

PERIODIC SOLUTIONS FOR NONAUTONOMOUS SECOND ORDER DIFFERENTIAL INCLUSIONS SYSTEMS WITH p -LAPLACIAN

DANIEL PAŞCA

ABSTRACT. Using the nonsmooth variant of minimax point theorems, some existence results are obtained for periodic solutions of nonautonomous second-order differential inclusions systems with p -Laplacian.

1. INTRODUCTION

Consider the second order system

$$(1) \quad \begin{aligned} \ddot{u}(t) &= \nabla F(t, u(t)) \text{ a.e. } t \in [0, T], \\ u(0) - u(T) &= \dot{u}(0) - \dot{u}(T) = 0 \end{aligned}$$

where $T > 0$, $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the following assumption:

(A) $F(t, x)$ is measurable in t for each $x \in \mathbb{R}^n$ and continuously differentiable in x for a.e. $t \in [0, T]$, and there exist $a \in C(\mathbb{R}_+, \mathbb{R}_+)$, $b \in L^1(0, T; \mathbb{R}_+)$ such that

$$|F(t, x)| + \|\nabla F(t, x)\| \leq a(\|x\|)b(t),$$

for all $x \in \mathbb{R}^n$ and a.e. $t \in [0, T]$.

Suppose that the nonlinearity $\nabla F(t, x)$ is bounded, that is, there exists $g \in L^1(0, T; \mathbb{R}_+)$ such that

$$\|\nabla F(t, x)\| \leq g(t),$$

for all $x \in \mathbb{R}^n$ and a.e. $t \in [0, T]$. In [3] the authors proved the existence of solutions for problem (1) under the condition that

$$\int_0^T F(t, x)dt \rightarrow +\infty \text{ as } \|x\| \rightarrow \infty,$$

or that

$$\int_0^T F(t, x)dt \rightarrow -\infty \text{ as } \|x\| \rightarrow \infty.$$

Tang in [5] proved the existence of solutions for problem (1) under more general conditions. He supposes that assumption (A) holds, that

$$\|\nabla F(t, x)\| \leq f(t)\|x\|^\alpha + g(t),$$

1991 *Mathematics Subject Classification.* 34C25, 49J24.

Key words and phrases. p -Laplacian, differential inclusions, the (PS) condition, periodic solutions, Rabinowitz's saddle point theorem.

[†] The author was supported by Ministerio de Educación y Ciencia, grant number SB2003-0337.

for all $x \in \mathbb{R}^n$ and a.e. $t \in [0, T]$, where $f, g \in L^1(0, T; \mathbb{R}_+)$, $\alpha \in [0, 1)$ and

$$\|x\|^{-2\alpha} \int_0^T F(t, x) dt \rightarrow +\infty \text{ as } \|x\| \rightarrow \infty,$$

or that

$$\|x\|^{-2\alpha} \int_0^T F(t, x) dt \rightarrow -\infty \text{ as } \|x\| \rightarrow \infty.$$

In order to prove the above results, Mawhin-Willem and Tang apply the classical (smooth) variant of minimax methods. In [4] we have considered the following problem which is a generalization of problem (1)

$$(2) \quad \begin{aligned} \ddot{u}(t) &\in \partial F(t, u(t)) \text{ a.e. } t \in [0, T], \\ u(0) - u(T) &= \dot{u}(0) - \dot{u}(T) = 0 \end{aligned}$$

where $T > 0$, $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$, ∂ denotes the Clarke subdifferential (see [2]) and $F(t, x)$ is measurable in t for each $x \in \mathbb{R}^n$, and locally Lipschitz and regular (see [2]) in x for each $t \in [0, T]$. Under some additional assumptions (see [4]) on F and ∂F we proved the existence of solutions for problem (2).

The aim of this paper is to consider the problem (2) in a more general sense. More exactly our results represent the extensions to systems with p -Laplacian.

Consider the second order differential inclusions system

$$(3) \quad \begin{aligned} \frac{d}{dt}(|\dot{u}(t)|^{p-2}\dot{u}(t)) &\in \partial F(t, u(t)) \text{ a.e. } t \in [0, T], \\ u(0) &= u(T), \dot{u}(0) = \dot{u}(T), \end{aligned}$$

where $p > 1$, $T > 0$, $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$, and ∂ denotes the Clarke subdifferential.

The corresponding functional $\varphi(u) : W_T^{1,p} \rightarrow \mathbb{R}$ is given by

$$\varphi(u) = \frac{1}{p} \int_0^T |\dot{u}(t)|^p dt + \int_0^T F(t, u(t)) dt.$$

2. MAIN RESULTS

Theorem 1. *Let $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that $F(t, x)$ is measurable in t for each $x \in \mathbb{R}^n$ and regular in x for each $t \in [0, T]$. We suppose that exist $k \in L^q(0, T; \mathbb{R})$ such that*

$$(4) \quad |F(t, x_1) - F(t, x_2)| \leq k(t) \|x_1 - x_2\|$$

for all $t \in [0, T]$ and all $x_1, x_2 \in \mathbb{R}^n$. If there exist $c_1, c_2 > 0$ and $\alpha \in [0, 1)$ such that

$$(5) \quad \zeta_1 \in \partial F(t, x) \Rightarrow \|\zeta_1\| \leq c_1 \|x\|^\alpha + c_2$$

for all $t \in [0, T]$ and all $x \in \mathbb{R}^n$, and if for $q = \frac{p}{p-1}$

$$(6) \quad \|x\|^{-q\alpha} \int_0^T F(t, x) dt \rightarrow +\infty \text{ as } \|x\| \rightarrow \infty$$

then problem (3) has at least one solution which minimizes the functional φ on $W_T^{1,p}$.

Theorem 2. *Let $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that $F(t, x)$ is measurable in t for each $x \in \mathbb{R}^n$ and locally Lipschitz and regular in x for each $t \in [0, T]$. We suppose that exist $a \in C(\mathbb{R}_+, \mathbb{R}_+)$ and $b \in L^1(0, T; \mathbb{R}_+)$ such that*

$$(7) \quad \|F(t, x)\| \leq a(\|x\|)b(t)$$

for all $t \in [0, T]$ and all $x \in \mathbb{R}^n$. If there exist $c_1, c_2 > 0$ and $\alpha \in [0, 1)$ such that

$$\zeta_1 \in \partial F(t, x) \Rightarrow \|\zeta_1\| \leq c_1 \|x\|^\alpha + c_2$$

for all $t \in [0, T]$ and all $x \in \mathbb{R}^n$, and if for $q = \frac{p}{p-1}$

$$(8) \quad \|x\|^{-q\alpha} \int_0^T F(t, x) dt \rightarrow -\infty \text{ as } \|x\| \rightarrow \infty$$

then problem (3) has at least one solution on $W_T^{1,p}$.

Remark 1. *Theorems 1 and 2 generalizes the corresponding Theorems 1 and 2 of [4]. In fact, it follows from these theorems letting $p = 2$.*

3. THE PRELIMINARY RESULTS

We introduce some functional spaces. Let T a positive real number and $1 < p < \infty$. We denote by $W_T^{1,p}$ the Sobolev space of functions $u \in L^p(0, T; \mathbb{R}^n)$ having a weak derivative $\dot{u} \in L^p(0, T; \mathbb{R}^n)$. The norm over X is defined by

$$\|u\|_{W_T^{1,p}} = \left(\int_0^T \|u(t)\|^p dt + \int_0^T \|\dot{u}(t)\|^p dt \right)^{\frac{1}{p}}.$$

We recall that

$$\|u\|_{L^p} = \left(\int_0^T \|u(t)\|^p dt \right)^{\frac{1}{p}} \text{ and } \|u\|_\infty = \max_{t \in [0, T]} \|u(t)\|.$$

For our aims it is necessary to recall some very well know results (for proof and details see [3]).

Proposition 3. *If $u \in W_T^{1,p}$ then*

$$\|u\|_\infty \leq c \|u\|_{W_T^{1,p}}.$$

If $u \in W_T^{1,p}$ and $\int_0^T u(t) dt = 0$ then

$$\|u\|_\infty \leq c \|\dot{u}\|_{L^p} \text{ (Sobolev inequality),}$$

$$\|u\|_{L^p} \leq c \|\dot{u}\|_{L^p} \text{ (Wirtinger's inequality).}$$

Proposition 4. *If the sequence $(u_k)_k$ converges weakly to u in $W_T^{1,p}$, then $(u_k)_k$ converges uniformly to u on $[0, T]$.*

Let X be a Banach space. Now follows [2], for each $x, v \in X$, we define the *generalized directional derivative* at x in the direction v of a given $f \in Lip_{loc}(X, \mathbb{R})$ as

$$f^0(x; v) = \limsup_{y \rightarrow x, \lambda \searrow 0} \frac{f(y + \lambda v) - f(y)}{\lambda}$$

and we denote by

$$\partial f(x) = \{x^* \in X^* : f^0(x; v) \geq \langle x^*, v \rangle, \text{ for all } v \in X\}$$

the *generalized gradient* of f at x (the Clarke subdifferential).

We recall the *Lebourg's mean value theorem* (see [2], Theorem 2.3.7).

Theorem 5. *Let x and y be points in X , and suppose that f is Lipschitz on open set containing the line segment $[x, y]$. Then there exists a point u in (x, y) such that*

$$f(y) - f(x) \in \langle \partial f(u), y - x \rangle.$$

Clarke consider in [2] the following abstract framework:

- let (T, \mathcal{T}, μ) be a positive complete measure space with $\mu(T) < \infty$, and let Y be a separable Banach space;
- let Z be a closed subspace of $L^p(T; Y)$ (for some p in $[1, \infty)$), where $L^p(T; Y)$ is the space of p -integrable functions from T to Y ;
- define a functional f on Z via

$$f(x) = \int_T f_t(x(t)) \mu(dt),$$

where $f_t : Y \rightarrow \mathbb{R}$, $(t \in T)$ is a given family of functions;

- suppose that for each y in Y the function $t \rightarrow f_t(y)$ is measurable, and that x is a point at which $f(x)$ is defined (finitely).

Hypothesis 1: There is a function k in $L^q(T, \mathbb{R})$, $\left(\frac{1}{p} + \frac{1}{q} = 1\right)$ such that, for all $t \in T$,

$$|f_t(y_1) - f_t(y_2)| \leq k(t) \|y_1 - y_2\|_Y \text{ for all } y_1, y_2 \in Y$$

Hypothesis 2: Each function f_t is Lipschitz (of some rank) near each point of Y , and for some constant c , for all $t \in T$, $y \in Y$, one has

$$\zeta \in \partial f_t(y) \Rightarrow \|\zeta\|_{Y^*} \leq c\{1 + \|y\|_Y^{p-1}\}.$$

Under this conditions described above Clarke prove (see [2], Theorem 2.7.5):

Theorem 6. *Under the conditions described above, under either of Hypothesis 1 or 2, f is uniformly Lipschitz on bounded subsets of Z , and one has*

$$\partial f(x) \subset \int_T \partial f_t(x(t)) \mu(dt).$$

Further, if each f_t is regular at $x(t)$ then f is regular at x and equality holds.

Remark 2. *f is globally Lipschitz on Z when Hypothesis 1 hold.*

Now we can prove the following result.

Theorem 7. *Let $F : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that $F(t, x)$ is measurable in t for each $x \in \mathbb{R}^n$, and locally Lipschitz and regular in x for each $t \in [0, T]$, and there exist $a \in C(\mathbb{R}_+, \mathbb{R}_+)$, $b \in L^1(0, T; \mathbb{R}_+)$, $c_1, c_2 > 0$ and $\alpha \in [0, p-1)$ such that*

$$(9) \quad |F(t, x)| \leq a(\|x\|)b(t),$$

$$(10) \quad \zeta_1 \in \partial F(t, x) \Rightarrow \|\zeta_1\| \leq c_1 \|x\|^\alpha + c_2,$$

for all $t \in [0, T]$ and all $x \in \mathbb{R}^n$. We suppose that $L : [0, T] \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$, is given by $L(t, x, y) = \frac{1}{p}\|y\|^p - F(t, x)$.

Then, the functional $f : Z \in \mathbb{R}$, where

$$Z = \left\{ (u, v) \in L^p(0, T; Y) : u(t) = \int_0^t v(s)ds + c, c \in \mathbb{R}^n \right\}$$

given by $f(u, v) = \int_0^T L(t, u(t), v(t))dt$, is uniformly Lipschitz on bounded subsets of Z and one has

$$(11) \quad \partial f(u, v) \subset \int_0^T \partial L(t, u(t), v(t))dt.$$

Proof. We can apply Theorem 6 under Hypothesis 2, with the following cast of characters:

- $(T, \mathcal{T}, \mu) = [0, T]$ with Lebesgue measure, $Y = \mathbb{R}^n \times \mathbb{R}^n$ be the Hilbert product space (hence is separable);
- $p > 1$ and

$$Z = \left\{ (u, v) \in L^p(0, T; Y) : u(t) = \int_0^t v(s)ds + c, c \in \mathbb{R}^n \right\}$$

be a closed subspace of $L^p(0, T; Y)$;

- $f_t(x, y) = L(t, x, y) = \frac{1}{p}\|y\|^p + F(t, x)$; in our assumptions it results that the integrand $L(t, x, y)$ is measurable in t for a given element (x, y) of Y , locally Lipschitz in (x, y) for each $t \in [0, T]$.

Proposition 2.3.15 from [2] implies

$$\partial L(t, x, y) \subset \partial_x L(t, x, y) \times \partial_y L(t, x, y) = \partial\{F(t, x)\} \times \{\|y\|^{p-2}y\}.$$

Using (3) and (4), if $\zeta = (\zeta_1, \zeta_2) \in \partial L(t, x, y)$ it results $\zeta_1 \in \partial\{F(t, x)\}$ and $\zeta_2 = \|y\|^{p-2}y$, and hence

$$\|\zeta\| = \|\zeta_1\| + \|\zeta_2\| \leq c_1\|x\|^\alpha + c_2 + \|y\|^{p-1} \leq \tilde{c}\{1 + \|(x, y)\|^{p-1}\}$$

for each $t \in [0, T]$, since $\alpha < p - 1$ and $p > 1$. The hypotheses of Theorem 6 are satisfied, therefore f is uniformly Lipschitz on the bounded subsets of Z and one has (11). \square

Remark 3. The interpretation of expression (11) is as follows: if (u_0, v_0) is an element of Z (so that $v_0 = \dot{u}_0$) and if $\zeta \in \partial f(u_0, v_0)$, we deduce the existence of a measurable function $(q(t), p(t))$ such that

$$(12) \quad q(t) \in \partial\{F(t, u_0(t))\} \text{ and } p(t) = \|v_0(t)\|^{p-2}v_0(t) \text{ a.e. on } [0, T]$$

and for any (u, v) in Z , one has

$$\langle \zeta, (u, v) \rangle = \int_0^T \{ \langle q(t), u(t) \rangle + \langle p(t), v(t) \rangle \} dt.$$

In particular, if $\zeta = 0$ (so that u_0 is critical point for $\varphi(u) = \int_0^T \left[\frac{1}{p}\|\dot{u}(t)\|^p + \right.$

$F(t, u(t)) \Big] dt$), it then follows easily that $q(t) = \dot{p}(t)$ a.e., or taking into account (12)

$$\frac{d}{dt} (\| \dot{u}_0(t) \|^{p-2} \dot{u}_0(t)) \in \partial F(t, u_0(t)) \text{ a.e. on } [0, T],$$

so that u_0 satisfies the inclusions system (3).

Remark 4. If $p = 2$ then the system (3) becomes system (2). If in addition F is continuously differentiable in x , then the system (3) becomes system (1).

In proving Theorem 2 we will invoke the following nonsmooth variant of the Rabinowitz's saddle point theorem (see [1], Theorem 3.3):

Theorem 8. Let X be a real Banach space, and let f be a locally Lipschitz function defined on X satisfies (PS) condition. Suppose $X = X_1 \oplus X_2$ with a finite-dimensional subspace X_1 , and there exist constants $b_1 < b_2$ and a bounded neighborhood N of θ in X_1 such that

$$f|_{X_2} \geq b_2, \quad f|_{\partial N} \leq b_1,$$

then f has a critical point.

The definitions of a critical point and the Palais-Smale condition are now recalled.

Definition 1. A point $u \in X$ is said to be a critical point of $f \in Lip_{loc}(X, \mathbb{R})$ if $\theta \in \partial f(u)$, namely $f^0(u, v) \geq 0$ for every $v \in X$. A real number c is called a critical value of f if there is a critical point $u \in X$ such that $f(u) = c$.

Definition 2. If $f \in Lip_{loc}(X, \mathbb{R})$, we say that f satisfies the Palais-Smale condition (in short (PS)) if each sequence (x_n) in X such that $(f(x_n))$ is bounded and $\lim_{n \rightarrow \infty} \lambda(x_n) = 0$ has a convergent subsequence. We denote $\lambda(x) = \min_{x^* \in \partial f(x)} \|x^*\|$.

4. PROOF OF THE THEOREMS

4.1. Proof of Theorem 1. For $u \in W_T^{1,p}$, let $\bar{u} = \frac{1}{T} \int_0^T u(t) dt$ and $\tilde{u} = u - \bar{u}$. From Lebourg's mean value theorem it follows that for each $t \in [0, T]$ there exist $z(t)$ in $(\bar{u}, u(t))$ and $\zeta \in \partial F(t, z(t))$ such that $F(t, u(t)) - F(t, \bar{u}) = \langle \zeta, \tilde{u}(t) \rangle$. It follows from (5) and Hölder's inequality that

$$\begin{aligned} \left| \int_0^T [F(t, u(t)) - F(t, \bar{u})] dt \right| &\leq \int_0^T |F(t, u(t)) - F(t, \bar{u})| dt \leq \\ &\leq \int_0^T |\zeta| |\tilde{u}(t)| dt \leq \int_0^T \left[2c_1 (|\bar{u}|^\alpha + |\tilde{u}(t)|^\alpha) + c_2 \right] |\tilde{u}(t)| dt \leq \\ &\leq C_1 \|\tilde{u}\|_\infty^{\alpha+1} + C_2 \|\tilde{u}\|_\infty \|\bar{u}\|^\alpha + C_3 \|\tilde{u}\|_\infty \leq \\ &\leq C_4 \|\dot{u}\|_{L^p}^{\alpha+1} + \frac{1}{2p} \|\dot{u}\|_{L^p}^p + C_5 \|\dot{u}\|_{L^p} + C_6 \|\bar{u}\|^{q\alpha} \end{aligned}$$

for all $u \in W_T^{1,p}$ and some positive constants C_4 , C_5 and C_6 . Hence we have

$$\varphi(u) \geq \frac{1}{p} \int_0^T |\dot{u}(t)|^p dt + \int_0^T F(t, \bar{u}) dt + \int_0^T [F(t, u(t)) - F(t, \bar{u})] dt \geq$$

$$\begin{aligned}
&\geq \frac{1}{2p} \|\dot{u}\|_{L^p}^p - C_4 \|\dot{u}\|_{L^p}^{\alpha+1} - C_5 \|\dot{u}\|_{L^p} - C_6 \|\bar{u}\|^{q\alpha} + \int_0^T F(t, \bar{u}) dt \geq \\
&\geq \frac{1}{2p} \|\dot{u}\|_{L^p}^p - C_4 \|\dot{u}\|_{L^p}^{\alpha+1} - C_5 \|\dot{u}\|_{L^p} + \|\bar{u}\|^{q\alpha} \left\{ \frac{1}{\|\bar{u}\|^{q\alpha}} \int_0^T F(t, \bar{u}) dt - C_6 \right\}
\end{aligned}$$

for all $u \in W_T^{1,p}$, which implies that $\varphi(u) \rightarrow \infty$ as $\|u\| \rightarrow \infty$ by (6) because $\alpha < p - 1$, and the norm $\|u\| = (\|\bar{u}\|^p + \|\dot{u}\|_{L^p}^p)^{\frac{1}{p}}$ is an equivalent norm on $W_T^{1,p}$. Now we write $\varphi(u) = \varphi_1(u) + \varphi_2(u)$ where

$$\varphi_1(u) = \frac{1}{p} \int_0^T |\dot{u}(t)|^p dt \text{ and } \varphi_2(u) = \int_0^T F(t, u(t)) dt.$$

The function φ_1 is weakly lower semi-continuous (w.l.s.c.) on $W_T^{1,p}$. From (4), (5) and Theorem 7, taking to account Remark 2 and Proposition 4, it follows that φ_2 is w.l.s.c. on $W_T^{1,p}$. By Theorem 1.1 in [3] it follows that φ has a minimum u_0 on $W_T^{1,p}$. Evidently $Z \simeq W_T^{1,p}$ and $\varphi(u) = f(u, v)$ for all $(u, v) \in Z$. From Theorem 7, it results that f is uniformly Lipschitz on bounded subsets of Z , and therefore φ possesses the same properties relative to $W_T^{1,p}$. Proposition 2.3.2 in [2] implies that $0 \in \partial\varphi(u_0)$ (so that u_0 is critical point for φ). Now from Theorem 7 and Remark 3 it follows that the problem (3) has at least one solution $u \in W_T^{1,p}$. ■

Remark 5. Evidently if $p = 2$ then we obtain the existence of solutions of problem (2). If in addition F is continuously differentiable in x , then we obtain the existence of solutions of problem (2).

4.2. Proof of Theorem 2. We will see that the functional

$$\varphi(u) : W_T^{1,p} \rightarrow \mathbb{R}, \quad \varphi(u) = \frac{1}{p} \int_0^T |\dot{u}(t)|^p dt + \int_0^T F(t, u(t)) dt.$$

verify the assumptions of Theorem 8. Evidently $Z \simeq W_T^{1,p}$ and $\varphi(u) = f(u, v)$ for all $(u, v) \in Z$. From Theorem 7, it results that f is uniformly Lipschitz on bounded subsets of Z and regular at each $(u, v) \in Z$, and therefore φ possesses the same properties relative to $W_T^{1,p}$. The functional φ is neither bounded from below, nor from above. Indeed, if $w \in W_T^{1,p}$ is a constant function, then

$$\varphi(w) = \int_0^T F(t, w) dt = \|w\|^{q\alpha} \left(\|w\|^{-q\alpha} \int_0^T F(t, w) dt \right) \rightarrow -\infty \text{ as } \|w\| \rightarrow \infty$$

and, if $v \in W_T^{1,p}$ has mean zero, by the proof of Theorem 1 one has

$$\begin{aligned}
\varphi(v) &= \frac{1}{p} \int_0^T |\dot{v}(t)|^p dt + \int_0^T F(t, 0) dt + \int_0^T [F(t, v(t)) - F(t, 0)] dt = \\
&= \frac{1}{p} \int_0^T |\dot{v}(t)|^p dt + \int_0^T F(t, 0) dt + \int_0^T \langle \zeta_1, v(t) \rangle dt \geq \\
&\geq \frac{1}{2p} \|\dot{u}\|_{L^p}^p - C_4 \|\dot{u}\|_{L^p}^{\alpha+1} - C_5 \|\dot{u}\|_{L^p} + \int_0^T F(t, 0) dt
\end{aligned}$$

where we applied the Lebourg's mean value theorem and Sobolev inequality, and where C_1 and C_2 are positive constants, so that φ is not bounded from above. We denote

$$X_1 = \{w \in W_T^{1,p} : w = \text{constant}\}$$

and

$$X_2 = \left\{v \in W_T^{1,p} : \int_0^T v(t) dt = 0\right\}.$$

Evidently $W_T^{1,p} = X_1 \oplus X_2$ with $\dim X_1 < \infty$. From the above observations, we see that there exists $R > 0$ such that

$$\sup_{S_R} \varphi < \inf_{X_2} \varphi$$

where $S_R = \{w \in X_1 : \|w\|_{W_T^{1,p}} = R\}$.

We shall show that φ satisfies the (PS) condition. Let (u_k) be a sequence in $W_T^{1,p}$ such that $\varphi(u_k)$ is bounded and $\lambda(u_k) \rightarrow 0$ as $k \rightarrow \infty$. Writing $u_k(t) = \tilde{u}_k(t) + \bar{u}_k$ with $\bar{u}_k = \frac{1}{T} \int_0^T u_k(t) dt$, and using the definition of $\lambda(u_k)$ it results that there is some k_0 such that for each $k \geq k_0$ there exist $u_k^* \in \partial\varphi(u_k)$ with

$$|\langle u_k^*, h \rangle| \leq \|h\|_{W_T^{1,p}}, \quad \text{for all } h \in W_T^{1,p}.$$

From Theorem 7, if $u_k^* \in \partial\varphi(u_k)$ it results that there exist $q_k(t) \in \partial F(t, u_k(t))$ such that

$$|\langle u_k^*, \tilde{u}_k \rangle| = \left| \int_0^T [\|\dot{u}_k(t)\|^p + \langle q_k(t), \tilde{u}_k(t) \rangle] dt \right| \leq \|\tilde{u}_k\|_{W_T^{1,p}}, \quad \text{for all } k \geq k_0.$$

In similar way to the proof of Theorem 1, we have

$$\left| \int_0^T \langle q_k(t), \tilde{u}_k(t) \rangle dt \right| \leq \frac{1}{2p} \|\dot{u}_k\|_{L^p}^p + C_4 \|\dot{u}_k\|_{L^p}^{\alpha+1} + C_5 \|\dot{u}_k\|_{L^p} + C_6 \|\bar{u}_k\|^{q\alpha}$$

for all k . Hence one has

$$\begin{aligned} \|\tilde{u}_k\|_{W_T^{1,p}} &\geq \langle u_k^*, \tilde{u}_k \rangle = \int_0^T [\|\dot{u}_k(t)\|^p + \langle q_k(t), \tilde{u}_k(t) \rangle] dt \geq \\ &\geq \frac{2p-1}{2p} \|\dot{u}_k\|_{L^p}^p - C_4 \|\dot{u}_k\|_{L^p}^{\alpha+1} - C_5 \|\dot{u}_k\|_{L^p} - C_6 \|\bar{u}_k\|^{q\alpha} \end{aligned}$$

for $k \geq k_0$. It follows from Wirtinger's inequality that

$$\|\tilde{u}_k\|_{W_T^{1,p}} \leq (1+c)^{\frac{1}{p}} \|\dot{u}_k\|_{L^p}$$

for all k . Hence we obtain

$$(1+c)^{\frac{1}{p}} \|\dot{u}_k\|_{L^p} \geq \frac{2p-1}{2p} \|\dot{u}_k\|_{L^p}^p - C_4 \|\dot{u}_k\|_{L^p}^{\alpha+1} - C_5 \|\dot{u}_k\|_{L^p} - C_6 \|\bar{u}_k\|^{q\alpha}$$

for $k \geq k_0$, and it follows that

$$C_6 \|\bar{u}_k\|^{q\alpha} \geq \frac{2p-1}{2p} \|\dot{u}_k\|_{L^p}^p - C_4 \|\dot{u}_k\|_{L^p}^{\alpha+1} - [(1+c)^{\frac{1}{p}} + C_5] \|\dot{u}_k\|_{L^p}$$

or

$$(13) \quad C_7 \|\bar{u}_k\|^{q\alpha} \geq \|\dot{u}_k\|_{L^p}^p$$

for some $C_7 > 0$ and for $k \geq k_0$. By the proof of Theorem 1 we have

$$\left| \int_0^T [F(t, u_k(t)) - F(t, \bar{u}_k)] dt \right| \leq \frac{1}{2p} \|\dot{u}_k\|_{L^p}^p + C_4 \|\dot{u}_k\|_{L^p}^{\alpha+1} + C_5 \|\dot{u}_k\|_{L^p} + C_6 \|\bar{u}_k\|^{q\alpha}$$

for all k . It follows from the boundedness of $(\varphi(u_k))$, (13) and the above inequality that

$$\begin{aligned} C_8 \leq \varphi(u_k) &= \frac{1}{p} \int_0^T |\dot{u}_k(t)|^p dt + \int_0^T [F(t, u_k(t)) - F(t, \bar{u}_k)] dt + \int_0^T F(t, \bar{u}_k) dt \leq \\ &\leq \frac{2p-1}{2p} \|\dot{u}_k\|_{L^p}^p + C_4 \|\dot{u}_k\|_{L^p}^{\alpha+1} + C_5 \|\dot{u}_k\|_{L^p} + C_6 \|\bar{u}_k\|^{q\alpha} + \int_0^T F(t, \bar{u}_k) dt \leq \\ &\leq \|\bar{u}_k\|^{q\alpha} \left(\|\bar{u}_k\|^{-q\alpha} \int_0^T F(t, \bar{u}_k) dt + C_9 \right) \end{aligned}$$

for $k \geq k_0$ and some positive constants C_8 and C_9 . The above inequality and (8) implies that $(\|\bar{u}_k\|)$ is bounded. Hence (u_k) is bounded by (13). Thus (u_k) is bounded in $W_T^{1,p}$ and hence contains a subsequence, relabeled (u_k) , which converge to some $u \in W_T^{1,p}$, weakly in $W_T^{1,p}$ and strongly in $C([0, T]; \mathbb{R}^n)$ (see Proposition 4). Therefore we have for $u_k^* \in \partial\varphi(u_k)$ and $u^* \in \partial\varphi(u)$

$$\langle u_k^* - u^*, u_k - u \rangle \rightarrow 0 \text{ as } k \rightarrow \infty.$$

But

$$\begin{aligned} \langle u_k^* - u^*, u_k - u \rangle &= \int_0^T \left[\langle q_k(t) - q(t), u_k(t) - u(t) \rangle + \|\dot{u}_k(t) - \dot{u}(t)\|^p \right] dt = \\ &= \|\dot{u}_k - \dot{u}\|_{L^p}^p + \int_0^T \langle q_k(t) - q(t), u_k(t) - u(t) \rangle dt \end{aligned}$$

where $q_k(t) \in \partial F(t, u_k(t))$ and $q(t) \in \partial F(t, u(t))$. It is easy to verify, that $\|\dot{u}_k - \dot{u}\|_{L^p} \rightarrow 0$ as $k \rightarrow \infty$, and hence $u_k \rightarrow u$ in $W_T^{1,p}$. We conclude that (PS) is satisfied and from Theorem 8, φ admits a critical point. Now from Theorem 7 and Remark 3 it follows that the problem (3) has at least one solution $u \in W_T^{1,p}$. \blacksquare

Remark 6. Evidently if $p = 2$ then we obtain the existence of solutions of problem (2). If in addition F is continuously differentiable in x , then we obtain the existence of solutions of problem (2).

ACKNOWLEDGEMENTS

The author thanks Centre de Recerca Matemàtica for the hospitality and facilities for doing this work.

REFERENCES

- [1] Kung-Ching Chang - *Variational Methods for Non-Differentiable Functionals and Their Applications to Partial Differential Equations*, J. Math. Anal. Appl. 80 (1981), 102–129.
- [2] F.H. Clarke - *Optimization and Nonsmooth Analysis*, SIAM, Classics in Applied Mathematics vol.5, Philadelphia, 1990.
- [3] J. Mawhin and M. Willem - *Critical Point Theory and Hamiltonian Systems*, Springer-Verlag, Berlin/New York, 1989.
- [4] Daniel Pașca - *Periodic Solutions for Second Order Differential Inclusions with Sublinear Nonlinearity*, PanAmerican Mathematical Journal, vol. 10, nr. 4 (2000) 35–45.
- [5] Chun-Lei Tang - *Periodic Solutions for Nonautonomous Second Order Systems with Sublinear Nonlinearity*, Proc. AMS, vol. 126, nr. 11 (1998), 3263–3270.

PERMANENT ADDRESS: DEPARTMENT OF MATHEMATICS AND INFORMATICS, UNIVERSITY OF ORADEA, UNIVERSITY STREET 1, 410087 ORADEA, ROMANIA

E-mail address: `dpasca@uoradea.ro`

CURRENT ADDRESS: CENTRE DE RECERCA MATEMÀTICA, 08193 BELLATERRA, BARCELONA, SPAIN

E-mail address: `dpasca@crm.es`