

POWER, HEART RATE AND PERCEIVED EXERTION RESPONSES TO 3X3 AND 4X4 BASKETBALL SMALL-SIDED GAMES

Jaime Sampaio, Catarina Abrantes & Nuno Leite

KEY WORDS: Training intensity; heart rate; rating of perceived exertion; basketball

ABSTRACT: Despite the interest drawn by game adaptations in players' performance development, no study examined the effects of these task constraints in basketball games exercise intensity. Therefore, the aim of this study is to identify differences in power, heart rate and perceived exertion responses to 3x3 and 4x4 basketball small-sided games. Eight young male basketball players participated in this study. Player's individual peak heart rate value and global perceived exertion was registered immediately after two small-sided games, 3x3 and 4x4. Additionally, squat jump and countermovement jump were used to assess power. Our results show that both small-sided games promoted high physiological demands, whereas the players performed the tasks above 80% of HRmax. Nevertheless, another interesting finding of this study is related to the fact that 3x3 contributed to higher physiological demands than the 4x4. The significant increase in the countermovement jump posttest jump results could suggest that the 4x4 were not played as quickly nor intensely as the 3x3. Decreases of the space and number of players in game allow greater self-recreation of players and greater intervention in game. Therefore, the heart rate response during the series displays a higher physiologic impact in 3x3 than in 4x4.

Jaime Sampaio, Sport Sciences, Exercise and Health Department, Universidade de Trás-os-Montes e Alto Douro, Quinta de Prados, Apartado 202, 5001-911 Vila Real, Portugal.

E-mail: ajaime@utad.pt.

Research Center in Sports Sciences, Health Sciences and Human Development (CIDESD), Portugal

Introduction

Challenges in sport sciences and coaching are not exclusively focused in clarifying the complex structures of performance and their associated parameters. The understanding of interactions between different capacities, contexts and other variables could delimit its influence on the performance of players and teams. Basketball is played by two teams of five players running in an area of approximately 450m². However, during the training process, it is common to reduce the number of players on each team and the size of the playable area. These small-sided games are present in the majority of methods used for basketball training. The players are more actively involved in these games and face with higher and continuous concentration all the demands, without breaks. Furthermore, with this reduce game format all players are encouraged to better develop awareness of basic attack and defense elements.

Technical and tactical dividends are evident, once players have more time in contact with the ball and therefore more success in decision-making. Additionally, at this point, available evidence suggests that these constraints (i.e., playing area and number of players) promote higher levels of enjoyment and dedication among players, who therefore, further develop at this level (Wall & Côté, 2007). Despite the interest drawn by game adaptations in players' performance development, to authors' knowledge, no study examined the effects of these task constraints on the exercise intensity of basketball games. Similarly, no study has evaluated the physiological, technical, tactical or psychological responses caused by this type of games. Such information could be useful for coaches and to help improve the design of effective physical training programs. Therefore, the aim of this study is to identify differences in

power, heart rate and perceived exertion responses to 3x3 and 4x4 basketball small-sided games.

Method

Participants and procedures

Eight young male basketball players participated in this study (weight: 65.2±5.9 kg, age: 15.5±0.6 years old, and height: 178.1±7.2 cm). Data collection was divided into three stages. First, in order to obtain the player's individual peak heart rate value (HRmax) during this period, all subjects completed the yo-yo intermittent recovery test level 2. Second and third stages corresponded to the application of 3x3 and 4x4 small-sided games. The games total duration was 25 min and consisted in 4×4 min bouts interspersed with 3 min active recovery. The court sizes used for both games kept the ratio player/area of 12m² and 16.8m², respectively for 3x3 and 4x4 games. The participants were well familiarized with all this protocols.

Instruments

Heart rate (HR) was measured (5 s recording intervals) via short-range radiotelemetry (Polar Team Sport System, Polar Electro, Finland). Perceived exertion was registered immediately after the games using the OMNI Perceived Exertion Scale (0-10, Robertson et al., 2005). The Ergojump (Globus Inc., Treviso, Italy) was used to measure the squat jump and countermovement jump flight time and jump height. Results were analyzed using Student t test. Statistical significance was set at 5%. Corresponding Cohen's effect sizes were also calculated. All data were analyzed with the statistical package SPSS for Windows, release 16.0 (SPSS Inc., Chicago, IL).

	Period	Heart Rate					RPE	
		Bout	Active Recovery	% maximum per period	% maximum per bout	% maximum in active recovery	Period	Bout
3X3	Effort 1	176,0±6,9		88,4			1,7±0,2	
	Effort 2	175,9±9,1		88,4			2,5±0,4	
	Effort 3	170,4±7,7	173,4	152,2		87,1	76,5	3,0
	Effort 4	171,1±9,6		86,0			4,2±1,0	
4X4	Effort 1	165,9±17,7		83,3			2,8±0,4	
	Effort 2	167,1±17,6		83,9			3,8±0,9	
	Effort 3	165,9±15,1	164,7	144,7		82,7	72,7	4,1
	Effort 4	160,1±14,2		80,4			5,0±1,0	

Table 1. Results of the descriptive and inferential statistics of heart rate and perceived exertion.

Test	3X3 (3x3)		4X4 (4x4)	
	M	SD	M	SD
<i>Squat Jump, on</i>				
Flight time				
Pre-tes	0,521	± 0,026	0,498	± 0,034
Post-tes	0,503	± 0,034	0,499	± 0,033
t	1,240		-0,054	
p	0,270		0,958	
Height				
Pre-tes	0,366	± 0,073	0,305	± 0,042
Post-tes	0,311	± 0,043	0,306	± 0,039
t	1,979		-0,046	
p	0,105		0,965	
<i>Countermovement jump, on</i>				
Flight time				
Pre-tes	0,553	± 0,040	0,549	± 0,035
Post-tes	0,549	± 0,037	0,570	± 0,041
t	0,486		-2,959	
p	0,648		0,021*	
ES			0,683	
Height				
Pre-teste	0,390	± 0,072	0,370	± 0,048
Post-teste	0,371	± 0,049	0,400	± 0,055
t	0,811		-3,046	
p	0,454		0,019*	
ES			0,694	

Table 2. Results of the descriptive and inferential statistics of ergojump tests

Results

The means, standard deviations, and univariate differences are presented in Tables 1 and 2. The heart rate responses were not significantly different between the two different games (3X3, 173.4 ± 8.3 beats min^{-1} ; 4X4, 164.7 ± 16.2 beats min^{-1}). These values corresponded to an exercise intensity of $87 \pm 4\%$ HRmax and $83 \pm 4\%$ for 3X3 and 4X4, respectively. These similarities in the heart rate responses to the small-sided game formats were maintained when the individual 4 min exercise and recovery periods were analyzed separately. Perceived exertion was also very similar between both games (3X3, 3.0 ± 0.5 ; 4X4, 4.1 ± 0.8). No significant differences were found between 3X3 and 4X4 neither between pre and post-tests in squat jump. Differences were, however, identified in 4X4 pre and post-test in countermovement jump, both in flight time ($p=0.021$, $d=0.683$) and height ($p=0.019$, $d=0.694$).

Discussion

The purpose of this study was to identify differences in power, heart rate and perceived exertion responses to 3x3 and 4x4 basketball small-sided games. Our results show that both games promoted high physiological demands, whereas the players performed the tasks above 80% of HRmax. Thus, one may argue that these small-sided games can be used for basketball-specific aerobic training with multifactorial training benefits. All these results confirm the recommendation made to coaches to use this training method as a specific physical training, also called integrated training. Indeed, sided games produce similar cardiovascular stress as other intermittent exercises specifically designed to improve athletes' endurance (Kelly & Drust, 2008). One

of the differences between sided games and short-duration intermittent running training methods is the presence of the ball, which imposes a specific activity and allows the improvement of technical and tactical skills with high player motivation. Nevertheless, another interesting finding of this study is related to the fact that the 3X3 game contributed to higher physiological demands than the 4X4 game.

Despite the inexistence of significant differences between both games it is important to enhance the fact that reducing the number of players and the playable area seems to increase the game intensity. Several studies have been confirming the physiological impact of small-sided games on diverse team sports (see Hill-Haas et al., 2008). The game area, the number of players, the game instructions, the number and duration of the series or the total duration of the session directly influence the activity of the players and the physiological impact. Indeed, Rampinini et al. (2007) showed that the intensity of sided games increases while the number of players decreases. Nevertheless, this is also dependent on the playing area, with the game intensity increasing when the available game area is decreased.

The results of the jump protocol reflected significant differences in 4X4 countermovement jump, both in flight time and height. The significant increase in the posttest jump results could suggest that the 4X4 were not played as intensively as the 3X3. Decreases of the space and number of players in game may allow greater self-recreation of players and greater intervention in game. Thus, the HR response during the series displays a higher physiological impact in 3X3 than 4X4.

References

- BALČIŪNAS, M., STONKUS, S., ABRANTES, C. et al. (2006). Long term effects of different training modalities on power, speed, skill and anaerobic capacity in young male basketball players. *Journal of Sports Science and Medicine* 5:163-170.
- KELLY, D. AND DRUST, B. (2008). The effect of pitch dimensions on heart rate responses and technical demands of small-sided soccer games in elite players. *J Sci Med Sport* 12(4):475-479.
- MCINNES, S., CARLSON, J., JONES, C. AND MCKENNA, M. (1995). The physiological load imposed on basketball players during competition. *Journal of Sport Science* 13:387-397.
- DELLAL, A., CHAMARI, K., PINTUS, A. ET AL. (2008). Heart rate responses during small-sided games and short intermittent running training in elite soccer players: a comparative study. *Journal of Strength and Conditioning Research* 22(5): 1449-1457.
- LEITE, N., SAMPAIO, J., ABRANTES, C. ET AL. (2007). Physical and technical componentes in boys' under-16 basketball by game quarter. *Iberian Congress on Basketball Research* 4:133-136.
- RAMPININI E, IMPELLIZZERI F, CASTAGNA C, et al. (2007). Factors influencing physiological responses to small-sided soccer games. *J Sports Sci* 25(6):659-66.
- IMPELLIZZERI, F., MARCORA, S., CASTAGNA, C. et al. (2006). Physiological and performance effects of generic versus specific aerobic training in soccer players. *Int J Sport Med* 27(6):483-492.
- IMPELLIZZERI, F., RAMPININI E., MARCORA S. (2005). Physiological assessment of aerobic training in soccer. *J Sports Sci* 23(6):583-92.
- ROBERTSON R., GOSS F., ANDREACCI J. et al. (2005). Validation of the children's OMNI RPE scale for stepping exercise. *Med Sci Sports Exer* 37(2):290-8.
- WALL, M. & CÔTÉ, J. (2007). Developmental activities that lead to drop out and investment in sport. *Physical Education and Sport Pedagogy* 12:77-87.

