

Accelerated drawdown of meridional overturning in the late-glacial Atlantic triggered by transient pre-H event freshwater perturbation

I. R. Hall,¹ S. B. Moran,² R. Zahn,^{1,3} P. C. Knutz,⁴ C.-C. Shen,⁵ and R. L. Edwards⁶

Received 8 March 2006; revised 30 June 2006; accepted 7 July 2006; published 31 August 2006.

[1] Abrupt decreases of the Atlantic meridional overturning circulation (MOC) during the Late Pleistocene have been directly linked to catastrophic discharges of glaciomarine freshwater, triggering disruption of northward marine heat transport and causing global climate changes. Here we provide measurements of excess sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ from a high-accumulation sediment drift deposit in the NE Atlantic that record a sequence of sudden variations in the rate of MOC, associated deep ocean ventilation and surface-ocean climatology. The data series reveal a sequential decrease in the MOC rate at ~ 18.0 ka BP ago that coincides with only transient and localized freshwater inputs. This change represents a substantial, though not total, cessation in MOC that predates the major Heinrich (H1) meltwater event by at least 1,200 years. These results highlight the potential of targeted freshwater perturbations in promoting substantial MOC changes without a direct linking with catastrophic freshwater surges. **Citation:** Hall, I. R., S. B. Moran, R. Zahn, P. C. Knutz, C.-C. Shen, and R. L. Edwards (2006), Accelerated drawdown of meridional overturning in the late-glacial Atlantic triggered by transient pre-H event freshwater perturbation, *Geophys. Res. Lett.*, 33, L16616, doi:10.1029/2006GL026239.

1. Introduction

[2] ^{231}Pa (half life $t_{1/2} = 32.5$ ka) and ^{230}Th ($t_{1/2} = 75.2$ ka) are particle-reactive radionuclides produced in the oceans at a constant initial $^{231}\text{Pa}/^{230}\text{Th}$ activity ratio of 0.093 and deposited in underlying sediments by attachment to settling particles. Interest in the application of excess, or unsupported, sediment $^{231}\text{Pa}/^{230}\text{Th}$ (decay-corrected to the time of deposition; herein referred to as $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$) as a palaeocirculation proxy in the Atlantic is based on the similar residence time of ^{231}Pa (~ 100 –200 yrs) [Nozaki and Nakanishi, 1985] and Atlantic deep waters [Broecker, 1979], whereby the rate of meridional overturning circulation (MOC) directly affects the lateral export of ^{231}Pa [Yu *et al.*, 1996; Marchal *et al.*, 2000]. Presently, approximately

half of the water column production of ^{231}Pa is exported in North Atlantic Deep Water (NADW) to the Southern Ocean [Yu *et al.*, 1996; Marchal *et al.*, 2000] by comparison, lateral transport of ^{230}Th is minimized due to its shorter residence time (~ 20 –40 yr) [Nozaki and Nakanishi, 1985; Yu *et al.*, 1996; Marchal *et al.*, 2000]. Thus, changes in Atlantic MOC are recorded down-core as variations in sediment $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$, with vigorous rates of MOC causing a reduction in $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ from the production ratio. The $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ proxy therefore enables the reconstruction of deep ocean circulation changes that goes beyond what traditional proxies such as $\delta^{13}\text{C}$ from benthic foraminifera can provide. Benthic $\delta^{13}\text{C}$ is driven by end-member variation, water mass chemical “aging” and mixing while $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ does depend more directly on lateral water mass advection, i.e., the physical vigor of the MOC.

2. Material and Methods

[3] We measured sedimentary $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ along sediment core DAPC2 recovered from a high-accumulation drift deposit located in the northern Rockall Trough ($58^{\circ}58.10'\text{N}$, $09^{\circ}36.75'\text{W}$, 1709 m water depth) (Figure 1). The site is presently influenced by Wyville-Thomson Overflow Water (WTOW) from the Norwegian Sea, a precursor to North Atlantic Deep Water (NADW), and recirculated upper NADW [New and Smythe-Wright, 2001]. This core provides high-resolution, multi-proxy, paleoceanographic records [Knutz *et al.*, 2002] of iceberg discharge and meltwater variability that indicate a rapid response of ice sheet surges and meltwater from NW Europe over the last deglaciation.

[4] Radiochemical analyses of ^{231}Pa and ^{230}Th were made by isotope dilution using a thermal ionization mass spectrometer for ^{231}Pa and a high-resolution magnetic sector inductively coupled plasma mass spectrometer for ^{230}Th [Shen *et al.*, 2002, 2003]. Diatom abundance as a possible modulator of $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ was quantified following Scherer [1995]. The chronology of DAPC2 is constrained by 10 calibrated accelerator mass spectrometry ^{14}C dates (age scale of Knutz *et al.* [2002] augmented by 4 further ^{14}C -AMS dates) that provide a tight temporal framework for the abrupt MOC changes indicated in the records and their linking with sudden changes of surface ocean climatology. The resulting time step along our stable isotope, faunal and ice rafted debris (IRD) records during the last glacial maximum (LGM) and H1 is less than 50 years, mean time step along the $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ is less than 120 years. At this temporal resolution, and because of its geographic location, DAPC2 provides an ideal sedimentary record in which to investigate the sequence of events

¹School of Earth, Ocean, and Planetary Sciences, Cardiff University, Cardiff, UK.

²Graduate School of Oceanography, University of Rhode Island, Narragansett, Rhode Island, USA.

³Now at Institut de Ciència i Tecnologia Ambientals, Universitat Autònoma de Barcelona, Bellaterra, Spain.

⁴Geological Survey of Denmark and Greenland, Copenhagen, Denmark.

⁵Department of Geosciences, National Taiwan University, Taipei, Taiwan.

⁶Department of Geology and Geophysics, University of Minnesota, Minneapolis, Minnesota, USA.




Figure 1. Study area and core DAPC2 data. (a) Map of the North Atlantic showing the location of core DAPC2 from Rockall Trough ($58^{\circ}58.10'N$, $09^{\circ}36.75'W$, 1709 m water depth). (b) ^{230}Th normalized biogenic silica flux (diatoms), note axis break (c) sedimentary record of $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$. Horizontal dashed line represents the $^{231}\text{Pa}/^{230}\text{Th}$ production ratio. AMS ^{14}C ages denoted by solid black triangles. Error bars represent uncertainties in ^{231}Pa and ^{230}Th data (2σ), which range from 3–7% and are generally $<5\%$. Shaded area highlights data not considered here for the reconstruction of paleocirculation due to the likely effects of preferential removal of ^{231}Pa by the increased biogenic silica flux.

surrounding a major past freshwater perturbation, the H1 meltwater event, and its consequences for the MOC.

3. Results and Discussion

[5] The $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ record displays (Figures 1 and 2) minimum values during the LGM and a multi-step transition to higher values that includes the Heinrich 1 (H1) meltwater event ($H-1_{\text{MW}}$). As non-carbonate clay and carbonate fluxes were high during the LGM (see auxiliary material¹ Figure S1), conditions typically associated with elevated $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ ratios [Bacon, 1988], the low $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ ratios prior to ~ 18 ka BP imply a vigorous MOC rate and export of ^{231}Pa from this location. At this time $\delta^{18}\text{O}$ values in DAPC2 of surface (*G. bulloides* [Schiebel et al., 1997]) and deep-dwelling (thermocline depths, *N. pachyderma* sinistral [Kohfeld et al., 1996]) planktonic foraminifera converge suggesting a weakly developed thermocline and decreased vertical stability, indicative of a well mixed ocean upper layer, plausibly marking conditions favorable for convective overturn in the region, in line with elevated benthic $\delta^{13}\text{C}$ in our core [Knutz et al., 2002]. In contrast, following the Younger Dryas stadial and throughout the Holocene, $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ ratios remain close to the production ratio which we attribute to preferential removal of ^{231}Pa by biogenic silica [Guo et al., 2002; Luo and Ku, 2004; Siddall et al., 2005], as evidenced by enhanced

diatom accumulation in the uppermost 50 cm (~ 12 ka BP) of DAPC2 (Figure 1c). The absence of diatom accumulation at the DAPC2 site during the LGM and early deglaciation indicates that $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ is not dominated by variations in biogenic silica flux during this period.

[6] Fine scale variability is observed in DAPC2 on centennial to sub-centennial time scales (Figure 2). These changes are directly coupled to atmospheric and surface




Figure 2. Paleoceanographic time-series from DAPC2 (a) *Neogloboquadrina pachyderma* sinistral (Nps) abundance. (b) Abundance of ice-raftered debris, quartz grains (Quartz, black) and detrital carbonate (DC, brown), in the $>250\text{ }\mu\text{m}$ fraction (c) *N. pachyderma* s. (Nps, cyan), *Globigerina bulloides* $\delta^{18}\text{O}$ (*G. bull.*, blue) and benthic $\delta^{18}\text{O}$ (*Cibicidoides wuellerstorfi*, Cw, orange). (d) Benthic $\delta^{13}\text{C}$ (Cw). (e) Sediment $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$. Acronyms are: H-1_{MW}, Heinrich 1 meltwater event; E-1_{a-c}, European sourced glacimarine events [Knutz et al., 2002]. These E-events and additional IRD pulses that do not display measurable planktonic $\delta^{18}\text{O}$ meltwater excursions at this location are highlighted by the grey bands. Heinrich Event 1 (H1) indicated by yellow band. Arrows between d and e mark onset of a multi-step MOC slowdown.

¹Auxiliary materials are available in the HTML. doi:10.1029/2006GL026239.

ocean (buoyancy) forcing revealed by the planktonic records that document a sequence of transient warming events followed by cooling/freshwater pulses, displayed in brief $\delta^{18}\text{O}$ and *N. pachyderma* (sin) abundance anomalies, prior to and during the onset of deglaciation (Figures 2a and 2c). Concomitant IRD peaks (Figures 2a–2c) suggest that the latter are linked with brief but intense pulses of icebergs discharged from the EIS (Figure 2b), previously labeled as E-events [Knutz et al., 2002], most likely triggered by short-lived collapses of marine-based ice margins caused by the slow but steady global warming at the end of the LGM. Similar lithic “precursor” events preceding Heinrich events have been reported from North Atlantic core sites and inferred to originate from European and Icelandic glacial sources [Grouset et al., 2000; Scourse et al., 2000]. The DAPC2 $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ record indicates that these events went along with measurable slowdown in MOC. In particular the event centered at ~ 19.1 ka BP stands out that is associated with a cooling and freshwater pulse, labeled as E-1_c event (Figure 2). The transient MOC slow-down during E-1_c that is evident in $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ is around half the magnitude of the major MOC drawdown starting at ~ 18 ka BP, although a value similar to those around H1 is reached in a single data point at 19.0 ka. MOC rapidly recovered from the slow down during the E-1_c event, within ~ 200 years returning to similar rates as before. A sustained step-wise increase in $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ then starts at ~ 18.0 ka BP, within ~ 170 yr after the warm pulse that followed the European sourced glacimarine E-1_b event [Knutz et al., 2002]. This signifies the onset of reduced lateral ^{231}Pa escape and a slow-down of deep water export from the region (Figures 2a and 2e). The first step of apparent MOC slow-down is coincident with a centennial scale IRD pulse that suggests a coeval EIS surge event. About one third of the total MOC drawdown, relative to the ~ 18.0 ka BP $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ maxima, is accounted for in this initial step. Approximately 270 years later, a second step of increasing $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ directly coincides with the incursion of freshwater/IRD pulse E-1_a. This phase of further MOC slow-down is accompanied by a coeval stepwise benthic $\delta^{13}\text{C}$ decrease that is also recorded at other core sites in the wider North Atlantic region and has been used, in conjunction with benthic trace element ratios, to infer decreasing chemical ventilation from northern sources and increased advection of southern hemisphere water masses [Zahn et al., 1997; Williams and Zahn, 2000; Zahn and Stüber, 2002; Rickaby and Elderfield, 2005]. The lack of equivalent negative benthic $\delta^{18}\text{O}$ excursions during the E-1_{b-a} events supports the view that these events did not cause globally significant changes in ice volume and sea level. The main H1 event lithologically identified in DAPC2 by the presence of a discrete detrital (dolomitic) carbonate IRD layer that is embedded within the quartz IRD deposit [Knutz et al., 2002] and the associated meltwater surge then occur as MOC rates approach peak minimum rates, some 1,200 yr after the onset of the MOC drawdown (Figure 2, H-1_{MW}). $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ ratios throughout the H1 interval remain below production ratio indicating that MOC never achieved a total cessation.

[7] The changes in deep water export traced by $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ in DAPC2 are also recorded at deep subtropical Atlantic sites (core OCE326-GGC5, lower Bermuda Rise, 4550 m water depth [McManus et al., 2004]; SU81-18, western Iberian margin, 3135 m water depth

[Gherardi et al., 2005]). The inference drawn at these sites was a significant slow-down in MOC coincident with the catastrophic iceberg discharge H1 [McManus et al., 2004; Gherardi et al., 2005] dated to 16.8 ka BP in the North Atlantic [Hemming, 2004]. The fine-scale time series of complementary proxies of ice rafted detritus, planktonic $\delta^{18}\text{O}$, benthic $\delta^{13}\text{C}$, and $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ (Figure 2) in DAPC2 from within the more immediate region of iceberg drift reveals that such MOC decrease occurred significantly before the H-1 event. The fact that all our records are from the same single sediment core ensures that the sequence of events and the temporal offsets between them are robust features independent of age modeling. In particular, we emphasize the similarity between the $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ and benthic $\delta^{13}\text{C}$ signals that underscores the significance of the changes in the rate of overturning inferred from $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$. Based on our age model the MOC decrease precedes the H1 event by as much as $\sim 1,200$ ($^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$) to $\sim 1,300$ yrs (benthic $\delta^{13}\text{C}$).

[8] Transient $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ increases associated with the E-1_{a-b} meltwater events suggest a direct link between circulation changes and local European sector derived meltwater perturbations. Taking the E-1_{a-b} planktonic $\delta^{18}\text{O}$ anomalies at face value suggests a salinity reduction of ~ 0.3 – 1.0 (freshwater endmember $\delta^{18}\text{O}$ $-35\text{\textperthousand}$ VSMOW), probably higher if accounting for the effects of sea surface cooling on planktonic $\delta^{18}\text{O}$. While salinity anomalies appear close to the magnitude of the H1_{MW} perturbation recorded at the DAPC2 site, at ~ 90 – 150 year duration these localized episodes are extremely short-lived. The possibility of additional sources of freshwater input at the time of MOC collapse remains for instance, from the small Iceland Ice Sheet and the much larger Laurentide and Greenland Ice Sheets [Hagen and Hald, 2002; Voelker et al., 2000] but the European provenance for the E-1_{a-b} meltwater perturbations is consistent with the reconstruction of a widespread salinity reduction west of Ireland at 15.0–17.7 ^{14}C ka BP (17.3–20.4 ka BP) that likewise has been linked to European melt sources [Sarnthein et al., 1995]. Moreover, the close linkage of the multi-step $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ increase documented in DAPC2 with the sequence of IRD and negative planktonic $\delta^{18}\text{O}$ events in the same core make a compelling case that while additional freshwater sources may have existed elsewhere in the region it were these brief freshwater pulses that played a decisive role in the sustained, but not total, drawdown of meridional overturning. Thus the H1 event and its associated major meltwater surge (H-1_{MW}) appear to have been embedded into an already slowed-down MOC and acted to sustain rather than initiate minimum MOC rates. Such lead of MOC slow down is supported by a similar temporal offset between the initial $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ increase after 19 ka BP in the deep subtropical Atlantic [McManus et al., 2004] and the initiation of surface ocean heat being retained in the tropical [Rühlemann et al., 1999] and subtropical [Flower et al., 2004] Atlantic that each significantly precede the age of H1 event. Continued surface ocean stratification is suggested in DAPC2 in the enhanced offset between the paired planktonic $\delta^{18}\text{O}$ records following H1 (Figure 2c) [Knutz et al., 2002] most likely indicating enhanced poleward transport of warm surface waters to the Nordic Sea, via the North Atlantic Drift, and that the location of deep convection has

shifted to north of the Iceland-Scotland Ridge [Rahmstorf, 1995], leaving the DAPC2 site under the influence of WTOW and an increasing recirculation of upper deep water [Knutz *et al.*, 2002].

4. Conclusions

[9] Our data demonstrate that catastrophic freshwater surges, e.g., during the past from the disintegrating Laurentide Ice Sheet, are not the sole component involved with major MOC change. Model simulations suggest that the MOC is sensitive to small-scale freshwater perturbations, on order of 0.1 Sv [Stocker and Wright, 1991; Manabe and Stouffer, 1997]. Our data from core DAPC2 indeed demonstrate that transient localized freshwater episodes occurred between 1,200–1,300 years prior to the large-scale H1 event and coincided with likewise transient phases of increasing $^{231}\text{Pa}_{\text{xs}}/^{230}\text{Th}_{\text{xs}}$ and decreasing benthic $\delta^{13}\text{C}$, both fully consistent with decreasing deep water ventilation and export. The events formed part of a multi-stepped shift toward a collapsed MOC, lending credence to numerical models that demonstrate the role of targeted freshwater events in forcing the MOC toward and across a threshold from where on convection enters a collapsed mode [Rahmstorf, 1994, 1995]. From this it appears that accelerated (yet non-catastrophic scale) melting of the Greenland Ice Sheet [Zwally *et al.*, 2002], may indeed bear significance for future MOC stability and climate in the wider North Atlantic region.

[10] **Acknowledgments.** We thank H. Medley, G. Bianchi, and J. Pike for technical assistance. The helpful comments of two anonymous reviewers improved this manuscript. This work was supported by funding from the U.K. National Environmental Research Council, U.S. National Science Foundation, Ministerio de Educación y Ciencia, Spain, and the Taiwan ROC National Science Council.

References

Bacon, M. P. (1988), Tracers of chemical scavenging in the ocean: Boundary effects and large scale chemical fractionation, *Philos. Trans. R. Soc. London, Ser. A*, 325, 147–160.

Broecker, W. S. (1979), A revised estimate of the radiocarbon age of the North Atlantic Deep Water, *J. Geophys. Res.*, 84, 3218–3226.

Flower, B. P., D. W. Hastings, H. W. Hill, and T. M. Quinn (2004), Phasing of deglacial warming and Laurentide Ice Sheet meltwater in the Gulf of Mexico, *Geology*, 32, 597–600.

Gherardi, J. M., L. Labeyrie, J. F. McManus, R. Francois, L. C. Skinner, and E. Cortijo (2005), Evidence from the Northeastern Atlantic basin for variability in the rate of the meridional overturning circulation through the last deglaciation, *Earth Planet. Sci. Lett.*, 240, 710–723.

Grousset, F. E., C. Pujol, L. Labeyrie, G. Auffret, and A. Boelaert (2000), Were the North Atlantic Heinrich events triggered by the behavior of the European ice sheets?, *Geology*, 28, 123–126.

Guo, L., M. Chen, and C. Gueguen (2002), Control of Pa/Th ratio by particulate chemical composition in the ocean, *Geophys. Res. Lett.*, 29(20), 1961, doi:10.1029/2002GL015666.

Hagen, S., and M. Hald (2002), Variation in surface and deep water circulation in the Denmark Strait, North Atlantic, during marine isotope stages 3 and 2, *Paleoceanography*, 17(4), 1061, doi:10.1029/2001PA000632.

Hemming, S. R. (2004), Heinrich events: Massive late Pleistocene detritus layers of the North Atlantic and their global climate imprint, *Rev. Geophys.*, 42, RG1005, doi:10.1029/2003RG000128.

Knutz, P. C., I. R. Hall, R. Zahn, T. L. Rasmussen, A. Kuijpers, M. Moros, and N. J. Shackleton (2002), Multidecadal ocean variability and NW European ice sheet surges during the last deglaciation, *Geochem. Geophys. Geosyst.*, 3(12), 1077, doi:10.1029/2002GC000351.

Kohfeld, K. E., R. G. Fairbanks, S. L. Smith, and I. D. Walsh (1996), *Neogloboquadrina pachyderma* (sinistral coiling) as paleoceanographic tracers in polar oceans: Evidence from Northeast Water Polynya plankton tows, sediment traps, and surface sediments, *Paleoceanography*, 11(6), 679–699.

Luo, S., and T.-L. Ku (2004), On the importance of opal, carbonate, and lithogenic clays in scavenging and fractionating ^{230}Th , ^{231}Pa , and ^{10}Be in the ocean, *Earth Planet. Sci. Lett.*, 220, 201–211.

Manabe, S., and R. J. Stouffer (1997), Coupled ocean-atmosphere model response to freshwater input: Comparison to Younger Dryas event, *Paleoceanography*, 12(2), 321–336.

Marchal, O., R. Francois, T. F. Stocker, and F. Joos (2000), Ocean thermohaline circulation and sedimentary $^{231}\text{Pa}/^{230}\text{Th}$ ratio, *Paleoceanography*, 15(6), 625–641.

McManus, J. F., R. Francois, J. M. Gherardi, L. D. Keigwin, and S. Brown-Leger (2004), Collapse and rapid resumption of Atlantic meridional circulation linked to deglacial climate changes, *Nature*, 428, 834–837.

New, A. L., and D. Smythe-Wright (2001), Aspects of the circulation in the Rockall Trough, *Cont. Shelf. Res.*, 21, 777–810.

Nozaki, Y., and T. Nakanishi (1985), ^{231}Pa and ^{230}Th in the open ocean water column, *Deep Sea Res., Part A*, 32, 1209–1220.

Rahmstorf, S. (1994), Rapid climate transitions in a coupled ocean-atmosphere model, *Nature*, 372, 82–85.

Rahmstorf, S. (1995), Bifurcation of the Atlantic thermohaline circulation in response to changes in the hydrological cycle, *Nature*, 378, 145–149.

Rickaby, R. E. M., and H. Elderfield (2005), Evidence from the high-latitude North Atlantic for variations in Antarctic Intermediate water flow during the last deglaciation, *Geochem. Geophys. Geosyst.*, 6, Q05001, doi:10.1029/2004GC000858.

Rühlmann, C., S. Mülitz, P. J. Müller, G. Wefer, and R. Zahn (1999), Warming of the tropical Atlantic Ocean and slowdown of thermohaline circulation during the last deglaciation, *Nature*, 402, 511–514.

Sarnthein, M., et al. (1995), Variations in Atlantic surface ocean paleoceanography, 50° – 80° N: A time-slice record of the last 30,000 years, *Paleoceanography*, 10(6), 1063–1094.

Scherer, R. P. (1995), A new method for the determination of absolute abundance of diatoms and other silt-sized sedimentary particles, *J. Paleolimnol.*, 12, 171–179.

Schiebel, R., J. Bijma, and C. Hemleben (1997), Population dynamics of the planktic foraminifer *Globigerina bulloides* from the eastern North Atlantic, *Deep Sea Res., Part I*, 44, 1701–1713.

Scourse, J. D., I. R. Hall, I. N. McCave, J. R. Young, and C. Sugdon (2000), The origin of Heinrich layers: Evidence from H2 for European precursor events, *Earth Planet. Sci. Lett.*, 182, 187–195.

Shen, C.-C., et al. (2002), Uranium and thorium isotopic and concentration measurements by magnetic sector inductively coupled plasma mass spectrometry, *Chem. Geol.*, 185, 165–178.

Shen, C.-C., et al. (2003), Measurement of attogram quantities of ^{231}Pa in dissolved and particulate fractions of seawater by isotope dilution thermal mass spectrometry, *Anal. Chem.*, 75, 1075–1079.

Siddall, M., G. M. Henderson, N. R. Edwards, S. A. Müller, T. F. Stocker, F. Joos, and M. Frank (2005), $^{231}\text{Pa}/^{230}\text{Th}$ fractionation by ocean transport, biogenic particle flux and particle type, *Earth Planet. Sci. Lett.*, 237, 137–155.

Stainforth, D. A., et al. (2005), Uncertainty in predictions of the climate response to rising levels of greenhouse gases, *Nature*, 433, 403–406.

Stocker, T. F., and D. G. Wright (1991), Rapid transitions of the ocean's deep circulation induced by changes in surface water fluxes, *Nature*, 351, 729–732.

Voelker, A. H. L., P. M. Grootes, M.-J. Nadeau, and M. Sarnthein (2000), Radiocarbon levels in the Iceland Sea from 25–53 kyr and their link to the Earth's magnetic field intensity, *Radiocarbon*, 42, 437–452.

Willamowski, C., and R. Zahn (2000), Upper ocean circulation in the glacial North Atlantic from benthic foraminiferal isotope and trace element fingerprinting, *Paleoceanography*, 15(5), 515–527.

Yu, E.-F., R. Francois, and M. P. Bacon (1996), Similar rates of modern and last-glacial ocean thermohaline circulation inferred from radiochemical data, *Nature*, 379, 689–694.

Zahn, R., and A. Stüber (2002), Suborbital intermediate water variability inferred from paired benthic foraminiferal Cd/Ca and $\delta^{13}\text{C}$ in the tropical West Atlantic and linking with North Atlantic climates, *Earth Planet. Sci. Lett.*, 200, 191–205.

Zahn, R., J. Schönfeld, H. Kudrass, M. Park, H. Erlenkeuser, and P. Grootes (1997), Thermohaline instability in the North Atlantic during meltwater events: Stable isotope and ice-rafted detritus records from core SO75-26KL, Portuguese margin, *Paleoceanography*, 12(5), 696–710.

Zwally, H. J., et al. (2002), Surface melt-induced acceleration of Greenland ice-sheet flow, *Science*, 297, 218–222.

R. L. Edwards, Department of Geology and Geophysics, University of Minnesota, 310 Pillsbury Drive SE, Minneapolis, MN 55455, USA.

I. R. Hall, School of Earth, Ocean and Planetary Sciences, Cardiff University, Main Building, Park Place, Cardiff CF10 3YE, UK. (hall@cardiff.ac.uk)

P. C. Knutz, Geological Survey of Denmark and Greenland, Øster Voldgade 10, 1350 Copenhagen, Denmark.

S. B. Moran, Graduate School of Oceanography, University of Rhode Island, South Ferry Road, Narragansett, RI 02882-1197, USA.

C.-C. Shen, Department of Geosciences, National Taiwan University, No 1 Sec 4, Roosevelt Road, Taipei 106, Taiwan.

R. Zahn, Institut de Ciencia i Tecnologia Ambientals, Universitat Autònoma de Barcelona, Edifici Cn - Campus UAB, Bellaterra E-08193, Spain.