

---

This is the **accepted version** of the journal article:

Martín i Pedret, Joaquim; Soria, Javier. «Characterization of rearrangement invariant spaces with fixed points for the Hardy-Littlewood maximal operator». *Annales Academiae Scientiarum Fennicae. Mathematica*, Vol. 31 (2006), p. 39-46.

---

This version is available at <https://ddd.uab.cat/record/271410>

under the terms of the  license

# Characterization of rearrangement invariant spaces with fixed points for the Hardy–Littlewood maximal operator

JOAQUIM MARTÍN<sup>§</sup> AND JAVIER SORIA<sup>¶</sup>

**Abstract.** We characterize the rearrangement invariant spaces for which there exists a non-constant fixed point, for the Hardy–Littlewood maximal operator (the case for the spaces  $L^p(\mathbb{R}^n)$  was first considered in [7]). The main result that we prove is that the space  $L^{\frac{n}{n-2},\infty}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  is minimal among those having this property.

## 1 Introduction

The centered Hardy–Littlewood maximal operator  $\mathcal{M}$  is defined on the Lebesgue space  $L^1_{\text{loc}}(\mathbb{R}^n)$  by

$$\mathcal{M}f(x) = \sup_{r>0} \frac{1}{|B_r|} \int_{B_r} |f(x-y)| dy,$$

where  $|B_r|$  denotes the measure of the Euclidean ball  $B_r$  centered at the origin of  $\mathbb{R}^n$ .

In this paper we study the existence of non-constant fixed points of the maximal operator  $\mathcal{M}$  (i.e.,  $\mathcal{M}f = f$ ) in the framework of the rearrangement invariant (r.i.) functions spaces (see Section 2 below). We will use some of the estimates proved in [7], where the case  $L^p(\mathbb{R}^n)$  was studied, and show that they can be sharpened to obtain all the rearrangement invariant norms with this property (in particular we extend Korry’s result to the end point case  $p = n/(n-2)$ , where the weak-type spaces have to be considered.) The main argument behind this problem is the existence of a minimal space  $L^{\frac{n}{n-2},\infty}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  contained in all the r.i. spaces with the fixed point property.

## 2 Background on Rearrangement Invariant Spaces

Since we work in the context of rearrangement invariant spaces it will be convenient to start by reviewing some basic definitions about these spaces.

A rearrangement invariant space  $X = X(\mathbb{R}^n)$  (r.i. space) is a Banach function space on  $\mathbb{R}^n$  endowed with a norm  $\|\cdot\|_{X(\mathbb{R}^n)}$  such that

$$\|f\|_{X(\mathbb{R}^n)} = \|g\|_{X(\mathbb{R}^n)}$$

---

<sup>§</sup>Research partially supported by Grants MTM2004-02299, 2001SGR00069 and by Programa Ramón y Cajal (MCYT).

<sup>¶</sup>Research partially supported by Grants MTM2004-02299, 2001SGR00069.

**Keywords:** Fixed points, fractional integrals, maximal operators, rearrangement invariant spaces, super-harmonic functions.

**MSC2000:** 42B25, 46E30.

whenever  $f^* = g^*$ . Here  $f^*$  stands for the non-increasing rearrangement of  $f$ , i.e., the non-increasing, right-continuous function on  $[0, \infty)$  equimeasurable with  $f$ .

An r.i. space  $X(\mathbb{R}^n)$  has a representation as a function space on  $\bar{X}(0, \infty)$  such that

$$\|f\|_{X(\mathbb{R}^n)} = \|f^*\|_{\bar{X}(0, \infty)}.$$

Any r.i. space is characterized by its **fundamental function**

$$\phi_X(s) = \|\chi_E\|_{X(\mathbb{R}^n)}$$

(here  $E$  is any subset of  $\mathbb{R}^n$  with  $|E| = s$ ) and the **fundamental indices**

$$\bar{\beta}_X = \inf_{s>1} \frac{\log M_X(s)}{\log s} \quad \text{and} \quad \underline{\beta}_X = \sup_{s<1} \frac{\log M_X(s)}{\log s},$$

where

$$M_X(s) = \sup_{t>0} \frac{\phi_X(ts)}{\phi_X(t)}, \quad s > 0.$$

It is well known that

$$0 \leq \underline{\beta}_X \leq \bar{\beta}_X \leq 1.$$

(We refer the reader to [2] for further information about r.i. spaces.)

### 3 Main result

Before formulating our main result, it will be convenient to start with the following remarks (see [7]):

**Remark 3.1** By the Lebesgue's differentiation theorem one easily obtains that

$$|f(x)| \leq \mathcal{M}f(x) \text{ a.e. } x \in \mathbb{R}^n,$$

thus  $f$  is a fixed point of  $\mathcal{M}$ , if and only if  $f$  is positive and

$$\frac{1}{|B(x, r)|} \int_{B(x, r)} f(y) dy \leq f(x) \text{ a.e. } x \in \mathbb{R}^n,$$

or equivalently  $f$  is a positive **super-harmonic** function (i.e.  $\Delta f \leq 0$ , where  $\Delta$  is the Laplacian operator).

**Remark 3.2** If  $f$  is a non-constant fixed point of  $\mathcal{M}$ , and  $\varphi \geq 0$  belongs to the Schwartz class  $\mathcal{S}(\mathbb{R}^n)$ , with  $\int_{\mathbb{R}^n} \varphi(x) dx = 1$ , then the function  $f_t(x) = (f * \varphi_t)(x)$ , with  $\varphi_t(x) = t^{-n} \varphi(x/t)$  is also a non-constant fixed point of  $\mathcal{M}$  which belongs to  $\mathcal{C}^\infty(\mathbb{R}^n)$  (notice that using the Lebesgue differentiation theorem, there exists some  $t > 0$  such that  $f_t$  is non-constant, since  $f$  is non-constant). In particular if  $X(\mathbb{R}^n)$  is an r.i. space and  $f \in X(\mathbb{R}^n)$  is a non-constant fixed point of  $\mathcal{M}$ , since  $\mathcal{S}(\mathbb{R}^n) \subset L^1(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$  we get that  $f_t \in X(\mathbb{R}^n) \cap \mathcal{C}^\infty(\mathbb{R}^n)$  is a non-constant fixed point of  $\mathcal{M}$ .

**Remark 3.3** Using the theory of weighted inequalities for  $\mathcal{M}$  (see [6]), if  $\mathcal{M}f = f$ , in particular  $f \in A_1$  (the Muckenhoupt weight class), and hence  $f(x) dx$  defines a doubling measure. Hence,  $f \notin L^1(\mathbb{R}^n)$ . Also, using the previous remark we see that if  $f \in L^p(\mathbb{R}^n)$  is a fixed point, then  $f \in L^q(\mathbb{R}^n)$ , for all  $p \leq q \leq \infty$ .

**Definition 3.4** Given an r.i. space  $X(\mathbb{R}^n)$ , we define

$$D_{I_2}(X(\mathbb{R}^n)) = \left\{ f \in L^0(\mathbb{R}^n) : \|I_2 f\|_{X(\mathbb{R}^n)} < \infty \right\},$$

where  $I_2$  is the Riesz potential,

$$(I_2 f)(x) = \int_{\mathbb{R}^n} |x - y|^{2-n} f(y) dy.$$

It is not hard to see that the space  $D_{I_2}(X(\mathbb{R}^n))$  is either trivial or is the largest r.i. space which is mapped by  $I_2$  into  $X(\mathbb{R}^n)$ , and is also related with the theory of the optimal Sobolev embeddings (see [4] and the references quoted therein).

**Theorem 3.5** Let  $X(\mathbb{R}^n)$  be an r.i. space. The following statements are equivalent:

1. There is a non-constant fixed point  $f \in X(\mathbb{R}^n)$  of  $\mathcal{M}$
2.  $n \geq 3$  and  $|x|^{2-n} \chi_{\{|x|>1\}}(x) \in X(\mathbb{R}^n)$ .
3.  $n \geq 3$  and  $\chi_{[0,1]}(t) + t^{2/n-1} \chi_{[1,\infty)}(t) \in \bar{X}(0, \infty)$ .
4.  $n \geq 3$  and  $(L^{\frac{n}{n-2}, \infty}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)) \subset X(\mathbb{R}^n)$ .
5.  $n \geq 3$  and  $D_{I_2}(X(\mathbb{R}^n)) \neq \{0\}$ .

**Proof.** (1  $\rightarrow$  2) Since if  $n = 1$  or  $n = 2$ , the only positive super-harmonic functions are the constant functions (see [8, Remark 1, p. 210]), necessarily  $n \geq 3$ . Moreover, it is proved in [7] that, if  $f \in \mathcal{C}^\infty(\mathbb{R}^n)$  is a non-constant fixed point of  $\mathcal{M}$ , then

$$f(x) \geq c |x|^{2-n} \chi_{\{|x|>1\}}(x).$$

Since  $f \in X(\mathbb{R}^n)$ , then  $|x|^{2-n} \chi_{\{|x|>1\}}(x) \in X(\mathbb{R}^n)$ .

(2  $\rightarrow$  3) Since if  $|x|^{2-n} \chi_{\{|x|>1\}}(x) \in X(\mathbb{R}^n)$ , then

$$F(x) = \chi_{\{|x|\leq 1\}}(x) + |x|^{2-n} \chi_{\{|x|>1\}}(x) \in X(\mathbb{R}^n).$$

An easy computation shows that

$$F^*(t) \simeq \chi_{[0,1]}(t) + t^{2/n-1} \chi_{[1,\infty)}(t).$$

(3  $\rightarrow$  4) Since  $f \in (L^{\frac{n}{n-2},\infty}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n))$  if and only if

$$\sup_{t>0} f^*(t)W(t) < \infty,$$

where  $W(t) = \max(1, t^{1-2/n})$ , we have that

$$f^*(t) \leq \|f\|_{L^{\frac{n}{n-2},\infty}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)} W^{-1}(t)$$

and since  $W^{-1}(t) = \chi_{[0,1]}(t) + t^{2/n-1}\chi_{[1,\infty)} \in \bar{X}(0, \infty)$  we have that

$$\|f\|_{X(\mathbb{R}^n)} = \|f^*\|_{\bar{X}(0,\infty)} \leq c \|f\|_{L^{\frac{n}{n-2},\infty}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)}$$

with  $c = \|W^{-1}\|_{\bar{X}(0,\infty)}$ .

(4  $\rightarrow$  5) Since (see [9] and [1])

$$(I_2 f)^*(t) \leq c_1 \left( t^{2/n-1} \int_0^t f^*(s) ds + \int_t^\infty f^*(s) s^{2/n-1} ds \right) \leq c_2 (I_2 f^0)^*(t)$$

where  $f^0(x) = f^*(c_n |x|^n)$ ,  $c_n = \text{measure of the unit ball in } \mathbb{R}^n$ . (Observe that  $(f^0)^* = f^*$ ). Rewriting the middle term in the above inequalities, using Fubini's theorem, we get

$$(I_2 f)^*(t) \leq d_1 \left( \frac{n}{n-2} \int_t^\infty f^{**}(s) s^{2/n-1} ds \right) \leq d_2 (I_2 f^0)^*(t),$$

where  $f^{**}(t) = t^{-1} \int_0^t f^*(s) ds$ . Thus,  $f \in D_{I_2}(X(\mathbb{R}^n))$  if and only if

$$\left\| \int_t^\infty f^{**}(s) s^{2/n-1} ds \right\|_{\bar{X}(0,\infty)} < \infty. \quad (1)$$

Since

$$F(t) = \int_t^\infty \chi_{[0,1]}^{**}(s) s^{2/n-1} ds = c(\chi_{[0,1]}(t) + t^{2/n-1}\chi_{[1,\infty)}(t))$$

is a decreasing function, and

$$F^0(x) = F(c_n |x|^n) \simeq (\chi_{\{x:|x|\leq 1\}}(x) + |x|^{2-n} \chi_{\{x:|x|>1\}}(x)) \in L^{\frac{n}{n-2},\infty}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$$

we get that  $\chi_{[0,1]}^\circ \in D_{I_2}(X(\mathbb{R}^n))$ .

Another argument to prove this part is the following:

Since, if  $n \geq 3$  (see [2, Theorem 4.18, p. 228])

$$I_2 : L^1(\mathbb{R}^n) \rightarrow L^{\frac{n}{n-2},\infty}(\mathbb{R}^n) \text{ and } I_2 : L^{\frac{n}{2},1}(\mathbb{R}^n) \rightarrow L^\infty(\mathbb{R}^n)$$

is bounded, we have that

$$I_2 : (L^1(\mathbb{R}^n) \cap L^{\frac{n}{2},1}(\mathbb{R}^n)) \rightarrow (L^{\frac{n}{n-2},\infty}(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)) \subset X(\mathbb{R}^n)$$

is bounded, and hence  $L^1(\mathbb{R}^n) \cap L^{\frac{n}{2},1}(\mathbb{R}^n) \subset D_{I_2}(X(\mathbb{R}^n))$ .

(5  $\rightarrow$  1) Since  $n \geq 3$ , we can use the classical formula of potential theory (see [10, p. 126])

$$-h = \Delta(I_2 h)$$

to conclude that there is a positive function  $f = I_2 \chi_{[0,1]}^\circ \in X(\mathbb{R}^n)$ . Then  $0 \leq f_t = I_2(\chi_{[0,1]}^\circ * \varphi_t) \in X(\mathbb{R}^n) \cap \mathcal{C}^\infty(\mathbb{R}^n)$  and  $\Delta f_t \leq 0$ . ■

We now consider particular examples, like the Lorentz spaces:

**Corollary 3.6** *Let  $1 \leq p < \infty$ , and assume  $\Lambda^p(\mathbb{R}^n, w)$  is a Banach space (i.e.,  $w \in B_p$  if  $1 < p < \infty$  or  $p \in B_{1,\infty}$  if  $p = 1$ , see [3]). Then, there exists a non-constant function  $f \in \Lambda^p(\mathbb{R}^n, w)$  such that  $\mathcal{M}(f) = f$  if and only if  $n \geq 3$  and*

$$\int_1^\infty \frac{w(t)}{t^{p(1-2/n)}} dt < \infty.$$

*In particular, this condition always holds, for  $p > 1$  and  $n$  large enough.*

**Proof.** The integrability condition follows by using the previous theorem. Now, if  $w \in B_p$ , then there exists an  $\varepsilon > 0$  such that  $w \in B_{p-\varepsilon}$ , and hence, it suffices to take  $n > 2/\varepsilon$ . Observe that if  $w = 1$  and  $p = 1$ , then  $\Lambda^1(\mathbb{R}^n, w) = L^1(\mathbb{R}^n)$ , which does not have the fixed point property for any dimension  $n$ . ■

**Corollary 3.7** *Let  $1 \leq p, q \leq \infty$  (if  $p = 1$  we only consider  $q = 1$ ). Then, there exists a non-constant function  $f \in L^{p,q}(\mathbb{R}^n)$  such that  $\mathcal{M}(f) = f$  if and only if  $n \geq 3$  and*

$$\begin{cases} n/(n-2) < p \leq \infty \\ \text{or} \\ p = n/(n-2) \text{ and } q = \infty. \end{cases}$$

**Corollary 3.8** (See [7]) *Let  $1 \leq p \leq \infty$ . There exists a non-constant function  $f \in L^p(\mathbb{R}^n)$  such that  $\mathcal{M}(f) = f$  if and only if  $n \geq 3$  and  $n/(n-2) < p \leq \infty$ .*

It is interesting to know when given an r.i. space  $X(\mathbb{R}^n)$ , the space  $D_{I_2}(X(\mathbb{R}^n))$  is not trivial, or equivalently

$$\overline{D_{I_2}(X(\mathbb{R}^n))} := \left\{ f \in L^0([0, \infty)) : \left\| \int_t^\infty f^{**}(s) s^{2/n-1} ds \right\|_{\bar{X}(0, \infty)} < \infty \right\} \quad (2)$$

is not trivial. This will be done in terms of the fundamental indices of  $X$ . We start by computing the fundamental function of  $D_{I_2}(X(\mathbb{R}^n))$ .

**Lemma 3.9** *Let  $X$  be an r.i. space on  $\mathbb{R}^n$ ,  $n \geq 3$ . Let  $Y$  be given by (2). Then*

$$\phi_Y(s) \simeq s^{n/2} \|P_{1-2/n} \chi_{[0,s]}\|_X$$

where  $P_{1-2/n} f(t) = t^{2/n-1} \int_0^t f(s) s^{-2/n} ds$ .

**Proof.**

$$\begin{aligned} s^{n/2} P_{1-2/n} \chi_{[0,s]}(t) &\simeq s^{n/2} (\chi_{[0,s]}(t) + \left(\frac{s}{t}\right)^{1-2/n} \chi_{[s,\infty)}(t)) \\ &\simeq \int_t^\infty \chi_{[0,s]}^{**}(r) r^{2/n-1} dr. \end{aligned}$$

■

**Theorem 3.10** *Let  $X$  be an r.i. space on  $\mathbb{R}^n$ ,  $n \geq 3$ . Let  $Y$  be given by (2). Then*

1. *If  $\bar{\beta}_X < 1 - 2/n$ , then  $Y \neq \{0\}$ .*
2. *If  $Y \neq \{0\}$  then  $\underline{\beta}_X \leq 1 - 2/n$ .*

**Proof.** 1.) Let  $\chi_r = \chi_{[0,r]}$ . Then

$$P_{1-2/n}\chi_r(t) = \int_0^1 \chi_r(\xi t) \frac{d\xi}{\xi^{n/2}} \leq c \sum_{k=0}^{\infty} 2^{-k(1-n/2)} \chi_{2^k r}(t).$$

Thus

$$\|P_{1-2/n}\chi_r\|_X \leq c \sum_{k=0}^{\infty} 2^{-k(1-n/2)} \phi_X(2^k r) \leq c \phi_X(r) \sum_{k=0}^{\infty} 2^{-k(1-n/2)} M_X(2^k).$$

Let  $\varepsilon > 0$  be such that  $\bar{\beta}_X + \varepsilon < 1 - 2/n$ . Then by the definition of  $\bar{\beta}_X$  it follows readily that there is a constant  $c > 0$  such that

$$M_X(2^k) \leq c 2^{k(\bar{\beta}_X + \varepsilon)},$$

and hence

$$\sum_{k=0}^{\infty} 2^{-k(1-n/2)} M_X(2^k) \leq \sum_{k=0}^{\infty} 2^{-k(1-n/2-\bar{\beta}_X-\varepsilon)} < \infty,$$

which implies that  $\chi_r \in Y$ .

2.) Since  $Y \neq \{0\}$  if and only if  $\|P_{1-2/n}\chi_{[0,1]}\|_X < \infty$  and

$$\sup_{t>0} (P_{1-2/n}\chi_{[0,1]})^{**}(t) \phi_X(t) \leq \|P_{1-2/n}\chi_{[0,1]}\|_X < \infty, \quad (3)$$

and easy computations show that (3) implies that

$$1 \leq \sup_{t \geq 1} \frac{\phi_X(t)}{t^{1-2/n}} = c < \infty, \quad (4)$$

then, by (4)

$$\begin{aligned} M_X(a) &= \max \left( \sup_{t \geq 1/a} \frac{\phi_X(ta)}{\phi_X(t)}, \sup_{t < 1/a} \frac{\phi_X(ta)}{\phi_X(t)} \right) \\ &= \max \left( \sup_{t \geq 1/a} \frac{\phi_X(ta)}{(at)^{1-2/n}} \frac{(at)^{1-2/n}}{\phi_X(t)}, \sup_{t < 1/a} \frac{\phi_X(ta)}{\phi_X(t)} \right) \\ &\simeq \max \left( a^{1-2/n} \sup_{t \geq 1/a} \frac{t^{1-2/n}}{\phi_X(t)}, \sup_{t < 1/a} \frac{\phi_X(ta)}{\phi_X(t)} \right). \end{aligned}$$

Thus, if  $a < 1$ , using again (4) we get

$$M_X(a) \geq a^{1-2/n} \sup_{t \geq 1/a} \frac{t^{1-2/n}}{\phi_X(t)} \geq a^{1-2/n}$$

which implies that

$$\underline{\beta}_X \leq 1 - 2/n.$$

■

Let us see that the converse in the previous theorem is not true.

**Proposition 3.11** *There are rearrangement invariant spaces  $X$  such that*

$$1. Y \neq \{0\} \text{ and } \overline{\beta}_X \geq 1 - 2/n.$$

$$2. Y = \{0\} \text{ and } \underline{\beta}_X < 1 - 2/n.$$

**Proof.** Let  $\varphi(t) = t^a \chi_{[0,1]}(t) + t^b \chi_{[1,\infty)}(t)$ , with  $0 \leq a, b \leq 1$ . Let

$$X = \left\{ f \in L^0([0, \infty)) : \sup_{t>0} f^{**}(t) \varphi(t) < \infty \right\}.$$

Since  $\varphi$  is a quasi-concave function, we have that

$$\varphi(t) = \phi_X(t)$$

and

$$\underline{\beta}_X = \min(a, b), \overline{\beta}_X = \max(a, b).$$

On the other hand, the space  $Y$  defined by (2) is not trivial if and only if

$$b \leq 1 - 2/n.$$

Now, to prove 1) take  $b \leq 1 - 2/n$  and  $a \geq 1 - 2/n$ . And to see 2) take  $b > 1 - 2/n$  and  $a \leq 1 - 2/n$ . ■

**Remark 3.12** If we consider

$$X_0 = \left\{ f \in L^0([0, \infty)) : \sup_{t>0} f^{**}(t) t^{1-2/n} (1 + \log^+ t) < \infty \right\}$$

and

$$X_1 = \left\{ f \in L^0([0, \infty)) : \sup_{t>0} f^{**}(t) \frac{t^{1-2/n}}{(1 + \log^+ t)} < \infty \right\}$$

then  $\underline{\beta}_{X_i} = \overline{\beta}_{X_i} = 1 - 2/n$ ,  $Y_0 = \{0\}$  and  $Y_1 \neq \{0\}$ .

**Remark 3.13** It was proved in [7] that if we consider the strong maximal function (i.e, the maximal operator associated to centered intervals in  $\mathbb{R}^n$ ), then there were no fixed points in any  $L^p(\mathbb{R}^n)$  space, regardless of the dimension. The same argument works to show that  $L^p(\mathbb{R}^n)$  cannot be replaced by any different r.i. space. Also, if we study this question for other kind of sets, like, e.g., Buseman–Feller differentiation bases (see [5]), then the only possible fixed points are the constant functions. This observation applies to any non-centered maximal operator (with respect to balls, cubes, etc.)



## References

- [1] C. Bennett and K. Rudnick, *On Lorentz-Zygmund Spaces*, Dissertationes Math. (Rozprawy Mat.) **175**, 1980.
- [2] C. Bennett and R. Sharpley, *Interpolation of Operators*, Academic Press, Boston, 1988.
- [3] M. Carro, L. Pick, J. Soria, and V. Stepanov, *On embeddings between classical Lorentz spaces*, Math. Inequal. Appl. **4** (2001), 397–428.
- [4] D. E. Edmunds, R. Kerman and L. Pick, *Optimal Sobolev imbeddings involving rearrangement-invariant quasinorms*, J. Funct. Anal. **170** (2000), 307–355.
- [5] M. de Guzmán, *Differentiation of Integrals in  $\mathbb{R}^n$* , Lecture Notes in Mathematics **481**, Springer-Verlag, Berlin-New York, 1975.
- [6] J. García-Cuerva and J.L. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland Mathematics Studies **116**, North-Holland Publishing Co., Amsterdam, 1985.
- [7] S. Korry, *Fixed points of the Hardy-Littlewood maximal operator*, Collect. Math. **52** (2001), 289–294.
- [8] E. H. Lieb and M. Loss, *Analysis*, American Mathematical Society, Providence, 2001.
- [9] E. Sawyer, *Boundedness of classical operators on classical Lorentz spaces*, Studia Math. **96** (1990), 145–158.
- [10] E. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, New Jersey, 1970.

Joaquim Martín  
Dept. Mathematics  
U. Autònoma de Barcelona  
E-08193 Bellaterra (Barcelona), SPAIN    *E-mail:* jmartin@mat.uab.es

Javier Soria  
Dept. Appl. Math. and Analysis  
University of Barcelona  
E-08071 Barcelona, SPAIN    *E-mail:* soria@mat.ub.es