

FLOW CURVATURE METHOD APPLIED TO CANARD EXPLOSION

JEAN-MARC GINOUX¹ AND JAUME LLIBRE²

ABSTRACT. The aim of this work is to establish that the bifurcation parameter value leading to a *canard explosion* in dimension two obtained by the so-called *Geometric Singular Perturbation Method* can be found according to the *Flow Curvature Method*. This result will be then exemplified with the classical Van der Pol oscillator.

1. INTRODUCTION

The classical geometric theory of differential equations developed originally by Andronov [1], Tikhonov [30] and Levinson [23] stated that *singularly perturbed systems* possess *invariant manifolds* on which trajectories evolve slowly, and toward which nearby orbits contract exponentially in time (either forward or backward) in the normal directions. These manifolds have been called asymptotically stable (or unstable) *slow invariant manifolds*¹. Then, Fenichel [11, 12, 13, 14] theory² for the *persistence of normally hyperbolic invariant manifolds* enabled to establish the *local invariance* of *slow invariant manifolds* that possess both expanding and contracting directions and which were labeled *slow invariant manifolds*.

During the last century, various methods have been developed to compute the *slow invariant manifold* or, at least an asymptotic expansion in power of ε .

The seminal works of Wasow [32], Cole [6], O’Malley [25, 26] and Fenichel [11, 12, 13, 14] to name but a few, gave rise to the so-called *Geometric Singular Perturbation Method*. According to this theory, existence as well as local invariance of the *slow invariant manifold* of *singularly perturbed systems* has been stated. Then, the determination of the *slow invariant manifold* equation turned into a regular perturbation problem in which one generally expected the asymptotic validity of such expansion to breakdown [26].

Recently a new approach of n -dimensional singularly perturbed dynamical systems of ordinary differential equations with two time scales, called *Flow Curvature Method* has been developed [17]. In dimension two and three, it consists in considering the *trajectory curves* integral of such systems as *plane* or *space* curves. Based on the use of local metrics properties of *curvature* (*first curvature*) and *torsion* (*second curvature*) resulting from the *Differential Geometry*, this method which does not require the use of asymptotic expansions, states that the location of the points

Key words and phrases. Geometric Singular Perturbation Method, Flow Curvature Method, singularly perturbed dynamical systems, canard solutions.

¹In other articles the *slow manifold* is the approximation of order $O(\varepsilon)$ of the *slow invariant manifold*.

²The theory of invariant manifolds for an ordinary differential equation is based on the work of Hirsch, *et al.* [19]

where the local *curvature* (resp. *torsion*) of *trajectory curves* of such systems, vanishes, directly provides an approximation of the *slow invariant manifold* associated with two-dimensional (resp. three-dimensional) *singularly perturbed systems* up to suitable order $O(\varepsilon^2)$ (resp. $O(\varepsilon^3)$). This method gives an implicit non intrinsic equation, because it depends on the euclidean metric.

Solutions of “canard” type have been discovered by a group of French mathematicians [2] in the beginning of the eighties while they were studying relaxation oscillations in the classical Van der Pol’s equation (with a constant forcing term) [31]. They observed, within a small range of the control parameter, a fast transition for the amplitude of the limit cycle varying suddenly from small amplitude to a large amplitude. Due to the fact that the shape of the limit cycle in the phase plane looks as a duck they called it “canard cycle”. Hence, they named this new phenomenon “canard explosion³” and triggered a “duck-hunting”.

Many methods have been developed to analyze “canard” solution such as non-standard analysis [2, 8], matched asymptotic expansions [10], or the blow-up technique [9, 22, 28] which extends the *Geometric Singular Perturbation Method* [11, 12, 13, 14].

Meanwhile, two other geometric approaches have been proposed. The first, elaborated by [4] involves *inflection curves*, while the second makes use of the *curvature* of the trajectory curve, integral of any n -dimensional singularly perturbed dynamical system [16, 17]. This latter, entitled *Flow Curvature Method* will be used in this work in order to compute the bifurcation parameter value leading to a *canard explosion*. Moreover, the total correspondence between the results obtained in this paper for two-dimensional singularly perturbed dynamical systems such as Van der Pol oscillator and those previously established by [2] will enable to highlight another link between the *Flow Curvature Method* and the *Geometric Singular Perturbation Method*.

2. SINGULARLY PERTURBED SYSTEMS

According to Tikhonov [30], Takens [29], Jones [20] and Kaper [21] *singularly perturbed systems* may be defined such as:

$$(1) \quad \begin{aligned} \vec{x}' &= \vec{f}(\vec{x}, \vec{z}, \varepsilon), \\ \vec{z}' &= \varepsilon \vec{g}(\vec{x}, \vec{z}, \varepsilon). \end{aligned}$$

where $\vec{x} \in \mathbb{R}^m$, $\vec{z} \in \mathbb{R}^p$, $\varepsilon \in \mathbb{R}^+$, and the prime denotes differentiation with respect to the independent variable t . The functions \vec{f} and \vec{g} are assumed to be C^∞ functions⁴ of \vec{x} , \vec{z} and ε in $U \times I$, where U is an open subset of $\mathbb{R}^m \times \mathbb{R}^p$ and I is an open interval containing $\varepsilon = 0$.

In the case when $0 < \varepsilon \ll 1$, i.e., is a small positive number, the variable \vec{x} is called *fast* variable, and \vec{z} is called *slow* variable. Using Landau’s notation: $O(\varepsilon^k)$ represents a function f of x and ε such that $f(u, \varepsilon)/\varepsilon^k$ is bounded for positive ε going to zero, uniformly for u in the given domain.

³According to Krupa and Szmolyan [22, p. 312] this terminology has been introduced in chemical and biological literature by Brøns and Bar-Eli [3, p. 8707] to denote a sudden change of amplitude and period of oscillations under a very small range of control parameter.

⁴In certain applications these functions will be supposed to be C^r , $r \geq 1$.

It is used to consider that generally \vec{x} evolves at an $O(1)$ rate; while \vec{z} evolves at an $O(\varepsilon)$ *slow* rate. Reformulating system (1) in terms of the rescaled variable $\tau = \varepsilon t$, we obtain

$$(2) \quad \begin{aligned} \varepsilon \dot{\vec{x}} &= \vec{f}(\vec{x}, \vec{z}, \varepsilon), \\ \dot{\vec{z}} &= \vec{g}(\vec{x}, \vec{z}, \varepsilon). \end{aligned}$$

The dot represents the derivative with respect to the new independent variable τ .

The independent variables t and τ are referred to the *fast* and *slow* times, respectively, and (1) and (2) are called the *fast* and *slow* systems, respectively. These systems are equivalent whenever $\varepsilon \neq 0$, and they are labeled *singular perturbation problems* when $0 < \varepsilon \ll 1$. The label “singular” stems in part from the discontinuous limiting behavior in system (1) as $\varepsilon \rightarrow 0$.

In such case system (2) leads to a differential-algebraic system called *reduced slow system* whose dimension decreases from $m + p = n$ to p . Then, the *slow* variable $\vec{z} \in \mathbb{R}^p$ partially evolves in the submanifold M_0 called the *critical manifold*⁵ and defined by

$$(3) \quad M_0 := \left\{ (\vec{x}, \vec{z}) : \vec{f}(\vec{x}, \vec{z}, 0) = \vec{0} \right\}.$$

When $D_x f$ is invertible, thanks to implicit function theorem, M_0 is given by the graph of a C^∞ function $\vec{x} = \vec{F}_0(\vec{z})$ for $\vec{z} \in D$, where $D \subseteq \mathbb{R}^p$ is a compact, simply connected domain and the boundary of D is an $(p - 1)$ -dimensional C^∞ submanifold⁶.

According to Fenichel theory [11, 12, 13, 14] if $0 < \varepsilon \ll 1$ is sufficiently small, then there exists a function $\vec{F}(\vec{z}, \varepsilon)$ defined on D such that the manifold

$$(4) \quad M_\varepsilon := \left\{ (\vec{x}, \vec{z}) : \vec{x} = \vec{F}(\vec{z}, \varepsilon) \right\},$$

is locally invariant under the flow of system (1). Moreover, there exist perturbed local stable (or attracting) M_a and unstable (or repelling) M_r branches of the *slow invariant manifold* M_ε . Thus, normal hyperbolicity of M_ε is lost via a saddle-node bifurcation of the *reduced slow system* (2).

Definition 1. A “*canard*” is a solution of a singularly perturbed dynamical system following the attracting branch M_a of the slow invariant manifold, passing near a bifurcation point located on the fold of the critical manifold, and then following the repelling branch M_r of the slow invariant manifold during a considerable amount of time.

Geometrically a *maximal canard* corresponds to the intersection of the attracting and repelling branches $M_a \cap M_r$ of the slow manifold in the vicinity of a non-hyperbolic point. Canards are a special class of solution of singularly perturbed dynamical systems for which normal hyperbolicity is lost.

⁵It corresponds to the approximation of the slow invariant manifold, with an error of $O(\varepsilon)$.

⁶The set D is overflowing invariant with respect to (2) when $\varepsilon = 0$.

3. GEOMETRIC SINGULAR PERTURBATION METHOD

Earliest geometric approaches to *singularly perturbed dynamical systems* have been developed by Cole [6], O'Malley [25, 26], Fenichel [11, 12, 13, 14] for the determination of the *slow manifold* equation.

Geometric Singular Perturbation Method is based on the following assumptions and theorem stated by Nils Fenichel in the middle of the seventies⁷.

3.1. Assumptions.

- (H₁) Functions \vec{f} and \vec{g} are C^∞ functions in $U \times I$, where U is an open subset of $\mathbb{R}^m \times \mathbb{R}^p$ and I is an open interval containing $\varepsilon = 0$.
- (H₂) There exists a set M_0 that is contained in $\{(\vec{x}, \vec{z}) : \vec{f}(\vec{x}, \vec{z}, 0) = 0\}$ such that M_0 is a compact manifold with boundary and M_0 is given by the graph of a C^1 function $\vec{x} = \vec{F}_0(\vec{z})$ for $\vec{z} \in D$, where $D \subseteq \mathbb{R}^p$ is a compact, simply connected domain and the boundary of D is an $(p - 1)$ -dimensional C^∞ submanifold. Finally, the set D is overflowing invariant with respect to (2) when $\varepsilon = 0$.
- (H₃) M_0 is normally hyperbolic relative to the *reduced fast system* and in particular it is required for all points $\vec{p} \in M_0$, that there are k (resp. l) eigenvalues of $D_{\vec{x}}\vec{f}(\vec{p}, 0)$ with positive (resp. negative) real parts bounded away from zero, where $k + l = m$.

Theorem 2 (Fenichel's Persistence Theorem). *Let system (1) satisfying the conditions (H₁) – (H₃). If $\varepsilon > 0$ is sufficiently small, then there exists a function $\vec{F}(\vec{z}, \varepsilon)$ defined on D such that the manifold $M_\varepsilon = \{(\vec{x}, \vec{z}) : \vec{x} = \vec{F}(\vec{z}, \varepsilon)\}$ is locally invariant under (1). Moreover, $\vec{F}(\vec{z}, \varepsilon)$ is C^r , and M_ε is C^r $O(\varepsilon)$ close to M_0 . In addition, there exist perturbed local stable and unstable manifolds of M_ε . They are unions of invariant families of stable and unstable fibers of dimensions l and k , respectively, and they are C^r $O(\varepsilon)$ close to their counterparts.*

Proof. See [11], [20] and [21]. □

3.2. Invariance. Generally, Fenichel theory enables to turn the problem for explicitly finding functions $\vec{x} = \vec{F}(\vec{z}, \varepsilon)$ whose graphs are locally *slow invariant manifolds* M_ε of system (1) into regular perturbation problem. Invariance of the manifold M_ε implies that $\vec{F}(\vec{z}, \varepsilon)$ satisfies:

$$(5) \quad \varepsilon D_{\vec{z}}\vec{F}(\vec{z}, \varepsilon) \vec{g}\left(\vec{F}(\vec{z}, \varepsilon), \vec{z}, \varepsilon\right) = \vec{f}\left(\vec{F}(\vec{z}, \varepsilon), \vec{z}, \varepsilon\right).$$

Then, plugging the perturbation expansion:

$$\vec{F}(\vec{z}, \varepsilon) = \sum_{i=0}^{N-1} \vec{F}_i(\vec{z}) \varepsilon^i + O(\varepsilon^N)$$

into (5) enables to solve order by order for $\vec{F}(\vec{z}, \varepsilon)$.

⁷For an introduction to Geometric Singular Perturbation Method see [21].

Taylor series expansion for $\vec{f}(\vec{F}(\vec{z}, \varepsilon), \vec{z}, \varepsilon)$ up to terms of order two in ε leads at order ε^0 to

$$(6) \quad \vec{f}(\vec{F}_0(\vec{z}), \vec{z}, 0) = \vec{0}$$

which defines $\vec{F}_0(\vec{z})$ due to the invertibility of $D_{\vec{x}}\vec{f}$ and the *Implicit Function Theorem*.

At order ε^1 we have:

$$(7) \quad D_{\vec{z}}\vec{F}_0\vec{g}(\vec{F}_0, \vec{z}, 0) = D_{\vec{x}}\vec{f}(\vec{F}_0, \vec{z}, 0)\vec{F}_1 + \frac{\partial\vec{f}}{\partial\varepsilon}(\vec{F}_0, \vec{z}, 0),$$

which yields $\vec{F}_1(\vec{z})$ and so forth.

$$(8) \quad D_{\vec{x}}\vec{f}(\vec{F}_0, \vec{z}, 0)\vec{F}_1 = D_{\vec{z}}\vec{F}_0\vec{g}(\vec{F}_0, \vec{z}, 0) - \frac{\partial\vec{f}}{\partial\varepsilon}(\vec{F}_0, \vec{z}, 0).$$

So, regular perturbation theory enables to build locally *slow invariant manifolds* M_ε . But for high-dimensional *singularly perturbed systems slow invariant manifold* asymptotic equation determination leads to tedious calculations.

Proof. For application of this technique see [14]. □

3.3. Slow invariant manifold and canards. A manifold of canards is an invariant manifold, where first approximation is M_0 . For two-dimensional singularly perturbed dynamical systems with just one fast variable (x) and one slow variable (y), canards are non generic according to Krupa and Szmolyan [22] and *maximal canards* can only occur in such systems only for discrete values of a control parameter μ . It means that in dimension two a one parameter family of singularly perturbed systems is needed to exhibit canard phenomenon. Because along a canard, the differential $D_x f$ is not always invertible, we can not write the manifold of canards as $x = F(y, \varepsilon)$. Thus, we will suppose that $D_y f$ is invertible and we will try to compute the canard as $y = F(x, \mu, \varepsilon)$. See [5] for a theory of this identification of formal series. We consider the following *singularly perturbed dynamical system*:

$$\begin{aligned} \varepsilon\dot{x} &= f(x, y, \mu, \varepsilon), \\ \dot{y} &= g(x, y, \mu, \varepsilon), \end{aligned}$$

with $x, y \in \mathbb{R}$, i.e. $(m, p) = (1, 1)$ and we suppose that due to the nature of the problem perturbation expansions of the canard and of the canard value read:

$$y = F(x, \varepsilon) = \sum_{i=0}^{N-1} F_i(x)\varepsilon^i + O(\varepsilon^N) \quad \text{and} \quad \mu(\varepsilon) = \sum_{i=0}^{N-1} \mu_i\varepsilon^i + O(\varepsilon^N).$$

According to Eq. (5) invariance of the manifold M_ε reads:

$$(9) \quad \left(\frac{\partial F}{\partial x}(x, \varepsilon) \right) f(x, F(x, \varepsilon), \mu(\varepsilon), \varepsilon) = \varepsilon g(x, F(x, \varepsilon), \mu(\varepsilon), \varepsilon).$$

To avoid technical complications in the computations below, we assume that, at order $O(\varepsilon^0)$, the critical manifold does not depend on the parameter μ .

Indeed,

$$\frac{\partial f}{\partial \mu}(x, F_0(x), \mu_0, 0) = 0$$

Then, solving equation (9) order by order provides at:

Order ε^0

$$(10) \quad \frac{\partial F_0}{\partial x}(x)f(x, F_0(x), \mu_0, 0) = 0 \quad \Leftrightarrow \quad f(x, F_0(x), \mu_0, 0) = 0.$$

because the function $\frac{\partial F_0}{\partial x}(x)$ is almost everywhere non zero. Indeed, the function F_0 is given by the implicit function theorem. In what follows f , g , and their derivatives are evaluated at $(x, F_0(x), \mu_0, 0)$, and F_0 , F_1 and F_2 are evaluated at x .

Order ε^1

$$F_0' \left(\frac{\partial f}{\partial y} F_1 + \frac{\partial f}{\partial \mu} \mu_1 + \frac{\partial f}{\partial \varepsilon} \right) + F_1' f = g.$$

Since according to what has been stated before, we have:

$$(11) \quad F_1 = \frac{\frac{g}{F_0'} - \frac{\partial f}{\partial \varepsilon}}{\frac{\partial f}{\partial y}}$$

A priori, this function is singular at the bifurcation point x_0 of the fast system, because F_0' vanishes at this point. To avoid this singularity in function F_1 , the relation $g(x_0, F(x_0), \mu_0, 0) = 0$ is needed. With an appropriate hypothesis on $\frac{\partial g}{\partial \mu}$, it gives a value for μ_0 .

Higher order The computation can be done with the same arguments. When condition of order k are studied, we have to fix F_k , and to avoid singularity in F_k we have to fix μ_{k-1} . An example will be done in the next paragraph.

3.4. Van der Pol's "canards". Van der Pol system

$$(12) \quad \begin{aligned} \varepsilon \dot{x} &= f(x, y) = x + y - \frac{x^3}{3}, \\ \dot{y} &= g(x, y) = \mu - x, \end{aligned}$$

satisfies Fenichel's assumptions (H_1) – (H_3) except on the points $(x, y) = \pm(1, -\frac{2}{3})$. The critical manifold is the cubic $y = x^3/3 - x$. Thus, the problem is to find a function $y = F(x, \varepsilon)$ whose graph is locally the *slow invariant manifold* M_ε of the Van der Pol system. We write:

$$F(x, \varepsilon) = F_0(x) + \varepsilon F_1(x) + \varepsilon^2 F_2(x) + O(\varepsilon^3)$$

and

$$\mu = \mu_0 + \varepsilon \mu_1 + \varepsilon^2 \mu_2 + O(\varepsilon^3).$$

The identification we have to perform is

$$\sum_{i=0} \frac{\partial F_i}{\partial x} \varepsilon^i \left(\sum_{i=0} F_i \varepsilon^i - \left(\frac{x^3}{3} - x \right) \right) = \sum_{i=0} \mu_i \varepsilon^{i+1} - \varepsilon x$$

Then, solving order by order provides at:

Order ε^0

$$F_0(x) = \frac{x^3}{3} - x.$$

Order ε^1

$$(13) \quad F_1(x) = -\frac{x - \mu_0}{x^2 - 1}$$

This function is singular at the fold point $x_0 = 1$ corresponding to the Hopf bifurcation point of the fast system⁸. So, to avoid this singularity in function $F_1(x)$ we pose: $\mu_0 = 1$ and thus we have: $F_1(x) = \frac{-1}{1+x}$.

Order ε^2

$$(14) \quad F_2(x) = \frac{\mu_1 + \frac{(x - \mu_0)(x^2 + 1 - 2\mu_0 x)}{(x^2 - 1)^3}}{x^2 - 1}$$

Taking into account that $\mu_0 = 1$ and, in order to avoid singularity in $F_2(x)$, we find that $\mu_1 = -\frac{1}{8}$ and so $F_2(x) = -\frac{x^2 + 4x + 7}{8(1+x)^4}$.

Order ε^3

Using the same process, a tedious computation (or, better a computation with the help of a computer) leads to $\mu_2 = -\frac{3}{32}$, $\mu_3 = -\frac{173}{1024}$. Thus, the bifurcation parameter value leading to canard solutions reads:

$$\mu = 1 - \frac{1}{8}\varepsilon - \frac{3}{32}\varepsilon^2 + O(\varepsilon^3)$$

Then, for $\varepsilon = 0.01$ one finds again the value obtained numerically by Benoît *et al.* [2, p.99] and Diener [8]:

$$\mu = 0.99874$$

⁸Due to the symmetry of the vector field: $(-x, -y, -\mu) \rightarrow (x, y, \mu)$ the same computation could have been done on the fold point $x_0 = -1$ in the vicinity of which a ‘‘canard explosion’’ also takes place.

The phenomenon of “canard explosion” of Van der Pol system (12) with $\varepsilon = 0.01$ is exemplified on Fig. 1. where the periodic solution has been plotted in red, the critical manifold in black and the positive fixed point in green. Double arrows indicate the *fast* motion while simple arrows indicate the *slow* motion. Exponentially small variations of the parameter value $\mu = 0.99874$ enable to exhibit the transition from relaxation oscillation (a) to small amplitude limit cycles (b) via canard cycles (c). Then, at the parameter value $\mu = 1$ corresponding to the Hopf bifurcation, the canard disappears (d).




FIGURE 1. Transition from relaxation oscillation to canard explosion.

4. FLOW CURVATURE METHOD

Recently, a new approach called *Flow Curvature Method* and based on the use of *Differential Geometry* properties of *curvatures* has been developed, see [16, 17]. According to this method, the highest *curvature of the flow*, i.e. the $(n-1)^{th}$ *curvature of trajectory curve* integral of n -dimensional dynamical system defines a *manifold* associated with this system and called *flow curvature manifold*. In the case of n -dimensional singularly perturbed dynamical system (1) for which with $\vec{x} \in \mathbb{R}^1$, $\vec{z} \in \mathbb{R}^{n-1}$, i.e. $(m, p) = (1, n-1)$ we have the following result.

Proposition 3. *The location of the points where the $(n-1)^{th}$ curvature of the flow, i.e. the curvature of the trajectory curve \vec{X} , integral of any n -dimensional singularly perturbed dynamical system vanishes, represents its $(n-1)$ -dimensional slow manifold M_ε the equation of which reads*

$$(15) \quad \phi(\vec{X}, \varepsilon) = \dot{\vec{X}} \cdot (\ddot{\vec{X}} \wedge \ddot{\vec{X}} \wedge \dots \wedge \overset{(n)}{\vec{X}}) = \det(\dot{\vec{X}}, \ddot{\vec{X}}, \ddot{\vec{X}}, \dots, \overset{(n)}{\vec{X}}) = 0$$

where $\overset{(n)}{\vec{X}}$ represents the time derivatives up to order n of $\vec{X} = (\vec{x}, \vec{z})^t$.

Proof. For proof of this proposition see [17, p. 185 and next] and below. \square

Remark 4. *First, let's notice that with the Flow Curvature Method the slow manifold is defined by an implicit equation. Secondly, in the most general case of n -dimensional singularly perturbed dynamical system (1) for which $\vec{x} \in \mathbb{R}^m$, $\vec{z} \in \mathbb{R}^p$ the Proposition 3 still holds. In dimension three, the example of a Neuronal Bursting Model (NBM) for which $(m, p) = (2, 1)$ has already been studied by Ginoux et al. [15]. In this particular case, one of the hypotheses of the Tihonov's theorem is not checked since the fast dynamics of the singular approximation has a periodic solution. Nevertheless, it has been established by Ginoux et al. [15] that the slow manifold can all the same be obtained while using the Flow Curvature Method. According to this method, the slow invariant manifold of a three-dimensional singularly perturbed dynamical system for which $(m, p) = (1, 2)$ is given by the 2nd curvature of the flow, i.e. the torsion. In the case of a Neuronal Bursting Model for which $(m, p) = (2, 1)$ it has been stated by Ginoux et al. [15] that the slow manifold is then given by the 1st curvature of the flow, i.e. the curvature. In such a case, the flow curvature manifold is defined by the location of the points where the three-dimensional pseudovector $\ddot{\vec{X}} \wedge \ddot{\vec{X}}$ vanishes. This condition leads to a nonlinear system of three equations two of which being linearly independent. These two equations define a curve corresponding to the slow invariant manifold⁹. Thus, one can deduce that for a three-dimensional singularly perturbed dynamical system for which (m, p) the slow manifold is given by the p^{th} curvature of the flow.*

4.1. Invariance. According to Schlomiuk [27] and Llibre et al. [24] the concept of *invariant manifold* has been originally introduced by Gaston Darboux [7, p. 71] in a memoir entitled: *Sur les équations différentielles algébriques du premier ordre et du premier degré* and can be stated as follows.

⁹See also Gilmore et al. [18]

Proposition 5. *The manifold defined by $\phi(\vec{X}, \varepsilon) = 0$ where ϕ is a C^1 in an open set U , is invariant with respect to the flow of (1) if there exists a C^1 function denoted by $\kappa(\vec{X}, \varepsilon)$ and called cofactor which satisfies*

$$(16) \quad L_{\vec{V}}\phi(\vec{X}, \varepsilon) = \kappa(\vec{X}, \varepsilon)\phi(\vec{X}, \varepsilon),$$

for all $\vec{X} \in U$, and with the Lie derivative operator defined as

$$L_{\vec{V}}\phi = \vec{V} \cdot \vec{\nabla}\phi = \sum_{i=1}^n \frac{\partial \phi}{\partial x_i} \dot{x}_i = \frac{d\phi}{dt}.$$

Proof. According to Fenichel's Persistence Theorem (see Th. 2) the slow invariant manifold M_ε may be written as an explicit function $\vec{x} = \vec{F}(\vec{z}, \varepsilon)$, the invariance of which implies that $\vec{F}(\vec{z}, \varepsilon)$ satisfies

$$(17) \quad \varepsilon D_{\vec{z}}\vec{F}(\vec{z}, \varepsilon) \vec{g}(\vec{F}(\vec{z}, \varepsilon), \vec{z}, \varepsilon) = \vec{f}(\vec{F}(\vec{z}, \varepsilon), \vec{z}, \varepsilon)$$

We write the slow manifold M_ε as an implicit function by posing

$$(18) \quad \phi(\vec{x}, \vec{z}, \varepsilon) = \vec{x} - \vec{F}(\vec{z}, \varepsilon) = \phi(\vec{X}, \varepsilon).$$

According to Darboux invariance theorem M_ε is invariant if its Lie derivative reads

$$(19) \quad L_{\vec{V}}\phi(\vec{X}, \varepsilon) = \kappa(\vec{X}, \varepsilon)\phi(\vec{X}, \varepsilon).$$

Plugging Eq. (18) into the Lie derivative (19) leads to

$$L_{\vec{V}}\phi(\vec{X}, \varepsilon) = \dot{\vec{x}} - D_{\vec{z}}\vec{F}(\vec{z}, \varepsilon)\dot{\vec{z}} = \kappa(\vec{X}, \varepsilon)\phi(\vec{X}, \varepsilon),$$

which may be written according to Eq. (2) as

$$L_{\vec{V}}\phi(\vec{X}, \varepsilon) = \frac{1}{\varepsilon}(\vec{f}(\vec{X}, \varepsilon) - \varepsilon D_{\vec{z}}\vec{F}(\vec{z}, \varepsilon)\vec{g}(\vec{X}, \varepsilon)) = \kappa(\vec{X}, \varepsilon)\phi(\vec{X}, \varepsilon).$$

Evaluating this Lie derivative in the location of the points where $\phi(\vec{X}, \varepsilon) = 0$, i.e. $\vec{x} = \vec{F}(\vec{z}, \varepsilon)$ leads to

$$L_{\vec{V}}\phi(\vec{F}(\vec{z}, \varepsilon), \vec{z}, \varepsilon) = \frac{1}{\varepsilon}(\vec{f}(\vec{F}(\vec{z}, \varepsilon), \vec{z}, \varepsilon) - \varepsilon D_{\vec{z}}\vec{F}(\vec{z}, \varepsilon)\vec{g}(\vec{F}(\vec{z}, \varepsilon), \vec{z}, \varepsilon)) = 0,$$

which is exactly identical to Eq. (18) used by Fenichel. \square

Remark 6. *This last equation for the invariance of the manifold M_ε may be written in a simpler way which implies that $\phi(\vec{x}, \vec{z}, \varepsilon)$ satisfies*

$$(20) \quad \frac{d}{dt}[\phi(\vec{x}, \vec{z}, \varepsilon)] = 0,$$

on the solutions of the differential system.

4.2. Slow invariant manifold. We consider again the following two-dimensional singularly perturbed dynamical system

$$\begin{aligned}\varepsilon \dot{x} &= f(x, y), \\ \dot{y} &= g(x, y),\end{aligned}$$

and we suppose that due to the nature of the problem perturbation expansion reads

$$y = F(x, \varepsilon) = F_0(x) + \varepsilon F_1(x) + \varepsilon^2 F_2(x) + O(\varepsilon^3).$$

According to the *Flow Curvature Method* each function $\vec{F}_i(\vec{z})$ of this perturbation expansion may be found again starting from the *slow manifold implicit equation* (16) as stated in the next result.

Proposition 7. *The functions $F_i(x)$ of the slow invariant manifold associated with a two-dimensional singularly perturbed dynamical system are given by the following expressions*

$$(21) \quad \begin{aligned} F_n'(x) &= \lim_{\varepsilon \rightarrow 0} \left[\frac{1}{n!} \frac{\partial^n a_{10}}{\partial \varepsilon^n} \right] \text{ with } n \geq 0, \\ F_n(x) &= \lim_{\varepsilon \rightarrow 0} \left[\frac{1}{n!} \frac{\partial^{n-1} a_{01}}{\partial \varepsilon^{n-1}} \right] \text{ with } n \geq 1, \end{aligned}$$

where

$$a_{10} = - \left. \frac{\frac{\partial \phi_i}{\partial x}}{\frac{\partial \phi_i}{\partial y}} \right|_{y=F(x, \varepsilon)} \quad \text{and} \quad a_{01} = - \left. \frac{\frac{\partial \phi_i}{\partial \varepsilon}}{\frac{\partial \phi_i}{\partial y}} \right|_{y=F(x, \varepsilon)},$$

and

$$\phi_i(x, y, \varepsilon) = \frac{d^{i-1}}{dt^{i-1}} [\phi_1(x, y, \varepsilon)] \quad \text{with } i = 1, 2, \dots, n,$$

where $\phi_i(x, y, \varepsilon)$ corresponds to the i^{th} order approximation in ε .

Proof. We have that

$$\phi(\vec{X}, \varepsilon) = \left\| \dot{\vec{X}} \wedge \ddot{\vec{X}} \right\| = \det(\dot{\vec{X}}, \ddot{\vec{X}}) = 0 \quad \Leftrightarrow \quad \phi(x, y, \varepsilon) = 0.$$

Since, for a two-dimensional singularly perturbed dynamical systems this *slow manifold equation* is defined by the *second order tensor of curvature*, i.e. by a determinant involving the first and second time derivatives of the vector field \vec{X} , it corresponds to the first order approximation in ε of the *slow manifold* obtained with the *Geometric Singular Perturbation Method*. So, we denote it by

$$\phi_1(\vec{X}, \varepsilon) = \left\| \dot{\vec{X}} \wedge \ddot{\vec{X}} \right\| = \det(\dot{\vec{X}}, \ddot{\vec{X}}) = 0.$$

A *third order tensor of curvature* can be easily given by the time derivative of $\phi_1(\vec{X}, \varepsilon)$. We denote it by

$$\phi_2(\vec{X}, \varepsilon) = \dot{\phi}_1(\vec{X}, \varepsilon) = \det(\dot{\vec{X}}, \ddot{\vec{X}}) = 0.$$

Thus, $\phi_2(\vec{X}, \varepsilon)$ corresponds to the second order approximation in ε . Using the same process, we consider the *slow manifold* $\phi_i(\vec{X}, \varepsilon)$ which corresponds to the i^{th} order approximation in ε .

Writing the *total differential* of the *slow manifold* we obtain

$$(22) \quad d\phi_i(x, y, \varepsilon) = \frac{\partial\phi_i}{\partial x} dx + \frac{\partial\phi_i}{\partial y} dy + \frac{\partial\phi_i}{\partial \varepsilon} d\varepsilon = 0.$$

Replacing in Eq. (23) dy by its *total differential* $dy = \frac{\partial F}{\partial x} dx + \frac{\partial F}{\partial \varepsilon} d\varepsilon$ yields

$$(23) \quad d\phi_i(x, y, \varepsilon) = \left(\frac{\partial\phi_i}{\partial x} + \frac{\partial\phi_i}{\partial y} \frac{\partial F}{\partial x} \right) dx + \left(\frac{\partial\phi_i}{\partial \varepsilon} + \frac{\partial\phi_i}{\partial y} \frac{\partial F}{\partial \varepsilon} \right) d\varepsilon.$$

According to Eq. (21) $\phi_i(x, y, \varepsilon)$ is *invariant* if and only if $d\phi_i(x, y, \varepsilon) = 0$, i.e. if

$$(24) \quad \begin{aligned} \frac{\partial\phi_i}{\partial x} + \frac{\partial\phi_i}{\partial y} \frac{\partial F}{\partial x} = 0 &\Leftrightarrow \frac{\partial F}{\partial x} = -\frac{\frac{\partial\phi_i}{\partial x}}{\frac{\partial\phi_i}{\partial y}}, \\ \frac{\partial\phi_i}{\partial \varepsilon} + \frac{\partial\phi_i}{\partial y} \frac{\partial F}{\partial \varepsilon} = 0 &\Leftrightarrow \frac{\partial F}{\partial \varepsilon} = -\frac{\frac{\partial\phi_i}{\partial \varepsilon}}{\frac{\partial\phi_i}{\partial y}}. \end{aligned}$$

By replacing $y = F(x, \varepsilon)$ by its expression in both parts of Eq. (25) and by setting

$$a_{10} = -\left. \frac{\frac{\partial\phi_i}{\partial x}}{\frac{\partial\phi_i}{\partial y}} \right|_{y=F(x,\varepsilon)} \quad \text{and} \quad a_{01} = -\left. \frac{\frac{\partial\phi_i}{\partial \varepsilon}}{\frac{\partial\phi_i}{\partial y}} \right|_{y=F(x,\varepsilon)},$$

we have that

$$(25) \quad \begin{aligned} \frac{\partial F(x, \varepsilon)}{\partial x} &= F'_0(x) + \varepsilon F'_1(x) + O(\varepsilon^2) = a_{10}, \\ \frac{\partial F(x, \varepsilon)}{\partial \varepsilon} &= F_1(x) + 2\varepsilon F_2(\varepsilon) + O(\varepsilon^2) = a_{01}. \end{aligned}$$

By using a *recurrence reasoning* it may be easily stated that the functions $F_i(x)$ of the *slow invariant manifold* associated with a two-dimensional singularly perturbed dynamical system are given by the expressions (21). \square

4.3. Van der Pol's "canards". We consider again the Van der Pol system (12). All functions $F_i(x)$ of the perturbation expansion may be deduced from the *slow manifold* equation defined by (16). But, since the determination of $F_3(x)$, i.e. the computation μ_2 requires a *third order tensor of curvature* we consider $\phi_3(\vec{X}, \varepsilon)$ the second time derivative of $\phi_1(\vec{X}, \varepsilon)$ which corresponds to the third order approximation in ε .

Thus, we find at

Order ε^0

$$F_0'(x) = \lim_{\varepsilon \rightarrow 0} [a_{10}] = -1 + x^2,$$

from which one deduces that

$$F_0(x) = \frac{x^3}{3} - x + C_0,$$

where the constant C_0 may be chosen in such a way that the *critical manifold* can be found again ($C_0 = 0$).

Order ε^1

$$(26) \quad F_1(x) = \lim_{\varepsilon \rightarrow 0} [a_{01}] = \frac{\mu_0 - x}{x^2 - 1}.$$

Thus, one find again exactly the same functions $F_1(x)$ as those given by *Geometric Singular Perturbation Method* (14) and of course the same value of the bifurcation parameter $\mu_0 = 1$.

Order ε^2

$$(27) \quad F_2(x) = \frac{\mu_1 + \frac{(x - \mu_0)(x^2 + 1 - 2\mu_0 x)}{(x^2 - 1)^3}}{x^2 - 1}.$$

Taking into account that $\mu_0 = 1$ we find again exactly the same functions $F_2(x)$ as those given by *Geometric Singular Perturbation Method* (15) and of course the same value of the bifurcation parameter $\mu_1 = -1/8$.

Order ε^3

A simple and direct computation leads to $\mu_2 = -\frac{3}{32}$. Thus, the bifurcation parameter value leading to canard solutions reads

$$\mu = 1 - \frac{1}{8}\varepsilon - \frac{3}{32}\varepsilon^2 + O(\varepsilon^3).$$

Then, for $\varepsilon = 0.01$ one finds again the value obtained by Benoît *et al.* [2, p.99] and Diener [8]

$$\mu = 0.99874\dots$$

A program made with Mathematica and available at: <http://ginoux.univ-tln.fr> enables to compute all order of approximations in ε of any two-dimensional singularly perturbed systems.

5. CONCLUSION

Thus, the bifurcation parameter value leading to a *canard explosion* in dimension two obtained by the so-called *Geometric Singular Perturbation Method* has been found again with the *Flow Curvature Method*. This result could be also extended to three-dimensional singularly perturbed dynamical systems such as the 3D-autocatalator in which canard phenomenon occurs.

ACKNOWLEDGMENTS

Authors would like to thank Professor Eric Benoît and the referees for their fruitful advices.

The second author is supported by the grants MIICIN/FEDER MTM 2008–03437, AGAUR 2009SGR410, and ICREA Academia.

REFERENCES

- [1] A.A. ANDRONOV & S.E. CHAIKIN, *Theory of Oscillators*, I. Moscow, 1937; English transl., Princeton Univ. Press, Princeton, N.J., 1949.
- [2] E. BENOÎT, J.L. CALLOT, F. DIENER & M. DIENER, *Collectanea Mathematica* 31-32 (1-3) (1981) 37-119.
- [3] M. BRØNS & K. BAR-ELI, *J. Phys. Chem.*, Vol. 95 (22) (1991) 8706-8713.
- [4] M. BRØNS & K. BAR-ELI, *Proc. R. Soc. Lond. A* 445 (1994) 305-322.
- [5] M. CANALIS-DURAND, F. DIENER AND M. GAETANO, Calcul des valeurs à canard à l'aide de Macsyma, in *Mathématiques finitaires et Analyse Non Standard*, Ed. M. Diener and G. Wallet, Publications mathématiques de l'Université Paris 7 (1985), pp "153–167".
- [6] J.D. COLE, *Perturbation Methods in Applied Mathematics*, Blaisdell, Waltham, MA, 1968.
- [7] G. DARBOUX, *Bull. Sci. Math.* 2(2) (1878) 60-96, 123-143 & 151-200.
- [8] M. DIENER, *The Mathematical Intelligencer* 6 (1984) 38-49.
- [9] F. DUMORTIER & R. ROUSSARIE, Canard cycles and center manifolds, *Memoires of the AMS*, 557, 1996.
- [10] W. ECKHAUS, Relaxation oscillations including a standard chase on French ducks; in *Asymptotic Analysis II*, Springer Lecture Notes Math. 985 (1983) 449-494.
- [11] N. FENICHEL, *Ind. Univ. Math. J.* 21 (1971) 193-225.
- [12] N. FENICHEL, *Ind. Univ. Math. J.* 23 (1974) 1109-1137.
- [13] N. FENICHEL, *Ind. Univ. Math. J.* 26 (1977) 81-93.
- [14] N. FENICHEL, *J. Diff. Eq.* 31 (1979) 53-98.
- [15] J.M. GINOUX & B. ROSSETTO, Slow manifold of a neuronal bursting model, in *Emergent Properties in Natural and Artificial Dynamical Systems* (eds M.A. Aziz-Alaoui & C. Bertelle), Springer-Verlag, 2006, pp. 119-128.
- [16] J.M. GINOUX, B. ROSSETTO & L.O. CHUA, *Int. J. Bif. & Chaos* 11(18) (2008) 3409-3430.
- [17] J.M. GINOUX, *Differential geometry applied to dynamical systems*, World Scientific Series on Nonlinear Science, Series A **66** (World Scientific, Singapore), 2009.
- [18] R. GILMORE, J.M. GINOUX, T. JONES, C. LETELLIER & U. S. FREITAS, *J. Phys. A: Math. Theor.* 43 (2010) 255101 (13pp).
- [19] M. W. HIRSCH, C.C. PUGH & M. SHUB, *Invariant Manifolds*, Springer-Verlag, New York, 1977.
- [20] C.K.R.T. JONES, *Geometric Singular Perturbation Theory*, in *Dynamical Systems*, Montecatini Terme, L. Arnold, Lecture Notes in Mathematics, vol. 1609, Springer-Verlag, 1994, pp. 44-118.
- [21] T. KAPER, *An Introduction to Geometric Methods and Dynamical Systems Theory for Singular Perturbation Problems*, in *Analyzing multiscale phenomena using singular perturbation methods*, (Baltimore, MD, 1998), pages 85-131. Amer. Math. Soc., Providence, RI, 1999.
- [22] M. KRUPA & P. SZMOLYAN, *J. Diff. Eq.* 174 (2001) 312-368.
- [23] N. LEVINSON, *Ann. Math.* 50 (1949) 127-153.
- [24] J. LLIBRE & J.C. MEDRADO, *J. Phys. A: Math. Theor.* 40 (2007) 8385-8391.
- [25] R.E. O'MALLEY, *Introduction to Singular Perturbations*, Academic Press, New York, 1974.
- [26] R.E. O'MALLEY, *Singular Perturbations Methods for Ordinary Differential Equations*, Springer-Verlag, New York, 1991.
- [27] D. SCHLOMIUK, *Expositiones Mathematicae* 11 (1993) 433-454.
- [28] SZMOLYAN, M. WECHSELBERGER, *J. Diff. Eq.* 177 (2001) 419-453.
- [29] F. TAKENS, *Constrained equations; A study of implicit differential equations and their discontinuous solutions*, in *Lecture Notes in Mathematics* volume 525 (1976), Springer Verlag.
- [30] A.N. TIKHONOV, *Mat. Sbornik N.S.*, 31 (1948) 575-586.
- [31] B. VAN DER POL, *Phil. Mag.*, 7(2) (1926) 978-992.

- [32] W.R. WASOW, *Asymptotic Expansions for Ordinary Differential Equations*, Wiley-Interscience, New York, 1965.

¹ LABORATOIRE PROTEE, I.U.T. DE TOULON, UNIVERSITÉ DU SUD, BP 20132, F-83957 LA GARDE CEDEX, FRANCE

E-mail address: ginoux@univ-tln.fr

² DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BELLATERRA, BARCELONA, SPAIN

E-mail address: jllibre@mat.uab.cat