

Additional file 2. Classification of patients with creatinine and albumin excretion rate available according to KDIGO 2009

	n (%)			Albuminuria stages (mg/g)				
				A1		A2	A3	All
				Optimal and high normal		High	Very high	
				< 10	10-29	30-299	≥ 300	
				GFR stages (ml/min per 1.73m ²)	G1	Increased and optimal	≥ 105	128 (8.6)
90-104	142 (9.6)	42 (2.8)	40 (2.8)				4 (0.3)	228 (15.4)
G2	Mild	75-89	269 (18.2)		109 (7.4)	71 (4.8)	5 (0.3)	454 (30.7)
		60-74	165 (11.1)		71 (4.8)	42 (2.8)	6 (0.4)	284 (19.2)
G3a	Mild-moderate	45-59	73 (5.0)		36 (2.4)	34 (2.3)	9 (0.6)	152 (10.3)
G3b	Moderate-severe	30-44	20 (1.3)		11 (0.7)	12 (0.8)	4 (0.3)	47 (3.2)
G4	Severe	15-29	13 (0.9)		20 (1.4)	8 (0.5)	2 (0.1)	43 (2.9)
G5	Kidney failure	<15	24 (1.6)		20 (1.4)	11 (0.7)	8 (0.5)	63 (4.3)
All			834 (56.4)		356 (24.1)	249 (16.8)	39 (2.6)	1478 (100)

Albuminuria alone

Nonalbuminuric RI

GFR: glomerular filtration rate