

POLYNOMIAL DIFFERENTIAL SYSTEMS WITH EXPLICIT NON-ALGEBRAIC LIMIT CYCLES

REBIHA BENTERKI¹ AND JAUME LLIBRE²

ABSTRACT. Up to know all the examples of polynomial differential systems for which non-algebraic limit cycles are known explicitly have degree ≥ 5 . Here we show that already there are polynomial differential systems of degree ≥ 3 exhibiting explicit non-algebraic limit cycles. It is well known that polynomial differential systems of degree 1 (i.e. linear differential systems) has no limit cycles. It remains the open question to determine if the polynomial differential systems of degree 2 can exhibit explicit non-algebraic limit cycles.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

Probably the existence of limit cycles is one of the more difficult objects to study in the qualitative theory of differential equations in the plane. There are hundreds of papers and some books dedicated to this topic, see for instance the book of Ye Yanqian et al. [12], or the famous Hilbert 16th problem [6] and [7]. More closed papers to the problem here considered are [4, 5, 1, 2, 8, 9].

A *polynomial differential system* is a system of the form

$$(1) \quad \begin{aligned} \dot{x} &= P(x, y), \\ \dot{y} &= Q(x, y), \end{aligned}$$

where $P(x, y)$ and $Q(x, y)$ are real polynomials in the variables x and y . The *degree* of the system is the maximum of the degrees of the polynomials P and Q . As usual the dot denotes derivative with respect to the independent variable t .

A *limit cycle* of system (1) is an isolated periodic solution in the set of all periodic solutions of system (1). If a limit cycle is contained in an algebraic curve of the plane, then we say that it is *algebraic*, otherwise it is called *non-algebraic*.

To distinguish when a limit cycle is algebraic or not, in general, it is not easy. Thus the well-known limit cycle of the van der Pol differential system discovered in 1927 (see [11]) was not proved until 1995 by Odani [10] that it was non-algebraic. The van der Pol system can be written as a polynomial differential system (1) of degree 3, but its limit cycles is not known explicitly.

These last years (from 2006 up to now) several papers have been published exhibiting polynomial differential systems for which non-algebraic limit cycles are known explicitly. This means that in some coordinates we have an explicit analytic expression of the curve containing the non-algebraic limit cycle. The first explicit

2010 *Mathematics Subject Classification.* 34C29, 34C25.

Key words and phrases. non-algebraic limit cycle, polynomial vector field.

The first author is partially supported by a MCYT/FEDER grant MTM2008-03437, by a CIRIT grant number 2009SGR-410 and by ICREA Academia. The second author is partially supported by the Algerian Ministry of Higher Education and Scientific Research under project B03320090002.

non-algebraic limit cycle, due to Gasull, Giacomini and Torregrosa [4], was for a polynomial differential system of degree 5. Of course, multiplying the right hand part of this polynomial differential system of degree 5 by $(ax + by + c)^n$ with n an arbitrary positive integer, where the straight line $ax + by + c = 0$ must be chosen in such a way that it does not intersect the explicit limit cycle of the system, we get a polynomial differential system of degree $5 + n$ exhibiting an explicit non-algebraic limit cycle.

Immediately after this first paper appeared the paper of Al-Dosary [1] inspired in [4] (note that this reference is quoted in [1]), providing a similar polynomial differential system of degree 5 exhibiting an explicit non-algebraic limit cycle.

Giné and Grau [5] provide a polynomial differential system of degree 9 exhibiting simultaneously two explicit limit cycles one algebraic and another non-algebraic. Note that the paper [4] is also quoted in [5].

The objective of this paper is to show that already there are polynomial differential systems of degree 3 exhibiting explicit non-algebraic limit cycles. Thus, our main result is the following one.

Theorem 1. *The differential polynomial system of degree 3*

$$(2) \quad \begin{aligned} \dot{x} &= x + (y - x)(x^2 - yx + y^2), \\ \dot{y} &= y - (y + x)(x^2 - yx + y^2), \end{aligned}$$

has a unique non-algebraic limit cycle whose expression in polar coordinates (r, θ) , defined by $x = r \cos \theta$ and $y = r \sin \theta$, is

$$(3) \quad r(\theta) = e^\theta \sqrt{r_*^2 - f(\theta)},$$

where

$$\begin{aligned} r_* &= e^{2\pi} \sqrt{\frac{f(2\pi)}{e^{4\pi} - 1}} \approx 1.1911644871948721\dots, \\ f(\theta) &= 4 \int_0^\theta \frac{e^{-2s}}{2 - \sin(2s)} ds. \end{aligned}$$

Moreover, this limit cycle is a stable hyperbolic limit cycle.

Theorem 1 is proved in section 2.

In short, since it is well known that the linear differential systems (or polynomial differential systems of degree 1) have no limit cycles, it remains the following open question:

Open question. *Are there or not polynomial differential systems of degree 2 exhibiting explicit non-algebraic limit cycles.*

2. PROOF OF THEOREM 1

The polynomial differential system (2) in polar coordinates becomes

$$(4) \quad \begin{aligned} \dot{r} &= r + \frac{1}{2}(\sin(2\theta) - 2)r^3, \\ \dot{\theta} &= \frac{1}{2}r^2(\sin(2\theta) - 2). \end{aligned}$$

Taking as independent variable the coordinate θ , this differential system writes

$$(5) \quad \frac{dr}{d\theta} = r + \frac{2}{r(\sin(2\theta) - 2)}.$$




FIGURE 1. The phase portrait in the Poincaré disc of the polynomial differential system (2).

Note that since $\dot{\theta} < 0$, the orbits $r(\theta)$ of the differential equation (5) has reversed their orientation with respect to the orbits $(r(t), \theta(t))$ or $(x(t), y(t))$ of the differential systems (4) and (2), respectively.

It is easy to check that the solution $r(\theta; r_0)$ of the differential equation (5) such that $r(0; r_0) = r_0$ is

$$(6) \quad r(\theta; r_0) = e^\theta \sqrt{r_0^2 - f(\theta)},$$

where $f(\theta)$ is the function defined in the statement of Theorem 1.

Clearly the unique equilibrium point of the differential system (2) is the origin of coordinates, which is an unstable node because its eigenvalues are 1 with multiplicity two, for more details see for instance Theorem 2.15 of [3]. This equilibrium point in polar coordinates become $r = 0$. This is the unique point of the plane where the differential equation (5) is not defined. But we can extend the flow of this differential equation to $r = 0$, assuming that at the origin of the plane in polar coordinates we have an unstable node.

The periodic orbits $r(\theta; r_0)$ of the differential equation (5) must satisfy $r(2\pi; r_0) = r_0$. The unique solution of this equation is $r_0 = r_*$, where r_* is defined in the statement of Theorem 1. So, if $r(\theta; r_*) > 0$ for all $\theta \in \mathbb{R}$, we shall have $r(\theta; r_*) > 0$ would be a periodic orbit, and consequently a limit cycle. In what follows it is proved that $r(\theta; r_*) > 0$ for all $\theta \in \mathbb{R}$. Indeed

$$\begin{aligned} r(\theta; r_*) &= e^\theta \sqrt{\frac{e^{4\pi}}{e^{4\pi} - 1} f(2\pi) - f(\theta)} \\ &\geq e^\theta \sqrt{f(2\pi) - f(\theta)} \\ &= 2e^\theta \sqrt{\int_\theta^{2\pi} \frac{e^{-2s}}{2 - \sin(2s)} ds} > 0, \end{aligned}$$

because $e^{-2s}/(2 - \sin(2s)) > 0$ for all $s \in \mathbb{R}$.

An easy computation shows that

$$\left. \frac{dr(2\pi; r_0)}{dr_0} \right|_{r_0=r_*} = e^{4\pi} > 1.$$

Therefore the limit cycle of the differential equation (5) is unstable and hyperbolic, for more details see section 1.6 of [3]. Consequently, this is a stable and hyperbolic limit cycle for the differential system (2).

Clearly the curve $(r(\theta) \cos \theta, r(\theta) \sin \theta)$ in the (x, y) plane with

$$r(\theta)^2 = e^{2\theta}(r_*^2 - f(\theta)),$$

is not algebraic, due the expression $e^{2\theta}r_*^2$. This completes the proof of Theorem 1.

Now we shall present the phase portrait of the differential system (2) in the Poincaré disc, see the Poincaré compactification in the Chapter 5 of [3].

Since the polynomial $\dot{x}y - \dot{y}x = (x^2 + y^2)(x^2 - xy + y^2)$ has no real linear factors, the compactification of Poincaré of the differential system (2) has no equilibrium points at infinity, i.e. the infinity is a periodic orbit. Doing the change of variables $r = 1/\rho$, the infinity of the differential equation (5) passes at the origin, and equation (5) becomes

$$\frac{d\rho}{d\theta} = -\rho - \frac{2\rho^3}{\sin(2\theta) - 2}.$$

Hence, clearly $\rho = 0$ is an stable equilibrium point of this differential equation, consequently the periodic orbit at infinity of the differential equation (3) is an unstable limit cycle. Then the phase portrait in the Poincaré disc of the polynomial differential system (2) is given in Figure 1.

REFERENCES

- [1] K.I.T. AL-DOSARY, *Non-algebraic limit cycles for parametrized planar polynomial systems*, Int. J. of Math. **18** (2007), 179–189.
- [2] C. DU AND Y. LIU, *The problem of general center-focus and bifurcation of limit cycles for a planar system of nine degrees*, J. Comput. Appl. Math. **223** (2009), 1043–1057.
- [3] F. DUMORTIER, J. LLIBRE AND J.C. ARTÉS, *Qualitative theory of planar differential systems*, UniversiText, Springer-Verlag, New York, 2006.
- [4] A. GASULL, H. GIACOMINI AND J. TORREGROSA, *Explicit non-algebraic limit cycles for polynomial systems*, J. Comp. and Appl. Math. **200** (2007), 448–457.
- [5] J. GINÉ AND M. GRAU, *Coexistence of algebraic and non-algebraic limit cycles, explicitly given, using Riccati equations*, Nonlinearity **19** (2006), 1939–1950.
- [6] D. HILBERT, *Mathematische Problem (lecture)*, Second Internat. Congress Math. Paris, 1900, Nachr. Ges. Wiss. Göttingen Math.-Phys. Kl. 1900, pp 253–297.
- [7] YU. S. ILYASHENKO, *Centennial history of Hilbert’s 16th problem*, Bull. Amer. Math. Soc. **39** (2002), 301–354.
- [8] J. LLIBRE AND M. LIMA *Limit cycles and invariant cylinders for a class of continuous and discontinuous vector field in dimension $2n$* , Appl. Math. Comput. **217** (2011), 9985–9996.
- [9] J. LLIBRE AND A. MAKHLOUF, *Bifurcation of limit cycles from a 4-dimensional center in 1:n resonance*, Appl. Math. Comput. **215** (2009), 140–146.
- [10] K. ODANI, *The limit cycle of the van der Pol equation is not algebraic*, J. Differential Equations **115** (1995), 146–152.
- [11] B. VAN DER POL, *On relaxation-oscillations*, The London, Edinburgh and Dublin Phil. Mag. & J. of Sci. **2**(7) (1927), 978–992.
- [12] YE YANQIAN ET AL., *Theory of Limit Cycles*, Translations of Math. Monographs, Vol. **66**, Amer. Math. Soc, Providence, 1986.

¹ DÉPARTEMENT DE MATHÉMATIQUES, CENTRE UNIVERSITAIRE DE BORDJ BOU ARRÉRIDJ, BORDJ BOU ARRÉRIDJ 34265, EL ANASSER, ALGERIA

E-mail address: **r-benterki@yahoo.fr**

² DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BEL-LATERRA, BARCELONA, CATALONIA, SPAIN

E-mail address: **jllibre@mat.uab.cat**