

## Fertilization capacity of cryopreserved Iberian ibex epididymal sperm in a heterologous *in vitro* fertilization assay

J. López-Saucedo<sup>2</sup>, J. Santiago-Moreno<sup>1</sup>, R. Fierro<sup>4</sup>, D. Izquierdo<sup>5</sup>, M.A. Coloma<sup>3</sup>, M.G. Catalá<sup>5</sup>, I. Jiménez<sup>4</sup> and M.T. Paramio<sup>5</sup>

Doctorado en Ciencias Biológicas y de la Salud de la Universidad Autónoma Metropolitana. Mexico City, Mexico; Departamento de Reproducción Animal, INIA, 28040 Madrid, Spain; Departamento de Ciencias de la Salud, Universidad Autónoma Metropolitana-Iztapalapa, Mexico City, Mexico; and Departamento de Ciencia Animal y de los Alimentos, Facultad de Veterinaria, Universidad Autónoma de Barcelona, Bellaterra, Barcelona, Spain

Date submitted: 23.12.2012. Date accepted: 23.04.2013

### Summary

*In vitro* fertilization (IVF) can be used to assess the fertilization capacity of sperm. Heterologous IVF may be useful when assessing that of wild animals as it is often difficult to obtain adequate numbers of naturally corresponding oocytes. The aim of the present study was to assess the fertilization capacity of frozen–thawed ibex epididymal spermatozoa via heterologous IVF involving the oocytes of prepubertal domestic goats. The effect on fertilization and embryo development of adding oestrous sheep serum (ESS) to the fertilization medium was also examined. Cumulus–oocyte complexes (COCs) were matured in TCM-199 for 24–27 h at 38.5°C in a 5% CO<sub>2</sub> in air atmosphere. Frozen–thawed epididymal spermatozoa were selected by density gradient centrifugation. After maturation, the oocytes were co-incubated with spermatozoa in synthetic oviductal fluid (SOF) with different concentrations of ESS: SOF-C (0%), SOF-2 (2%) and SOF-20 (20%). At 17 h post-insemination (hpi), zygotes with one female and one male pronucleus (2PN) were categorised as normal; zygotes with 3PN were recorded as polyspermic, and oocytes with 1PN as asynchronous. Cleavage and blastocyst development were assessed at 48 and 168 hpi respectively. The percentage of zygotes with 2PN was higher in the SOF-2 than in the SOF-20 treatment group (27.7% versus 2.9%  $P < 0.05$ ). The percentage of blastocysts formed with the SOF-C, SOF-2 and SOF-20 treatments were 1.1%, 7.5% and 0% respectively. These results show that the presence of 2% ESS achieves better results than the use of no serum or the standard 20% concentration. Heterologous IVF may be an effective method for predicting the fertilization capacity of ibex spermatozoa, and therefore perhaps that of other wild mountain ungulates.

**Keywords:** Domestic goat, Embryo development, Epididymal sperm, Heterologous *in vitro* fertilization, Ibex

### Introduction

Reproductive technologies may be of help in preventing the loss of threatened wild species. Genetic resources can be built up by the cryopreservation of their gametes and the establishment of genetic resource banks (Watson & Holt, 2001). Certainly, the use of stored gametes in artificial insemination (AI) programmes should help in the dispersal of genes across isolated gene pools. Unfortunately, the collection of gametes can present many difficulties. Indeed, even sperm collection can be complex in some wild species, although viable epididymal spermatozoa can sometimes be retrieved *post mortem* (Soler *et al.*, 2003; Cocchia *et al.*, 2010; Anel *et al.*, 2011; Keeley *et al.*,

<sup>1</sup>All correspondence to: J. Santiago-Moreno. <sup>2</sup>Departamento de Reproducción Animal, INIA, 28040 Madrid, Spain. Tel: +34 91 347 40 20. Fax: +34 91 347 40 14. e-mail address: moreno@inia.es

<sup>2</sup>Doctorado en Ciencias Biológicas y de la Salud de la Universidad Autónoma Metropolitana. 09340 Mexico City, Mexico.

<sup>3</sup>Departamento de Reproducción Animal, INIA, 28040 Madrid, Spain.

<sup>4</sup>Departamento de Ciencias de la Salud, Universidad Autónoma Metropolitana-Iztapalapa, 09340 Mexico City, Mexico.

<sup>5</sup>Departamento de Ciencia Animal y de los Alimentos, Facultad de Veterinaria, Universidad Autónoma de Barcelona, 08193 Bellaterra, Barcelona, Spain.

2012). Furthermore, assisted reproduction typically produces few offspring (Andrabi & Maxwell, 2007). One reason for this situation is the variation in the fertilization capacity of cryopreserved sperm. This aspect should always be checked before using stored sperm for conservation purposes.

The Iberian ibex (*Capra pyrenaica*) is a wild caprine endemic to the Iberian Peninsula. Phylogenetically it is related to the domestic goat (*Capra hircus*). Studies on the cryopreservation of its sperm and assisted reproduction in the species (Santiago-Moreno *et al.*, 2007, 2009a; Coloma *et al.*, 2011) have led to its recognition as a good model for other wild mountain ungulates (Santiago-Moreno *et al.*, 2010). Heterologous *in vivo* insemination of domestic goats has been used to determine the fertilization capacity of Iberian ibex frozen-thawed epididymal spermatozoa, permitting the selection of sperm doses most likely to produce live offspring via AI (Santiago-Moreno *et al.*, 2006). However, the fertility rates obtained in both normal and interspecies AI are profoundly influenced by the fertility of the females involved, the quality of the corpus luteum, early embryonic mortality, and the response to hormonal synchronisation treatments (Salamon & Maxwell, 2000; Maurel *et al.*, 2003). The ability to assess fertilization capacity *in vitro* would, therefore, be very useful. Indeed, it is an important goal for understanding epididymal sperm biology and may be critical for developing optimised assisted reproduction protocols.

*In vitro* fertilization (IVF) can be used to predict the *in vivo* fertilization capacity of sperm. Ideally, IVF procedures should use homologous oocytes. However, in wild animals, obtaining oocytes in sufficient quantities for IVF can be very challenging. Consequently, heterologous sperm-oocyte interaction assays, such as the generic oocyte zona-pellucida-free hamster assay (Soler & Garde, 2003), or IVF that involves oocytes from related species (Comizzoli *et al.*, 2001; Soler *et al.*, 2008; García-Álvarez *et al.*, 2009), must be used.

The aim of the present study was to assess the use of prepubertal domestic goat oocytes to determine the fertilization capacity of frozen-thawed epididymal Iberian ibex spermatozoa. As the optimal concentration of oestrous sheep serum (ESS) in the fertilization medium has not been determined for domestic sheep and goat IVF protocols that involve cryopreserved epididymal sperm, the effect of different concentrations on fertilization and embryo development was also examined.

## Materials and methods

Dulbecco's phosphate-buffered saline (PBS) was purchased from Gibco (Paisley, UK), and OviPure®

from Nidacon International AB (Mönldal, Sweden). The antibiotics for sperm extenders were purchased from Calbiochem® (La Jolla, California, USA). All other reagents were purchased from Sigma-Aldrich Chemical Co. (Madrid, Spain).

### Sperm collection

Testes were obtained from five mature Iberian ibexes that had been legally hunted during the rutting season (November/December) in the Tejeda and Almiraj Game Reserve in southern Spain (36°N latitude). The testes, with their scrotal sac, were transported to the laboratory immediately after removal. All materials were kept at ambient temperature (about 11°C) during transport and laboratory processing. To reduce the death-to-sperm collection time, a small laboratory was set up in the mountains of the hunting reserve. All epididymal spermatozoa were collected between 2 and 8 h after death; no sperm variables are affected within 8–9 h of death (Santiago-Moreno *et al.*, 2006). The testes and epididymes were removed from the scrotal sac and the caudae epididymides isolated from the testes and surrounding connective tissue. Epididymal spermatozoa were collected by the retrograde flushing method (Santiago-Moreno *et al.*, 2009b) using 1 ml of Tris-citric acid-glucose medium (TCG) at ambient temperature. This TCG was composed of 3.8% Tris (w/v), 2.2% citric acid (w/v), 0.6% glucose (w/v), 5% glycerol (v/v), and 6% egg yolk (v/v). Two antibiotics – penicillin G potassium salt (1000 IU/ml) and streptomycin sulphate (1 mg/ml) – were added to this medium. Sperm counts were recorded using a Neubauer chamber (Marienfeld, Lauda-Königshofen, Germany). The final samples were diluted with TCG medium to a concentration of  $800 \times 10^6$  sperm/ml, placed in 15 ml centrifuge tubes (Falcon™) set in a beaker with 100 ml of water at 12°C, and then transferred to a refrigerator at 5°C. Cooling to 5°C took about 1 h; the suspension was then maintained at this temperature for an additional 2 h. At this point, aliquots of samples were loaded into 0.25 ml French straws (IMV®; L'Aigle Cedex, France), sealed by ultrasound using an Ultra Seal 31 apparatus (Minitub, Germany), and frozen by placing them in the vapour 5 cm above the surface of a liquid nitrogen container for 10 min before plunging them into the liquid nitrogen itself.

### Sperm analysis and processing

Sperm motility, morphological abnormalities, acrosome integrity and plasma membrane integrity were assessed for each sample – both fresh and frozen-thawed – to determine *in vitro* sperm quality. The percentage of motile spermatozoa and the quality of sperm motility were evaluated subjectively using a

phase-contrast microscope (Zeiss, Germany) at  $\times 400$ , examining samples incubated for 20 min at  $37^{\circ}\text{C}$ . The vigour with which the sperm cells moved was scored on a scale from 0 (lowest) to 5 (highest). Sperm viability was assessed by staining an aliquot of sperm suspension with nigrosin-eosin (Campbell *et al.*, 1956). Simultaneously, plasma membrane integrity was assessed using the hypo-osmotic swelling test (Jeyendran *et al.*, 1984). Morphological abnormalities were assessed by phase-contrast microscopic examination of glutaraldehyde-fixed samples. Spermatozoa with cytoplasmic droplets were considered morphologically normal as these are commonly seen in epididymal sperm cells. The percentage of spermatozoa with intact acrosomes was assessed by observing spermatozoa in samples fixed in buffered 2% glutaraldehyde solution at  $37^{\circ}\text{C}$ , using phase-contrast microscopy (magnification  $\times 1000$ ) (Pursel & Johnson, 1974). Each analysis required the observation of 200 cells. In addition, the motility variables of the frozen-thawed sperm samples were assayed using a computer-aided sperm analysis system (CASA) running Sperm Class Analyzer<sup>®</sup> v.4.0. software (Microptic S.L., Barcelona, Spain), and their membrane and acrosome integrity examined by fluorescence staining (Santiago-Moreno *et al.*, 2013; Soler *et al.*, 2005) (the necessary equipment was not available in the field laboratory to test the fresh sperm).

For IVF, the straws were thawed for 1 min at  $37^{\circ}\text{C}$  and highly motile spermatozoa selected by density gradient (OviPure<sup>®</sup>) centrifugation. This procedure is the first time that ibex sperm has been purified by density gradient; a protocol for domestic rams was therefore followed. Briefly, 1 ml of Bottom Layer Medium was placed in a 15 ml Falcon<sup>®</sup> tube, and then carefully overlaid with 1 ml of Top Layer Medium. Aliquots (50  $\mu\text{l}$ ) of frozen-thawed sperm were placed at the top of the gradient and centrifuged at 300 g for 10 min. After centrifugation, the supernatant was carefully removed using a pipette. The sperm pellet was resuspended in 4 ml of synthetic oviductal fluid (SOF), centrifuged at 300 g for 5 min, and the supernatant again carefully removed. The remaining pellet was resuspended in SOF for use in IVF procedures.

### Oocyte collection and *in vitro* maturation

Ovaries from prepubertal domestic goats were obtained from a local abattoir and transported to the laboratory in sterile Dulbecco's PBS at  $37\text{--}39^{\circ}\text{C}$ . Once in the laboratory, the ovaries were washed three times with PBS. Cumulus-oocyte complexes (COCs) were recovered by slicing the ovary in HEPES-buffered TCM-199 medium supplemented with 1.1 mg/ml sodium bicarbonate and 50  $\mu\text{g}/\text{ml}$  gentamicin. Only

COCs with two or more complete and compact layers of cumulus cells and homogenous cytoplasm were chosen for use.

The selected COCs were washed in TCM-199. Groups of 25–30 COCs were matured in 100  $\mu\text{l}$  TCM-199 supplemented with 275  $\mu\text{g}/\text{ml}$  sodium pyruvate, 146  $\mu\text{g}/\text{ml}$  L-glutamine, 10% (v/v) donor bovine serum (DBS), 10  $\mu\text{g}/\text{ml}$  oLH, 10  $\mu\text{g}/\text{ml}$  oFSH, 1  $\mu\text{l}/\text{ml}$   $17\beta$ -estradiol, 100  $\mu\text{M}$  cysteamine and 50  $\mu\text{g}/\text{ml}$  gentamicin. Their maturation was completed by incubating these samples under mineral oil for 24–27 h at  $38.5^{\circ}\text{C}$  in a humidified atmosphere containing 5%  $\text{CO}_2$  in air (Jiménez-Macedo *et al.*, 2005).

### IVF, assessment of nuclear stage, and embryo development

After maturation, the COCs were partially denuded by gentle pipetting and transferred into SOF. Fertilization was performed in 50  $\mu\text{l}$  microdrops of medium supplemented with ESS at different concentrations: SOF-C (0% ESS), SOF-2 (2% ESS) or SOF-20 (20% ESS) (five replicates). Groups with a maximum of 15 COCs per microdrop were co-incubated with  $1 \times 10^6$  spermatozoa/ml for 20–24 h at  $38.5^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , 5%  $\text{O}_2$  and 90%  $\text{N}_2$  in a humidified atmosphere (Comizzoli *et al.*, 2001).

At 17 h post-insemination (hpi), a sample of half of the presumptive zygotes was separated from the cumulus cells by gently pipetting, and fixed in acetic acid:ethanol (1:3, v/v) for 24 h at  $4^{\circ}\text{C}$ . The presumptive zygotes were stained with 1% lacmoid and observed under a phase-contrast microscope. Zygotes were classified as normally fertilized if one female and one male pronucleus (2PN) was observed. The male pronucleus was detected when one sperm tail was close to the pronucleus. Zygotes with 3PN were considered polyspermic; oocytes with 1PN were classified as asynchronous (Mogas *et al.*, 1997).

The remaining presumptive zygotes were completely denuded, washed by gentle pipetting and cultured in groups of 10–15 in 20  $\mu\text{l}$  culture drops of SOF medium at  $38.5^{\circ}\text{C}$  in a humidified 5%  $\text{CO}_2$ , 5%  $\text{O}_2$  and 90%  $\text{N}_2$  atmosphere for 7 days. On day 5, the embryos were transferred to microdrops with fresh medium. Cleavage and blastocyst formation rates were evaluated at 48 and 168 hpi respectively (Jiménez-Macedo *et al.*, 2005).

### Statistical analysis

The data for the sperm variables were not normally distributed (as determined by the Shapiro-Wilk's test); they were therefore arcsine-transformed before analysis. The *t*-test for matched pairs was used to compare the values of sperm variables before and after thawing. The nuclear status after fertilization

**Table 1** Sperm variable (mean  $\pm$  S.E.) values for fresh and frozen–thawed epididymal spermatozoa collected within 8 h of death

	Fresh samples	Frozen–thawed samples
% Motile sperm	86.0 $\pm$ 1.8 <sup>a</sup>	57.3 $\pm$ 6.4 <sup>b</sup>
Quality of motility (0–5)	3.5 $\pm$ 0.1 <sup>a</sup>	3.5 $\pm$ 0.1 <sup>a</sup>
% Viable sperm	87.2 $\pm$ 1.4 <sup>a</sup>	73.4 $\pm$ 3.7 <sup>b</sup>
% Positive endosmosis	86.1 $\pm$ 2.4 <sup>a</sup>	50.4 $\pm$ 5.6 <sup>b</sup>
Acrosome integrity (%NAR)	91.8 $\pm$ 2.0 <sup>a</sup>	82.9 $\pm$ 2.6 <sup>b</sup>
% Morphological abnormalities	2.6 $\pm$ 0.3 <sup>b</sup>	15.6 $\pm$ 3.2 <sup>a</sup>

<sup>a,b</sup>Different superscript letters between columns indicate significant differences ( $P < 0.05$ ).

**Table 2** Sperm motility variables (mean  $\pm$  S.E.) analysed by CASA, and membrane and acrosome integrity fluorescence staining (fluorochrome combination of propidium iodide and fluorescein isothiocyanate-conjugated peanut [*Arachis hypogea*] agglutinin) in frozen–thawed epididymal spermatozoa collected within 8 h of death

	Frozen–thawed samples
% Immotile sperm	44.0 $\pm$ 4.7
% Non-progressive motility	20.1 $\pm$ 2.4
% Progressive motility	35.6 $\pm$ 3.3
% Live sperm with an intact acrosome	60.7 $\pm$ 7.4
% Live sperm with a damaged acrosome	1.4 $\pm$ 0.8
% Dead sperm with an intact acrosome	21.6 $\pm$ 5.4
% Dead sperm with a damaged acrosome	16.3 $\pm$ 3.2

**Table 3** Nuclear status of zygotes at 17 h following insemination of prepubertal domestic goat oocytes with frozen–thawed ibex epididymal sperm

Group	Total presumptive zygotes	Nuclear status (mean %)			Total penetrated (mean %)
		1PN asynchrony	2PN	3PN polyspermic	
SOF-C	150	3 (1.9) <sup>a</sup>	21 (14.5) <sup>a,b</sup>	0 (0) <sup>a</sup>	24 (16.4) <sup>a,b</sup>
SOF-2	140	7 (8.8) <sup>a</sup>	39 (27.7) <sup>a</sup>	8 (10.5) <sup>a</sup>	54 (47.1) <sup>a</sup>
SOF-20	151	0 (0) <sup>a</sup>	4 (2.9) <sup>b</sup>	0 (0) <sup>a</sup>	4 (2.9) <sup>b</sup>

SOF: synthetic oviductal fluid medium. ESS: oestrous sheep serum. SOF-C: 0% ESS; SOF-2: 2% ESS; SOF-20: 20% ESS. 1PN: asynchrony, i.e. oocytes with one female pronucleus and one condensed sperm head; 2PN: zygotes with two pronuclei (normal fertilization); 3PN: polyspermic zygotes.

<sup>a,b</sup>Values with different subscripts within the same column indicate significant differences ( $P < 0.05$ ). Values in parentheses are the mean (%) of five replicates.


and embryonic development data were assessed using general linear model (GLM) ANOVA. When this analysis revealed the ESS concentration to have a significant effect, differences between ESS treatment groups were examined using Tukey's *post hoc* test. All calculations were performed using SAS v.9.1 software. Data are presented as means  $\pm$  standard error (SE). Significance was set at  $P < 0.05$ .

## Results

The freezing–thawing process significantly reduced the percentage of motile spermatozoa ( $P < 0.001$ ), the percentage of viable sperm according to nigrosin–eosin

staining ( $P < 0.05$ ), the membrane integrity according to the hypo-osmotic swelling test ( $P < 0.001$ ), the percentage of sperm showing acrosome integrity ( $P < 0.05$ ), and the percentage of morphological sperm abnormalities ( $P < 0.001$ ) (Table 1). Table 2 shows the results for the motility variables for frozen–thawed sperm as analysed objectively by CASA, plus the percentage of sperm showing plasma membrane and acrosome integrity as assessed by fluorescence staining. The results are similar to those obtained by the previous methods.

Table 3 shows the recorded nuclear stage data for oocytes at 17 h after IVF. The SOF-C treatment provided 14.5% 2PN zygotes (Fig. 1 shows a representative 2PN zygote). In the SOF-2 treatment, the percentage of 2PN



**Figure 1** Normal zygote obtained via heterologous IVF involving prepubertal domestic goat oocytes and frozen–thawed ibex epididymal sperm (recovered after death). Arrows indicate the female and male pronucleus (2PN).

**Table 4** Embryonic development at 48 and 168 h post-fertilization with frozen–thawed ibex epididymal sperm

Group	Total oocytes	Cleavage (mean %) 48 hpi	Blastocyst formation (mean %) 168 hpi
SOF-C	165	29 (18.3) <sup>a</sup>	2 (1.1) <sup>a</sup>
SOF-2	162	36 (25.6) <sup>a</sup>	12 (7.5) <sup>a</sup>
SOF-20	153	4 (2.8) <sup>a</sup>	0 (0) <sup>a</sup>

<sup>a</sup>Values with different subscripts within the same column indicate significant differences ( $P < 0.05$ ). SOF: synthetic oviductal fluid medium. ESS: oestrous sheep serum. SOF-C: 0% ESS; SOF-2: 2% ESS; SOF-20: 20% ESS. Values in parentheses are the mean (%) of five replicates.

zygotes increased (but not significantly) to 27.7%. In the SOF-20 group, the percentage decreased to 2.9% ( $P < 0.05$  with respect to the SOF-2 result). Polyspermic fertilization (3PN) was only observed with the SOF-2 treatment (10.5%). The percentage of asynchronous oocytes (1PN) was 1.9% with the SOF-C treatment, and 8.8% with the SOF-2 treatment (no significant difference).


Table 4 shows the recorded embryo development data after 7 days of culture. The percentage of cleaved embryos at 48 hpi was higher with the SOF-2 than with the SOF-C and SOF-20 treatments (25.6% versus 18.3 and 2.8% respectively); however, none of these differences was significant. At 168 hpi, the blastocyst formation rate was higher with the SOF-2 than with the SOF-C or SOF-20 treatments (7.5 versus 1.1 and 0% respectively); but again, the differences were not significant. Figure 2 shows a blastocyst stage embryo.

## Discussion

Overall, the values for the fresh and frozen–thawed sperm variables were similar to those previously reported for this species in samples collected within 8 h of death (Santiago-Moreno *et al.*, 2006; Fernández-Santos *et al.*, 2011). The percentage of live spermatozoa with damaged acrosomes was very low; such damage was usually seen in dead sperm. Both the motility and viability variables measured and the CASA and fluorescence results showed the quality of the present frozen–thawed epididymal sperm to be greater than that of electro-ejaculated sperm from the same species (Coloma *et al.*, 2010). This finding suggests that epididymal sperm cells better resist freezing and thawing.

This report describes the first use of heterologous IVF in a threatened ibex species. The number of penetrated oocytes and the embryo development achieved indicated that heterologous IVF may be an effective method for predicting the fertilization capacity of ibex spermatozoa.

While the fertilization capacity of ibex sperm doses can be determined via interspecific AI *in vivo*, the results can be influenced by factors that are difficult to control, such as sub-clinical ovarian and uterine pathologies (Smith & Sherman, 2009), the quality and early regression of the corpus luteum, early embryonic mortality, and the response of females to hormonal treatments for synchronisation and the induction of ovulation (Salamon & Maxwell, 2000; Maurel *et al.* 2003). Fertility also depends on the presence of a heterogeneous population of sperm (i.e. with sperm subpopulations) (Holt, 2009), the quality



**Figure 2** Blastocyst stage embryo obtained by heterologous IVF of a prepubertal domestic goat oocyte by a frozen-thawed ibex epididymal sperm.

of the oocyte vestments, and the quality of the oocyte itself. Therefore, neither a simple spermogram nor routine post-thaw evaluation can adequately determine the fertilization capacity of cryopreserved sperm (Rodríguez-Martínez, 2003). Furthermore, correlations between sperm motility and fertility may vary greatly in domestic ruminant species (Kjaestad *et al.*, 1993; Bailey *et al.*, 1994; Januskauskas *et al.*, 2003) and the relationship between the degree of damage to the sperm plasma membrane and fertility is not always clear (Garner *et al.*, 1997; Rodríguez-Martínez & Larsson, 1998). Functional *in vitro* assays that involve IVF, however, seem able to determine the ability of sperm cells to accomplish specific steps in fertilization and the triggering of early embryo development. This aspect is important when valuable sperm doses from non-domestic and endangered species are used, as in these species definitive correlations between normally recorded sperm variables and fertility are even more difficult to establish (Durrant, 2009).

The success of the present heterologous fertilization assay is underpinned by the fact that both domestic goats and Iberian ibex have 60 chromosomes. Other authors report heterologous IVF assays involving mouflon (*Ovis gmelini musimon*) sperm and sheep (*Ovis aries*) zona-pellucida-intact oocytes – both species with 54 chromosomes – to be successful (Berlinguer *et al.*, 2003).

We previously reported the usefulness and adequacy of prepubertal domestic goat oocytes in IVF also involving goat sperm (Romaguera *et al.*, 2010).

The present results also support the idea that they may be used in heterologous IVF assays of ibex sperm fertilization capacity. The mean percentage of oocytes appropriately fertilized (2PN) in the SOF-2 treatment group was 27.7%; this finding is similar to the 20% reported in earlier *in vivo* work (Santiago-Moreno *et al.*, 2006).

The concentration of ESS in the culture medium may be a key factor in the success of IVF. The present results showed a significantly higher percentage of 2PN zygotes with the SOF-2 treatment than with SOF-20. How higher concentrations of ESS might prevent oocyte penetration is unknown, but the possibilities include the induction of sperm agglutination and alterations of the factors controlling oocyte-sperm interactions, such as sperm-ZP3 binding affinity (Cummins, 1995; Oehninger & Franken, 2006). The percentage of cleaved embryos at 48 hpi was higher with the SOF-2 than the SOF-C and SOF-20 treatments (25.6 versus 18.3 and 2.8% respectively), although the differences were not significant. A larger number of oocytes and/or replicates might be needed to detect significant differences in embryo development. While the SOF-2 treatment seemed to have some beneficial effect on fertilization, this effect was not so visible at the blastocyst stage. Using both fresh and frozen-thawed goat semen, Velilla *et al.* (2004) found 22% of oocytes to be appropriately fertilized (2PN), similar to the figure obtained with the present SOF-2 treatment (27.7%). In another study that used freshly ejaculated sperm and oocytes from follicles of more than 3 mm

diameter, an 18.1% blastocyst formation rate was achieved (Romaguera *et al.*, 2011), greater than that achieved with the present SOF-2 treatment (7.5%) for frozen-thawed sperm. It has been reported that the ESS concentration does not affect the fertilization rates of electro-ejaculated deer sperm (Comizzoli *et al.*, 2001), but these authors did not follow embryo development to the blastocyst stage. In addition, some authors (Huneau *et al.*, 1994) have reported a beneficial effect of adding 20% ESS to the fertilization medium for sheep, but while high rates of fertilization were obtained, these authors did not use cryopreserved epididymal sperm. The present results may differ from those of previous studies for this reason. In addition, little information is known about the effects of ESS on ibex sperm. Further studies are needed to clarify the effects of ESS. The effect of fetal bovine serum should also be examined (Ghasemzadeh-Nava & Tajik, 2000; Tibary *et al.*, 2005).

IVF usually involves the selection of sperm via a density or swim-up gradient (Mortimer, 1994). The literature contains no report regarding the selection of functional ibex sperm, so a density gradient designed for rams (OviPure®) was used in this study. As 27.7% of oocytes were appropriately fertilized (2PN) even in the SOF-2 treatment group, this method of sperm selection can be recommended. However, density gradients more specific for caprines (e.g., CapriPure® [Batista *et al.*, 2011]) should be tested.

In conclusion, heterologous IVF assays using prepubertal domestic goat oocytes can be used to assess the fertilization capacity of cryopreserved ibex epididymal sperm, and therefore perhaps that of other wild mountain ungulates. The use of 2% ESS in the fertilization medium may be a better option than using none at all or the standard 20% concentration as, in the present work, it increased the percentage of appropriately fertilized oocytes (2PN).

## Acknowledgements

This study was funded by grants from MINECO (AGL2011-23784 and AGL2011-25495), and CONACYT Mexico (0105961/10110/194/09). J. Lopez was the recipient of a CONACYT scholarship number 240847. The authors thank Dr Cecilia Esquivelzeta and Dr Manuel Castillo for statistical assistance.

## References

- Andrabi, S. & Maxwell, W. (2007). A review on reproductive biotechnologies for conservation of endangered mammalian species. *Anim. Reprod. Sci.* **99**, 223–43.
- Anel, L., Alvarez, M., Anel, E., Martinez-Pastor, F., Martinez, F., Chamorro, C. & de Paz, P. (2011). Evaluation of three different extenders for use in emergency salvaging of epididymal spermatozoa from a Cantabric brown bear. *Reprod. Domest. Anim.* **46**, e85–90.
- Bailey, J.L., Robertson, L. & Buhr, M.M. (1994). Relations among *in vivo* fertility, computer-analysed motility and  $\text{Ca}^{++}$  influx in bovine spermatozoa. *Can. J. Anim. Sci.* **74**, 53–8.
- Batista, A.M., Silva, S.V., Soares, A.T., Monteiro, P.L.J., Wischral, A. & Guerra, M.M.P. (2011). Comparison of CapriPure® and Percoll® density gradients for sperm separation of frozen-thawed goat spermatozoa. *Anim. Prod.* **8**, 81–4.
- Berlinguer, F., Ledda, S., Rosati, I., Bogliolo, L., Leoni, G. & Naitana, S. (2003). Superoxide dismutase affects the viability of thawed European mouflon (*Ovis g. musimon*) semen and the heterologous fertilization using both IVF and intracytoplasmic sperm injection. *Reprod. Fertil. Dev.* **15**, 19–25.
- Campbell, R.C., Dott, H.M. & Glover, T.D. (1956) Nigrosin eosin as a stain for differentiating live and dead spermatozoa. *J. Agric. Sci.* **48**, 1–8.
- Cocchia, N., Ciani, F., El-Rass, R., Russo, M., Borzacchiello, G., Esposito, V., Montagnaro, S., Avallone, L., Tortora, G. & Lorizio, R. (2010). Cryopreservation of feline epididymal spermatozoa from dead and alive animals and its use in assisted reproduction. *Zygote* **18**, 1–8.
- Coloma, M.A., Gómez-Brunet, A., Velázquez, R., Toledano-Díaz, A., López-Sebastián, A. & Santiago-Moreno, J. (2010). Freezability of Iberian ibex (*Capra pyrenaica*) spermatozoa according to the glycerolization temperature and plasma testosterone concentration. *Cryobiology* **61**, 204–10.
- Coloma, M.A., Toledano-Díaz, A., Castaño, C., Velázquez, R., Gómez-Brunet, A., López-Sebastián, A. & Santiago-Moreno, J. (2011) Seasonal variation in reproductive physiological status in the Iberian ibex (*Capra pyrenaica*) and its relationship with sperm freezability. *Theriogenology* **76**, 1695–705.
- Comizzoli, P., Mauget, R. & Mermillod, P. (2001). Assessment of *in vitro* fertility of deer spermatozoa by heterologous IVF with zona-free bovine oocytes. *Theriogenology* **56**, 261–74.
- Cummins, J.M. (1995). Test of sperm function. In *Gametes. The Spermatozoon. Cambridge Reviews in Human Reproduction*. (eds J.G. Grudzinskas & J.L. Yovich), pp. 70–103. Cambridge, UK: Cambridge University Press.
- Durrant, B.S. (2009). The importance and potential of artificial insemination in CANDES (companion animals, non-domestic, endangered species). *Theriogenology* **71**, 113–22.
- Fernández-Santos, M.R., Soler, A.J., Ramon, M., Ros-Santaella, J.L., Maroto-Morales, A., Garcia-Alvarez, O., Bisbal, A., Garde, J.J., Coloma, M.A. & Santiago-Moreno, J. (2011). Effect of post-mortem time on post-thaw characteristics of Spanish ibex (*Capra pyrenaica*) spermatozoa. *Anim. Reprod. Sci.* **129**, 56–66.
- García-Álvarez, O., Maroto-Morales, A., Martínez-Pastor, F., Fernández-Santos, M., Estesio, M., Pérez-Guzmán, M. & A.J. Soler. (2009). Heterologous *in vitro* fertilization is a good procedure to assess the fertility of thawed ram spermatozoa. *Theriogenology* **71**, 643–50.



- Garner, D.L., Thomas, C.A., Allen, C.H., Senger, P.L. & Sasser, R.G. (1997). Effect of cryopreservation on bovine sperm viability as determined by dual DNA staining. *Reprod. Domest. Anim.* **32**, 279–83.
- Ghasemzadeh-Nava, H. & Tajik, P. (2000). *In vitro* maturation of ovine follicular oocytes in different concentrations of fetal calf serum and estrous sheep serum. *Theriogenology* **53**, 453 (Abstr.).
- Holt, W.V. (2009). Is semen analysis useful to predict the odds that the sperm will meet the egg? *Reprod. Domest. Anim.* **44**, 31–8.
- Huneau, D., Crozet, N. & Ahmed-Ali, M. (1994). Estrous sheep serum as a potent agent for ovine IVF: effect on cholesterol efflux from spermatozoa and the acrosome reaction. *Theriogenology* **42**, 1017–28.
- Januskauskas, A., Johannisson, A. & Rodríguez-Martínez, H. (2003). Subtle membrane changes in cryopreserved bull semen in relation to sperm viability, chromatin structure and field fertility. *Theriogenology* **60**, 743–58.
- Jeyendran, R., Van Der Ven, H., Perez-Pelaez, M., Crabo, B. & Zaneveld, L. (1984). Development of an assay to assess the functional integrity of the human sperm membrane and its relation to other semen characteristics. *J. Reprod. Fertil.* **70**, 219–28.
- Jiménez-Macedo, A.R., Izquierdo, D., Anguita, B. & Paramio, M.T. (2005). Comparison between intracytoplasmic sperm injection and *in vitro* fertilization employing oocytes derived from prepubertal goats. *Theriogenology* **64**, 1249–62.
- Keeley, T., McGreevy, P.D. & O'Brien, J.K. (2012). Cryopreservation of epididymal sperm collected postmortem in the Tasmanian devil (*Sarcophilus harrisii*). *Theriogenology* **78**, 315–25.
- Kjaestad, H., Ropstad, E. & Andersen Berg, K. (1993). Evaluation of spermatological parameters used to predict the fertility of frozen bull semen. *Acta Vet. Scand.* **34**, 299–303.
- Maurel, M., Roy, F., Hervé, V., Bertin, J., Vaiman, D., Crihiu, E., Manfredi, E., Bouvier, F., Lantier, I., Boue, P. & Guillou, F. (2003). Immune response to equine chorionic gonadotropin used for the induction of ovulation in goats and ewes. *Ginecol Obstet. Fertil.* **31**, 766–9.
- Mogas, T., Palomo, M., Izquierdo, M. & Paramio, M. (1997). Morphological events during *in vitro* fertilization of prepubertal goat oocytes matured *in vitro*. *Theriogenology* **48**, 815–29.
- Mortimer, D. (1994). Sperm washing. In *Practical Laboratory Andrology* (ed. D. Mortimer), pp. 267–86. New York: Oxford University Press Inc.
- Oehninger, S. & Franken, D. (2006). Testing sperm manufacturing quality: the sperm–zona binding assay. In *The Sperm Cell. Production, Maturation, Fertilization, Regeneration* (eds C.J. De Jonge & C.L. Barratt), pp. 194–216. New York, USA: Cambridge University Press.
- Pursel, V. & Johnson, L. (1974). Glutaraldehyde fixation of boar spermatozoa for acrosome evaluation. *Theriogenology* **1**, 63–8.
- Rodríguez-Martínez, H. (2003). Laboratory semen assessment and prediction of fertility: still Utopia? *Reprod. Domest. Anim.* **38**, 312–8.
- Rodríguez-Martínez, H. & Larsson, B. (1998). Assessment of sperm fertilizing ability in farm animals. *Acta Agr. Scand. Sect. A Animal Sci. Suppl.* **29**, 12–8.
- Romaguera, R., Casanovas, A., Morató, R., Izquierdo, D., Catalá, M., Jiménez-Macedo, A.R., Mogas, T. & Paramio, M.T. (2010). Effect of follicle diameter on oocyte apoptosis, embryo development and chromosomal ploidy in prepubertal goats. *Theriogenology* **74**, 364–73.
- Romaguera, R., Moll, X., Morató, R., Roura, M., Palomo, M., Catalá, M., Jiménez-Macedo, A.R., Hammami, S., Izquierdo, D., Mogas, T., Paramio, M.T. (2011). Prepubertal goat oocytes from large follicles result in similar blastocyst production and embryo ploidy than those from adult goats. *Theriogenology* **76**, 1–11.
- Salamon, S., & Maxwell, W. (2000). Storage of ram semen. *Anim. Reprod. Sci.* **62**, 77–112.
- Santiago-Moreno, J., Toledano-Díaz, A., Pulido-Pastor, A., Gómez-Brunet, A. & López-Sebastián, A. (2006). Birth of live Spanish ibex (*Capra pyrenaica hispanica*) derived from artificial insemination with epididymal spermatozoa retrieved after death. *Theriogenology* **66**, 283–91.
- Santiago-Moreno, J., Toledano-Díaz, A., Dorado, J., Pulido-Pastor, A., Coloma, M.A. & López-Sebastián, A. (2007). Recovery and cryopreservation of Spanish ibex epididymal spermatozoa. *Arch. Androl.* **53**, 309–16.
- Santiago-Moreno, J., Coloma, M., Dorado, J., Pulido-Pastor, A., Gómez-Guillamon, F., Salas-Vegas, R., Gómez-Brunet, A. & López-Sebastián, A. (2009a). Cryopreservation of Spanish ibex (*Capra pyrenaica*) sperm obtained by electroejaculation outside the rutting season. *Theriogenology* **71**, 1253–60.
- Santiago-Moreno, J., Astorga, R.J., Luque, I., Coloma, M.A., Toledano-Díaz, A., Pulido-Pastor, A., Gómez-Guillamon, F., Salas-Vega, R. & López-Sebastián, A. (2009b). Influence of recovery method and microbial contamination on the response to freezing-thawing in ibex (*Capra pyrenaica*) epididymal spermatozoa. *Cryobiology* **59**, 357–62.
- Santiago-Moreno, J., Coloma, M., Toledano-Díaz, A., Castaño, C., Gómez-Brunet, A., López-Sebastián, A. (2010). Assisted reproduction in Mediterranean wild ruminants: lessons from the Spanish ibex (*Capra pyrenaica*). *Soc. Reprod. Fertil. Suppl.* **67**, 431–41.
- Santiago-Moreno, J., Castaño, C., Toledano-Díaz, A., Esteso, M.C., López-Sebastián, A., Guerra, R., Ruiz, M.J., Mendoza, N., Luna, C., Cebrián-Pérez, J.A. & Hildebrandt, T.B. (2013). Cryopreservation of aoudad (*Ammotragus lervia sahariensis*) sperm obtained by transrectal ultrasound-guided massage of the accessory sex glands and electroejaculation. *Theriogenology* **79**, 383–91.
- Smith, M.C. & Sherman, D.M. (eds) (2009). *Goat Medicine*, 2nd edn. Iowa, USA: Wiley-Blackwell.
- Soler, A.J., & Garde, J.J. (2003). Relationship between the characteristics of epididymal red deer spermatozoa and penetrability into zona-free hamster ova. *J. Androl.* **24**, 393–400.
- Soler, A.J., Pérez-Guzmán, M.D. & Garde, J.J. (2003). Storage of red deer epididymides for four days at 5°C: effects on sperm motility, viability, and morphological integrity. *J. Exp. Zool.* **295A**, 188–99.



- Soler, A.J., Esteso, M.C., Fernández-Santos, M.R. & Garde, J.J. (2005). Characteristics of Iberian red deer (*Cervus elaphus hispanicus*) spermatozoa cryopreserved after storage at 5°C in the epididymis for several days. *Theriogenology* **64**, 1503–17.
- Soler, A., Poulin, N., Fernández-Santos, M., Cognie, Y., Esteso, M., Gardé, J.J. & Mermillod, P. (2008). Heterologous *in vitro* fertility evaluation of cryopreserved Iberian red deer epididymal spermatozoa with zona intact sheep oocytes and its relationship with the characteristics of thawed spermatozoa. *Reprod. Domest. Anim.* **43**, 293–98.
- Tibary, A., Anouassi, A. & Khatir, H. (2005). Update on reproductive biotechnologies in small ruminants and camelids. *Theriogenology* **64**, 618–38.
- Velilla, E., Izquierdo, D., Rodríguez-González, E., López-Béjar, M., Vidal, F. & Paramio, M. (2004). Distribution of prepubertal and adult goat oocyte cortical granules during meiotic maturation and fertilization: ultrastructural and cytochemical study. *Mol. Reprod. Dev.* **8**, 507–14.
- Watson, P.F & Holt, W.V. (eds) (2001) *Cryobanking the Genetic Resource*. London: Taylor and Francis.