

This is the **accepted version** of the journal article:

Jiménez-Murcia, Susana; Granero, Roser; Stinchfield, Randy; [et al.]. «Typologies of young pathological gamblers based on sociodemographic and clinical characteristics». *Comprehensive Psychiatry*, Vol. 54 (2013), p. 1153-1160. DOI 10.1016/j.comppsy.2013.05.017

This version is available at <https://ddd.uab.cat/record/319689>

under the terms of the  **COPYRIGHT** license

Jiménez-Murcia S, Granero R, Stinchfield R, Fernández-Aranda F, Penelo E, Savvidou LG, Fröberg F, Aymamí N, Gómez-Peña M, Moragas L, del Pino-Gutiérrez A, Fagundo AB, Menchón JM. (2013). Typologies of young pathological gamblers based on sociodemographic and clinical characteristics. *Comprehensive Psychiatry*, 54(8), 1153-1160. doi: 10.1016/j.comppsy.2013.05.017.

*This is the author manuscript, which has undergone full peer review but has not yet been copyedited, typeset, paginated, or proofread. Consequently, this version may differ from the final Version of Record.

Abstract

Objective: The aim of this study is to explore empirical clusters within the population of young Spanish individuals attending outpatient pathological gambling treatment. **Method:** The South Oaks Gambling Screen (SOGS), the Symptom Checklist (SCL-90-R), the Temperament and Character Inventory-R (TCI-R) and other clinical and psychopathological measures were administered to 154 patients (between 17 and 25 years old). The two-step cluster analysis explored the presence of empirical heterogeneous groups based on clinical and socio-demographic characteristics. **Results:** Three clusters of young pathological gambling patients emerged. Type I showed less psychopathology and more functional personality traits. Type II showed a profile characterized by major emotional distress, shame, immaturity, hostility and negative feelings. Type III showed the most severe psychopathological profile and most psychopathological disturbances and schizotypal traits. **Conclusions:** These results suggest that three distinct endophenotypes exist, and that environmental factors have a stronger influence in the first, while in the second and third, individual factors related to deficits of emotional regulation stand out. © 2013 Elsevier Inc. All rights reserved.

1. Introduction

Pathological gambling (PG) is defined by uncontrollable gambling behavior that results in severe negative effects on the patient's occupation, relationships, psychological health and other relevant areas of life [1]. Moreover, although people suffering from this disorder recognize that gambling is the reason of severe impairments in the family and on a community level (many patients indicate that gambling was the reason they lost significant relationships and jobs), they cannot resist the impulse to gamble.

According to Chiu & Storm [2], youth are overly sensitive to norms that contribute to the maintenance of high-risk behaviors such as gambling. Further studies consistently report that youth problem gambling is associated with other psychosocial problems, such as depression, substance abuse, and delinquency [3,4]. The first exposure to gambling usually happens early in life, in the form of electronic and/or internet gambling, lotteries, slot-machines, card games, casinos, and many other types of games [5]. This early exposure represents one of the critical risk factors for the onset of PG. Furthermore, the main steps or actions that form the addictive behavior are located in the same social atmosphere, involving gambling as well as the excitement of risk-taking (strong characteristic of youth).

It is crucial to identify PG behaviors while still in their earliest manifestations, since the persistence to gamble involves the accumulation of negative social and psycho-pathological consequences, such as unemployment, debts, disrupted family relationships, and search for money by illegal means, substance abuse and affective-anxiety disorders [6].

Even though there is strong evidence about the heterogeneity of PG and the existence of different subgroups based on socio demographic and clinical characteristics among adults [7–11], few studies have attempted to identify clusters of young pathological gamblers. An exception is the recent work published by Gupta et al. [12], which was conducted with students of English-speaking schools in Quebec and Ontario. This study explores the “Pathways Model” proposed by Blaszczynski & Nower [8], in a sample of 109 adolescents that meet the criteria for problem gambling. The results suggest that, in addition to the three subtypes described in the “Pathways Model” (behaviorally conditioned, emotionally vulnerable and antisocial impulsive problem gamblers) there are two more subtypes, one characterized by depressive symptoms and the other by externalizing and internalizing symptoms. Furthermore, Goldstein et al. [13] examined subgroups of gamblers among adolescent emergency patients, finding two groups with few or many consequences of gambling. The group with many consequences also exhibited substance abuse, delinquency and violence, and resembled Pathway 3 in the model by Blaszczynski & Nower [8].

The aim of this study is to explore empirical clusters in a sample of Spanish young people (17–25 years old) attending outpatient PG treatment. Based on empirical evidence, we hypothesized that it would be possible to identify clinically relevant subgroups of young PG patients, based on psychopathology and personality traits. We expected to identify one subgroup characterized by low psychopathology and non-dysfunctional personality traits, a second with moderate psychopathology and melancholic personality

traits (following character types described by Cloninger et al. [14] and a third formed by a subgroup of patients with severe psychopathology and disorganized or schizotypal personality configuration (as defined by Cloninger et al. [14]).

2. Method

2.1. Participants

The sample included 154 pathological gambling patients, 17–25 years old, who presented for assessment and outpatient treatment at the Pathological Gambling Unit in the Psychiatric Department of Bellvitge University Hospital (HUB) in Barcelona. All participants were diagnosed by experienced psychologists and psychiatrists in PG, using the Diagnostic Questionnaire for Pathological Gambling according to DSM-IV criteria [15].

The first column of Table 1 shows the socio demographic and clinical characteristics of the sample. The mean age was 22.4 years old ($SD = 2.2$), 94.2% were male and 61.3% of them were employed at the intake. More than half of the sample had achieved no more than primary educational level and only 11.3% were married or lived with a partner.

2.2. Measures

South Oaks Gambling Screen (SOGS) [16], Spanish validation by Echeburúa et al. [17]. This is a 20-item screening questionnaire that identifies probable pathological gambling. The Spanish validation of this questionnaire shows high reliability and validity. Test–retest reliability is .98 ($p < 0.001$) and internal consistency .94 (Cronbach's alpha). Convergent validity with respect to DSM-III-R criteria for pathological gambling (APA, 1987) was estimated .92 ($p < 0.001$). The total score was used in this study. Additionally, several studies have reported the use and utility of this test as an index of gambling severity [18–20].

Diagnostic questionnaire for Pathological Gambling according to DSM-IV criteria [15]. Spanish adaptation by Jimenez-Murcia et al. [21]. This 19-item questionnaire assesses the DSM-IV diagnostic criteria for PG. Internal consistency ranged between .81 for the general population and .77 for gambling treatment samples. Convergent validity with the SOGS total score was very good: $r = .77$ for the general population and $r = .75$ for gambling treatment groups [15].

Symptom Check List-90 items-Revised (SCL-90-R, Spanish version) [22]. We administered the SCL-90-R to evaluate a broad range of psychological problems and symptoms of psychopathology. Containing 90 items, the SCL-90-R is used to measure nine primary symptom dimensions: Somatization, Obsession–Compulsion, Interpersonal Sensitivity, Depression, Anxiety, Hostility, Phobic Anxiety, Paranoid Ideation, and Psychoticism. In addition, it includes three global indices: a global severity index (GSI), which measures overall psychological distress; a positive symptom distress index (PSDI) to measure the intensity of symptoms; and a positive symptom total (PST), which reports the total self-reported symptoms. The GSI can be used as a summary of the test. This scale has been validated in a Spanish population, obtaining an internal consistency (coefficient alpha) of the items ranging between .81 and .90.

Temperament and Character Inventory-Revised (TCI-R) [23]. This is a 240-item questionnaire. Like the original TCI version [14], this questionnaire is a reliable and valid measure of seven dimensions of personality: four temperament dimensions (harm avoidance, novelty seeking, reward dependence and persistence) and three character dimensions (self-directedness, cooperativeness and self-transcendence). The performance of the Spanish revised version of this questionnaire [24] has been well-documented. The reliability of the different temperament and character dimensions in the Spanish adaptation ranged between .77 and .84 (Cronbach's alpha).

Additional demographic, clinical and socio-familial variables related to gambling were measured using a semi-structured clinical interview, described elsewhere [25,26].

2.3. Procedure

This study was carried out according to the latest version of the Declaration of Helsinki. The Ethics Committee of the Bellvitge University Hospital (Barcelona) approved this study and written informed consent was obtained from all final participants.

The assessment was conducted prospectively at base- line and it involved a single session (with a mean duration of 90 min), during which the above mentioned tests were administered by trained clinical psychologists. In addition to the assessment battery, the patients were explored through a semi-structured face-to-face interview regarding their PG, psychopathological symptoms and personality traits (Jiménez-Murcia et al., 2007). The same interview also assessed sociodemographic data (e.g., education, occupation, marital status) and additional clinical information.

2.4. Statistical analysis

A two-step cluster analysis explored the presence of empirical groups based on the socio demographic and clinical variables (number of problematic games, co- morbidity, SCL-90-R and TCI-R scores, SOGS and DSM Total scores). The two-step method allows discriminating natural groups from a set of variables stabilizing the nearness criterion, with a hierarchical agglomerative clustering whose centres are far apart. Likelihood was the distance measure selected, defining the normal density for continuous variables and the multinomial probability mass function for categorical variables [27]. Average Silhouette Coefficient (a measure of how tightly grouped all the data in the cluster are) measured the goodness-of-fit [28]. This index combines both, cohesion (based on the average distances between all the objects into a cluster) and separation (based on the average distance of any object to all the other objects not contained into the same cluster), and can range between -1 and $+1$; values over 0 are indicative of inappropriate fit, between 0 and 0.2 are considered poor, between 0.2 and 0.5 fair and indices above 0.5 good.

Next, binary logistic regressions (for categorical criteria) and analysis of variance (ANOVA, for quantitative outcomes) compared the empirical clusters on all the measures of the study. Cohen's d measured the effect size of proportions and mean differences (d -coefficients over 0.50 were considered good). Due to the multiple statistical comparisons, Bonferroni-Finner's correction was applied to avoid bias due to Type-I error.

3. Results

Cluster analysis was carried out with 110 participants (71.4% of sample), who had completed all the measures of the study and were considered valid for the analysis. There were no statistical differences between those included (with complete information) and excluded (due to missing data) into the two-step cluster analysis in the set of variables available. Three clusters emerged, with sample sizes of 52 (47.3%), 44 (40.0%) and 14 (12.7%) (the ratio of sizes comparing the largest to smallest cluster was 3.71). Goodness-of-fit was achieved, with fair average Silhouette Coefficient equal to 0.30.

Table 1 shows the sociodemographic composition of clusters, as well as the distribution of clinical variables associated with gambling. No variable differed significantly between the empirical clusters.

Table 2 shows the comparison of SOGS and DSM-IV total scores between clusters. Patients in cluster 3 had the highest intensity level of gambling, followed by cluster 2 and 1 whereas considering the individual items of the SOGS, there were no statistical differences comparing the three clusters. However, post-hoc analyses comparing specific groups and Cohen's d -coefficients achieved relevant differences for: playing slot machines, other bets, spent more than 300 Euros, returning to win back lost money, claiming winning when losing, gambling more than planned, being criticized, being unable to stop gambling, discussions with family or friends, borrowing money and not paying back, skipping school or work due to gambling, obtaining money from family or banks, using credit cards for gambling and obtaining money for gambling from property sales (1.7% vs. 20.8%).

Table 3 shows the comparison of general psychopathology (SCL-90-R) and personality traits (TCI-R) mean scores between clusters. ANOVA tests obtained significant mean differences between clusters for all the scales, except for TCI-R novelty seeking, reward dependence and persistence. Post-hoc comparisons showed that the three clusters differed in all the measures (except for the TCI-R novelty seeking, reward dependence and persistence scales), with the highest mean scores for patients in cluster 3 followed by cluster 2. Effect sizes were good to very good (Cohen's d clearly over 0.50), except for the three TCI-R scores that did not achieve statistical significant differences.

4. Discussion

The heterogeneity of PG has led to several attempts to establish different subtypes within the spectrum of the disorder. This study found evidence for three different clusters of young people, aged 17 to 25 years attending outpatient PG treatment.

The sociodemographic features were equally distributed between clusters, with the exception of educational level. Lack of education was associated with greater PG severity. The first cluster (or Type I) was composed of cases with high educational level (secondary or university studies) and who reported fewer negative consequences of gambling according to the questions in the SOGS (spending more money gambling, returning to win back lost money, gambling more than planned, being criticized due to the gambling addiction, being unable to stop gambling, borrowing money and not paying back, skipping school or work due to gambling, using credit cards to gamble and obtaining money destined to gambling from property sales). This cluster also achieved the lowest mean scores in the severity of the disorder (SOGS- Total score and DSM-IV-Total score), lower levels of general psychopathology (assessed with the SCL-90-R questionnaire) and healthier personality traits (lower scores on Harm Avoidance and Self-Transcendence and higher on Self-Directedness and Cooperativeness). This subtype, which we named High General Functioning (Type I), coincides with the “Behaviorally Conditioned Problem Gamblers” Pathway 1 type among adult PGs, as described by Blaszczynski & Nower

[8]. Our High General Functioning group (Type I) was the healthiest group compared to the other two in terms of psychopathology. If anxiety or depressive symptoms were present among these patients, they would be treated quickly once treatment is initiated. Type I also coincides with cluster I described in a study by Lesieur [29] named “normal problem gambler”: a group that presented low levels of psychopathology, impulsivity, depression, and anxiety, amongst other clinical characteristics. Likewise, our Type I resembles Class 2 in the study by Gupta et al. [12] conducted among adolescents.

The second cluster (Type II) that we call Depressive Type was characterized by major emotional distress, shame, immaturity, hostility and negative feelings. This cluster resembles the Pathway 2 type among adults described by Blaszczynski & Nower [8] as “Emotionally Vulnerable Problem Gamblers”, and would include those PGs that have suffered premorbid states of anxiety and depression, as well as significant deficits in coping strategies. This type feels the need to regulate and modulate their negative emotional states through gambling behavior. Similarly, our group partially coincides with the cluster II identified by Lesieur [29], which he labeled as “moderately-impulsive action seeker” patients with moderate levels of psychopathology and impulsivity, as well as those with more severe gambling behavior. However, Lesieur’s cluster also includes individuals with early age of onset and high levels of search for excitement.

Our Type II coincides with those called Class 4 and 5 in the study of Gupta et al. [12] among adolescents. While Class 4 is associated with depression, suicidal tendencies, childhood abuse and family conflicts, Class 5 is strictly associated with depressive symptoms (which the authors suggest as a unique subtype in adolescents). According to the Cloninger et al. [14] model of personality, three character dimensions (Self-directedness, Cooperativeness and Self-Transcendence) can be subsyndromic indicators of depressive or psychotic episodes. In our Type II, we observe low scores in Self-Directedness and Cooperativeness, but moderate scores in Self-Transcendence. This profile would fit the melancholy character, described by Cloninger et al. [14], in which immature traits, emotional reactivity, selfishness, competition and rivalry feelings toward others are present. Basically, the affective state of these individuals would be negative, rarely experiencing positive emotions. In these cases, and as suggested by Blaszczynski & Nower [8], gambling would be a maladaptive strategy to escape from and/or modulate these symptoms and negative emotions.

We identified a third subgroup, which we called Disorganized Type, or Type III, with the most severe psychopathological profile, including schizotypal traits, and most severe gambling behavior (assessed by the SOGS and DSM-IV questionnaires). Consistent with Cloninger et al. [14], these subjects, with low scores in Self-Directedness and Cooperativeness and high scores in Self-Transcendence, could be considered as disorganized or schizotypal. They would appear suspicious, illogical and immature, prone to imagination and fantasy, as well as demonstrating peculiar and unconventional behavior. In addition, this configuration is associated with the temperament traits Novelty Seeking and Harm Avoidance (as in the case of this subtype), as well as borderline personalities (explosive and overly-sensitive). This type

coincides with the Pathway 3 among adults described by Blaszczynski & Nower [8] as "Antisocial and Impulsivist Problem Gamblers": a subgroup with major alterations in psychopathology, major psychosocial interference due to their gambling behavior and more dysfunctional personality profiles. To these authors, this subgroup would represent an etiology of the disorder strongly associated with both neurobiological and psychological risk factors.

Both the subgroup described in the explanatory model of Blaszczynski & Nower [8] and our Type III, are equivalent with the Class 1 in the study by Gupta et al. [12]. Furthermore, Goldstein et al. [13] found a subgroup of adolescent gamblers with many consequences of gambling that was associated with substance abuse, delinquency and violence, which resembles the "Antisocial Impulsivist"/ Pathway 3 in the model by Blaszczynski & Nower [8] and our Type III which scored high on Novelty seeking. One may speculate that some of the adolescents in the high consequence group in the study by Goldstein et al. [13] represent our Type III at a younger age.

Lesieur's [29] cluster 3, composed of "impulsive escape seekers", only partially coincides with our results. In this group, Lesieur [29] included individuals with the most elevated levels of impulsivity and psychopathology, although also those with later age of onset and with low levels

of excitement seeking. In other words, this group would better meet the Type II and Type III in our study and the clusters 2 and 3 obtained by Blaszczynski and Nower [8]. However, it is important to highlight that Lesieur's [29] results were based on an inpatient sample, with elevated levels of severity, which could explain the partial concordance with the subtypes observed in other studies.

The impact that the identification of homogeneous subgroups in PG has in the design of therapeutic approaches is crucial. Although studies exploring the treatment response suggest that this disorder can be successfully treated [30], there are high rates of dropouts and relapses throughout the rehabilitation process of these patients [11,31]. Studies about the response to treatment programs show that mood disorders or substance dependence [32] and dysfunctional personality traits such as Sensation-seeking traits [33] Harm Avoidance and Self-Directedness [34] are associated with poor response to treatment. Based on research considering the subgroups, including our present findings, we might consider that the Type II and III patients would be those at risk for a more torpid and complex response to treatment. Therefore, including techniques and strategies for specific treatment of cognitive strategies for inhibitory control or regulation of negative emotions, among other aspects, could be crucial to improve the results of treatment programs. It is of great scientific interest to further investigate the effectiveness of new tools aimed at treating the underlying neurocognitive PG processes such as cognitive remediation [35], serious games [36,37] or mindfulness training based treatments [38].

4.1. Limitations

The first limitation concerns generalization: the results of this study must only be extrapolated to populations of young male gamblers who seek treatment due to problems related to gambling. Another limitation is related to the sample size: the small number of participants attenuated the power of the cluster analysis to detect more differences for the set of variables. Finally, although our main objective was to obtain an early identification of the subtypes of pathological gamblers, the average age of the sample was 22 years. The facts that gambling is illegal under the age of 18 in our country and that only the most severe cases seek treatment [39] are factors that may be related to the average age of the subjects studied. Several studies have shown that from the start of gambling behavior to loss of control it usually takes 6–7 years [5,40,41] which could also justify that although gambling starts earlier for most people (in some cases younger than 18), they do not seek treatment until they are older (e.g. 20–22 years of age).

4.2. Implications

To our knowledge, this is the first study examining subgroups of PG in a treatment-seeking sample of youth, so the findings have high relevance from a clinical point of view. Overall, the results of this work indicate that PG constitutes a complex disorder with heterogeneous clusters even among adolescents and young adults. Identifying the specific earliest manifestations of this problem is essential in order to develop adequate therapeutic programs and to prevent the disorder from progressing to the most severe stages. Finally, research should test for the empirical clusters that emerged in this study with larger and more

diverse samples particularly in relation to gender and response to treatment. Future research on PG types should also consider examining biological, neuropsychological, biochemical and genetic variables. The goal is to better understand this disorder and provide more effective assessment and treatment.

References

- [1] American Psychiatric Association, APA. Diagnostic and statistical manual of mental disorders (4th ed. revised). Washington, DC: American Psychiatric Association; 2000.
- [2] Chiu J, Storm L. Personality, perceived luck and gambling attitudes as predictors of gambling involvement. *J Gambl Stud* 2010;26:205-27.
- [3] Blinn-Pike L, Worthy SL, Jonkman JN. Adolescent gambling. A review of an emerging field of research. *J Adolesc Health* 2010;47:223-36.
- [4] Hardoon KK, Gupta R, Derevensky JL. Psychosocial variables associated with adolescent gambling. *Psychol Addict Behav* 2004; 18:170-9.
- [5] Jiménez-Murcia S, Álvarez-Moya E, Stinchfield R, Fernández-Aranda F, Granero R, Aymamí N, et al. Age of onset in pathological gambling: clinical, therapeutic and personality correlates. *J Gambl Stud* 2010;26(2):235-48, <http://dx.doi.org/10.1007/s10899-009-9175-3>.
- [6] Raylu N, Oei TPS. Pathological gambling: a comprehensive review. *Clin Psychol Rev* 2002;22:1009-61.
- [7] Álvarez-Moya E, Jimenez-Murcia S, Aymami MN, Gomez-Peña M, Granero R, Santamaria JJ, et al. Subtyping study of a male pathological gambling sample. *Can J Psychiatry* 2010;55(8):498-506.
- [8] Blaszczynski A, Nower L. A pathways model of problem and pathological gambling. *Addiction* 2002;97(5):487-99.
- [9] Cunningham-Williams RM, Hong SI. A latent class analysis (LCA) of problem gambling among a sample of community-recruited gamblers. *J Nerv Ment Dis* 2007;195:939-47.
- [10] von Ranson KM, Wallace LM, Holub A, Hodgins DC. Eating Disorders, Substance Use Disorders, and Impulsiveness among Disordered Gamblers in a Community Sample. *Eur Eat Disorders Rev* 2013;21:148-54, <http://dx.doi.org/10.1002/erv.2207>.
- [11] Ledgerwood DM, Petry NM. Psychological experience of gambling and subtypes of pathological gamblers. *Psychiatry Res* 2006;144: 17-27.
- [12] Gupta R, Nower L, Derevensky JL, Blaszczynski A, Faregh N, Temcheff C. Problem gambling in adolescents: an examination of the pathways model. *J Gambl Stud* 2012, <http://dx.doi.org/10.1007/s10899-012-9322-0>.
- [13] Goldstein AB, Faulkner B, Cunningham RM, Zimmerman MA, Chermack S, Walton MA. A latent class analysis of adolescent gambling: application of resilience theory. *Int J Ment Health Addiction* 2013;11:13-30.
- [14] Cloninger CR, Svrakic DM, Przybeck TR. A psychobiological model of temperament and character. *Arch Gen Psychiatry* 1993;50(12): 975-90.
- [15] Stinchfield R. Reliability, validity, and classification accuracy of a measure of DSM-IV diagnostic criteria for pathological gambling. *Am J Psychiatry* 2003;160:180-2.
- [16] Lesieur HR, Blume SB. The South Oaks Gambling Screen (SOGS): a new instrument for the identification of pathological gamblers. *Am J Psychiatry* 1987;144(9):1184-8.
- [17] Echeburúa E, Báez C, Fernández J, Páez D. Cuestionario de juego patológico de South Oaks (SOGS): Validación española (South Oaks Gambling Screen (SOGS): Spanish validation). *Análisis de Modifica- ción de Conducta* 1994;20:769-91.
- [18] Alessi SM, Petry NM. Pathological gambling severity is associated with impulsivity in a delay discounting procedure. *Behav Process* 2003;64(3):345-54.

- [19] Stinchfield R. Reliability, validity, and classification accuracy of the South Oaks Gambling Screen (SOGS). *Addict Behav* 2002; 27(1):1-19.
- [20] Strong DR, Daughters SB, Lejuez CW, Breen RB. Using the Rasch model to develop a revised Gambling Attitudes and Beliefs Scale (GABS) for use with male college student gamblers. *Subst Use Misuse* 2004;39(6):1013-24.
- [21] Jimenez-Murcia S, Stinchfield R, Alvarez-Moya E, Jaurieta N, Bueno B, Granero R, et al. Reliability, validity, and classification accuracy of a Spanish translation of a measure of DSM-IV diagnostic criteria for pathological gambling. *J Gambl Stud* 2009;25(1):93-104.
- [22] Derogatis LR. SCL-90-R. Cuestionario de 90 síntomas. [SCL-90-R. 90-Symptoms Questionnaire]. Madrid: TEA; 1994.
- [23] Cloninger CR. The Temperament and Character Inventory-Revised. St. Louis, MO: Center for Psychobiology of Personality, Washington University; 1999.
- [24] Gutierrez-Zotes JA, Bayon C, Montserrat C, Valero J, Labad A, Cloninger CR, et al. Temperament and Character Inventory Revised (TCI-R). Standardization and normative data in a general population sample. *Actas Esp Psiquiatr* 2004;32(1):8-15.
- [25] Jiménez-Murcia S, Bove FI, Vergé B, Álvarez-Moya E, Granero R, Penelo E, et al. Cognitive-behavioral therapy for pathological gambling in Parkinson's disease: a pilot controlled study. *Eur Addict Res* 2012;18(6):265-74, <http://dx.doi.org/10.1159/000337442>.
- [26] Jiménez-Murcia S, Aymamí MN, Gómez-Peña M, Álvarez-Moya EM, Vallejo J. *Protocols de tractament cognitivoconductual pel joc patològic i d'altres addiccions no tòxiques (Guidelines of cognitive- behavioral treatment of pathological gambling and other non-toxic addictions)*. Barcelona, Spain: Hospital Universitari de Bellvitge, Departament de Salut, Generalitat de Catalunya; 2006.
- [27] Fraley C, Raftery AE. How many clusters? Which clustering method? Answers via model-based cluster analysis. *The Computer Journal* 1998;41(8):578-88.
- [28] Rousseeuw PJ. Silhouettes: a graphical aid to the interpretation and validation of cluster analysis. *J Comput Appl Math* 1998;20:53-65.
- [29] Lesieur HR. Cluster analysis of types of inpatient pathological gamblers. *Dissertation Abstracts International: Section B: The Sciences and Engineering* 2001; 62: 4-B.
- [30] Gooding P, Tarrier N. A systematic review and meta-analysis of cognitive-behavioral interventions to reduce problem gambling: hedging our bets? *Behav Res Ther* 2009;47(7):592-607.
- [31] Melville KM, Casey LM, Kavanagh DJ. Psychological treatment dropout among pathological gamblers. *Clin Psychol Rev* 2007;27(8): 944-58.
- [32] Hodgins DC, el-Guebaly N. The influence of substance dependence and mood disorders on outcome from pathological gambling: five-year follow-up. *J Gambl Stud* 2010;26(1):117-27.
- [33] Smith D, Harvey P, Battersby M, Pols R, Oakes J, Baigent M. Treatment outcomes and predictors of drop out for problem gamblers in South Australia: a cohort study. *Aust N Z J Psychiatry* 2010;44(10): 911-20.
- [34] Nordin C, Nylander PO. Temperament and character in pathological gambling. *J Gambl Stud* 2007;23(2):113-20.
- [35] Pedrero-Perez EJ, Rojo-Mota G, de Leon JM Ruiz-Sanchez, Llanero- Luque M, Puerta-Garcia C. Rehabilitación cognitiva en el tratamiento de las adicciones (Cognitive remediation in addictions treatment). *Rev Neurol* 2011;52(3):163-72.
- [36] Fernández-Aranda F, Jiménez-Murcia S, Santamaría JJ, Gunnard K, Soto A, Kalapanidas E, et al. Video games as a complementary therapy tool in mental disorders: Playmancer, a European multicentre study. *J Ment Health* 2012;21(4):364-74.
- [37] Jiménez-Murcia S, Fernández-Aranda F, Kalapanidas E, Konstantas D, Ganchev T, Kocsis O, et al.

Playmancer project: a serious videogame as an additional therapy tool for eating and impulse control disorders. *Stud Health Technol Inform* 2009;144:163-6.

- [38] Witkiewitz K, Lustyk MK, Bowen S. Retraining the addicted brain: a review of hypothesized neurobiological mechanisms of mindfulness- based relapse prevention. *Psychol Addict Behav* 2012.
- [39] Slutske WS. Natural recovery and treatment-seeking in pathological gambling: results of two U.S. national surveys. *Am J Psychiatry* 2006;163(2):297-302.
- [40] Breen RB, Zimmerman M. Rapid onset of pathological gambling in machine gamblers. *J Gambl Stud* 2002;18(1):31-43.
- [41] Tavares H, Zilberman ML, Beites FJ, Gentil V. Gender differences in gambling progression. *J Gambl Stud* 2001;17(2):151-9.

Table 1

	Total sample (<i>n</i> = 154)	Cluster 1 (<i>n</i> = 52)	Cluster 2 (<i>n</i> = 44)	Cluster 3 (<i>n</i> = 14)	<i>p</i>
Socio demographic and socio-economic characteristics					
Age (years); <i>mean (SD)</i>	22.4 (2.2)	22.6 (2.1)	23.0 (2.3)	22.6 (1.8)	.692
Gender: male; %	94.2	100	93.2	100	.374
Employment status (employed); %	61.3	73.1	63.6	57.1	.625
Educational level; % <i>Primary or less</i>	56.4	48.1	72.7	78.6	.252
Civil status; % <i>Without partner</i>	89.0%	94.2	86.0	78.6	.478
Own incomes (euros); <i>mean (SD)</i>	704 (603)	907 (579)	751 (524)	771 (841)	.625
Family incomes (euros); <i>mean (SD)</i>	2482 (1560)	2613 (1664)	2322 (1643)	3067 (1679)	.625
Smoker (yes); %	75.2	74.4	86.5	69.2	.597
Clinical history					
Number of problematic games; <i>mean (SD)</i>	1.5 (0.8)	1.33 (0.57)	1.53 (.86)	2.00 (1.15)	.252
Evolution (years); <i>mean (SD)</i>	2.8 (2.5)	2.55 (2.04)	3.32 (2.80)	2.54 (2.23)	.597
Age of onset; <i>mean (SD)</i>	20.0 (2.6)	20.42 (2.12)	19.76 (2.86)	20.31 (2.27)	.625
Maximum bets (euros); <i>mean (SD)</i>	741 (1107)	688 (1394)	659 (798)	1454 (1148)	.374
Mean bets (euros); <i>mean (SD)</i>	86 (143)	80 (110)	82 (125)	131 (286)	.646
Cumulate debts (euros); <i>mean (SD)</i>	2842 (5723)	3144 (6931)	2635 (5157)	2620 (2721)	.942

Table 2

SOGS	Prevalence (%)			<i>p</i>	Logistic regression/ANOVA					
	Cluster 1 (<i>n</i> = 52)	Cluster 2 (<i>n</i> = 44)	Cluster 3 (<i>n</i> = 14)		Contrasts: OR/MD (Cohens' <i>d</i>) ^a					
					CI2 vs CI1		CI3 vs CI1		CI3 vs CI2	
1a-Playing cards	36.54%	38.64%	57.14%	.381	1.094	(0.04)	2.31	(0.42)	2.12	(0.38)
1b-Playing horse races	1.92%	0.00%	0.00%	.998	—	(0.20)	—	(0.20)	1.00	(0.00)
1c-Playing sports	3.85%	6.98%	0.00%	.898	1.875	(0.14)	—	(0.28)	—	(0.39)
1d-Playing numbers-lotteries	75.00%	84.09%	78.57%	.587	1.762	(0.23)	1.22	(0.08)	0.69	(0.14)
1e-Playing casino	34.62%	34.09%	50.00%	.545	0.977	(0.01)	1.89	(0.32)	1.93	(0.33)
1f-Playing bingo	50.00%	68.18%	50.00%	.553	2.143	(0.38)	1.00	(0.00)	0.47	(0.38)
1g-Stock market	5.77%	0.00%	7.14%	.733	—	(0.35)	1.26	(0.06)	—	(0.39)
1h-Playing slot machines	84.31%	100.00%	100.00%	.998	—	(0.61)	—	(0.61)	1.00	(0.00)
1i-Other bets	17.31%	22.73%	57.14%	.064	1.405	(0.14)	6.37*	(0.90)	4.52*	(0.75)
2-Amount money spent: ≥ 300 euros	51.92%	54.55%	92.86%	.083	1.111	(0.05)	12.05*	(1.03)	10.87*	(0.97)
3-Family antecedents of gambling	26.92%	38.64%	42.86%	.331	1.709	(0.25)	2.04	(0.34)	1.19	(0.09)
4-Going back to win back lost money	86.54%	100.00%	100.00%	.998	—	(0.56)	—	(0.56)	1.00	(0.00)
5-Claimed winning when loosing	34.62%	59.09%	71.43%	.064	2.728*	(0.51)	4.72*	(0.79)	1.73	(0.26)
6-Problem recognition	90.38%	97.73%	100.00%	.248	4.574	(0.31)	1.00	(0.46)	1.00	(0.22)
7-Gambling more than planned	82.69%	97.73%	100.00%	.085	9.000*	(0.52)	—	(0.65)	—	(0.22)
8-Being criticized	53.85%	84.09%	85.71%	.064	4.531*	(0.69)	5.15*	(0.74)	1.14	(0.05)
9-Feeling guilty	86.54%	97.73%	100.00%	.144	6.689	(0.42)	—	(0.56)	—	(0.22)
10-Unable to stop gambling	76.47%	93.18%	100.00%	.064	4.205*	(0.50)	—	(0.78)	—	(0.38)
11-Hiding signs of gambling	64.71%	77.27%	78.57%	.331	1.855	(0.28)	2.00	(0.31)	1.08	(0.03)
12-Discussions with family/friends	86.54%	88.64%	100.00%	.360	1.213	(0.06)	—	(0.56)	—	(0.51)
13-Discussions and fights	62.75%	65.12%	78.57%	.490	1.108	(0.05)	2.18	(0.35)	1.96	(0.30)
14-Borrowing money, not paying back	44.23%	59.09%	92.86%	.064	1.821	(0.30)	16.39*	(1.23)	9.01*	(0.86)
15-Skipping work due to gambling	42.31%	43.18%	100.00%	.064	1.036	(0.02)	—	(1.65)	—	(1.62)
16a-Money from home	56.86%	63.64%	69.23%	.497	1.328	(0.14)	1.71	(0.26)	1.29	(0.12)
16b-Money from couple	19.61%	34.09%	30.77%	.331	2.121	(0.33)	1.82	(0.26)	0.86	(0.07)
16c-Money from family	25.00%	40.91%	61.54%	.064	2.077	(0.34)	4.81*	(0.79)	2.31	(0.42)
16d-Money from banks	25.00%	47.73%	61.54%	.064	2.739*	(0.50)	4.81*	(0.79)	1.73	(0.28)
16e-Credit cards	26.92%	45.45%	69.23%	.064	2.262	(0.39)	6.10*	(0.93)	2.70	(0.50)
16f-Money from money lender	3.92%	4.55%	0.00%	.733	1.167	(0.03)	—	(0.29)	—	(0.31)
16 g-Money from financial papers	1.96%	0.00%	0.00%	.998	—	(0.20)	—	(0.20)	—	(0.00)
16 h-Money from property sales	1.96%	20.45%	7.69%	.193	12.86*	(0.61)	4.17	(0.27)	0.32	(0.37)
16i-Money from falsified checks	0.00%	2.27%	0.00%	.701	—	(0.22)	1.00	(0.00)	—	(0.22)
16j-Money from credit account casino	1.92%	2.27%	7.69%	.501	1.186	(0.02)	4.26	(0.27)	3.58	(0.25)
SOGS: total score; <i>mean (SD)</i>	8.58 (3.7)	11.14 (2.8)	12.71 (1.8)	<.001	2.56*	(0.78)	4.14*	(1.41)	1.58	(0.68)
DSM-IV: total criteria; <i>mean (SD)</i>	6.27 (2.4)	7.98 (1.4)	8.93 (1.1)	<.001	1.71*	(0.80)	2.66*	(1.42)	0.95	(0.63)

Table 3

	Mean; standard deviation						Comparison between clusters: contrasts (Cohen's d)						
	Cluster 1 (n = 52)		Cluster 2 (n = 44)		Cluster 3 (n = 14)		p	C11 vs C12		C11 vs C13		C12 vs C13	
SCL-90-R scores													
Somatization	0.34;	0.36	0.95;	0.51	2.18;	0.87	<.001	-0.61*	(1.38)	-1.84*	(2.76)	-1.23*	(1.72)
Obsessive/compulsive	0.52;	0.38	1.23;	0.50	2.43;	0.69	<.001	-0.71*	(1.60)	-1.91*	(3.43)	-1.20*	(1.99)
Interpersonal sensitivity	0.30;	0.28	1.10;	0.52	2.19;	0.87	<.001	-0.80*	(1.92)	-1.89*	(2.92)	-1.09*	(1.52)
Depressive	0.58;	0.44	1.59;	0.58	2.45;	0.71	<.001	-1.01*	(1.96)	-1.87*	(3.17)	-0.86*	(1.33)
Anxiety	0.36;	0.36	1.13;	0.54	2.43;	0.54	<.001	-0.78*	(1.68)	-2.07*	(4.51)	-1.29*	(2.41)
Hostility	0.44;	0.36	1.22;	0.71	2.38;	1.03	<.001	-0.79*	(1.39)	-1.95*	(2.51)	-1.16*	(1.31)
Phobic anxiety	0.15;	0.22	0.46;	0.49	1.56;	0.92	<.001	-0.31*	(0.82)	-1.41*	(2.11)	-1.10*	(1.49)
Paranoid Ideation	0.34;	0.34	1.02;	0.59	2.06;	0.67	<.001	-0.68*	(1.41)	-1.72*	(3.24)	-1.04*	(1.65)
Psychotic	0.27;	0.31	0.96;	0.43	1.98;	0.47	<.001	-0.69*	(1.84)	-1.71*	(4.30)	-1.02*	(2.26)
GSI score	0.40;	0.24	1.13;	0.32	2.22;	0.43	<.001	-0.73*	(2.58)	-1.82*	(5.23)	-1.09*	(2.88)
PST score	24.6;	13.8	53.5;	12.5	71.6;	8.55	<.001	-28.9*	(2.19)	-47.0*	(4.09)	-18.03*	(1.69)
PSDI score	1.41;	.29	1.90;	0.31	2.80;	0.46	<.001	-0.49*	(1.63)	-1.39*	(3.61)	-0.90*	(2.29)
TCI-R scores													
Novelty seeking	112.2;	13.1	115.0;	11.7	117.0;	13.1	.362	-2.78	(0.22)	-4.81	(0.37)	-2.02	(0.16)
Harm avoidance	92.4;	13.8	101.6;	16.1	106.3;	7.0	.001	-9.15*	(0.61)	-13.8*	(1.27)	-4.69	(0.38)
Reward dependence	100.7;	15.7	94.5;	15.2	99.6;	9.8	.137	6.75	(0.40)	1.07	(0.08)	-5.10	(0.40)
Persistence	110.4;	19.2	108.6;	22.8	113.3;	15.8	.749	1.75	(0.08)	-2.90	(0.16)	-4.65	(0.24)
Self-directedness	139.1;	18.0	119.5;	16.0	107.4;	14.4	<.001	19.6*	(1.15)	31.7*	(1.94)	12.12*	(0.79)
Cooperativeness	132.2;	18.9	124.4;	17.5	111.9;	16.1	.001	7.79	(0.43)	20.3*	(1.16)	12.5*	(0.74)
Self-Transcendence	55.7;	14.1	65.5;	14.2	75.4;	11.7	<.001	-9.74*	(0.69)	-19.7*	(1.52)	-9.90*	(0.76)