

EXPLICIT UPPER AND LOWER BOUNDS FOR THE TRAVELING WAVE SOLUTIONS OF FISHER-KOLMOGOROV TYPE EQUATIONS

A. GASULL*, H. GIACOMINI, AND J. TORREGROSA*

ABSTRACT. It is well-known that the existence of traveling wave solutions for reaction-diffusion partial differential equations can be proved by showing the existence of certain heteroclinic orbits for related autonomous planar differential equations. We introduce a method for finding explicit upper and lower bounds of these heteroclinic orbits. In particular, for the classical Fisher-Kolmogorov equation we give rational upper and lower bounds which allow to locate these solutions analytically and with very high accuracy.

1. INTRODUCTION AND MAIN RESULTS

Consider the adimensionalized reaction-diffusion partial differential equation

$$u_t = u_{xx} + f(u), \quad (1)$$

of Fisher-Kolmogorov type, where $f(u)$ is a smooth function satisfying certain hypotheses.

The usual Fisher-Kolmogorov equation corresponds to $f(u) = u(1-u)$ and models the spreading of biological populations, see [2, 5]. Other well-known cases are the Newell-Whitehead-Segel equation, $f(u) = u(1-u^2)$, for describing Rayleigh-Benard convection, see [6, 9], and the Zeldovich equation, $f(u) = u(1-u)(u-\alpha)$ with $0 < \alpha < 1$, that appears in combustion theory, see [11]. See also [4, 7, 8, 10].

It is known that the traveling wave solutions $u = u(x-ct)$ of (1), satisfying

$$\lim_{s \rightarrow -\infty} u(s) = 0 \quad \text{and} \quad \lim_{s \rightarrow \infty} u(s) = 1,$$

appear when $u(s)$ is a special solution of the second order equation

$$\ddot{u} + c\dot{u} + f(u) = 0,$$

where the dot indicates the derivative with respect to s . This solution can be seen as the heteroclinic orbit H_c of the planar system


$$\begin{cases} \dot{u} &= v, \\ \dot{v} &= -cv - f(u), \end{cases} \quad (2)$$

that connects the origin with the saddle point at $(u, v) = (1, 0)$, see Figure 1. For instance for the classical Fisher-Kolmogorov equation it exists only when $c \geq 2$.

2010 *Mathematics Subject Classification.* 34C37, 35C07, 37C29.

Key words and phrases. Traveling wave, heteroclinic orbit, Fisher-Kolmogorov equation, reaction-diffusion partial differential equation, invariant manifold.

*The first and third authors are partially supported by the MCYT/FEDER grant number MTM2008-03437 by the CIRIT grant number 2009SGR410.

FIGURE 1. Phase portrait showing the unstable separatrix H_c .

The goal of this paper is to give analytic upper and lower bounds of the heteroclinic orbit H_c as well as of their time parametrization. We will approach to this question with similar tools to those introduced in [3]. A key point consists in using the local behavior of the separatrices of the critical points to guess global algebraic bounds for the actual orbits.

First we prove a general result for system (2). It is illustrated in Figure 2.

Theorem 1.1. *Consider system (2), with f satisfying $f(0) = f(1) = 0$, $f'(0) > 0$, $f'(1) < 0$, $f''(u) < 0$ for all $u \in \mathbb{R}$ and $c \geq 2\sqrt{f'(0)}$. Let H_c be its heteroclinic orbit and define*

$$\underline{\lambda} := \frac{c - \sqrt{c^2 - 4f'(0)}}{2f'(0)} \quad \text{and} \quad \bar{\lambda} := \frac{c - \sqrt{c^2 - 4f'(1)}}{2f'(1)}.$$

Then H_c can be parametrized as $H_c = \{(u, h_c(u)), u \in [0, 1]\}$ and for all $u \in (0, 1)$,


$$-\underline{\lambda}f(u) < h_c(u) < -\bar{\lambda}f(u) < 0.$$

Moreover, if $(u_c(s), v_c(s))$ is the parametrization of H_c such that $u_c(0) = 1/2$ it holds that

$$u_c(s) \in \langle z_{\underline{\lambda}}(s), z_{\bar{\lambda}}(s) \rangle,$$

where $\langle a, b \rangle$ denotes the smallest closed interval containing a and b and $u = z_{\lambda}(s)$ is the solution of the Cauchy problem

$$\begin{cases} \frac{du}{ds} = -\lambda f(1-u), \\ u(0) = 1/2. \end{cases}$$

FIGURE 2. (i) Upper and lower bounds of the unstable separatrix H_c (dotted line). (ii) Bounds for the time parametrization of H_c .

For instance when we particularize the above theorem to the classical Fisher-Kolmogorov system we obtain that

$$u_c(s) \in \left\langle \frac{1}{1 + e^{\lambda s}}, \frac{1}{1 + e^{\bar{\lambda} s}} \right\rangle. \quad (3)$$

The results obtained for system (2) can be improved when we study this case. We prove:

Theorem 1.2. *Let $H_c = \{(u, h^c(u)), u \in [0, 1]\}$ be a parametrization of the heteroclinic solution of system (2) when $f(u) = u(1 - u)$. For $u \in (0, 1)$ and $c \geq 2$ it holds that*

$$h_2^c(u) < h_3^c(u) < \dots < h_{100}^c(u) < h^c(u) < R_{10}^c(u) < R_9^c(u) < \dots < R_1^c(u),$$

where $h_n^c(u)$ is the Taylor polynomial of degree n of the unstable separatrix of the saddle point $(1, 0)$ of the system and each R_m^c is a rational function whose numerator has degree $m + 2$ and its denominator degree m , constructed from the Padé approximants of $v = h_{2m+2}^c(u)$. See Figure 3.




FIGURE 3. Upper and lower bounds of the separatrix H_c of the Fisher-Kolmogorov equation, which is plotted as a dotted line.

For a given value of c it is also possible to study the maximum distance between two of the above functions. For instance when $c = 99/100$, we can prove that $R_9^{99/100}(x) - h_{20}^{99/100}(x) < 2 \times 10^{-19}$. It is also possible to go further in the computations. For example we get that

$$h_{82}^{99/100}(u) < h^{99/100}(u) < R_{40}^{99/100}(u) \quad \text{and} \quad R_{40}^{99/100}(u) - h_{82}^{99/100}(u) < 2 \times 10^{-47},$$

for all $u \in [0, 1]$ and the maximum error is at $u = 0$. For bigger values of c it is needed to compute approximations of higher degree to arrive to similar bounds of the error.

The s -parametrization of H_c can also be obtained with more accuracy for the classical Fisher-Kolmogorov case. We only present here a first result. Sharper approximations are detailed in Section 4.

Theorem 1.3. *Let $(u^c(s), v^c(s))$ be the time-parametrization of the heteroclinic orbit H^c of system (2) with $f(u) = u(1 - u)$ and such that $u^c(0) = 1/2$. Define*

$$U^c(s) = \frac{3 + 2\sqrt{2}}{(1 + \sqrt{2} + e^{(\sqrt{c^2+4}-c)s/2})^2}.$$

Then:

- (i) When $2 \leq c < 5/\sqrt{6}$ it holds that $\text{sgn}(u^c(s) - U^c(s)) = -\text{sgn}(s)$.

- (ii) When $c = 5/\sqrt{6}$ it holds that $u^c(s) = U^c(s)$.
- (iii) When $c > 5/\sqrt{6}$ it holds that $\text{sgn}(u^c(s) - U^c(s)) = \text{sgn}(s)$.

The above inequalities improve the bounds given in (3). Notice also that when $c = 5/\sqrt{6}$ we have obtained the exact expression of the traveling wave solution of the classical Fisher-Kolmogorov equation, $u_t = u_{xx} + u(u - 1)$,

$$u(x, t) = \frac{3 + 2\sqrt{2}}{\left(1 + \sqrt{2} + e^{\frac{1}{\sqrt{6}}\left(x - \frac{5}{\sqrt{6}}t\right)}\right)^2},$$

which coincides with the one given in [1]. The novelty of our result is that a similar expression gives a bound of the traveling wave for all the values of c .

The methods developed to study the classical Fisher-Kolmogorov equation can also be applied for the Newell-Whitehead-Segel and the Zeldovich equations.

2. PROOF OF THEOREM 1.1

For computational reasons it is more convenient to locate the saddle point of system (2) at the origin. So we introduce the new variables $x = 1 - u$, $y = v$ and $t = s$ and it writes as

$$\begin{cases} x' &= -y, \\ y' &= -cy + g(x), \end{cases} \quad (4)$$

where $g(x) = -f(1 - x)$ and the prime denotes derivative with respect to t . Observe that the variable x introduced above does not coincide with the one used in equation (1). Notice that g satisfies the following set of hypotheses

H : $g(0) = g(1) = 0$, $g'(0) < 0$, $g'(1) > 0$ and $g''(x) > 0$ for all $x \in \mathbb{R}$.

The above system has only two critical points $(0, 0)$ and $(1, 0)$. The origin is a saddle point with eigenvalues

$$\lambda_s^\pm = \frac{-c \pm \sqrt{c^2 - 4g'(0)}}{2} \quad (5)$$

and corresponding eigenvectors $(1, -\lambda_s^+)$ and $(1, -\lambda_s^-)$. Similarly, when $c^2 - 4g'(1) \geq 0$ the point $(1, 0)$ is an attracting node and its eigenvalues are

$$\lambda_n^\pm = \frac{-c \pm \sqrt{c^2 - 4g'(1)}}{2} < 0.$$

A sketch of the phase portrait of system (4) is given in Figure 4. There we can see the heteroclinic connection that we are interested to locate. Indeed it is given by ones of the branches of the unstable separatrix of the saddle point. In the new coordinates we will call it Γ_c .

Proposition 2.1. *Consider system (4), with g satisfying hypotheses **H** and $c \geq 2\sqrt{g'(1)}$. Set*

$$\underline{\lambda} := \frac{c - \sqrt{c^2 - 4g'(1)}}{2g'(1)} \quad \text{and} \quad \bar{\lambda} := \frac{c - \sqrt{c^2 - 4g'(0)}}{2g'(0)}.$$

Let Γ_c be its heteroclinic orbit. Then Γ_c can be parametrized as $\Gamma_c = \{(x, \gamma_c(x)), x \in [0, 1]\}$ and it holds for all $x \in (0, 1)$ that

$$\underline{\lambda}g(x) < \gamma_c(x) < \bar{\lambda}g(x) < 0.$$




FIGURE 4. Phase portrait showing the unstable separatrix Γ_c of system (4).

Proof. Consider the 1-parameter family of maps $G_\lambda(x, y) = y - \lambda g(x)$. We compute

$$\langle \nabla G_\lambda(x, y), (-y, -cy + g(x)) \rangle \Big|_{y=\lambda g(x)} = g(x)(1 - c\lambda + g'(x)\lambda^2) =: g(x)N_\lambda(x).$$

Since on $(0, 1)$ it holds that $g(x) < 0$, if we choose λ such that N_λ does not vanish on the same interval we will have that the corresponding curve $y = \lambda g(x)$ is without contact. Notice that $N'_\lambda(x) = \lambda^2 g''(x)$ and so for $\lambda \neq 0$, the hypotheses **H** imply that the function N_λ is increasing. Therefore:

- (a) If for some $\lambda > 0$, $N_\lambda(0) \geq 0$ then $N_\lambda(x) > 0$ for all $x > 0$.
- (b) If for some $\lambda > 0$, $N_\lambda(1) \leq 0$ then $N_\lambda(x) < 0$ for all $x < 1$.

The conditions $N_\lambda(j) = 0$ for $j = 0, 1$, write as $1 - c\lambda + g'(j)\lambda^2 = 0$. Their solutions are

$$\lambda_j^\pm = \frac{c \pm \sqrt{c^2 - 4g'(j)}}{2g'(j)}.$$

It is easy to prove that

$$\lambda_0^+ < 0 < \lambda_0^- =: \bar{\lambda} < \underline{\lambda} := \lambda_1^- \leq \lambda_1^+.$$

Therefore, taking $G_{\bar{\lambda}}(x, y) = 0$ and $G_{\underline{\lambda}}(x, y) = 0$ as an upper and lower boundary, respectively, we have constructed a subset of the strip $\{(x, y), 0 \leq x \leq 1\}$, that contains the heteroclinic orbit Γ_c , see Figure 5. Then, on it, $x' = -y > 0$, Γ_c can be parametrized as a function of x , say $y = \gamma_c(x)$, and the inequalities of the statement follow. \square




FIGURE 5. Upper and lower bounds of the unstable separatrix (dotted line) Γ_c .

Notice that $y = \bar{\lambda}g(x) = -\lambda_s^+ x + O(x^2)$ and therefore this curve is tangent to the unstable separatrix of the saddle point.

Using the above proposition we can also approach the parametrization of Γ_c with respect the actual time t . We need to introduce some new functions. Given the Cauchy problem

$$\begin{cases} \frac{dx}{dt} = -\lambda g(x), \\ x(0) = 1/2, \end{cases} \quad (6)$$

we will denote by $x = w_\lambda(t)$ its corresponding solution.

Theorem 2.2. *Under the same hypotheses and notations of Proposition 2.1, if $(x_c(t), y_c(t))$ is the parametrization of Γ_c such that $x_c(0) = 1/2$ it holds that*

$$x_c(t) \in \langle w_\lambda(t), w_{\bar{\lambda}}(t) \rangle,$$

where $w_\lambda(t)$ is defined in (6), see Figure 6.




FIGURE 6. Upper and lower bounds of the t -parametrization of Γ_c (dotted curve).

Proof. By Proposition 2.1 we know that

$$\underline{\lambda} g(x) < \gamma_c(x) < \bar{\lambda} g(x) < 0.$$

Recall that Γ_c is parametrized as $y = \gamma_c(x)$ and therefore since in system (4), $x'(t) = -y(t)$, it holds that $x'_c(t) = -\gamma_c(x_c(t))$. Therefore

$$-\underline{\lambda} g(x_c(t)) > x'_c(t) > -\bar{\lambda} g(x_c(t)) > 0.$$

Since $x_c(0) = w_{\bar{\lambda}}(0) = 1/2$, $x_c(t)$ satisfies the differential inequality

$$x'_c(t) > -\bar{\lambda} g(x_c(t))$$

and $w_{\bar{\lambda}}(t)$ the equality, it holds that $x_c(t) > w_{\bar{\lambda}}(t)$ for all $t > 0$, as we wanted to see. The other cases follow similarly. \square

Corollary 2.3. *If in system (4), $g(x) = x^m - x$, $2 \leq m \in \mathbb{N}$, then*

$$x_c(t) \in \left\langle \frac{1}{\sqrt[m-1]{1 + (2^{m-1} - 1)e^{-(m-1)\underline{\lambda}t}}}, \frac{1}{\sqrt[m-1]{1 + (2^{m-1} - 1)e^{-(m-1)\bar{\lambda}t}}} \right\rangle.$$

In particular for the Fisher-Kolmogorov case, $m = 2$,

$$x_c(t) \in \left\langle \frac{1}{1 + e^{-\underline{\lambda}t}}, \frac{1}{1 + e^{-\bar{\lambda}t}} \right\rangle. \quad (7)$$

Proof. It suffices to solve the Cauchy problem (6) and then apply Theorem 2.2. When $g(x) = x^m - x$ we obtain

$$w_\lambda(t) = \frac{1}{\sqrt[m-1]{1 + (2^{m-1} - 1)e^{-(m-1)\lambda t}}}$$

and so the result follows. \square

Theorem 1.1 is a reformulation for system (2) of the results of this section obtained for system (4).

3. SHARPER UPPER AND LOWER BOUNDS FOR Γ_c IN THE FISHER-KOLMOGOROV CASE

This section will be devoted to find sharper upper and lower bounds for Γ_c in the Fisher-Kolmogorov system (4) when $g(x) = x(x - 1)$. We will use dynamical tools inspired in [3]. One of the key points will be to find algebraic curves constructed by imposing that these curves coincide as much as possible with the unstable separatrix of the saddle point.

To avoid the appearance of square roots during the computations it is convenient to include a new parameter r in such a way that

$$c = \frac{1}{r} - r.$$

Then system (4) writes as

$$\begin{cases} x' &= -y, \\ y' &= \left(r - \frac{1}{r}\right)y + x(x - 1), \end{cases} \quad r \in (0, \sqrt{2} - 1]. \quad (8)$$

Notice that the condition on r implies that $c = r - 1/r \geq 2$. One advantage of introducing this new parameter is that the eigenvalues of the saddle are now $-1/r < 0 < r$. In the notation of the previous section $\lambda_s^+ = \bar{\lambda} = r$. Hence, if we denote $\Gamma^r := \Gamma_{1/r-r}$ the searched heteroclinic trajectory, from Proposition 2.1 we know that $y = rx(x - 1)$ is an upper bound for Γ^r .

First we need to know the local expansion of the unstable manifold of the saddle point.

Lemma 3.1. *The local unstable manifold of the origin of system (8) writes as the analytic function*

$$\begin{aligned} y = h^r(x) &= -rx + \frac{r}{2r^2 + 1}x^2 + \frac{2r^3}{(2r^2 + 1)^2(3r^2 + 1)}x^3 \\ &+ \frac{10r^5}{(2r^2 + 1)^3(3r^2 + 1)(4r^2 + 1)}x^4 \\ &+ \frac{12r^7(19r^2 + 6)}{(2r^2 + 1)^4(3r^2 + 1)^2(4r^2 + 1)(5r^2 + 1)}x^5 + O(x^5), \end{aligned}$$

and the subsequent terms can be determined recurrently and are positive for $r > 0$.

Proof. Let $y = h^r(x) = h(x)$ be the local expression of any of the separatrices of the saddle point, being h an analytic function at zero. Then $y' - h'(x)x'|_{y=h(x)} \equiv 0$, or equivalently,

$$\left(r - \frac{1}{r}\right)y + x(x-1) + h'(x)y \Big|_{y=h(x)} = \left(r - \frac{1}{r} + h'(x)\right)h(x) - x + x^2 \equiv 0. \quad (9)$$

Writing $h(x) = h_1x + h_2x^2 + \dots$ and plugging this expression in the above one we get that $h_1^2 + (r - 1/r)h_1 - 1 = 0$. So we choose $h_1 = -r$. Then the right hand identity in (9) writes as

$$\left(-\frac{1}{r} + 2h_2x + 3h_3x^2 + \dots\right)(-rx + h_2x^2 + h_3x^3 + \dots) - x + x^2 \equiv 0.$$

So $-2h_2rx^2 - h_2x^2/r + x^2 \equiv 0$, which implies $h_2 = r/(2r^2 + 1)$. In general, for $n > 2$ it holds that

$$-\frac{1}{r}h_n + 2h_2h_{n-1} + 3h_3h_{n-2} + \dots + (n-1)h_{n-1}h_2 - nh_nr = 0.$$

Therefore,

$$h_n = \frac{2h_2h_{n-1} + 3h_3h_{n-2} + \dots + (n-1)h_{n-1}h_2}{nr + 1/r}$$

and by induction $h_n > 0$, for $n > 2$, as we wanted to prove. \square




FIGURE 7. Upper and lower bounds of the separatrix Γ_c plotted as a dotted thin line. Here $2 \leq n \leq 100$ and $1 \leq m \leq 10$.

Proposition 3.2. *Let $\Gamma^r = \{(x, \gamma^r(x)), x \in [0, 1]\}$, be a parametrization of the heteroclinic solution of system (8). Then for all $x \in (0, 1)$ it holds that*

$$h_2^r(x) < h_3^r(x) < \dots < h_{99}^r(x) < h_{100}^r(x) < \gamma^r(x),$$

where $h_n^r(x)$ is the Taylor polynomial of degree n in powers of x of the function $h^r(x)$ defined in Lemma 3.1. See Figure 7.

Proof. We will give first the details for $n = 3$. It is evident that for $x > 0$ sufficiently small we have $h^r(x) > h_3^r(x)$. First we evaluate the polynomial $h_3^r(x)$ at $x = 1$. We obtain

$$h_3^r(1) = -\frac{2r^5(5 + 6r^2)}{(1 + 2r^2)^2(1 + 3r^2)}.$$

This quantity is negative for $r > 0$ and it is a monotonous decreasing function of r . Its value at $r = \sqrt{2} - 1$ is approximately -0.054 . Moreover, $h_3^r(0) = 0$, $h_3^r(x)'' > 0$ for $x > 0$ and $h_3^r(x) < 0$ for $x > 0$ and sufficiently small. Then we

conclude that $h_3^r(x) < 0$ for all $x \in (0, 1]$. We will show now that the flow of the vector field $Z^r(x, y) = (-y, (r - 1/r)y + x(x - 1))$ associated to system (8) crosses the curve $F_3^r(x, y) = y - h_3^r(x) = 0$ upwards. We compute

$$M_3^r(x) := \langle \nabla F_3^r(x, y), Z^r(x, y) \rangle|_{y=h_3^r(x)}.$$

We obtain

$$M_3^r(x) = \frac{10r^4x^4}{(1 + 2r^2)^3(1 + 3r^2)} + \frac{12r^6x^5}{(1 + 2r^2)^4(1 + 3r^2)^2}.$$

This expression is positive for $x > 0$.

All the other cases can be studied by using the same method. The key point is that all the monomials in the corresponding expression of $M_n^r(x)$ are positive for $x > 0$. \square

In Proposition 3.2, we have chosen to stop at the value 100 because the computation of the function M_{100}^r used in its proof together with the testing that all its coefficients are positive takes more than four hours of CPU time in our computer. In any case, for practical uses, it suffices to consider small values of n .

Remark 3.3. *It is clear that if we could prove that the radius of convergence of the series given in Lemma 3.1, when $r \in (0, \sqrt{2} - 1]$, is 1 we would have obtained an infinite monotonous sequence of polynomials tending to the actual separatrix. Unfortunately we have not succeeded in our attempts. In any case, computing an approximation of the radius of convergence, by using several hundreds of terms of the series, seems to show that the result is true.*

We give now upper bounds for Γ^r . Before to state the next proposition we introduce some definitions. Consider the rational function

$$R^r(x) = \frac{h_{22}^r(x)}{rx(x - 1)}.$$

We define the sequence of rational functions $R_n^r(x)$ as follows:

$$R_n^r(x) = rx(x - 1)P_{n,n}^r(x), \quad n = 1, \dots, 10,$$

where the $P_{n,n}^r(x)$ are the Padé approximants of order (n, n) of the function $R^r(x)$. For instance

$$R_1^r(x) = rx(x - 1) \frac{(2r^2 + 1)(3r^2 + 1) - 3r^2x}{(2r^2 + 1)(3r^2 + 1) - r^2(6r^2 + 5)x}.$$

Recall that the Padé approximants $P_{n,n}^r(x)$ are rational functions whose numerators and denominators are polynomials of degree n and their Taylor expansions in powers of x are the same that the Taylor expansion of $R^r(x)$ up to order $2n$. We write

$$R_n^r(x) = rx(x - 1) \frac{A_n^r(x)}{C_n^r(x)}, \quad \text{with } C_n^r(0) > 0.$$

The rational functions $R_n^r(x)$ vanish at $x = 0$ and $x = 1$ and their Taylor expansions coincide with the Taylor expansion of $h^r(x)$ up to order $2n$. As we will see in the proof of next proposition they are well defined for all $x \in [0, 1]$ because $C_n^r(x) \neq 0$.

Proposition 3.4. *Let $\Gamma^r = \{(x, \gamma^r(x)), x \in [0, 1]\}$, be a parametrization of the heteroclinic solution of system (8). For $x \in (0, 1)$ and $r \in (0, \sqrt{2} - 1]$ it holds that*

$$\gamma^r(x) < R_{10}^r(x) < R_9^r(x) < \cdots < R_2^r(x) < R_1^r(x),$$

see Figure 7.

Proof. We will show that for each n , $1 \leq n \leq 10$, the flow of the vector field $Z^r(x, y) = (-y, (r - 1/r)y + x(x - 1))$ associated to system (8) crosses each curve $G_n^r(x, y) = y - R_n^r(x) = 0$ forwards. We compute

$$N_n^r(x) := \langle \nabla G_n^r(x, y), Z^r(x, y) \rangle|_{y=R_n^r(x)}.$$

We obtain

$$N_n^r(x) = r^{4n+2} x^{2n+2} (x - 1) \frac{B_n^r(x)}{C_n^r(x)^3},$$

where $B_n^r(x)$ and $C_n^r(x)$ are polynomials of degree n in x . The coefficients of $B_n^r(x)$ and $C_n^r(x)$ are polynomials in r .

We want to prove that for $r > 0$ both polynomials $B_n^r(x)$ and $C_n^r(x)$ are positive for $x \in (0, 1)$. We will approach the problem varying the parameter r and studying which are the possible bifurcations for the number of zeros of these polynomials when $x \in (0, 1)$. The values of $B_n^r(0)$, $C_n^r(0)$, $B_n^r(1)$ and $C_n^r(1)$, are polynomials in r whose coefficients are natural numbers. Then we have $B_n^r(0) > 0$, $C_n^r(0) > 0$, $B_n^r(1) > 0$ and $C_n^r(1) > 0$ for $r > 0$. For $n \geq 2$, we evaluate now the discriminants of the polynomials $B_n^r(x)$ and $C_n^r(x)$, $\text{Dis}(B_n^r(x), x)$ and $\text{Dis}(C_n^r(x), x)$. We obtain a polynomial in r of degree b_n with positive integers numbers as coefficients for $\text{Dis}(B_n^r(x), x)$ and a polynomial in r of degree c_n with also positive integer numbers as coefficients for $\text{Dis}(C_n^r(x), x)$, where the degrees b_n and c_n are given in Table 3. Then, the two discriminants do not vanish for $r > 0$. In consequence, the number of real roots in $(0, 1)$ of each polynomial, $B_n^r(x)$ or $C_n^r(x)$, does not change for $r > 0$. Picking a concrete value of r , for instance $r = 1/10$, we found, by applying the Sturm algorithm, that they have no roots in $(0, 1)$. Then, we deduce that $B_n^r(x) > 0$ and $C_n^r(x) > 0$ for $x \in (0, 1)$ and $r > 0$. Therefore, we have proved that for $n \geq 2$, $N_n^r(x) < 0$ for $x \in (0, 1)$ and $r > 0$, as we wanted to see. The case $n = 1$ is much easier because we do not need to compute the discriminants and we omit the details.

n	2	3	4	5	6	7	8	9	10
b_n	40	212	624	1480	2900	5028	8260	12560	18180
c_n	24	100	264	584	1100	1860	2996	4496	6444

TABLE 1. Degrees of the numerator and the denominator of the resultants computed in the proof of Proposition 3.4.

Finally for $k = 2, 3, \dots, 10$ we obtain by a direct computation

$$R_k^r(x) - R_{k-1}^r(x) = \frac{x^{2k}(x-1)r^{4k-1}D_k(r^2)}{C_k^r(x)C_{k-1}^r(x)},$$

where $D_k(x)$ is a polynomial with positive coefficients. Hence $R_k^r(x) < R_{k-1}^r(x)$ as we wanted to show. \square

As a corollary of Propositions 3.2 and 3.4 we obtain the following result:

Theorem 3.5. *Let $\Gamma^r = \{(x, \gamma^r(x)), x \in [0, 1]\}$, be a parametrization of the heteroclinic solution of system (8). For $x \in (0, 1)$ and $r \in (0, \sqrt{2} - 1]$ it holds that*

$$h_2^r(x) < h_3^r(x) < \cdots < h_{100}^r(x) < \gamma^r(x) < R_{10}^r(x) < R_9^r(x) < \cdots < R_1^r(x).$$

If we were interested in obtaining more precise upper bounds for the heteroclinic orbit, we could simply increase n and apply the same procedure. For instance for $r = 1/10$ we have performed all the computations and proved that for $x \in (0, 1)$,

$$\begin{aligned} h_{42}^{1/10}(x) &< \gamma^{1/10}(x) < R_{20}^{1/10}(x) & \text{and} & & R_{20}^{1/10}(x) - h_{42}^{1/10}(x) < 9 \times 10^{-32}, \\ h_{62}^{1/10}(x) &< \gamma^{1/10}(x) < R_{30}^{1/10}(x) & \text{and} & & R_{30}^{1/10}(x) - h_{62}^{1/10}(x) < 2 \times 10^{-40}, \\ h_{82}^{1/10}(x) &< \gamma^{1/10}(x) < R_{40}^{1/10}(x) & \text{and} & & R_{40}^{1/10}(x) - h_{82}^{1/10}(x) < 2 \times 10^{-47}, \end{aligned}$$

where the functions $R_m^r(x)$, $m > 10$, are defined similarly to the ones given in the above theorem. We remark that in all the cases the maximum error is at $x = 1$. It is also important to notice that for bigger values of r we need bigger values of n in $R_n^r(x)$ and $h_n^r(x)$ to obtain similar bounds.

Theorem 1.2 is simply a reformulation of the above theorem.

4. ON THE TIME-PARAMETRIZATION OF Γ_c FOR THE FISHER-KOLMOGOROV CASE

By using normal forms theory it is well-known that in a neighborhood of the origin of system (8) its unstable manifold can be parametrized as $F(e^{rt})$ for some analytic function F . So it seems natural to find bounds of the actual heteroclinic orbit which are rational functions of e^{rt} . As far as we know this idea is new.

First, we consider the family of rational functions

$$X(t) = \frac{\beta e^{rt} + \alpha e^{2rt}}{1 + (\alpha + 2\beta - 1)e^{rt} + \alpha e^{2rt}}$$

to try to approximate the function $x^r(t)$, where we denote by $(x^r(t), y^r(t))$ the time-parametrization of the heteroclinic orbit Γ^r . Notice that the parameters are taken in such a way that

$$\lim_{t \rightarrow -\infty} X(t) = 0, \quad X(0) = \frac{1}{2}, \quad \lim_{t \rightarrow \infty} X(t) = 1,$$

properties that are also satisfied by $x^r(t)$. We do not care about the y component because $y(t) = -x'(t)$.

Recall that the vector field associated to system (8) is

$$Z^r(x, y) = (Z_1^r(x, y), Z_2^r(x, y)) = (-y, (r - 1/r)y + x(x - 1)).$$

To study the behavior of the flow of this system on the curve $(X(t), Y(t))$, where $Y(t) = -X'(t)$ we compute

$$M(t) := -Y'(t)Z_1^r(X(t), Y(t)) + X'(t)Z_2^r(X(t), Y(t)). \quad (10)$$

If we introduce the compact notation $\Phi := e^{rt}$ we obtain that

$$M(t) = \frac{\Phi^3 (\alpha(\alpha + \beta - 1)\Phi^2 + 2\alpha\Phi + \beta) P_3(\Phi)}{(\alpha\Phi^2 + (\alpha + 2\beta - 1)\Phi + 1)^5},$$

where P_3 is a polynomial of degree 3 with coefficients depending also polynomially on α and β .

In order to simplify the expression of $M(t)$ we take $\alpha = 1 - \beta$. Then

$$M(t) = \frac{\Phi^3 (2(1 - \beta)\Phi + \beta) P_3(\Phi)}{((1 - \beta)\Phi^2 + \beta\Phi + 1)^5}.$$

Finally, taking $\beta = 2\sqrt{2} - 2$, the numerator and the denominator of the above fraction have a common zero. Then it writes as

$$M(t) = \frac{2(17 + 12\sqrt{2})r(1 - 6r^2)\Phi^3}{(1 + \sqrt{2} + \Phi)^7}.$$

In short we have proved the following:

Theorem 4.1. *Let $(x^r(t), y^r(t))$ be the time-parametrization of the heteroclinic orbit Γ^r of system (8) that satisfies $x^r(0) = 1/2$. Define*

$$X^r(t) = \frac{(2 + 2\sqrt{2} + e^{rt})e^{rt}}{(1 + \sqrt{2} + e^{rt})^2}.$$

Then:

- (i) *When $r < 1/\sqrt{6}$ it holds that $\text{sgn}(x^r(t) - X^r(t)) = -\text{sgn}(t)$.*
- (ii) *When $r = 1/\sqrt{6}$ it holds that $x^r(t) = X^r(t)$.*
- (iii) *When $r > 1/\sqrt{6}$ it holds that $\text{sgn}(x^r(t) - X^r(t)) = \text{sgn}(t)$.*

Theorem 1.3 follows from the above result, simply using that $u = 1 - x$ and $c = 1/r - r$.

Theorem 3.5 can also be used to obtain an explicit bound for $x^r(t)$. Since $h_2(x) = -rx + \frac{r}{2r^2 + 1}x^2 < \gamma^r(x)$, where $(x, \gamma^r(x))$ is the parametrization of Γ^r , solving the Cauchy problem

$$x' = rx - \frac{r}{2r^2 + 1}x^2, \quad x(0) = \frac{1}{2},$$

we obtain a function

$$U^r(t) = \frac{2r^2 + 1}{1 + (4r^2 + 1)e^{-rt}},$$

such that $\text{sgn}(x^r(t) - U^r(t)) = -\text{sgn}(t)$. The other bounds given in Theorem 3.5 give rise to implicit bounds of the form $H(t, x) = 0$ of the curve $(t, x^r(t))$.

4.1. Sharper bounds when $r \in (0, 1/\sqrt{6})$. This subsection improves the results of the previous section when $r < 1/\sqrt{6}$.

Fixed $r \in (0, 1/\sqrt{6})$ and given any natural n , following similar techniques that the ones used to prove Lemma 3.1, we can compute a function $x_n^r(t) = \sum_{j=1}^{2n} a_j \Phi^j$, with $\Phi = e^{rt}$ and $a_1 = 1$, that coincides with the solution $x^r(t)$ until order $2n$ in Φ . For instance

$$a_2 = -\frac{1}{2r^2 + 1} \quad \text{and} \quad a_3 = \frac{1}{(2r^2 + 1)(3r^2 + 1)}.$$

Looking Φ as an independent variable, we compute the associated Padé approximant of $\sum_{j=1}^{2n} a_j \Phi^j$ of order (n, n) , obtaining

$$Z_n^r(t) = \tilde{Z}_n^r(\Phi) = \frac{\sum_{j=1}^n b_j(r) \Phi^j}{\sum_{j=0}^n c_j(r) \Phi^j},$$

where $b_j(r)$ and $c_j(r)$ are polynomials on r . For instance

$$\tilde{Z}_2^r(\Phi) = \frac{3(2r^2 + 1)(3r^2 + 1)(4r^2 + 1)\Phi - 2(r - 1)(r + 1)(3r^2 + 1)\Phi^2}{3(2r^2 + 1)(3r^2 + 1)(4r^2 + 1) + 5(2r^2 + 1)(3r^2 + 1)\Phi + (3r^2 + 2)\Phi^2}.$$

Notice that it satisfies that $\lim_{t \rightarrow -\infty} Z_n^r(t) = 0$ but we do not impose neither that $Z_n^r(0) = 1/2$ nor that $\lim_{t \rightarrow \infty} Z_n^r(t) = 1$.

As in the proof of Theorem 4.1, to study the behavior of the flow associated to system (8) on the curve $(Z_n^r(t), -Z_n^r(t)')$ we compute the corresponding function (10). It writes as

$$M_n(t) = \widetilde{M}_n(\Phi) = \frac{\Phi^{2n+2} r^3 (1 - 6r^2) P_{3n-3}(\Phi)}{(Q_n(\Phi))^5},$$

where P_k and Q_k are polynomials of degree k with coefficients depending also polynomially on r . It can be seen that all these coefficients, as functions of r , take positive values on $(0, 1/\sqrt{6})$. We prove this fact introducing a new variable z satisfying $r = z^2/(\sqrt{6}(1 + z^2))$ in each of the coefficients and then applying the Descartes' Theorem to the resulting polynomials. Then $M_n(t) > 0$ for all t and for all r in $(0, 1/\sqrt{6})$. Moreover, this also shows that $Z_n^r(t)$ is well defined because $Q_n(\Phi) = \sum_{j=0}^n c_j(r) \Phi^j > 0$.

Since we want that $Z_n^r(0) = 1/2$ we need to modify $Z_n^r(t)$. To do this, for $\rho > 0$, we define the new family of functions $W_n^r(t, \rho) = \tilde{Z}_n^r(\rho\Phi)$. It is not difficult to see that following the above procedure the same results hold. Moreover it can be proved that there exists a unique value $\rho_0(r) \in (0, 1)$ such that $W_n^r(0, \rho_0(r)) = 1/2$. So we define $X_n^r(t) = W_n^r(t, \rho_0(r))$ and it satisfies $X_n^r(0) = 1/2$. Since the corresponding function (10) is positive, for each $n = 2, \dots, 8$, it holds that

$$\text{sgn}(x^r(t) - X^r(t)) = -\text{sgn}(t).$$

Moreover $\lim_{t \rightarrow \infty} X_n^r(t) = b_n(r)/c_n(r)$ and it can be seen that these limits, which are rational functions in r , satisfy

$$1 < b_8(r)/c_8(r) < b_7(r)/c_7(r) < \dots < b_2(r)/c_2(r).$$

Furthermore

$$E_k := \max_{r \in (0, 1/\sqrt{6})} \left(\frac{b_k(r)}{c_k(r)} - 1 \right)$$

decreases with k . For instance $E_2 = 3 - 4\sqrt{5}/3 \approx 0.02$ and $E_8 \approx 0.0007$.

CONCLUSIONS

In spite of the great interest of studying the traveling wave solutions of reaction-diffusion equations, $u_t = u_{xx} + f(u)$, there are no methods for obtaining explicit bounds for them. In this paper we present an approach that allows to obtain this type of bounds in the general case. Moreover we introduce more elaborated tools, based on the control of the heteroclinic trajectories of an associated planar ordinary differential equation, that allow to improve the general results when we deal with a particular function f . We study with detail the classical Fisher-Kolmogorov case $f(u) = u(1 - u)$. The methods developed can be easily adapted to treat the Newell-Whitehead-Segel and the Zeldovich equations

REFERENCES

- [1] M. J. Ablowitz and A. Zeppetella, *Explicit solutions of Fisher's equation for a special wave speed*, Bull. Math. Biol. **41** (1979), 835–840.
- [2] R. A. Fisher, *The wave of advance of advantageous genes*, Ann. Eugenics **7** (1937), 355–369.
- [3] A. Gasull, H. Giacomini and J. Torregrosa, *Some results on homoclinic and heteroclinic connections in planar systems*, Nonlinearity **23** (2010), 2977–3001.
- [4] P. Grindrod, *Patterns and Waves: The Theory and Applications of Reaction-Diffusion Equations*, Clarendon Press (1991).
- [5] A. Kolmogorov, I. Petrovskii and N. Piscounov, *A study of the diffusion equation with increase in the amount of substance, and its application to a biological problem*, In V. M. Tikhomirov, editor, *Selected Works of A. N. Kolmogorov I*, 248–270. Kluwer 1991. Translated by V. M. Volosov from Bull. Moscow Univ., Math. Mech. **1** (1937), 1–25.
- [6] A. C. Newell and J. A. Whitehead, *Finite bandwidth, finite amplitude convection*, J. Fluid Mech. **38** (1969), 279–303.
- [7] F. Sánchez-Garduño and P. K. Maini, *Travelling wave phenomena in some degenerate reaction-diffusion equations*, J. Differential Equations **117** (1995), 281–319.
- [8] F. Sánchez-Garduño and P. K. Maini, *Travelling wave phenomena in non-linear diffusion degenerate Nagumo equations*, J. Math. Biol. **35** (1997), 713–728.
- [9] L. A. Segel, *Distant sidewalls cause slow amplitude modulation of cellular convection*, J. Fluid Mech. **38** (1969), 203–224.
- [10] J. Xin, *Front propagation in heterogeneous media*, SIAM Rev. **42** (2000), 161–230.
- [11] Y. B. Zeldovich and D. A. Frank-Kamenetskii, *A theory of thermal propagation of flame*, Acta Physicochimica URSS **9** (1938), 341–350. English translation: In Dynamics of curved fronts, editor R. Pelcé, Perspectives in Physics Series, Academic Press, New York (1988), 131–140.

DEPT. DE MATEMÀTIQUES. UNIVERSITAT AUTÒNOMA DE BARCELONA. EDIFICI C. 08193 BELLATERRA, BARCELONA. SPAIN
E-mail address: gasull@mat.uab.cat

LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE. FACULTÉ DES SCIENCES ET TECHNIQUES. UNIVERSITÉ DE TOURS. 37200 TOURS. FRANCE
E-mail address: Hector.Giacomini@lmpt.univ-tours.fr

DEPT. DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, EDIFICI C. 08193 BELLATERRA, BARCELONA. SPAIN
E-mail address: torre@mat.uab.cat