

1 **Potassium: a neglected nutrient in global change**

2

3 Jordi Sardans<sup>1,2\*</sup> & Josep Peñuelas<sup>1,2</sup>

4 <sup>1</sup>CSIC, Global Ecology Unit CREAM-CEAB-CSIC-UAB. 08913 Cerdanyola del Vallès.

5 Catalonia. Spain.

6 <sup>2</sup>CREAF. 08913 Cerdanyola del Vallès. Catalonia. Spain.

7 Running title: Potassium stoichiometry and global change

8 \*Corresponding author. Email: [j.sardans@creaf.uab.es](mailto:j.sardans@creaf.uab.es) Tel: 34 93 851 2934 Fax: 34 93

9 581 4151

10

11

12 Keywords: climate change, drought, ecological stoichiometry, eutrophication, global

13 change, invasiveness, N:K, plant growth

14

15

16

17

18

19

20

21

22

23

24

25

26 **ABSTRACT**

27 **Aim:** Potassium (K) is the second most abundant nutrient after nitrogen (N) in plant  
28 photosynthetic tissues. Thousands of physiological and metabolic studies in recent  
29 decades have established the fundamental role of K in plant function, especially in  
30 water use efficiency and economy, and yet macroecological studies have mostly  
31 overlooked this nutrient.

32 **Methods:** We have reviewed available studies on the contents, stoichiometries and  
33 roles of potassium in the soil-plant system and in terrestrial ecosystems. We have also  
34 reviewed the impacts of global change drivers on K contents, stoichiometries, and roles.

35 **Conclusions:** The current literature indicates that K, at a global level, is as limiting as N  
36 and phosphorus (P) for plant productivity in terrestrial ecosystems. K limitation has  
37 been seen up to some degree in 70% of all studied terrestrial ecosystems. However,  
38 atmospheric K deposition from human activities represents higher amounts than that  
39 from natural sources in some areas. We are far from understanding the K fluxes  
40 between the atmosphere and land, and the role of anthropogenic activities in these  
41 fluxes. The increasing aridity expected in wide areas of the world makes K more critical  
42 through its role in water use efficiency. N deposition exerts a strong impact on the  
43 ecosystem K-cycle, decreasing K availability and increasing K limitation. Plant invasive  
44 success is enhanced by higher soil K availability, especially in environments without  
45 strong abiotic stresses. The impacts of other global change drivers, such as increasing  
46 atmospheric CO<sub>2</sub> or changes in land use remain to be elucidated. Current models of the  
47 responses of ecosystems and carbon storage to projected global climatic and  
48 atmospheric changes are now starting to consider N and P, but they should also consider  
49 K, mostly in arid and semiarid ecosystems.

50

51 **Introduction: Potassium in plant-soil system**

52

53 Potassium (K) is the most abundant cation in plant cells and is the second most  
54 abundant nutrient after nitrogen (N) in leaves. K is thus more abundant than phosphorus  
55 (P) (Leigh & Jones, 1984; Sardans *et al.*, 2006; Zheng & Shangguan, 2007; Sardans *et*  
56 *al.*, 2011, 2012a). K represents 2.6% of the weight of the Earth's crust. The  
57 concentrations of P and N in rocks are lower; P represents only 0.1% of the weight of  
58 the Earth's crust, and N is only present in trace amounts in magmatic rocks (< 200 ppm  
59 in granite). The most N-rich sedimentary rocks have only 0.1% N (Greenwood &  
60 Earnshaw, 1997; Holloway & Dahlgren, 2002). Young soils tend to have high  
61 phosphorus availability and low nitrogen availability (Houlton *et al.*, 2008). However,  
62 as soils age, nitrogen gradually accumulates in the soils (Hedin *et al.*, 2003), while they  
63 become more depleted in phosphorus (Walker & Syers, 1976; Vitousek & Howarth,  
64 1991). K can be more easily leached than N or P (Boxman *et al.*, 1994; Neiryneck *et al.*,  
65 1998). However, terrestrial ecosystems, such as forests and shrublands, have a great  
66 biological capacity for K-retention by several processes, such as plant "pumping" and  
67 resorption of K (Nowak *et al.*, 1991; Jobbáry & Jackson, 2001).

68 K "pumping" describes the plant K uptake from deep soil layers (with K  
69 provided in part by mineral weathering release) to upper soil layers. This can imply  
70 higher K losses by runoff (Barré *et al.*, 2009), but despite this runoff the pumping  
71 activity allows maintenance of the high levels of available-K in the surface of most soils  
72 of the globe (Jobbáry & Jackson, 2001). This biological control of ecosystem K-cycle  
73 strongly suggests a great evolutionary pressure to retain K in terrestrial ecosystems.  
74 Moreover, this "pumping" capacity can allow a long-term conservation of K in soils  
75 (Stone & Kszystyniak, 1977). The potential of parent material together with a

76 conservative biogeochemical cycle are underlying the maintenance of K supply to  
77 plants (Nowak *et al.*, 1991; Jobbáry & Jackson, 2001).

78 K in soil can be allocated to three pools of availability for uptake by roots. It is  
79 dissolved in soil water, adsorbed onto particles of clay and organic matter and held  
80 within the crystal structures of feldspar and mica. The organic matter in soils contains a  
81 negligible amount of K because it is not a constituent of biomolecules, hence easily and  
82 quickly leached from leaves due to its high solubility. However, the fraction of soil K  
83 directly available to plants (i.e., in soil solution) is usually a small proportion (0.1-0.2%)  
84 of the total soil K, whereas both the immediate available-water soluble and the  
85 exchangeable fraction are 1-2%, and the soil unavailable fractions are 96-99% (Wang *et al.*  
86 *et al.*, 2010; Britzke *et al.*, 2012). The soil available-K pool is easily leached by runoff.  
87 Soil-plant available K concentrations can thus frequently be lower than the soil-plant  
88 available N and P concentrations, even though the total soil K concentrations are  
89 generally higher than those of N and, especially, P (Sardans & Peñuelas, 2007; Sardans  
90 *et al.*, 2008). We have estimated the global K contents of the various K fractions in soil  
91 from the data in Soil Taxonomy (Soil Survey Staff, 1999) and other bibliographic  
92 sources (Table 1, Figure 1).

93 On the other hand, in terrestrial ecosystems, K shows a greater loss from the  
94 plant canopy than other nutrients such as N and P by foliar leaching (Duchesne *et al.*,  
95 2001; Kopacek *et al.*, 2009). Thus, this poses the question that if K is potentially  
96 limiting, why there has not been a greater natural selection to minimize leaching losses  
97 from leaves?. We must take into account that differently from N and P, that mostly  
98 constitute complex molecular structures, K occurs exclusively in soluble K<sup>+</sup> form. For  
99 this reason, from a physical point-of-view perspective, K can be more easily leached  
100 through cell membranes due to its smaller size. Probably, the need of maintaining its

101 cationic soluble form for its function has not allowed to improve the plant capacity to  
102 retain it. Moreover, mechanisms, such as reduction of stomatal density or thickening of  
103 the cuticle, which could help in preventing K losses, very probably present several  
104 evolutionary trade-offs with other important plant functions such as gas exchanges and  
105 water economy.

106         However, soil-plant system has a notable control on the ecosystem K-cycle they  
107 contribute in its retention. Current available data show a notable plant capacity to retain  
108 K by resorption. Diverse results have been reported in the studies comparing N, P, and  
109 K resorption efficiency (RE). Some researchers have observed  $N=P<K$  RE (van den  
110 Driessche, 1985; Nieminen & Helmisaari, 1996; Sardans *et al.*, 2005), others have  
111 observed  $P<N=K$  RE (Ozbucak *et al.* 2008),  $N<K=P$  RE (Chuyong *et al.*, 2000),  
112  $N>K>P$  RE (Yin *et al.*, 2009),  $P>K>N$  RE (Sahehi *et al.*, 2013),  $K<N=P$  RE  
113 (Helmisaari, 1992; Duchesne *et al.*, 2001; Tartachnyk & Blanke, 2004; Hagen-Thorn *et*  
114 *al.*, 2006 ), and  $N=P=K$  RE (Gallardo *et al.*, 1999; Trémolières *et al.*, 1999). A recent  
115 review of a global dataset has observed that K RE on a global scale (70.1%) seems to be  
116 higher than the RE of N (62.1%) and P (64.9%) (Vergutz *et al.*, 2012), although most of  
117 the studies reviewed by these authors did not take into account the role of leaf leaching  
118 when calculating RE. In fact, K leaches more easily from leaves than N and P; hence it  
119 is difficult to know the role of K foliar leaching in the studies that have calculated RE of  
120 these three nutrients, and some of these studies might be biased. Nieminen &  
121 Helmisaari (1996) observed in *Pinus sylvestris* that after correcting RE for nutrient  
122 losses by net throughfall, K RE was higher than N and P RE. Some results add even  
123 more complexity in the interpretation of the comparison of RE among these three  
124 nutrients. Some studies showed that the RE of N, P and K varied considerably  
125 depending on the phase of RE, and that this variance depended of each studied species,

126 but that this K RE was generally higher than N and P RE (Nieminen & Helmisaari,  
127 1996; Fife & Nambiar, 1997; Fife et al., 2008). . In summary, the available data allow to  
128 deduce that the internal plant RE of K is important, and similar, on average, to the REs  
129 of N and P.

130 The availability of K in several terrestrial ecosystems can also strongly depends  
131 on the input of K by atmospheric deposition (Table 2), which can vary from 0.07 to  
132 more than 100 kg ha<sup>-1</sup> yr<sup>-1</sup> (with a mean of the total, dry plus wet, atmospheric K  
133 deposition of 4.1 kg ha<sup>-1</sup> yr<sup>-1</sup> for the available studies, Table 2). The available data  
134 suggest that the current levels of K fertilization ( $0.025 \times 10^9$  t yr<sup>-1</sup>) and K atmospheric  
135 deposition ( $0.066 \times 10^9$  t yr<sup>-1</sup>) are very low compared with the global amount of  
136 available soil K ( $57.7 \times 10^9$  t). The current data for atmospheric K deposition also  
137 suggest that the amounts of K from atmospheric deposition in some areas of the world  
138 could have a large impact on soil K concentrations and on the stoichiometric  
139 relationships between K and other elements. For example, the deposition of 281 kg K  
140 ha<sup>-1</sup> reported for southern Nigeria (Muoghalu & Oakhunen, 2000) or the 6-23 kg K ha<sup>-1</sup>  
141 of wet deposition observed in some areas of India (Prathibha *et al.*, 2010) are notable. In  
142 the last few decades, the National Atmospheric Deposition Program database  
143 (<http://nadp.sws.uiuc.edu/>) has determined wet K deposition by analyzing rainfall of  
144 237 plots throughout North-America; the data show a great variability depending on the  
145 site (from 0.022 to 3.0 kg K ha<sup>-1</sup> yr<sup>-1</sup>). In this regard, we must highlight that the data of  
146 wet K deposition from precipitation have been collected for some world areas, the data  
147 of total K deposition are scarce despite the current knowledge suggest that K dry  
148 deposition could also be important (Table 2). It is surprising that studies on the potential  
149 effects of high amounts of K in terrestrial ecosystems have been apparently neglected,

150 though there are so many on the causes and consequences of N- and, even, P  
151 depositions.

152         The atmospheric deposition of K originates from natural processes, such as sea  
153 spray (Table 2) but also from agricultural activities and human industrial activities such  
154 as fertilizer and cement production, petrochemistry, steel industry, and coal combustion  
155 (Klumpp *et al.*, 2002; Walker *et al.*, 2003; Urban *et al.*, 2012). Human activities can  
156 represent a higher proportion than the natural processes (Ferm & Hultberg, 1999;  
157 Golobocanin *et al.*, 2008). We are, however, far from understanding the K fluxes  
158 between the atmosphere and the land and the role of anthropogenic activities in these  
159 fluxes (Quennehen *et al.*, 2012).

160         Ecological stoichiometric studies of terrestrial ecosystems have focused mostly  
161 on N and P (Vitousek & Howarth 1991; Elser *et al.*, 2000; Sterner & Elser, 2002;  
162 Sardans *et al.*, 2012b). Similarly, studies on the impacts of the drivers of global change  
163 on terrestrial biogeochemical cycles have focused mainly on these two elements  
164 (Vitousek *et al.*, 2010; Sardans *et al.*, 2012c; Peñuelas *et al.*, 2013a,b). Models of global  
165 carbon (C) circulation frequently do not include the role of nutrients in the  
166 establishment of C balances and the capacity of sinks to store C (Lé Quéré *et al.*, 2009).  
167 However, those that do, usually include only N (Schimel *et al.*, 2001; Mack *et al.*, 2004;  
168 Langley & Megonigal, 2010) or more recently P (Peñuelas *et al.*, 2012, 2013a, 2013b).

169         There is an increase in number of reports that highlight the importance of K in  
170 plant functions (Box 1) and responses to environmental changes, and its frequent  
171 limiting role for plant growth and C storage in terrestrial ecosystems. In the present  
172 study, we aim to demonstrate the urgent necessity to include K in the studies of  
173 biogeochemistry and stoichiometry in terrestrial ecosystems, and of the consequences of  
174 global change on nutrient cycles and their mutual feedbacks.

175

176 **Plant K uptake**

177 The importance of K in terrestrial plants is reflected by the sophisticated mechanisms of  
178 K uptake, redistribution, and homeostasis involving several families of cell-wall and  
179 membrane proteins (Hirsch *et al.*, 1998; Armengaud *et al.*, 2004; Szczerba *et al.*, 2009;  
180 Pyo *et al.*, 2010). Several regulatory mechanisms have been identified for K transporters,  
181 which are activated by various environmental factors such as water availability and soil  
182 concentrations of K<sup>+</sup>, Na<sup>+</sup> or Ca<sup>2+</sup> (Zhang *et al.*, 2006; Szczerba *et al.*, 2009; Pyo *et al.*,  
183 2010). Low-affinity and high-affinity K-transport systems have been identified. Briefly,  
184 the low-affinity transport system, believed to be channel mediated, acts at high external  
185 K concentrations and is isothermodynamically passive (Leigh, 2001; Szczerba *et al.*,  
186 2009). The high-affinity transport system is a saturable system, which catalyzes the  
187 thermodynamically active uptake of K at low external K concentrations (< 1 mM)  
188 (Schachtman & Schroeder, 1994; Szczerba *et al.*, 2009; Cuéllar *et al.*, 2010). The high-  
189 affinity transport system, as opposed to the low-affinity transport system, is down-  
190 regulated at high external K concentrations and up-regulated at low external K  
191 concentrations (Schroeder & Fang, 1991; Maathuis & Sanders, 1994; Hirsch *et al.*, 1998;  
192 Szczerba *et al.*, 2009). Both the systems, though, represent an electrogenic entry of net  
193 positive charge, which requires the active removal of protons to maintain electrical  
194 neutrality (Leigh, 2001; Szczerba *et al.*, 2009).

195         The capacity of terrestrial plants to take up K and to maintain their homeostatic  
196 properties is determined by sophisticated mechanisms of genetic expression  
197 (Armengaud *et al.*, 2004; Yin *et al.*, 2011; El-Mesbahi *et al.*, 2012). Moreover, K  
198 uptake is closely linked to water economy. Some studies have observed that  
199 transmembrane channels for K and water are likely to be co-regulated and may function

200 in a coordinated manner to maintain the appropriate cytosolic osmolarity (Patrick *et al.*,  
201 2001; Liu *et al.*, 2006; Osakabe *et al.*, 2013). Reports on the co-regulation of K and  
202 water channels in plants, however, are still scarce.

203

#### 204 **A widespread limiting role of K in plant growth and ecosystem productivity**

205 In contrast to the many physiological studies demonstrating the importance of K in  
206 terrestrial plants, few ecological studies have investigated the possible role of K  
207 limitation in terrestrial plant productivity, particularly in comparison to the studies on N  
208 and P limitation. The existing studies suggest that the fraction of vegetation that is K  
209 limited or K co-limited with N and/or P is high (Figure 2, Table 3). Tripler *et al.* (2006)  
210 conducted a review and metadata analysis of K limitation in forest ecosystems with all  
211 the data available until 2004. This meta-analysis, including 38 articles (a subset that is  
212 included in Figure 2 and Table 3), showed an overall limiting effect of K in tree growth,  
213 with 69% of the studies showing some level of K limitation. We have now also included  
214 data from studies published up to July 2013 and from other terrestrial ecosystems such  
215 as grasslands, observing that 69% of 56 studies have detected at least some significant  
216 limiting role of K (Figure 2, Table 3).

217

#### 218 **K stoichiometry and water availability in terrestrial ecosystems**

219 Most of the studies on relationship between terrestrial plant stoichiometry and climatic  
220 gradients exclusively focused on N and P and their ratio (Reich & Oleksyn, 2004;  
221 Kerkhoff *et al.*, 2005). Some recent studies observed that foliar K concentrations are  
222 negatively correlated with mean annual precipitation (MAP) (Fyllas *et al.*, 2009; Zhang  
223 *et al.*, 2012), whereas few others observed contrary relationships (Sardans *et al.*, 2011;  
224 2012a). On the other hand, some studies positively related plant K uptake capacity with  
225 soil water availability (Fernandez *et al.*, 2011; Ge *et al.*, 2012); Tomlinson *et al.* (2013),

226 while comparing leaf traits of different plant species growing in wet and arid  
227 environments, observed that leaves of species adapted to arid sites are small with high K  
228 concentrations. Furthermore, it was observed that tree species at the driest sites, such as  
229 in Mediterranean evergreen and dry tropical forests, have a higher capacity to change  
230 their seasonal internal allocation of K, with a higher allocation of K to leaves during  
231 summer (the driest season) than the species at wetter sites (Milla *et al.*, 2005; Rivas-  
232 Ubach *et al.*, 2012; Sardans *et al.*, 2012a).

233         Thus a trade-off is expected because K uptake is strongly related to soil water  
234 content (Watson *et al.*, 2001; Oliveira *et al.*, 2004), while, as previously commented, K  
235 is specially determinant to increase water use efficiency and to limit water losses.  
236 Therefore, at a global scale in wet and more productive environments, we should expect  
237 that plants are able to uptake more K than in dry environments, but, on the other hand,  
238 in drier environments plants can also increase their K concentrations by conservative  
239 mechanisms, such as increasing K resorption. Thus, current data suggest not only that K  
240 uptake is strongly determined by water availability but also that K is crucial to the  
241 strategies of water economy, leading to the paradox that the more necessary K is (under  
242 drought), the more difficult is its uptake.

243         Some studies have observed that K accumulation in soil is greatest in arid  
244 ecosystems rather than in mesic ecosystems (Singer, 1989), despite studies of natural  
245 gradients have observed that K accumulates and concentrates more in biomass with  
246 increasing annual mean precipitation (Sardans *et al.*, 2011). However, RE of K remains  
247 considerable and could be expected to be higher in dry ecosystems, compensating the  
248 low capacity of K uptake in dry conditions (in comparison with plants at wetter sites as  
249 reported in some studies, see e.g. Fayyaz *et al.*, 2013; Tomlinson *et al.*, 2013).  
250 Moreover, the K retention capacity in plants in arid environments could be improved by

251 lower leaching (Singer, 1989). Thus at this moment, given all these previous results, it  
252 is difficult to reach any consistent conclusion on whether K is more actively retained in  
253 dry or wet ecosystems.

254         Recent stoichiometric studies have observed that K is even more variable among  
255 dry plant ecotypes than N and P (Sardans *et al.*, 2011) and is also more responsive to  
256 environmental pressure such as drought or herbivore attack (Sardans *et al.*, 2013; Rivas-  
257 Ubach *et al.*, 2012; 2014). Foliar K concentrations have been found to co-vary with the  
258 entire metabolome, suggesting the importance of K in global plant function, despite its  
259 absence from biomolecules (Rivas-Ubach *et al.*, 2012; 2014). The strong link between  
260 plant K concentrations and water availability justifies the study of K and its  
261 stoichiometric relationships with other nutrients. Focusing on K would more strongly  
262 integrate the dimension of water availability in the study of terrestrial ecological  
263 stoichiometry. Recent ecological stoichiometric studies have observed that K plays a  
264 more fundamental role than does N or P in the differences in elemental composition  
265 among and within species in response to different water availability.

266         Some studies have shown that different grass species have different optimal N:K  
267 ratios of supply (Lawniczak *et al.*, 2009). A reduction in soil K availability is frequently  
268 observed as a consequence of N deposition. As commented below, this is due to large  
269 losses of K by leaching that causes nutrient imbalances. Moreover, other studies have  
270 observed that K concentrations in the soil solution and in runoff are more sensitive to  
271 changes in land use and fertilization than N or P concentrations (Watmough *et al.*, 2005;  
272 Anguelov *et al.*, 2011), and also that K is more quickly leached from litter than N or P,  
273 and it has a much shorter residence time in soil organic matter (Schreeg *et al.*, 2013;  
274 Florez-Florez *et al.*, 2013).

275

276 **Anthropogenic impacts on the K cycle**

277 **K fertilizers:** K is required in large amounts for growing crops, but farmers in poor  
278 countries often do not fertilize with K due to high costs, which leads to K deficiencies in  
279 large areas of farmland (Hoa *et al.*, 2006; Andrist-Rangel *et al.*, 2007). The situation is  
280 especially critical in Africa. Many African croplands are dramatically under-fertilized,  
281 especially with K. Despite covering 25% of global potential cropland area (FAO 2003,  
282 2011), the African share of global fertilizer consumption is 2.7% for N, 2.4% for P and  
283 1.7% for K (FAO, 2011). The fertilizer nutrient supply/demand balance indicates that  
284 Africa will remain as an exporter of phosphate and nitrogen, but for potash, the region  
285 would continue to depend solely on import (FAO, 2011). This fact, along with the great  
286 economical deficits in most African states, makes K a critical limiting factor for food  
287 production in this continent.

288 Croplands in China, Egypt, Bulgaria and parts of southeastern Asia also have  
289 negative balances of K, mainly due to socio-economic factors that decrease crop  
290 productivity (Magen, 2008; Ryan & Sommer, 2012; Ryan *et al.*, 2012). This negative  
291 balance leads to an increasing impoverishment of K in cropland soils (Vanderpol &  
292 Traore, 1993; Singh & Bansai, 2009). Moreover, the increasing demand for food and  
293 increasing aridity in several areas of the world, such as the Mediterranean Basin (mainly  
294 the Maghreb) and the Sahel, make K fertilization more critical given the important role  
295 of K in water use efficiency (Ruan *et al.*, 1997; Römheld & Kirkby, 2010). A recent  
296 study has concluded that K is currently the most important limiting nutrient in  
297 Mediterranean croplands (Ryan & Sommer, 2012). Agriculture practices that decrease  
298 soil pH favor K leaching and thus decrease availability for plants (Sharifi *et al.*, 2013).  
299 For example, despite that sometimes N fertilization can enhance soil K-availability in a  
300 short term by decreasing soil pH (Mitchell & Smethurst, 2008), but in absence of K

301 fertilization, it finally leads to a decrease in soil K-availability because of leaching (He  
302 *et al.*, 1999; Mitchell & Smethurst, 2008; Yavitt *et al.*, 2011; Glab & Gondek, 2014).

303 Demands and uses of fertilizers are expected to increase in the coming decades  
304 (Roberts, 2008). Mineable reserves of K are sufficient to meet the projected demand for  
305 centuries (Roberts, 2008). Some results suggest that the increasing use of K fertilizer in  
306 some regions in recent decades has increased the atmospheric deposition of K in areas  
307 adjacent to cropland (Gélinas *et al.*, 2000). The deposition of K from agricultural  
308 activities is very high in some developing countries such as Brazil, reaching 55% of  
309 total K deposition (Allen *et al.*, 2010). In contrast, the deposition of K from energy  
310 production and industrial plants has decreased in some European countries (Ruoho-  
311 Airola & Salminen, 2003). Some studies have observed that K deposition, linked to  
312 industrial and, frequently, agricultural activities as well, can represent a higher  
313 proportion than that from natural processes (Ferm & Hultberg, 1999; Golobocanin *et al.*,  
314 2008) such as sea spray (Poor *et al.*, 2006; Prathibha *et al.*, 2010).

315 Inputs of K from atmospheric deposition have a determinant role in K balances  
316 of forests, counteracting, such as observed in some areas, the loss of K from leaching,  
317 which is favored by N deposition (Friedland & Miller, 1999; Watmough & Dillon,  
318 2004). These inputs of K contribute to K availability in tropical forest ecosystems  
319 (Vitousek & Sanford, 1986; Muoghalu & Oakhumen, 2000). All these findings justify  
320 the study of the impacts of an increased use of K fertilizers on the rates of K  
321 atmospheric deposition and thereafter on ecosystem function at least in some parts of  
322 the world.

323

324 **K in a drier world:** Climate models project an increase in the extent of drought over  
325 large areas of the world that are already dry such as the Mediterranean Basin and the

326 Sahel (IPCC, 2007). Despite the intensive study of drought resistance under field  
327 conditions, only a few field studies have tested the impacts of drier conditions on the  
328 status of K in terrestrial plant communities (Table 4). These studies show that drought  
329 increases total soil K but decreases its soluble soil fraction. Moreover, drought increases  
330 the retention of K in the wood of some species and the concentration of K in the  
331 photosynthetic tissues of others (Table 4). The effects of drought are usually  
332 asymmetrical in different plant organs, because plants tend to allocate and retain more  
333 K in stems, and a fraction of this K is transported to leaves during the driest seasons  
334 (Table 4). These few studies suggest a possible increase in biomass K concentrations in  
335 dry areas but also a possible decrease in total biomass K content as a result of reduced  
336 growth. This decrease of K in biomass and total soil K increases could enhance the risk  
337 of higher losses of K from soil.

338         As previously stated, the availability and uptake of K are essential for  
339 improving drought resistance in various types of plant communities (Harvey & van den  
340 Driessche, 1999; Ashraf *et al.*, 2002; Egilla *et al.*, 2005; Sardans & Peñuelas, 2012c).  
341 Globally, plant K uptake, however, appears to be associated with water availability  
342 (Sardans *et al.*, 2011; 2012a; Ge *et al.*, 2012), wherein less K uptake in turn negatively  
343 affects water uptake by reducing the activity of aquaporins (Kanai *et al.*, 2011). All  
344 these interconnected relationships strongly suggest a cascade of higher water deficit,  
345 lower K uptake and a reduced capacity to avoid drought, which could become a serious  
346 problem in currently dry areas that are threatened by future drier scenarios. These  
347 scenarios could even be more negative if torrential rain increases and MAP decreases,  
348 as projected for areas such as the Mediterranean Basin and the Sahel (ICPP, 2007).  
349 Moreover, the capacity of different plant species to use and remobilize K can be crucial  
350 in their competitive relationships (Dunlop *et al.*, 1979; Tilman *et al.*, 1999). Thus, due

351 to the high relevance of K under conditions of water stress, we can expect that K will  
352 become an important factor in the shifts in species composition of plant communities  
353 under drought. Plants can also respond to drought by improving efficiencies of K and  
354 water uptake. Some phenomena linked to an improved resistance to drought stress, such  
355 as soil patchiness or the capacity to redistribute water, can favor K uptake. Plants under  
356 drought can improve the use of water for the uptake of nutrients by their capacity to  
357 redistribute soil water (Sardans & Peñuelas, 2013b). Hydraulic lift and its opposite,  
358 downward siphoning (Smith *et al.*, 1999), can favor water recirculation, efficiency of  
359 water use and the capacity of plants to take up nutrients (Sardans & Peñuelas, 2013b). A  
360 higher capacity to redistribute water to increase K uptake can be a key factor for the  
361 survival of plants in drier conditions. In this context, due to the lack of studies, it is not  
362 possible to assess the importance of water redistribution in enhancing not only water use  
363 efficiency but also the capacity of nutrient uptake.

364         Moreover, the variability of water and nutrient availability on slopes in the drier  
365 areas of the Mediterranean Basin is related to the distribution of patches of vegetation  
366 relative to the patches of bare soil that can be considered as runoff sinks and sources,  
367 respectively (Gabarrón-Galeote *et al.*, 2012; Merino-Martin *et al.*, 2012; Sardans &  
368 Peñuelas, 2013c). This effect of patches is crucial in the conservation of vegetation at  
369 drier sites. Gómez-Aparicio *et al.* (2005) have observed that the presence of large  
370 woody plants improves the availability of soil nutrients and in particular of K in  
371 Mediterranean ecosystems. The role of soil patches in the changes in the complete K  
372 cycle has not been specifically studied, despite the possibility that in patches the higher  
373 availability of water, the better soil texture and the horizontal redistribution of water  
374 could strongly affect the K cycle.

375

376 **Impacts of N and acid atmospheric deposition on the K cycle:** The impacts of N  
377 deposition on decreasing C:N in plants and soils have been extensively reported  
378 (Mulder *et al.*, 2013). However, only a relatively small proportion of the studies on the  
379 impacts of N deposition on terrestrial ecosystems stoichiometry have taken K into  
380 account (Figure 3). Our review of current literature shows that atmospheric N  
381 deposition increases the leaching of K from soils, thus decreases K concentrations in the  
382 soils (and decreases K concentrations and increases N:K ratios in plants. Despite that  
383 plants respond to increasing K uptake, an ultimate enhancement of K limitation is  
384 frequently observed. This loss of K under N deposition is frequently related to K  
385 deficiencies in trees (Tzvetkova & Haddjiivanova, 2006). Moreover, atmospheric S  
386 deposition has also contributed to K leaching thus decreasing soil K availability in  
387 several European (Britton *et al.*, 2008; Sutton *et al.*, 2011) and North-American  
388 (Mitchell *et al.*, 1996; Navratil *et al.*, 2010) areas. However, this K leaching has been  
389 reduced in the last decades as a result of the dramatic reduction of S emissions in  
390 Europe since the early 1980s, after the reduction of coal combustions and the  
391 implementation of policies to improve air quality (Eurostat, 2009; Navratil *et al.*, 2010;  
392 Robinson *et al.*, 2013).

393         The N eutrophication effects on the K cycle and N:K ratios may have large flow-  
394 on impacts (Figure 3). Decreases in soil K contents and increases in soil N:K ratios are  
395 associated with the loss of species richness (Huang *et al.*, 2013). Moreover, P:K soil  
396 ratios can increase because the solubility and mobility of K are higher than those of P,  
397 therefore K leaches more easily than P (Olde Venterink *et al.*, 2009). Changes in N:K  
398 and P:K ratios can favor species tolerance to water stress or competitively faster-  
399 growing species, depending on the circumstances. For example, a decrease in N:K ratio  
400 under increasing drought would probably favor the stress-tolerant species, which are

401 better adapted to allocate K to their leaves and to use it to improve drought avoidance,  
402 more than favoring the faster-growing competitors.

403

404 **K and success of plant invasion:** Most studies have focused on the roles of N and P in  
405 the success of species invasion (Sardans & Peñuelas, 2012). The few studies  
406 investigating relationships between invasive success and the K availability of soils have  
407 found the relationships to be important. In most cases, the success of alien plants  
408 depends on a high availability of soil K (Figure 4) and on a higher capacity of K uptake  
409 by the invasive species compare to the native plants (Pieters & Baruch, 1997), but it has  
410 also been associated with low soil K availability after the establishment of invasive  
411 species (Figure 4). Other studies, however, have observed an opposite pattern. The  
412 invasive success of *Taraxacum officinalis* in the Andes has been linked to low soil K  
413 availability (Cavieres *et al.* 2008). By examining the impacts of seven widespread  
414 invasive plant species in northern Europe, Dassonville *et al.* (2008) observed that  
415 invasive species reduced soil K availability in K-rich soils, but increased soil K  
416 availability in K-poor soils. The success of alien plants thus appears to depend on high  
417 K availability in productive environments and on low K availability in less productive  
418 environments; however, the low number of studies precludes formulation of a definitive  
419 conclusion on this dependence (Funk & Vitousek, 2007; Peng *et al.*, 2011; Sardans &  
420 Peñuelas, 2012). Alien plants with a more efficient use of K may be able to increase  
421 their spread in arid areas. Islands are among the most threatened areas by plant invasive  
422 success (Denslow *et al.*, 2009; Atwood & Meyerson, 2011). Current literature relates  
423 alien success in islands to several variables linked to humans, such as population  
424 density (Atwood & Meyerson, 2011; Traveset *et al.*, 2014), level of human impact  
425 (MacDonald *et al.*, 1991; Monty *et al.*, 2013), ecological traits such as genetic diversity

426 (Hardman *et al.*, 2012; Traveset *et al.*, 2014), differences in phylogeny with native  
427 species (Bezeng *et al.*, 2013), island area (Traveset *et al.*, 2014), isolation and distance  
428 to nearest continent (Traveset *et al.*, 2014), and climate change (Yamashita *et al.*, 2000).  
429 The possible relationship between bedrock K<sub>2</sub>O concentrations and plant alien success  
430 remains mostly unexplored. However, Denslow *et al.* (2009) observed a significant  
431 relationship between the main substrate of the island (coralline limestone islands,  
432 basaltic with metamorphic, ultrabasic, and continental plutonic islands) and the number  
433 of invasive plants.

434

#### 435 **Final remarks**

436 1. The data available from current studies on the role of K availability in the growth of  
437 plants indicate that K is frequently a limiting factor for plant productivity in terrestrial  
438 ecosystems.

439 2. Apart from the well-known physiological processes linked to water economy, such as  
440 enzymatic function, internal transport systems and cell-membrane integrity, K has  
441 recently been associated with an increasing number of plant ecological processes, such  
442 as resistance to diseases, herbivory, salinity, cold, frost or waterlogging, with large  
443 implications on ecosystem function and capacity of global crop production.

444 3. Recent ecological stoichiometric studies of terrestrial ecosystems that have included  
445 K with the classically considered C, N and P have shown that the stoichiometric  
446 changes involving K are even larger than those involving N and P in plant responses to  
447 the environmental changes. These findings justify the inclusion of K in ecological  
448 stoichiometric studies.

449 4. Drought can be more intense in current arid areas. Given the importance of K in  
450 water use efficiency and the difficulty of K uptake under drought conditions, K is

451 expected to have an increasingly important role in determining the function and survival  
452 of plant communities.

453 5. N deposition has been linked to several processes, such as forest decline and soil  
454 imbalances, but the impacts on K are not well known.

455 6. Current studies suggest that the success of invasive species in environments that are  
456 not strongly limited by nutrients is associated with a high capacity of K uptake and with  
457 the consequent decrease in K availability of soils.

458

459

#### 460 **Acknowledgements**

461 This research was supported by the Spanish Government grant CGL2013-48074-P, the  
462 Catalan Government grant SGR 2014-274, and the European Research Council Synergy  
463 grant ERC-2013-SyG-610028, IMBALANCE-P.

464

465

466

467

468

469

470

471

472

473

474

475

476 BOX 1. Main Ecophysiological roles of K

477 **Water economy:** K is involved in many mechanisms that mitigate drought stress.  
478 Studies in greenhouses and with potted plants have demonstrated the importance of K  
479 supply in the resistance of plants toward drought. K plays many fundamental  
480 physiological and metabolic roles in terrestrial plants related to water economy and  
481 osmotic homeostasis such as maintenance of cellular turgor and osmotic pressure  
482 (Ashraf *et al.*, 2002; Levi *et al.*, 2011), control of water conductance and transpiration  
483 (Harvey & van Driessche, 1999; Arquero *et al.*, 2006; Benloch-González *et al.*, 2010),  
484 improvement of root hydraulic conductivity (El-Mesbahi *et al.*, 2012), regulation of sap  
485 flow (Oddo *et al.*, 2011), regulation of membrane potentials (Schroeder & Fang, 1991),  
486 stomatal control (Talbot & Zeiger, 1996; Benloch-González *et al.*, 2010), and  
487 maintenance of ionic homeostasis or transmembrane potential (Su *et al.*, 2001; Waraich  
488 *et al.*, 2011). Moreover, field studies have observed that K, apart from its role in  
489 stomatal control and water conductance, can further affect water economy as an  
490 osmolyte (Babita *et al.*, 2010; Levi *et al.*, 2011) directly linked to improving the  
491 capacity to retain water (Nandwal *et al.*, 1998; Rao *et al.*, 2012).

492 In support of these physiological observations, some recent metabolomic studies  
493 have observed that K concentrations are related to increases in molecular osmolytes  
494 under drought (Talbot & Zeiger, 1996; Levi *et al.*, 2011; Rivas-Ubach *et al.*, 2012).  
495 The Mediterranean evergreen *Olea europea* decreases its capacity to regulate stomatal  
496 control when growing under moderate K deficiency (Arquero *et al.*, 2006); this suggests  
497 that even drought-resistant tree species can suffer detrimental effects in their growth  
498 capacity and survival under drought conditions through a reduction in their capacity to  
499 take up K. A few studies, though, have observed no effects of K availability on drought  
500 avoidance (e.g., Ashraf *et al.*, 2002).

501 K also contributes to the detoxification of reactive oxygen species (ROS) (Wang  
502 *et al.*, 2013). The impairment of stomatal regulation in K-deficient plants under drought  
503 stress produces a consequent impairment of CO<sub>2</sub> fixation, and molecular oxygen is  
504 activated leading to the generation of ROS (Cakmak, 2000). Under drought conditions,  
505 an increase of extrachloroplastic K<sup>+</sup> concentrations in plant cells with an adequate K  
506 supply prevents the inhibition of photosynthesis (Egilla *et al.*, 2005). This effect is  
507 related to the role K plays in enhancing photosynthetic CO<sub>2</sub> fixation, improving the  
508 transport of photosynthates into sink organs and inhibiting the transfer of photosynthetic  
509 electrons to O<sub>2</sub>, thereby reducing ROS production (Cakmak, 2005). Many field studies  
510 have thus observed correlations between K concentrations in plants and soils with  
511 mechanisms of drought avoidance (Nandwal *et al.*, 1998; Arquero *et al.*, 2006; Teixeira  
512 *et al.*, 2008; Fayyaz *et al.*, 2013; Tomlinson *et al.*, 2013).

513 Many physiological and metabolomic studies in garden and potted plants (Babita  
514 *et al.*, 2010; Cuéllar *et al.*, 2010) and in crops (Römheld & Kirkby, 2010; Waraich *et al.*,  
515 2011; Wang *et al.*, 2013), and some studies across environmental gradients under field  
516 conditions (Kadar & Foldesi, 2001), have observed a strong link between K  
517 concentrations and the resistance and adaptation of plants to drought. Under drought K  
518 deficiency, plant defoliation (Harvey & van den Driessche, 1999) and mortality (Del  
519 Campo *et al.*, 2011; El-Mesbahi *et al.*, 2012) increases. A few studies, though, have  
520 observed no effects of K availability on drought avoidance (e.g., Ashraf *et al.*, 2002).

521

522 **Other physiological functions:** Many plant functions that are not directly related to  
523 water economy have recently been associated with K, such as enzymatic activity and  
524 protein synthesis (Armengaud *et al.*, 2004; 2009), internal transport and allocation of  
525 amino acids and nitrates (Jarvis *et al.*, 1989), photosynthetic activity and retranslocation

526 of fixed C (Zhao *et al.*, 2001; Teixeira *et al.*, 2008), signal transduction and energy  
527 transport (Gajdanowicz *et al.*, 2011), cellular, foliar and root expansion (Elumalai *et al.*,  
528 2002; Yin *et al.*, 2011) and foliar movement (Travlos *et al.*, 2008), among several other  
529 metabolic and physiological functions essential for plant life. Evidences, moreover, are  
530 increasing that adequate K fertilization plays an important role in defending crops not  
531 only against drought but also against many other biotic and abiotic stresses such as  
532 diseases, pests, salinity, cold, frost, and waterlogging (see recent review of Wang *et al.*,  
533 2013). Gajdanowicz *et al.* (2011) recently observed that K<sup>+</sup> circulating in the phloem  
534 creates a mobile gradient serving as a decentralized energy store that can be used to  
535 overcome local energy limitations through the activation of diverse membrane proteins.  
536 The authors observed that posttranscriptional modification of the phloem-expressed  
537 *Arabidopsis* K<sup>+</sup> channel AKT2 taps this “potassium battery”, which then efficiently  
538 assists the plasma membrane H<sup>+</sup>-ATPase in energizing the transmembrane phloem  
539 reloading processes. Thus K<sup>+</sup> will be loaded into the phloem from source tissues where  
540 sufficient chemical energy is available. The established transmembrane K<sup>+</sup> gradient is  
541 then transported with the phloem stream to other parts of the plant. The energy stored in  
542 the K<sup>+</sup> membrane gradient can be harvested to fuel transport processes by opening a  
543 gateway (for example the AKT2-like channels) for the passage of K<sup>+</sup> ions through the  
544 membrane being this process related to sucrose leakage from the phloem that enhances  
545 energy status.

546

547

548

549

550

551

552 **References**

553

554 Allen, A.G., Cardoso, A.A., Wiata, A.G., Machado, C.M.D., Paterlini, W.C. & Baker, J.  
555 (2010) Influence of intensive agriculture on dry deposition of aerosol nutrients.  
556 *Journal of the Brazilian Chemical Society*, **21**, 87-97.

557 Andrist-Randgel, Y., Edwards, A.C., Hillier, S. & Oborn, I. (2007) Long-term K  
558 dynamics in organic and conventional mixed cropping systems as related to  
559 management and soil properties. *Agriculture Ecosystems and Environment*, **122**,  
560 413-426.

561 Anguelov, G., Anguelova, I. & Bailey, N.O. (2011) Land-use impact on soil solution  
562 constituents from an Ultisol of North Florida. *Nutrient Cycling in Agroecosystems*,  
563 **90**, 171-187.

564 Armengaud, P., Breiting, R. & Amtmann, A. (2004) The potassium-dependent  
565 transcriptome of *Arabidopsis* reveals a prominent role of Jasmonic acid in nutrient  
566 signaling. *Plant Physiology*, **136**, 2556-2576.

567 Armengaud, P., Sulpice, R., Miller, A.J., Stitt, M., Amtmann, A. & Gibon, Y. (2009)  
568 Multilevel analysis of primary metabolism provides new insights into the role of  
569 potassium nutrition for glycolysis and nitrogen assimilation in *Arabidopsis* roots.  
570 *Plant Physiology*, **150**, 772-785.

571 Arquero, O., Barranco, D. & Benlloch, M. (2006) Potassium starvation increases  
572 stomatal conductance in olive trees. *HortScience*, **41**, 433-436.

573 Ashraf, M., Ashfaq, M. & Ashraf, M.Y. (2002) Effects of increased supply of  
574 potassium on growth and nutrient content in pearl millet under water stress.  
575 *Biologia Plantarum*, **45**, 141-144.

576 Atwood, J.P. & Meyerson, L.A. (2011) Island biogeography extends to small-scale  
577 habitats: low competitor density and richness on islands may drive trait in  
578 nonnative plants. *Biological Invasions*, **13**, 2035-2043.

579 Babita, M., Maheswari, M., Rao, L.M., Shanker, A.K. & Rao, D.G. (2010) Osmotic  
580 adjustment, drought tolerance and yield in castor (*Ricinus communis* L.) hybrids.  
581 *Environmental and Experimental Botany*, **69**, 243-249.

582 Barré, P., Berger, G. & Velde, B. (2009) How element translocation by plants may  
583 stabilize illitic clays in the surface of temperate soils. *Geoderma*, **151**, 22-30.

584 Becker, D., Hoth, S., Ache, P., Wenkel, S., Roelfsema, M.R.G., Meyerhoff, O., Hartung,  
585 W. & Hedrich, R. (2003) Regulation of the ABA-sensitive *Arabidopsis* potassium  
586 channel gene GORK in response to water stress. *FEMS Letters*, **554**, 119-126.

587 Benlloch-González, M., Romera, J., Cristescu, S., Harren, F., Fournier, J.M. & Benlloch,  
588 M. (2010) K<sup>+</sup> starvation inhibits water-stress-induced stomatal closure via  
589 ethylene synthesis in sunflower plants. *Journal of Experimental Botany*, **61**, 1139-  
590 1145.

591 Bezeng, B.S., Savolainen, V., Yessoufou, K., Papadopoulos, A.S.T., Maurin, O. & van  
592 der Bank, M. (2013) A phylogenetic approach towards understanding the drivers  
593 of plant invasiveness on Robben island, South Africa. *Botanical Journal of the*  
594 *Linnean Society*, **172**, 142-152.

595 Boxman, A.W., Cobben, P.L.W. & Roelofs, J.G.M. (1994) Does (K+Mg+Ca+P)  
596 fertilization lead to recovery of tree health in a nitrogen stressed *Quercus rubra* L.  
597 stand? *Environmental Pollution*, **85**, 297-303.

598 Britton, A.J., Helliwell, R.C., Fisher, J.M. & Gibbs, F.S. (2008) Interactive effects of  
599 nitrogen deposition and fire on plant and soil chemistry in an alpine heathland.  
600 *Environmental Pollution*, **156**, 409-416.

- 601 Britzke, D., da Silva, L.S., Moterle, D.F., dos Santos Rheinheimer, D. & Bortoluzzi,  
602 E.C. (2012) A study of potassium dynamics and mineralogy in soils from  
603 subtropical Brazilian lowlands. *Journal of Soils and Sediments*, **12**, 185-197.
- 604 Cakmak, I. (2000) Possible roles of zinc in protecting plant cells from damage by  
605 reactive oxygen species. *New Phytologist*, **146**, 185-205.
- 606 Cakmak, I. (2005) The role of potassium in alleviating detrimental effects of abiotic  
607 stresses in plants. *Journal of Plant Nutrition and Soil Science*, **168**, 521-530.
- 608 Cavieres, L.A., Quiroz, C.L. & Molina-Montenegro, M.A. (2008) Facilitation of the  
609 non-native *Taraxacum officinale* by native nurse cushion species in the high  
610 Andes of central Chile: are there differences between nurses? *Functional Ecology*,  
611 **22**, 148-156.
- 612 Chuyong, G.B., Newbery, D.M. & Songwe, N.C. (2000) Litter nutrients and  
613 retranslocation in a central African rain forest dominated by ectomycorrhizal trees.  
614 *New Phytologist*, **148**, 493-510.
- 615 Cuéllar, T., Pascaud, F., Verdeil, J.L., Torregrosa, L., Adam-Blondon, A.F.A., Thibaud,  
616 J.B., Santenac, H. & Gaillard, I. (2010) A grapevine shaker inward K<sup>+</sup> channel  
617 activated by the calcineurin B-like calcium sensor 1-protein kinase CIPK23  
618 network is expressed in grape berries under drought stress conditions. *The Plant*  
619 *Journal*, **61**, 58-69.
- 620 Da Silva, I.R., Furtini, Neto, A.E., Fernandes, L.A., Curi, N. & Do Vale, F.B. (2000)  
621 Formas, relação quantidade/intensidade e biodisponibilidade de potássio em  
622 diferentes latossolos. *Pesquisa agropecuária brasileira*, **35**, 2065-2073.
- 623 Dassonville, N., Vanderhoeven, S., Vanparys, V., Hayez, M., Gruber, W. & Meerts, P.  
624 (2008) Impacts of alien invasive plants on soil nutrients are correlated with initial  
625 site conditions in NW Europe. *Oecologia*, **157**, 131-140.
- 626 Del Campo, A.D., Hermoso, J., Flors, J., Lidón, A. & Navarro-Cerrillo, R.M. (2011)  
627 Nursey location and potassium enrichment in Aleppo pine stock 2. Performance  
628 under real and hydrogel-mediated drought conditions. *Forestry*, **84**, 235-245.
- 629 Denslow, J.S., Space, J.C. & Thomas, P.A. (2009) Invasive Exotic Plants in the  
630 Tropical Pacific Islands: Patterns of Diversity. *Biotropica*, **41**, 162-170.
- 631 Duchesne, L., Ouimet, R., Camire, C. & Houle, D. (2001) Seasonal nutrient transfers by  
632 foliar resorption, leaching, and litter fall in a northern hardwood forest at Lake  
633 Clair watershed, Quebec, Canada. *Canadian Journal of Forest Research*, **31**, 333-  
634 344.
- 635 Dunlop, J., Glass, A.D.M. & Tomkins, B.D. (1979) Regulation of K<sup>+</sup> uptake by  
636 ryegrass and white clover roots in relation to their competition for potassium. *New*  
637 *Phytologist*, **83**, 365-370.
- 638 Egilla, J.N., Davies, Jr F.T. & Boutton, T.W. (2005) Drought stress influences leaf  
639 water content, photosynthesis, and water-use efficiency of *Hibiscus rosa-sinensis*  
640 at three potassium concentrations. *Photosynthetica*, **43**, 135-140.
- 641 El-Mesbahi, M.N., Azcón, R., Ruiz-Lozano, J.M. & Aroca, R. (2012) Plant potassium  
642 content modifies the effects of arbuscular mycorrhizal symbiosis on root  
643 hydraulic properties in maize plants. *Mycorrhiza*, **22**, 555-564.
- 644 Elser, J.J., Fagan, W.F., Denno, R.F., Dobberfuhl, D.R. Folarin, A., Huberty, A.,  
645 Interlandi, S., Kilham, S.S., McCauley, E., Schulz, K.L., Siemann, E.H. & Sterner,  
646 R.W. (2000) Nutritional constraints in terrestrial and freshwater ecosystems.  
647 *Nature*, **408**, 578-580.
- 648 Elumalai, R.P., Nagpal, P. & Reed, J.W. (2002) A mutation in the Arabidopsis  
649 KTA/KUP2 potassium transporter gene affects shoot cell expansion. *The Plant*  
650 *Cell*, **14**, 119-131.

- 651 Eurostat (2009) Statistical books. Panorama of Energy. Energy statistics to support EU  
652 policies and solutions. Edited by: Office for Official Publications of the European  
653 Communities, Luxembourg.
- 654 FAO (2011) Current world fertilizer trends and outlook to 2015. Roma.
- 655 Fayyaz, P., Etemadi, E., Julaiee-Manesh, N. & Zolfaghari, R. (2013) Sodium and  
656 potassium allocation under drought stress in Atlas mastic tree (*Pistacia atlantica*  
657 subsp. *mutica*). *Forest-Biogeosciences and Forestry*, **6**, 90-94.
- 658 Ferm, M. & Hultberg, H. (1999) Dry deposition and internal circulation of nitrogen,  
659 sulphur and base cations to a coniferous forest. *Atmospheric Deposition*, **33**,  
660 4421-4430.
- 661 Fernandez, F.G., Brouder, S.M., Volenec, J.J., Beyrouy, C.A. & Hoyum, R. (2011)  
662 Soybean shoot and root response to localized water and potassium in a split-pot  
663 study. *Plant and Soil*, **344**, 197-212.
- 664 Fife, D.N., Nambiar, E.K.S. & Saur, E. (2008) Retranslocation of foliar nutrients in  
665 evergreen tree species planted in a Mediterranean environment. *Tree Physiology*,  
666 **28**, 187-196.
- 667 Filella, I. & Peñuelas, J. (2003) Indications of hydraulic lift by *Pinus halepensis* and its  
668 effects on the water relations of neighbour shrubs. *Biologia Plantarum*, **47**, 209-  
669 214.
- 670 Florez-Florez, C.P., Leon-Pelaez, J.D., Osorio-Vega, N.W. & Restrepo-Llano, M.F.  
671 (2013) Nutrient dynamics in forest plantations of *Azadirachta indica* (Meliaceae)  
672 established for restoration of degraded lands in Colombia. *Revista de Biología*  
673 *Tropical*, **61**, 515-529.
- 674 Friedland, A.J. & Miller, E.K. (1999) Major-element cycling in a high-elevation  
675 Adirondack forest: patterns and changes, 1986-1996. *Ecological Applications*, **9**,  
676 958-967.
- 677 Funk, J.L. & Vitousek, P.M. (2007) Resource-use efficiency and plant invasion in low-  
678 resource systems. *Nature*, **446**, 1079-1081.
- 679 Fyllas, N.M., Patino, S., Baker, T.R., Nardoto, G.B., Martinelli, L.A., Quesada, C.A.,  
680 Paiva, R., Schwarz, M., Horna, V., Mercado, L.M., Santos, A., Arroyo, L.,  
681 Jimenez, E.M., Luizao, F.J., Neill, D.A., Silva, N., Prieto, A., Rudas, A., Silveira,  
682 M., Vieira, I.C.G., Lopez-Gonzalez, G., Malhi, Y., Phillips, O.L. & Lloyd, J.  
683 (2009) Basin-wide variations in foliar properties of Amazonian forest: phylogeny,  
684 soils and climate. *Biogeosciences*, **6**, 2677-2708.
- 685 Gabarrón-Galeote, M.A., Martínez-Murillo, J.F. & Ruiz-Sinoga, J.D. (2012) Relevant  
686 effects of vegetal cover and litter on the soil hydrological response of two  
687 contrasting Mediterranean hill slopes at the end of the dry season (south of Spain).  
688 *Hydrological Proceedings*, **26**, 1729-1738.
- 689 Gajdanowicz, P., Michard, E., Sandmann, M., Rocha, M., Correa, L.G.G., Ramirez-  
690 Aguilar, S.J., Gomez-Porrás, J.L., Gonzalez, W., Thibaud, J.B., van Dongen, J.T.  
691 & Dreyer, I. (2011) Potassium (K<sup>+</sup>) gradients serve as a mobile energy source in  
692 plant vascular tissues. *Proceedings of the National Academy of Sciences USA*, **108**,  
693 864-869.
- 694 Gallardo, J.F., Martín, A. & Moreno, G. (1999) Nutrient efficiency and resorption in  
695 *Quercus pyrenaica* oak coppices under different rainfall regimes of the Sierra de  
696 gata mountains (central western Spain). *Annals of Forest Sciences*, **56**, 321-331.
- 697 Ge, T.D., Sun, N.B., Bai, L.P., Tong, C.L. & Sui, F.G. (2012) Effects of drought stress  
698 on phosphorus and potassium uptake dynamics in summer maize (*Zea mays*)  
699 through the growth cycle. *Acta Physiologiae Plantarum*, **34**, 2179-2186.

- 700 Gélinas, Y., Lucotte, M. & Schmit, J.P. (2000) History of the atmospheric deposition of  
701 major and trace elements in the industrialized St. Lawrence Valley, Quebec,  
702 Canada. *Atmospheric Environment*, **34**, 1797-1810.
- 703 Glab, T. & Gondek, K. (2014) The influence of soil compactation and fertilization on  
704 physic-chemical properties of mollic fluvisol under red clover/grass mixture.  
705 *Geoderma*, **226**, 204-212.
- 706 Golobocanin, D., Zujic, A., Milenkovic, A. & Miljevic, N. (2009) Precipitation  
707 composition and wet deposition temporal pattern in Central Serbia for the period  
708 from 1998 to 2004. *Environmental Monitoring and Assessment*, **142**, 185-198.
- 709 Gómez-Aparicio, L., Gómez, J.M., Zamora, R. & Boettinger, J.L. (2003) Canopy vs.  
710 soil effects of shrubs facilitating tree seedlings in Mediterranean montane  
711 ecosystems. *Journal of Vegetation Science*, **16**, 191-198.
- 712 Greenwood, N.N. & Earnshaw, A. (1997) *Chemistry of The Elements*. (Second Edition).  
713 Butterworth-Heinemann, Oxford.
- 714 Hagen-Thorn, A., Varmagiryte, I., Nihlgard, B. & Armolatis, K. (2006) Autumn  
715 nutrient resorption and losses in four deciduous forest tree species. *Forest  
716 Ecology and Management*, **228**, 33-39.
- 717 Hardman, C.J., Williams, S., Manco, B.N. & Hamilton, M.A. (2012) Predicting the  
718 potential threat of *Casuarina equisetifolia* to three endemic plant species on the  
719 Turks and Caicos Islands. *Oryx*, **46**, 204-212.
- 720 Harvey, H.P. & van den Driessche, R. (1999) Nitrogen and potassium effects on xylem  
721 cavitation and water-use efficiency in poplars. *Tree Physiology*, **19**, 943-950.
- 722 He, Z.L., Alva, A.K., Calvert, D.V., Li, Y.C. & Banks, D.J. (1999) Effects of nitrogen  
723 fertilization of grapefruit trees on soil acidification and nutrient availability in a  
724 Rivera fine sand. *Plant and Soil*, **206**, 11-19.
- 725 Hedin, L.O., Vitousek, P.M. & Matson, P.A. (2003) Nutrient losses over four million  
726 years of tropical forest development. *Ecology*, **84**, 2231-2255.
- 727 Hemmisaari, H.S. (1992) Nutrient retranslocation in three *Pinus sylvestris* stands.  
728 *Forest Ecology and Management*, **51**, 347-367.
- 729 Hirsch, R.E., Lewis, B.D., Spalding, E.P. & Sussman, M.R. (1998) A role for the AKT1  
730 potassium channel in plant nutrition. *Science*, **280**, 918-921.
- 731 Hoa, N.M., Janssen, B.H., Oenema, O. & Dobermann, A. (2006) Comparison of partial  
732 and complete soil K budgets under intensive rice cropping in the Mekong Delta,  
733 Vietnam. *Agriculture Ecosystems and Environment*, **116**, 121-131.
- 734 Holloway, J.A.M. & Dahlgren, R.A. (2002) Nitrogen in rock: Occurrences and  
735 biogeochemical implications. *Global Biogeochemical Cycles*, **16**, 1118.
- 736 Houlton, B.Z., Wang, Y.P., Vitousek, P.M., Christopher, B. & Field, C.B. (2008) A  
737 unifying framework for dinitrogen fixation in the terrestrial biosphere. *Nature*,  
738 **454**, 327-330.
- 739 Hultine, K.R., Scott, R.L., Cable, W.L., Goodrich, D.C. & Williams, D.G. (2004)  
740 Hydraulic redistribution by a dominant, warm-desert phreatophyte: seasonal  
741 patterns and response to precipitation pulses. *Functional Ecology*, **18**, 530-538.
- 742 Huang, S., Pan, X.H., Sun, Y.N., Zhang, Y., Hang, X.N., Yu, X.C. & Zhang, W.J.  
743 (2013) Effects of long-term fertilization on the weed growth and community  
744 composition in a double-rice ecosystem during the fallow period. *Weed Biology  
745 and Management*, **13**: 10-18.
- 746 <http://nadp.sws.uiuc.edu/> (2012) National Atmospheric Deposition Program. U.S.  
747 Department of Agriculture.
- 748 IPCC (2007) Climate Change 2007: The Physical Science Basis. Contribution of  
749 Working Group I. p. 1-32 In: S. Solomon, et al.. (ed.). Fourth Assessment Report

- 750 of the Intergovernmental Panel on Climate Change. Cambridge University Press,  
751 Cambridge, United Kingdom and New York, NY, USA.
- 752 Jobbágy, E.G. & Jackson, R.B. (2001) The distribution of soil nutrients with depth:  
753 global patterns and the imprint of plants. *Biogeochemistry*, **53**, 51-77.
- 754 Kadar, I. & Foldesi, D. (2001) Mineral fertilization of poppy (*Papaver somniferum* L.)  
755 on calcareous loamy chernozem soil. I. *Nevenytermeles*, **50**, 453-465.
- 756 Kanai, S., Moghaieb, R.E., El-Shemy, H.A., Panigrahi, R., Mohapatra, P.K., Ito, J.,  
757 Nguyen, N.T., Saneoka, H. & Fujita, K (2011) Potassium deficiency affects water  
758 status and photosynthetic rate of the vegetative sink in green house tomato prior to  
759 its effects on source activity. *Plant Science*, **180**, 368-374.
- 760 Kerkhoff, A.J., Enquist, B.J., Elser, J.J. & Fagan, W.F. (2005) Plant allometry,  
761 stoichiometry and the temperature-dependence of primary productivity. *Global*  
762 *Ecology and Biogeography*, **14**, 585-598.
- 763 Kopacek, J., Turek, J., Hejzlar, J. & Santruckova, H. (2009) Canopy leaching of  
764 nutrients and metals in a mountain spruce forest. *Atmospheric Environment*, **43**,  
765 5443-5453.
- 766 Klumpp, A., Domingos, M. & Klumpp, G. (2002) Foliar nutrient contents in tree  
767 species of the Atlantic rain forest as influenced by air pollution from the  
768 industrial complex of Cubatao, SE-Brazil. *Water Air and Soil Pollution*, **133**, 315-  
769 333.
- 770 Langley, J.A. & Megonigal, J.P. (2010) Ecosystem response to elevated CO<sub>2</sub> levels  
771 limited by nitrogen-induced plant species shift. *Nature*, **466**, 96-99.
- 772 Lawniczak, A.E., Güsewell, S. & Verhoeven, J.T.A. (2009) Effect of N:K supply ratios  
773 on the performance of three grass species from herbaceous wetlands. *Basic and*  
774 *Applied Ecology*, **10**, 715-725.
- 775 Lee, J.A. & Caporn, S.J.M. (1998) Ecological effects of atmospheric reactive nitrogen  
776 deposition on seminatural terrestrial ecosystems. *New Phytologist*, **139**, 127-134.
- 777 Lee, J.E., Oliveira, R.S., Dawson, T.E. & Fung, I. (2005) Root functioning modifies  
778 seasonal climate. *Proceedings of the National Academy of Sciences USA*, **102**,  
779 17576-17581.
- 780 Leigh, R.A. & Jones, R.G.W. (1984) A hypothesis relating critical potassium  
781 concentrations for growth to the distribution and functions of this ion in the plant-  
782 cell. *New Phytologist*, **97**, 1-13.
- 783 Leigh, R.A. (2001) Potassium homeostasis and membrane transport. *Journal of Plant*  
784 *Nutrition and Soil Science*, **164**, 193-198.
- 785 Lé Quére, C., Raupach, M.R., Canadell, J.G., Marland, G., Bopp, L., Ciais, P., Conway,  
786 T.J., Doney, S.C., Feely, R.A., Foster, P., Friedlingstein, P., Gurney, K.,  
787 Houghton, R.A., House, J.I., Huntingford, C., Levy, P.E., Lomas, M.R., Majkut,  
788 J., Metzl, N., Ometto, J.P., Peters, G.P., Prentice, I.C., Randerson, J.T., Running,  
789 S.W., Sarmiento, J.L., Schuster, U., Sitch, S., Takahashi, T., Viovy, N., van der  
790 Werf, G.R. & Eoodward, F.I. (2009) Trends in the sources and sinks of carbon  
791 dioxide. *Nature Geoscience*, **2**, 831-836.
- 792 Levi, A., Paterson, A.H., Cakmak, I. & Saranga, Y. (2011) Metabolite and mineral  
793 analyses of cotton near-isogenic lines introgressed with QTLs for productivity and  
794 drought-related traits. *Physiologia Plantarum*, **141**, 265-275.
- 795 Liu, H.Y., Sun, W.N., Su, W.A. & Tang, Z.C. (2006) Co-regulation of water channels  
796 and potassium channels in rice. *Physiologia Plantarum*, **128**, 58-69.
- 797 Lucas, R.W., Klaminder, J., Futter, M.N., Bishop, K.H., Egnell, G., Laudon, H. &  
798 Högberg, P. (2011) A meta-analysis of the effects of nitrogen additions on base

- 799 cations: implications for plants, soils, and streams. *Forest Ecology and*  
800 *Management*, **262**, 95-104.
- 801 Maathuis, F.J.M. & Sanders, D. (1994) Mechanism of high-affinity potassium uptake in  
802 roots of *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences*  
803 *USA*, **91**, 9272-9276.
- 804 MacDonald, I.A.W., Thebaud, C., Strahm, W.A. & Strasberg, D. (1991) Effects of alien  
805 plant invasions on native vegetation remnants on La Reunion (Mascarene-Islands,  
806 Indian-Ocean). *Environmental Conservation*, **18**, 51-61.
- 807 Mack, M.C., Schuur, E.A.G., Bret-Harte, M.S., Shaver, G.R. & Chapin III, F.S. (2004)  
808 Ecosystem carbon storage in arctic tundra reduced by long-term nutrient  
809 fertilization. *Nature*, **431**, 440-443.
- 810 Magen, H. (2008) Balanced crop nutrition: fertilizing for crop and food quality. *Turkish*  
811 *Journal of Agriculture*, **32**, 183-193.
- 812 Marchi, G., Silva, V.A., Guilherme, L.R.G., Lima, J.M., Nogueira, F.D. & Guimaraes,  
813 P.T.G. (2012) Potassium extractability from soils of Brazilian coffee regions.  
814 *Bioscience Journal*, **28**, 913-919.
- 815 Merino-Martin, L., Moreno-de las Heras, M., Pérez-Domingo, S., Espigarea, T. &  
816 Nicolau, J.M. (2012) Hydrological heterogeneity in Mediterranean reclaimed  
817 slopes: runoff and sediment yield at the patch and slope scales along a gradient of  
818 overland flow. *Hydrological Earth Systems Science*, **16**, 1305-1320.
- 819 Milla, R., Castro-Díaz, P., Maestro-Martínez, M. & Montserrat-Martí, G. (2005)  
820 Relationships between phenology and the remobilization of nitrogen, phosphorus  
821 and potassium in branches of eight Mediterranean evergreens. *New Phytologist*,  
822 **168**, 167-178.
- 823 Mitchell, M.J., Raynal, D.J. & Driscoll, C.T. (1996) Biogeochemistry of a forested  
824 watershed in the central Adirondack Mountains: Temporal changes and mass  
825 balances. *Water, Air and Soil Pollution*, **88**, 355-369.
- 826 Mitchell, A.D. & Smethurst, P.J. (2008) Base cation availability and leaching after  
827 nitrogen fertilization of a eucalypt plantation. *Australian Journal of Soil*  
828 *Research*, **46**, 445-454.
- 829 Monty, M.L.F., Florens, F.B.V. & Baider, C. (2013) Invasive alien plants elicit reduced  
830 production of flowers and fruits in various native forest species on the tropical  
831 island of Mauritius (Mascarenes, Indian Ocean). *Tropical Conservation Science*, **6**,  
832 35-49.
- 833 More, J.D., Camiré, C. & Ouimet, R. (2000) Effects of liming on the nutrition, vigor,  
834 and growth of sugar maple at the Lake Clair watershed, Québec, Canada.  
835 *Canadian Journal of Forest Research*, **30**, 725-732.
- 836 Mulder, C., Ahrestani, F.S., Bahn, M., Bohan, D.A., Bonkowski, M., Griffiths, B.S.,  
837 Guicharnaud, R.A., Kattge, J., Krogh, P.H., Lavorel, S., Lewis, O.T., Mancinelli,  
838 G., Naeem, S., Peñuelas, J., Poorter H., Reich, P.B., Rossi, L., Rusch, G.M.,  
839 Sardans, J. & Wright, I.J. (2013) Connecting the green and brown worlds:  
840 allometric and stoichiometric predictability of above- and below-ground networks.  
841 *Advances in Ecological Research*, **49**, 69-175.
- 842 Muoghalu, J.I. & Oakhumen, A. (2000) Nutrient content of incident rainfall, throughfall  
843 and stemflow in a Nigerian secondary lowland rainforest. *Applied Vegetation*  
844 *Science*, **3**, 181-188.
- 845 Nandwal, A.S., Hooda, A. & Datta, D. (1998) Effect of substrate moisture and  
846 potassium on water relations and C, N and K distribution in *Vigna radiate*.  
847 *Biologia Plantarum*, **41**, 149-153.

- 848 Neiryck, J., Maddelein, D., de Keersmaeker, L., Lust, N. & Muys, B. (1998) Biomass  
849 and nutrient cycling of a highly productive Corsican pine stand on former  
850 heathland in northern Belgium. *Annales des Sciences Forestieres*, **55**, 389-405.
- 851 Nieminen, T. & Helmisaari, H.S. (1996) Nutrient retranslocation in the foliage of *Pinus*  
852 *sylvestris* L. growing along a heavy metal pollution gradient. *Tree Physiology*, **16**,  
853 825-831.
- 854 Navratil, T., Norton, S.A., Fernandez, I.J. & Nelson, S.J. (2010) Twenty-year inter-  
855 annual trends and seasonal variations in precipitation and stream water chemistry  
856 at the Bear Brook Watershed in Maine, USA. *Environmental Monitoring and*  
857 *Assessment*, **171**, 23-45.
- 858 Nowak, C.A., Downard, R.B. & White, E.H. (1991) Potassium trends in red pine  
859 plantations at pack-forest, New York. *Soil Science Society of America Journal*, **55**,  
860 847-850.
- 861 Oddo, E., Inzerillo, S., La Bella, F., Grisafi, F., Salleo, S. & Nardini, A. (2011) Short-  
862 term effects of potassium fertilization on the hydraulic conductance of *Laurus*  
863 *nobilis* L. *Tree Physiology*, **31**, 131-138.
- 864 Olde Venterink, H., Kardel, I., Kotowski, W., Peeters, W. & Wassen, M.J. (2009)  
865 Long-term effects of drainage and hay-removal on nutrient dynamics and  
866 limitation in the Biebrza mires, Poland. *Biogeochemistry*, **93**, 235-252.
- 867 Osakabe, Y., Arinaga, N., Umezawa, T., Katsura, S., Nagamachi, K., Tanaka, H.,  
868 Ohiraki, H., Yamada, K., Seo, S.U., Abo, M., Yoshimura, E., Shinozaki, K. &  
869 Yamaguchi-Shinozaki, K. (2013) Osmotic stress responses and plant growth  
870 controlled by potassium transporters in *Arabidopsis*. *Plant Cell*, **25**, 609-624
- 871 Ozbucak, T.B., Kutbay, H.G., Kilic, D., Korkmaz, H., Bilgin, A., Yalcin, E. & Apaydin,  
872 Z. (2008) Foliar resorption of nutrients in selected sympatric tree species in  
873 gallery forest (Black Sea region). *Polish Journal of Ecology*, **56**, 227-237.
- 874 Peng, R.H., Fang, C.M., Li, B. & Chen, J.K. (2011) *Spartina alterniflora* invasion  
875 increases soil inorganic nitrogen pools through interactions with tidal subsides in  
876 the Yangtze Estuary, China. *Oecologia*, **165**, 797-807.
- 877 Peñuelas, J., Sardans, J., Rivas-Ubach, A. & Jansens, I. (2012) The human induced  
878 imbalance between C, N and P in Earth's life-system. *Global Change Biology*,  
879 **189**, 5-8.
- 880 Peñuelas, J., Sardans, J., Estiarte, M., Ogaya, R., Carnicer, J., Coll, M., Barbeta, A.,  
881 Rivas-Ubach, A., Llusia, J., Garbuslsky, M., Filalla, I. & Jump, A.S. (2013a)  
882 Evidence of current impact of climate change on life: a walk from genes to the  
883 biosphere. *Global Change Biology*, **19**, 2303-2338.
- 884 Peñuelas, J., Poulter, B., Sardans, J., Ciais, P., van der Velde, M., Bopp, L., Boucher, O.,  
885 Godderis, Y., Hinsinger, P., Llusia, J., Nardin, E., Vicca, S., Obersteiner, M. &  
886 Janssens, I.A. (2013b) Human-induced nitrogen-phosphorus imbalances alter  
887 natural and managed ecosystems across the globe. *Nature Communications*, **4**,  
888 3934.
- 889 Pieters, A. & Baruch, Z. (1997) Soil depth and fertility effects on biomass and nutrient  
890 allocation in jaragua grass. *Journal of Range Management*, **50**, 268-273.
- 891 Poor, N., Pollman, C., Tate, P., Begum, M., Evans, M. & Campbell, S. (2006) Nature  
892 and magnitude of atmospheric fluxes of total inorganic nitrogen and other  
893 inorganic species to the Tampa Bay watershed, FL, USA. *Water Air and Soil*  
894 *Pollution*, **170**, 267-283.
- 895 Potter, C.S., Ragsdale, H.L. & Swank, W.T. (1991) Atmospheric deposition and foliar  
896 leaching in a regenerating southern Appalachian forest canopy. *Journal of*  
897 *Ecology*, **79**, 97-115.

- 898 Prathibha, P., Kothai, P., Saradhi, I.V., Pandit, G.G. & Puranik, V.D. (2010) Chemical  
899 characterization of precipitation at a coastal site in Trombay, Mumbai, India.  
900 *Environmental Monitoring and Assessment*, **168**, 45-53.
- 901 Pyo, Y.J., Gierth, M., Schroeder, J.I. & Cho, M.H. (2010) High-affinity K<sup>+</sup> transport in  
902 *Arabidopsis*: AtHAK5 and AKT1 are vital for seedling establishment and post  
903 germination growth under low-potassium conditions. *Plant Physiology*, **153**, 863-  
904 875.
- 905 Quennehen, B., Schwarzenboeck, A., Matsuki, A., Burkhart, J.F., Stohl, A., Ancellet, G.  
906 & Law, K.S. (2012) Anthropogenic and forest fire pollution aerosol transported to  
907 the Arctic: observations from the POLARCAT-France spring campaign.  
908 *Atmospheric Chemistry and Physics*, **12**, 6437–6454.
- 909 Rao, S.R., Qayyum, A., Razzaq, A., Ahmad, M., Mahmood, I. & Sher, A. (2012) Role  
910 of foliar application of salicylic acid and L-tryptophan in drought tolerance of  
911 maize. *Journal of Animal and Plant Sciences*, **22**, 768-772.
- 912 Reich, P.B. & Oleksyn, J. (2004) Global patterns of plant leaf N and P in relation to  
913 temperature and latitude. *Proceedings National Academy of Sciences USA*, **101**,  
914 11001-11006.
- 915 Rivas-Ubach, A., Sardans, J., Pérez-Trujillo, M., Estiarte, M. & Peñuelas, J. (2012)  
916 Strong relationship between elemental stoichiometry and metabolome in plants.  
917 *Proceedings of the National Academy of Sciences USA*, **109**, 4181-4186.
- 918 Rivas-Ubach, A., Gargallo-Garriga, A., Sardans, J., Oravec, M., Pérez-Trujillo, M.,  
919 Ogaya, R., Urban, O. & Peñuelas, J. (2014) Drought stress enhances folivory rates  
920 by shifting foliar metabolomes. *New Phytologist*, **202**, 874-885..
- 921 Roberts, T.L. (2008) Global potassium reserves and potassium fertilizer use. IPNI  
922 (International Plant Nutrition Institute). Symposium-Global Nutrient Cycling.  
923 Georgia. USA.
- 924 Robinson, A.L., Scanlon, T.M., Cosby, B.J., Webb, J.R. & Galloway, J.N. (2013) Roles  
925 of sulfate adsorption and base cation supply in controlling the chemical response  
926 of streams of western Virginia to reduced acid deposition. *Biogeochemistry*, **116**,  
927 119-130.
- 928 Römheld, V. & Kirkby, E.S. (2010) Research on potassium in agriculture: needs and  
929 prospects. *Plant and Soil*, **335**, 155-180.
- 930 Ruan, J., Wu, X. & Härdter, R. (1997) The interaction between soil water regime and  
931 potassium availability on the growth of tea. *Communications in Soil Science and*  
932 *Plant Analysis*, **28**, 89-98.
- 933 Ruoho-Airola, T. & Salminen, K. (2003) Trends in the base cation deposition in Finland.  
934 *Air Pollution XI. Advances in Air Pollution Series*, **13**, 273-282.
- 935 Ryan, J. & Sommer, R. (2012) Soil fertility and crop nutrition research at an  
936 international center in the Mediterranean region: achievements and future  
937 perspective. *Archives of Agronomy and Soil Science*, **58**, S41-S54.
- 938 Ryan, J., Sommer, R. & Ibrikci, H. (2012) Fertilizer best management practices: a  
939 perspective from the dryland west Asia-North Africa region. *Journal of Agronomy*  
940 *and Crop Science*, **198**, 57-67.
- 941 Ryel, R.J., Leffler, A.J., Peek, M.S., Ivans, C.Y. & Caldwell, M.M. (2004) Water  
942 conservation in *Artemisia tridentate* through redistribution of precipitation.  
943 *Oecologia*, **141**, 335-345.
- 944 Salehi, A., Ghorbanzadeh, N. & Salehi, M. (2013) Soil nutrient status, nutrient return  
945 and retranslocation in poplar species and clones in northern Iran. *Forest-*  
946 *Biogeosciences and Forestry*, **6**, 336-341.

- 947 Sardans, J. & Peñuelas, J. (2007) Drought changes phosphorus and potassium  
948 accumulation patterns in an evergreen Mediterranean forest. *Functional Ecology*,  
949 **21**, 191-201.
- 950 Sardans, J. & Peñuelas, J. (2012) The role of plants in the effects of global change on  
951 nutrient availability and stoichiometry in the Plant-Soil system. *Plant Physiology*,  
952 **160**, 1741-1761.
- 953 Sardans, J., Peñuelas, J. & Rodà, F. (2005) Changes in nutrient use efficiency, status  
954 and retranslocation in young post-fire regeneration *Pinus halepensis* in response  
955 to sudden N and P input, irrigation and removal of competing vegetation. *Trees*  
956 *Structure and Function*, **19**, 233-250.
- 957 Sardans, J., Rodà, F. & Peñuelas, J. (2006) Effects of a nutrient pulse supply on nutrient  
958 status of the Mediterranean trees *Quercus ilex* subsp. *ballota* and *Pinus halepensis*  
959 on different soils and under different competitive pressure. *Trees Structure and*  
960 *Function*, **20**, 619-632.
- 961 Sardans, J., Peñuelas, J., Prieto, P. & Estiarte, M. (2008) Drought and warming induced  
962 changes in P and K concentration and accumulation in plant biomass and soil in a  
963 Mediterranean shrubland. *Plant Soil*, **306**, 261-271.
- 964 Sardans, J., Rivas-Ubach, A. & Peñuelas, J. (2011) Factors affecting nutrient  
965 concentration and stoichiometry of forest trees in Catalonia (NE Spain). *Forest*  
966 *Ecology and Management*, **262**, 2024-2034.
- 967 Sardans, J., Peñuelas, J., Coll, M., Vayreda, J. & Rivas-Ubach, A. (2012a)  
968 Stoichiometry of potassium is largely determined by water availability and growth  
969 in Catalanian forests. *Functional Ecology*, **26**, 1077-1089.
- 970 Sardans, J., Rivas-Ubach, A. & Peñuelas, J. (2012b) The elemental stoichiometry of  
971 aquatic and terrestrial ecosystems and its relationships with organismic lifestyle  
972 and ecosystem structure and function: a review and perspectives. *Biogeochemistry*,  
973 **111**, 1-39.
- 974 Sardans, J., Rivas-Ubach, A. & Peñuelas, J. (2012c) The C:N:P stoichiometry of  
975 organisms and ecosystems in a changing world: A review and perspectives.  
976 *Perspectives in Plant Ecology, Evolution and Systematics*, **14**, 33-47.
- 977 Sardans, J. & Peñuelas, J. (2013) Plant-soil interactions in Mediterranean forest and  
978 shrublands: impacts of climatic change. *Plant and Soil*, **365**, 1-33.
- 979 Sardans, J. & Peñuelas, J. (2014) Hydraulic redistribution by plants and nutrient  
980 stoichiometry. Shifts under global change. *Ecohydrology*, **7**, 1-20.
- 981 Schachtman, D.P. & Schroeder, J.I. (1994) Structure and transport mechanism of a  
982 high-affinity potassium uptake transporter from higher plants. *Nature*, **370**, 655-  
983 658.
- 984 Schimel, D.S., House, J.I., Hibbard, K.A. Bousquet, P., Ciais, P., Peylin, P., Braswell,  
985 B.H., Apps, M.J., Baker, D., Bondeau, A., Canadell, J., Churkina, G., Cramer, W.,  
986 Denning, A.S., Field, C.B., Friedlingstein, P., Goodale, C., Heimann, M.,  
987 Houghton, R.A., Melillo, J.M., Moore III, B., Murdiyarso, D., Noble, I., Pacala,  
988 S.W., Prentice, I.C., Raupach, M.R., Rayner, P.J., Scholes, R.J., Steffen, W.L., &  
989 Wirth, C. (2001) Recent patterns and mechanisms of carbon exchange by  
990 terrestrial ecosystems. *Nature*, **414**, 169-172.
- 991 Schreeg, L.A., Mack, M.C. & Turner, B.L. (2013) Nutrient-specific patterns of leaf  
992 litter across 41 lowland tropical woody species. *Ecology*, **94**, 94-105.
- 993 Schroeder, J.I. & Fang, H.H. (1991) Inward-rectifying K<sup>+</sup> channels in guard cells  
994 provide a mechanism for low-affinity K<sup>+</sup> uptake. *Proceedings of the National*  
995 *Academy of Sciences USA*, **88**, 11583-11587.

- 996 Sharifi, M., Cheema, M., McVicar, K., LeBlanc, L. & Fillmore, S. (2013) Evaluation of  
 997 liming properties and potassium bioavailability of three Atlantic Canada Wood  
 998 ash sources. *Canadian Journal of Plant Science*, **93**, 1209-1216.
- 999 Silva, V.A., Marchi, G., Guimaraes Guilherme, L.R., de Lima, J.M., Dias Nogueira, F.  
 1000 & Gontijo Guimaraes, P.T. (2008) Kinetics of K release from soils of Brazilian  
 1001 coffee regions: effect of organic acids. *Revista Brasileira de Ciencia do Solo*, **28**:  
 1002 597-603.
- 1003 Singer, A. (1989) Illite in the hot-aridic soil environment. *Soil Science*, **147**, 126-133.
- 1004 Singh, K. & Bansai, S.K. (2009) Potassium indexing of crops grown on eight  
 1005 Benchmark soil series of India. *Communications in Soil Science and Plant  
 1006 Analysis*, **40**, 1369-1379.
- 1007 Smith, D.M., Jackson, N.A., Roberts, J.M. & Ong, C.K. (1999) Reverse flow of sap in  
 1008 tree roots and downward siphoning of water by *Grevillea robusta*. *Functional  
 1009 Ecology*, **13**, 256-264.
- 1010 Soil Survey Staff (1999) Soil Taxonomy. A Basic System of Soil Classification for  
 1011 Making Classification for making and interpreting Soil Surveys.
- 1012 Sterner, R.W. & Elser, J.J. (2002) Ecological stoichiometry: the biology of elements  
 1013 from molecules to the biosphere. Princeton University Press, Princeton.
- 1014 Stone, E.L. & Kszystyniak, R. (1977) Conservation of potassium in the *Pinus resinosa*  
 1015 ecosystem. *Science*, **198**, 192-194.
- 1016 Su, H., Gollack, D., Katsuhara, M., Zhao, C. & Bohnert, H.J. (2001) Expression and  
 1017 stress-dependent induction of potassium channel transcripts in the common ice  
 1018 plant. *Plant Physiology*, **125**, 604-614.
- 1019 Sutton, M.A., Howard, C.M., Erisman, J.W., Billen, G., Bleeker, A., Grennfelt, P.,  
 1020 Grinsven, H.V. & Grizzetti, B. (eds) (2011) The European Nitrogen Assessment.  
 1021 Cambridge University Press.
- 1022 Szczerba, M.W., Britto, D.T. & Kronzucker, H.J. (2009) K<sup>+</sup> transport in plants:  
 1023 physiology and molecular biology. *Journal of Plant Physiology*, **166**, 447-466.
- 1024 Talbott, L.D. & Zeiger, E. (1996) Central roles for potassium and sucrose in guard-cell  
 1025 osmoregulation. *Plant physiology*, **111**, 1051-1057.
- 1026 Tartachnyk, I.I. & Blanke, M.M. (2004) Effect of delayed fruit harvest on  
 1027 photosynthesis, transpiration and nutrient remobilization of apple leaves. *New  
 1028 Phytologist*, **164**, 441-450.
- 1029 Teixeira, P.C., Gonçalves, J.L.M., Arthur, J.C. & Dezordi, C. (2008) *Eucalyptus* sp.  
 1030 seedling response to potassium fertilization and soil water. *Ciencia Forestal*, **18**,  
 1031 47-63.
- 1032 Tilman, E.A., Tilman, D., Crawley, M.J. & Johnston, A.E. (1999) Biological weed  
 1033 control via nutrient competition: potassium limitation of dandelions. *Ecological  
 1034 Monographs*, **9**, 103-111.
- 1035 Tomlinson, K.W., Pooter, L., Sterck, F.J., Borghetti, F., Ward, D., de Bie, S. & van  
 1036 Langevelde, F. (2013) Leaf adaptations of evergreen and deciduous trees of semi-  
 1037 arid and humid savannas on three continents. *Journal of Ecology*, **101**, 430-440.
- 1038 Traveset, A., Kueffer, C. & Daehler, C.C. (2014) Global and regional nested patterns of  
 1039 non-native invasive floras on tropical islands. *Journal of Biogeography*, **41**, 823-  
 1040 832.
- 1041 Travlos, I.S., Liakopoulos, G., Karabourniotis, G., Fasseas, C. & Karamanos, A.J. (2008)  
 1042 Ciscadian leaflet movements of *Tylosema esculatum* (Burch) A. schreib, and the  
 1043 abolishment of these diurnal movements by potassium deficiency. *Journal of Arid  
 1044 Environments*, **72**, 1745-1750.

- 1045 Trémolières, M., Schnitzler, A., Sánchez-Pérez, J.M. & Schmitt, D. (1999) Changes in  
 1046 foliar nutrient content and resorption in *Fraxinus excelsior* L., *Ulmus minor* Mill.  
 1047 and *Clematis vitalba* L. after prevention of floods. *Annals of Forest Science*, **56**,  
 1048 641-650.
- 1049 Tripler, C.E., Kaushal, S.S., Linkens, G.E. & Walter, M.T. (2006) Patterns in potassium  
 1050 dynamics in forest ecosystems. *Ecology Letters*, **9**, 451-466.
- 1051 Tzvetkova, N. & Hadjiivanova, C.H. (2006) Chemical composition and biochemical  
 1052 changes in Needles of Scots pine (*Pinus sylvestris* L.) stands at different stages of  
 1053 decline in Bulgaria. *Trees Structure and Function*, **20**, 405-409.
- 1054 Urban, R.C., Lima-Souza, M., Caetano-Silva, L., Queiroz, M.E.C., Nogueira, R.F.P.,  
 1055 Allen, A.G., Cardoso, A.A., Held, G. & Campos, M.L.A.M. (2012) Use of  
 1056 levoglucosan, potassium, and water-soluble organic carbon to characterize the  
 1057 origins of biomass-burning aerosols. *Atmospheric Environment*, **61**, 562-569.
- 1058 Van den Driessche, R. (1985) Late-season fertilization mineral nutrient reserves and  
 1059 retranslocation in planted douglass-fir *Pseudotsuga menziessi* seedlings. *Forest  
 1060 Science*, **31**, 485-496.
- 1061 Vanderpol, F. & Traore, B. (1993) Soil nutrient depletion by agricultural production in  
 1062 southern Mali. *Fertilizer Research*, **36**, 79-90.
- 1063 Vegas-Vilarrúbia, T., Baritto, F., López, P., Meléan, G., Ponce, M.E., Mora, L. &  
 1064 Gómez, O. (2010) Tropical Histosols of the lower Orinoco Delta, features and  
 1065 preliminary quantification of their carbon storage. *Geoderma*, **155**, 280-288.
- 1066 Vergutz, L., Manzoni, S., Porporato, A., Novais, R.F. & Jackson, R.B. (2012) Global  
 1067 resorption efficiencies and concentrations of carbon and nutrients in leaves of  
 1068 terrestrial plants. *Ecological Monographs*, **82**, 205-220.
- 1069 Vitousek, P.M. & Sanford, R.L. (1986) Nutrient cycling in moist tropical forest. *Annual  
 1070 Review of Ecological Systems*, **17**, 585-594.
- 1071 Vitousek, P.M. & Howarth, R.W. (1991) Nitrogen limitation on land and in the sea:  
 1072 How can it occur? *Biogeochemistry*, **13**, 87-115.
- 1073 Vitousek, P.M., Porder, S., Houlton, B.Z. & Chadwick, O.A. (2010) Terrestrial  
 1074 phosphorus limitation: mechanisms, implications, and nitrogen-phosphorus  
 1075 interactions. *Ecological Applications*, **20**, 5-15.
- 1076 Vogt RD, Seip HM, Larssen, T., Zhao, D., Xiang, R., Xiao, J., Luo, J. & Zhao, Y.  
 1077 (2006) Potential acidifying capacity of deposition experiences from regions with  
 1078 high NH<sub>4</sub><sup>+</sup> and dry deposition in China. *Science of the Total Environment*, **367**,  
 1079 394-404.
- 1080 Walker, T.W. & Syers, J.K. (1976). The fate of phosphorus during pedogenesis.  
 1081 *Geoderma*, **15**, 1-19.
- 1082 Walker, T.R., Young, S.D., Crittenden, P.D. & Zhang, H. (2003) Anthropogenic metal  
 1083 enrichment of snow and soil in north-eastern European Russia. *Environmental  
 1084 Pollution*, **121**, 11-21.
- 1085 Walna, B., Drzymala, S. & Siepak, J. (2000) The impact of acid rain on potassium and  
 1086 sodium status in typical soils of the Wielkopolski National Park (Poland). *Water  
 1087 Air and Soil Pollution*, **121**, 31-41.
- 1088 Wang, H.Y., Zhou, J.M., Du, C.W. & Chen, X.Q. (2010) Potassium fractions in soils as  
 1089 affected by monocalcium phosphate, ammonium sulfate and potassium chloride  
 1090 application. *Pedosphere*, **20**, 368-377.
- 1091 Wang, M., Zheng, Q., Shen, Q. & Guo, S. (2013) The critical role of potassium in plant  
 1092 stress response. *International Journal of Molecular Science*, **14**, 7370-7390.

- 1093 Waraich, E.A., Ahmad, R., Saifullah & Ehsanullah, M.Y.A. (2011) Role of mineral  
 1094 nutrition in alleviation of drought stress in plants. *Australian Journal of Crop*  
 1095 *Science*, **5**, 764-777.
- 1096 Watmough, S.A. & Dillon, P.J. (2004) Major element fluxes from a coniferous  
 1097 catchment in central Ontario, 1983-1999. *Biogeochemistry*, **67**, 369-398.
- 1098 Watmough, S.A., Aherne, J., Alewell, C., Arp, P., Bailey, S., Clair, T., Dillon, P.,  
 1099 Duchesne, L., Eimers, C., Fernandez, I., Foster, N., Larssen, T., Miller, E.,  
 1100 Mitchell, M. & Page, S. (2005) Sulphate, nitrogen and base cation budgets at 21  
 1101 forested catchments in Canada, the United States and Europe. *Environmental*  
 1102 *Monitoring and Assessment*, **109**, 1-36.
- 1103 Yamashita, N., Ishida, A., Kushima, H. & Tanaka, N. (2000) Acclimatation to sudden  
 1104 increase in light favoring an invasive over native trees in subtropical islands, Japan.  
 1105 *Oecologia*, **125**, 412-419.
- 1106 Yavitt, J.B., Harms, K.E., Garcia, M.N., Mirabello, M.J. & Wright, S.J. (2011) Soil  
 1107 fertility and fine root dynamics in response to 4 years of nutrient (N, P, K)  
 1108 fertilization in a lowland tropical forest, Panama. *Austral Ecology*, **36**, 433-445.
- 1109 Yin, L.M., Wang, L.H. & Liu, B. (2009) Dynamics variation and resorption of nutrient  
 1110 elements in the leaves of *Xanthoceras sorbifolia bunge*. *Bulletin of Botanical*  
 1111 *Research*, **29**, 685-691.
- 1112 Yin, X.M., Rocha, P.S.C.F., Wang, M.L., Zhu, Y.X., Li, L.Y., Song, S.F. & Xia, X.  
 1113 (2011) Rice gene OsDSR-1 promotes lateral root development in *Arabidopsis*  
 1114 under high-potassium conditions. *Journal of Plant Biology*, **54**, 180-189.
- 1115 Zhang, Y., Wang, Z., Zhang, L., Cao, Y., Huang, D. & Tang, K. (2006) Molecular  
 1116 cloning and stress-dependent regulation of potassium channel gene in Chinese  
 1117 cabbage (*Brassica rapa* ssp. *pekinensis*). *Journal of Plant Physiology*, **163**, 968-  
 1118 978.
- 1119 Zhang, S.B., Zhang, J.L., Slik, J.W.F. & Cao, K.F. (2012) Leaf element concentrations  
 1120 of terrestrial plants across China are influenced by taxonomy and the environment.  
 1121 *Global Ecology and Biogeography*, **21**, 809-818.
- 1122 Zhao, D., Oosterhuis, D.M. & Bednarz, C.W. (2001) Influence of potassium deficiency  
 1123 on photosynthesis, chlorophyll content, and chloroplast ultrastructure of cotton  
 1124 plants. *Photosynthetica*, **39**, 103-109.
- 1125 Zheng, S. & Shangguan, Z. (2007) Spatial patterns of leaf nutrient traits of the plants in  
 1126 the Loess Plateau of China. *Trees Structure and Function*, **21**, 357-370.

1127  
 1128  
 1129 Additional references to data sources can be found in Appendix S5 at [URL].

1130  
 1131  
 1132  
 1133

1134 **SUPPORTING INFORMATION**

1135 Additional supporting information may be found in the online version of this article at  
 1136 publisher's website.

1137

1138 **Appendix S1.** Annual atmospheric rates of K deposition ( $\text{kg ha}^{-1} \text{yr}^{-1}$ ).

1139

1140 **Appendix S2.** The limiting role of K on plant growth.

1141

1142 **Appendix S3.** Effects of water availability on ecosystemic K cycle.

1143

1144 **Appendix S4.** Impacts of increases of N supply on ecosystem K cycle.

1145

1146 **Appendix S5.** Plant invasion and soil K availability.

1147

1148 **Appendix S6.** References for the supporting information.

1149

1150

1151 **BIOSKETCH**

1152 Dr. Jordi Sardans and Prof. Josep Peñuelas are researchers focusing on the field of plant  
1153 ecophysiology, biogeochemical cycles and atmosphere-biosphere interactions. JS's  
1154 primary focus is on the elemental stoichiometry relationships with terrestrial ecosystems  
1155 structure and function and in the metabolomic plant responses to abiotic and biotic  
1156 changes. JP's research covers a wide range of ecological disciplines: chemical ecology,  
1157 biogeochemistry, global change, climate change, atmospheric pollution, biogenic VOCs  
1158 emissions, remote sensing, and functioning and structure of terrestrial plants and  
1159 ecosystems.

1160 Table 1. Global K contents of various soils. The data, except those indicated below the table, were obtained from Soil Taxonomy (Soil Survey  
 1161 Staff, 1999).

Soil type	Global surface area (x 10 <sup>6</sup> km <sup>2</sup> )	Average available K in soil profile (mg g <sup>-1</sup> ) (range of soil depth, cm)	Average bulk density (g cm <sup>-3</sup> )	Global total available K (x 10 <sup>9</sup> t)
Alfisol	12.62	0.123 (0-130)	1.73	3.51
Andisol	0.91	0.119 (0-200)	0.8	0.17
Aridisol	15.73	0.620 (0-279)	1.5*	40.1
Entisol	21.14	0.040 (0-152)	1.5*	1.70
Gelisol	11.3	0.072 (0-110)	1.41	1.57
Histosol	1.53	0.147 (0-100)***	1.5*	0.337
Inceptisol	12.83	0.005 (0-73)	1.2	0.059
Mollisol	9.0	0.706 (0-165)	1.48	21.8
Oxisol	9.81	0.024 (0-100)**	1.32	0.310
Spodosol	3.35	0.125 (0-100)	1.5*	0.210
Ultisol	11.05	0.064 (0-280)	1.65	3.25
Vertisol	3.35	0.311 (0-234)	1.95	4.69
Global total (continental areas covered by rocks or ice)	130.8			57.7

1162  
 1163 \* No data provided in Soil Taxonomy; estimated from the average of other soil types.

1164 \*\* Data from Marchi *et al.* (2012), Silva *et al.* (2008) and Da Silva *et al.* (2000).  
1165 \*\*\* Data from Vegas-Vilarrúbia *et al.* (2010).  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193


## Figure legends

Figure 1. K cycle among soils, atmosphere and oceans.

Figure 2. Number of studies reporting limiting and non-limiting roles of K availabilities for plant growth under field conditions in terrestrial ecosystems. (Detailed information regarding these studies is presented in Table S2, Supplementary material).

Figure 3. The reported effects of N loadings (by N deposition or by N fertilization simulating N deposition) in field conditions. The numbers within brackets correspond to the number of the studies that have observed the corresponding effect. (Detailed information of the studies is shown in Table S4, Supplementary material).

Figure 4. The reported relationships between Soil K-availability and alien success. The numbers within brackets correspond to the number of the studies that have observed the corresponding effect (Detailed information of the studies is shown in Table S5, Supplementary material).



1. Soil Survey Staff (1999) Soil Taxonomy. A Basic System of Soil Classification for Making and Interpreting Soil Surveys. Second Edition. 870 pages.
2. Hoelt RG, Nafziger ED, Johnson RR, Aldrich SR (2000) Modern Corn and Soybean Production. MCSP Publications. Champaign, IL. 353p.
3. FAO (2008) Current world fertilizer trends and outlook to 2014. <ftp://ftp.fao.org/ag/agp/docs/cwfto14.pdf>.
4. Eakins BW, Sharman GF (2010) Volumes of the World's Oceans from ETOPO1, NOAA National Geophysical Data Center, Boulder, CO.
5. USGS (2009). Potash. U.S. Geological Survey, Mineral commodity summaries. Online at <http://minerals.er.usgs.gov/minerals/pubs/mcs>.
6. Pawlowicz R (2013) Key physical variables in Ocean: temperature, salinity, and density. Nature Education Knowledge 4 (4): 13.
7. Chow V, Maidment D, Mays L (2000) *Hidrología Aplicada*. Ed. Nomos, S.A. Colombia. 584 p.
8. Cole GA (1983) Textbook of Limnology. 3rd Ed. The C.V. Mosby Co., St. Louis, Toronto, London, 401 pp.




Figure 2




Figure 3




Figure 4



