

Insulin-like Growth Factor 2 Overexpression Induces β -Cell Dysfunction and Increases Beta-cell Susceptibility to Damage*

Received for publication, February 3, 2015, and in revised form, April 29, 2015. Published, JBC Papers in Press, May 13, 2015, DOI 10.1074/jbc.M115.642041

Alba Casellas^{‡§¶}, Cristina Mallol^{‡§¶1}, Ariana Salavert^{‡2}, Veronica Jimenez^{‡§¶}, Miquel Garcia^{‡§¶}, Judith Agudo^{‡§¶}, Mercè Obach^{‡¶}, Virginia Haurigot^{‡§¶}, Laia Vilà^{‡§¶13}, Maria Molas^{‡§¶}, Ricardo Lage^{‡§¶}, Meritxell Morro^{‡§¶4}, Estefania Casana^{‡§¶1}, Jesús Ruberte^{‡¶||}, and Fatima Bosch^{‡§¶15}

From the [‡]Center of Animal Biotechnology and Gene Therapy, Departments of [§]Biochemistry and Molecular Biology and ^{||}Animal Health and Anatomy, School of Veterinary Medicine, Universitat Autònoma Barcelona, 08193-Bellaterra and [¶]Centro de Investigación Biomédica en Red de Diabetes y Enfermedades Metabólicas Asociadas, 08036-Barcelona, Spain

Background: Human and animal studies have shown associations between insulin growth factor 2 (IGF2) and diabetes.

Results: Overexpression of insulin growth factor 2 in β -cells leads to β -cell dysfunction and makes islets more vulnerable to β -cell damage and immune attack.

Conclusion: IGF2 may play an important role in the predisposition and development of diabetes.

Significance: This study unravels an unprecedented role of IGF2 on physiology of β -cells.

The human insulin-like growth factor 2 (IGF2) and insulin genes are located within the same genomic region. Although human genomic studies have demonstrated associations between diabetes and the insulin/IGF2 locus or the IGF2 mRNA-binding protein 2 (IGF2BP2), the role of IGF2 in diabetes pathogenesis is not fully understood. We previously described that transgenic mice overexpressing IGF2 specifically in β -cells (Tg-IGF2) develop a pre-diabetic state. Here, we characterized the effects of IGF2 on β -cell functionality. Overexpression of IGF2 led to β -cell dedifferentiation and endoplasmic reticulum stress causing islet dysfunction *in vivo*. Both adenovirus-mediated overexpression of IGF2 and treatment of adult wild-type islets with recombinant IGF2 *in vitro* further confirmed the direct implication of IGF2 on β -cell dysfunction. Treatment of Tg-IGF2 mice with subdiabetogenic doses of streptozotocin or crossing these mice with a transgenic model of islet lymphocytic infiltration promoted the development of overt diabetes, suggesting that IGF2 makes islets more susceptible to β -cell damage and immune attack. These results indicate that increased local levels of IGF2 in pancreatic islets may predispose to the onset of diabetes. This study unravels an unprecedented role of IGF2 on β -cells function.

β -Cell dysfunction and reductions in β -cell mass are pathological alterations central to the development of both type 1 (T1D)⁶ and type 2 diabetes (T2D) (1). In particular, T2D may begin with insulin resistance, but overt T2D only develops when β -cells fail to compensate for the increased insulin demand. Thus, the alteration in the function of β -cells would be a primary defect in the pathogenesis of diabetes (2).

The concept of loss of β -cell differentiation was first introduced by Jonas *et al.* (3) in 1999 to describe a change in the phenotype of β -cells by which functions of the fully differentiated cells necessary for optimal functioning, such as insulin secretion, are lost (4). The progressive changes in β -cell phenotypes observed during dedifferentiation include the down-regulation of many genes that are highly expressed in β -cells, *e.g.* insulin (INS), solute carrier family member 2 (SLC2A2 or GLUT2), glucokinase (GCK), and pancreatic and duodenal homeobox 1 (PDX1) (5). The term dedifferentiation has been used, for example, to describe the phenotype observed in knock-out mice for the transcription factor *Foxo1* specifically in β -cells. In this animal model that develops overt hyperglycemia, β -cells were described to have dedifferentiated to a state of immaturity characterized by positivity for markers of multipotency and plasticity and for markers of endocrine progenitor cells and by lack of insulin production (6). Other authors have also used the term dedifferentiation to describe the loss of β -cell identity due to the inactivation of transcription factors that define β -cell fate (7, 8). Importantly, recent evidence attributes to dedifferentiation of β -cells a causative role in β -cell failure in T2D (6, 9, 10).

In parallel to β -cell dedifferentiation, many animal models of T1D also show endoplasmic reticulum (ER) stress. Given that

* This work was supported in part by Ministerio de Economía y Competitividad, Plan Nacional I+D+I, Grants SAF2008-00962 and SAF2011-24698, and Generalitat de Catalunya Grants 2009 SGR-224 and ICREA Academia, Spain. The authors declare that they have no conflicts of interest with the contents of this article.

¹ Recipients of predoctoral fellowship from Ministerio de Ciencia e Innovación.

² Recipient of a predoctoral fellowship from Generalitat de Catalunya.

³ Recipient of Postdoctoral Fellowship JCI-2010-06388 from Ministerio de Ciencia e Innovación, Spain.

⁴ Recipient of Predoctoral Fellowship SAF2008-00962 from Ministerio de Ciencia e Innovación.

⁵ To whom correspondence should be addressed: Center of Animal Biotechnology and Gene Therapy. Edifici H. Universitat Autònoma de Barcelona. E-08193 Bellaterra, Spain. Tel.: 34-93-581-4182; Fax: 34-93-581-4180; E-mail: fatima.bosch@uab.es.

⁶ The abbreviations used are: T1D, type 1 diabetes; T2D, type 2 diabetes; INS, insulin; ER, endoplasmic reticulum; qPCR, quantitative real time PCR; STZ, streptozotocin; TRITC, tetramethylrhodamine isothiocyanate; bw, body weight; VNTR, variable number of tandem repeats; IR, insulin release; rIGF2, recombinant IGF2.

β -cells need to produce insulin in large quantities, they probably represent a cell type vulnerable to ER stress (11). In fact, ER stress has been proposed as a mechanism responsible for β -cell dysfunction and death in type 2 diabetes (12), and it has been reported to precede the onset of T1D in the nonobese diabetic mouse model (13). The major indicators of the ER stress response, such as X-box binding protein 1 (*Xbp1*) and DNA-damage inducible transcript 3 (*Ddit3/Chop*), are up-regulated in islets from type 1 and type 2 diabetic models (12, 14, 15).

Insulin-like growth factor 2 (IGF2) is a growth-promoting polypeptide that shares a high degree of structural homology with insulin. IGF2 is synthesized primarily by the liver, but it is also produced locally by many tissues, where it acts in an autocrine/paracrine manner (16). IGF2 is expressed at high levels during embryonic development, but its expression is progressively shut down in most tissues after birth. In humans, the *IGF2* gene is located on chromosome 11p15.5, in close linkage with the *INS* and tyrosine hydroxylase (*TH*) genes. Because the *IGF2* and *INS* genes share their promoter, the human pancreas can express *INS-IGF2* hybrid transcripts (17). Indeed, in human β -cells insulin is the most abundant transcript, followed by *INS-IGF2* hybrids and *IGF2* transcripts (18). Genome-wide association studies performed in humans from Caucasian and Asian populations have demonstrated associations between the *IDDM2* locus containing the *INS* and *IGF2* genes and diabetes (19). In particular, variations in the size and sequence of a regulatory element of the insulin promoter that includes a variable number of tandem repeats (VNTR) have been found to confer susceptibility to T1D (the shortest version of VNTR or class I) or to T2D (the longest version of VNTR or class III) (20, 21). Pugliese *et al.* (20) suggested that although there is clear evidence supporting the concept that the *INS* VNTR element is the main factor conferring susceptibility to diabetes to the *IDDM2* locus, there is also the possibility that, at least in some instances, both the *INS* and *IGF2* locus modulate *IDDM2*-conferred susceptibility. In contrast, the *IGF2BP2* (*IMP2*) gene, involved in the regulation of IGF2 expression by binding to the 5'-UTR of the *IGF2* mRNA to promote its translation (22), has also been associated with genome-wide association studies with an increased risk of T2D and gestational diabetes in Caucasian, Asian, and Arabic populations (23–25). *IGF2BP2* is expressed in β -cells and in insulin-sensitive tissues in postnatal life (22), and an altered expression or regulation of the *IGF2BP2* gene may end up impacting the levels of IGF2 in these cells. Furthermore, old Goto-Kakizaki rats, a nonobese model of mild T2D, show increased *IGF2* mRNA in their islets (26).

We previously showed that transgenic mice overexpressing IGF2 specifically in β -cells (Tg-IGF2) display several primary defects characteristic of T2D, such as hyperinsulinemia, mild hyperglycemia, and altered glucose and insulin tolerance tests (27). These observations together with the evidence from human genome-wide association studies suggested that altered levels of IGF2 in β -cells could confer susceptibility to diabetes. Here, we studied the mechanisms by which IGF2 exerted its action on β -cells of Tg-IGF2 to help unravel the role that this growth factor plays in the development of overt diabetes in humans. To this end, we studied in detail the phenotype of transgenic β -cells at the initial stages of the diabetic process.

Local IGF2 overexpression led to β -cell dysfunction as well as down-regulation of typical markers of β -cell differentiation. Up-regulation of markers of ER stress and immune response-related genes was also documented in islets of Tg-IGF2 mice, and this agreed with the observed increased susceptibility to β -cell death and diabetes development. Thus, in this study we provide evidence that local pancreatic overexpression of IGF2, besides causing islet hyperplasia and hyperinsulinemia (27), directly affects β -cells leading to cell dysfunction and susceptibility to damage and as a consequence predisposes to the onset of diabetes.

Experimental Procedures

Animals—Heterozygous male transgenic mice expressing either mouse *Igf2* or human *IFN β* under the control of the rat insulin promoter-I (RIP-I) were used (27, 28). Tg-IGF2 mice were in C57Bl6/SJL background, and Tg-IFN β mice, a model of islet lymphocytic infiltration with increased susceptibility to autoimmune diabetes (29), were in CD1 background, and double Tg-IFN β /IGF2 mice were obtained by crossing the two lines (C57Bl6/SJL + CD1). Diabetes was induced as described previously (29). All mice were fed *ad libitum* with a standard chow diet (2018S Teklad Global, Harlan) and maintained under conditions of controlled temperature and light (12 h light/dark cycles). When stated, mice were fasted for 16 h. Animal care and experimental procedures were approved by the Ethics Committee in Animal and Human Experimentation of the Universitat Autònoma de Barcelona.

Immunohistochemistry and Histopathology—For immunohistochemical detection of INS, GLUT2, PDX1, MAFA, and p62, pancreata were fixed for 12–24 h in formalin, embedded in paraffin, and sectioned. Sections were then incubated overnight at 4 °C with the following antibodies: guinea pig anti-porcine insulin (Sigma); rabbit anti-human glucagon (Signet Labs, Dedham, MA); rabbit polyclonal anti-GLUT2 (AB1342, Chemicon International Inc., EEUU); rabbit anti-human IGF2 ((PAC1) GroPep Ltd., Adelaide, Australia); rabbit polyclonal anti-MAFA (Santa Cruz Biotechnology, sc-66958); rabbit polyclonal anti-PDX1 (Millipore, AB3243); rat anti-mouse MAC-2 (Cedarlane CL8942AP); and mouse anti-human p62 (BD Biosciences 610833) as a autophagy marker. As secondary antibodies, peroxidase-conjugated rabbit anti-guinea pig IgG (Dako, Denmark), biotinylated goat anti-rabbit (Pierce), TRITC-conjugated goat anti-guinea pig (Molecular Probes Leiden, The Netherlands), biotinylated goat anti-rabbit (Molecular Probes), biotinylated rabbit anti-rat (Dako, Glostrup, Denmark), and biotinylated horse anti-mouse (Vector Laboratories, Burlingame, CA) antibodies were used. Streptavidin-conjugated Alexa 488 (Molecular Probes) or streptavidin-conjugated Alexa 568 (Molecular Probes) were used as fluorochromes. Images were obtained with a Nikon Eclipse E800 microscope (Nikon, Tokyo, Japan). For laser-scanning confocal analysis, a TCS SP2 microscope (Leica Microsystems, Heidelberg, Germany) was used.

Morphometric Analysis— β -Cell mass determination was performed as described previously (29). Macrophage infiltration was measured as the percentage of MAC-2-positive area

TABLE 1
Primers used for murine islets and INS-1 cells (rat)

Gene	Forward primer (5' to 3')	Reverse primer (5' to 3')
Mouse		
<i>Foxa2</i>	5'GAG-CAG-CAA-CAT-CAC-CAC-G3'	5'CGT-AGG-CCT-TGA-GGT-CCA-T3'
<i>Pdx1</i>	5'GAA-ATC-CAC-CAA-AGC-TAC-G3'	5'CGG-GTT-CCG-CTG-TGT-AAG3'
<i>RplpO</i>	5'TCC-CAC-CTT-GTC-TCC-AGT-CT3'	5'ACT-GGT-CTA-GGA-CCC-GAG-AAG3'
<i>Ins</i>	5'GCG-ATT-GTG-GAT-CAG-TGC-T3'	5'AGG-TGG-GCC-TTA-GTT-GCA-C3'
<i>Gck</i>	5'ATG-ACA-GAG-CCA-GGA-TGG-AG	5'CGG-CTC-ATC-ACC-TTC-TTC-AG3'
<i>Glut2</i>	5'CTG-GAG-CCC-TCT-TGA-CGG-GA3'	5'CCA-GTC-CTG-AAA-TTA-GCC-CAC-A3'
<i>Serca2</i>	5'GTC-CTA-ACT-GTG-GTG-TTT-TCC-TC3'	5'GTT-TAG-GAA-GCC-GTT-ACT-CCA-G3'
<i>Chop</i>	5'GCG-ACA-GAG-CCA-GAA-TAA-CA3'	5'GAT-GCA-CTT-CCT-TCT-GGA-ACA3'
B2-microglobulin	5'ACT-CCA-AGA-CCC-AGA-AAC-TGT-C3'	5'ACT-GGT-AGG-AGT-AGG-GAT-GCA-C3'
<i>H2-Aa</i>	5'CTC-TGA-TTC-TGG-GGG-TCC-T 3'	5'ACC-ATA-GGT-GCC-TAC-GTG-GT3'
β 2/NeuroD1	5'CTC-TGA-TTC-TGG-GGG-TCC-T3'	5'ACC-ATA-GGT-GCC-TAC-GTG-GT3'
<i>Hnf4a</i>	5'TGC-CAA-CCT-CAA-TTC-ATC-CA3	5'GCT-CGA-GGC-TCC-GTA-GTG-TT3'
<i>Mafa</i>	5'AGG-AGG-AGG-TCA-TCC-GAC-TG3'	5'CTT-CTC-GCT-CTC-CAG-AAT-GTG3'
<i>Xbp1s</i>	5'GCT-GAG-TCC-GCA-GCA-GGT3'	5'ACA-GGG-TCC-AAC-TTG-TCC-AG3'
<i>XBp1</i>	5'TCA-GTT-TCC-TCC-GCA-GCG-CTT-T3	5'GCT-GCC-GCT-CAT-GGT-ACC-CG3'
<i>Igf2r</i>	5'TCT-GTG-TTG-GCT-CGT-CAC-TC3'	5'CCG-GTG-ACA-GAC-GTT-GAT-GA3'
<i>Insr-B</i>	5'GCC-AGT-GAG-TGC-TGC-TCA-T3'	5'TAC-TGT-CCT-CGG-CAC-CAT-T3'
<i>Insr-A</i>	5'CCC-ACC-CTT-TGA-GTC-TGA-TG3'	5'GCT-TTC-GGG-ATG-GCC-T3'
<i>Gcg</i>	5'GGC-ACA-TTC-ACC-AGC-GAC-TA3'	5'GTC-CCT-TCA-GCA-TGC-CTC-TC3'
Rat		
<i>Foxa2</i>	5'CCA-GAC-AAC-GCG-AGT-CCT3'	5'ACG-GCT-CCC-AGC-ATA-CTT-T3'
<i>Pdx1</i>	5'CTG-TCG-TGC-CAT-GTG-AAC-C3'	5'TTC-TCT-AAA-TTG-GTC-CCA-GGA-A3'
<i>RplpO</i>	5'GAT-GCC-CAG-GGA-AGA-CAG3	5'CAC-AAT-GAA-GCA-TTT-TGG-GTA-G3'
<i>INS</i>	5' GCT-CTG-TAC-CTG-GTG-TGT-GG 3'	5' CCA-AGG-TCT-GAA-GAT-CCC-CG 3
<i>Gck</i>	5'GCC-CAG-TTG-TTG-ACT-CTG-GT	5'CAT-CAC-CTT-CTT-CAG-GTC-TTC 3'
<i>GLUT2</i>	5'TGA-AGG-ATC-TGC-TCA-CAT-AGT-CA 3'	5'CCA-ACA-TGG-CTT-TGA-TCC-TT 3'
<i>Serca</i>	5'TGT-TAA-TCA-AGA-CAA-AAA-GAA-CAT-GC3'	5'GGA-TCT-TGC-CGA-TCT-CAG-TAT-T3'

per islet area in islets from three different sections per mouse, with four animals per group.

Islets Isolation and Culture—Pancreatic islets were isolated as described previously (29) and then cultured in RPMI 1640 medium (11 mM glucose), supplemented with 1% BSA, 2 mM glutamine, penicillin/streptomycin at 37 °C in an atmosphere of 95% humidified air, 5% CO₂. To study the effects of recombinant IGF2 on β -cells, after overnight culture, pools of islets were treated with 13.5 nM recombinant IGF2 (792-MG, R&D Systems) and then cultured for 48 h. To express *Igf2* in islets, after isolation and overnight culture, pools of islets were infected with *Igf2*-expressing adenoviruses (Ad5/CMV-IGF2) or with noncoding adenoviruses as control (Ad5/CMV-Null) and incubated for 48 h. After treatments islets were hand-picked and processed. For qPCR analysis, islets were cultured overnight to recuperate from isolation stress and then hand-picked and processed to obtain RNA.

Cell Culture—INS-1 cells were cultured in RPMI 1640 medium (11 mM glucose), 2 mM glutamine, 10 mM Hepes, 1 mM sodium pyruvate, 50 μ M 2-mercaptoethanol, and 10% FBS at 37 °C in an atmosphere of 95% humidified air, 5% CO₂. For cell culture studies, medium was modified, and 10% FBS was substituted for 1% BSA, and cells were treated with 200 nM wortmannin (W1628, Sigma) followed an hour later by 100 ng/ml recombinant IGF2, or with either alone, and then cultured for 48 h.

Gene Expression Analysis—For qPCR analysis, total RNA was extracted from isolated islets using Tripure Isolation Reagent (Roche Applied Science) and RNeasy mini kit (Qiagen, Hilden, Germany). Total RNA (1 μ g) was retrotranscribed using the Transcriptor First Strand cDNA synthesis kit (Roche Applied Science). RT-PCR was performed using LightCycler 480 SYBR Green I MasterMix (Roche Applied Science). The primers used for murine islets and INS-1 cells (rat) are listed in the Table 1.

Results were analyzed using the mathematical model of Pfaffli (30). All samples were processed in triplicate, and a mean C_t was calculated. The C_t for each transcript of interest was normalized by the C_t obtained for the reference gene *RplpO*. Then, the relative expression of each gene in each sample was calculated by dividing the value obtained for each sample by the mean of wild-type/control values.

Hormone and Metabolite Assays—Blood glucose levels and serum insulin concentrations were measured as described previously (29). The concentration of serum glucagon was determined by radioimmunoassay with the Millipore glucagon radioimmunoassay kit (EMD Millipore Corp., Billerica, MA). Glucose tolerance was evaluated as described previously (29). For insulin release determination, glucose (3 g/kg body weight) was injected intraperitoneally, and venous blood was collected from tail vein at 0, 2, 5, 15, and 30 min in pre-chilled tubes (Microvette® CB 300, SARSTEDT) and immediately centrifuged to separate plasma, which was stored at -20 °C. Insulin levels were measured by ELISA (Crystal Chemical, Chicago).

Microarray Analysis—RNA samples were obtained following Affymetrix recommendations (Expression Analysis Technical Manual). Microarray analysis was performed by Progenika (Bilbao, Spain). The final gene list contained only those probe sets with a $p < 0.05$. For the interpretation, cross-checking and visualization of the data, FatiGO term enrichment (release date 11/20/2010), Database for Annotation, Visualization and Integrated Discovery (DAVID; david.abcc.ncifcrf.gov), Kyoto Encyclopedia of Genes and Genomes, and GeneCodis were used. Array data have been submitted to MIAMExpress (E-MEXP-3786).

Statistical Analysis—All values are expressed as the mean \pm S.E. Differences between groups were compared by Student's *t*




FIGURE 1. Local overexpression of IGF2 alters islet functionality. *A*, Tg-IGF2 mice showed altered glucose tolerance test at 4 months of age. Three-month-old WT (white circle) and Tg-IGF2 (black circle) mice were given an intraperitoneal injection of 2 mg of glucose/g bw, and blood glucose was measured at the indicated times as described under "Experimental Procedures." Results are expressed as mean \pm S.E. of 10 animals/group. *, $p < 0.05$. *B*, *in vivo* insulin release test after intraperitoneal glucose injection (3 g/kg bw) to 4-month-old WT mice (white circles) and Tg-IGF2 mice (black circles), $n = 10$ animals/group (*, $p < 0.05$). *C*, analysis of the expression of β -cell glucose sensor genes, *Gck* and *Slc2a2/Glut2*, by qPCR on WT (white bars) and Tg-IGF2 (black bars) islets from 4-month-old mice. Data are mean \pm S.E. of at least three pools of islets from eight animals/group. **, $p < 0.01$. *D*, immunohistochemical detection of INS (red) and GLUT2 (green) in pancreatic sections of 3-month-old WT and Tg-IGF2 mice. Original magnification $\times 400$ (scale bar, 50 μ m). *E*, electron microscopy of β -cells in pancreatic islets isolated from 4-month-old WT and Tg-IGF2 mice. Tg-IGF2 β -cells presented altered ultrastructure ($\times 10,000$) (scale bar, 2 μ m). Inset images show secretory granules in more detail ($\times 80,000$) (scale bar, 0.2 μ m).

test. A p value less than 0.05 was considered statistically significant.

Results

Local IGF2 Overexpression Impairs Islet Function—Transgenic mice overexpressing IGF2 specifically in β -cells (Tg-IGF2) have disrupted islet structure and islet hyperplasia and develop a pre-diabetic state (27). To study the role of IGF2 in β -cell function at the initial stages of the diabetic process, we performed a glucose tolerance test and *in vivo* insulin release (IR) test on 4-month-old mice. Tg-IGF2 mice showed glucose intolerance (Fig. 1A) (27) and were accompanied by reduced insulin release (Fig. 1B). In particular, the first phase of insulin

secretion in response to glucose was greatly diminished in these mice (Fig. 1B), suggesting an impaired secretory response by β -cells. Tg-IGF2 mice were, however, hyperinsulinemic (WT, 1.2 ± 0.16 ng/ml; Tg-IGF2, 2.2 ± 0.19 ng/ml, $p < 0.05$ (27)), which reflected the fact that these animals have a β -cell mass 3-fold higher than WT littermates (27). In contrast, Tg-IGF2 mice presented a normal α -cell mass (27), and no differences were observed in the levels of islet glucagon mRNA (1 ± 0.35 versus 0.81 ± 0.19 relative units for wild type and Tg-IGF2, respectively, $p \leq 0.31$) or serum glucagon (WT = 100.9 ± 8.2 versus Tg-IGF2 = 131.2 ± 26.5 pg/ml, $p \leq 0.21$). In agreement with the abnormal insulin release shown by Tg-IGF2 upon glu-

IGF2 Overexpression in β -Cells




FIGURE 2. **IGF2 overexpression leads to changes in β -cell phenotype.** A–F, confirmation of the differential gene expression of β -cell markers in WT (white bars) and Tg-IGF2 (black bars) islets from 4-month-old mice by qPCR. Data are expressed as mean \pm S.E. of at least three pools of islets from eight animals/group. *, $p < 0.05$. G, immunohistochemical detection of INS (red) and MAFa (green) in pancreatic sections from 3-month-old mice. Original magnification $\times 400$ (scale bar, 50 μ m). H, immunohistochemical detection of INS (red) and PDX1 (green) in pancreatic sections from 3-month-old mice. Original magnification $\times 400$ (scale bar, 50 μ m).

glucose stimulation, the β -cell glucose sensors, solute carrier family member 2 (*Slc2a2* or *Glut2*) and glucokinase (*Gck*), were down-regulated in transgenic islets (Fig. 1C). Immunohistochemical analysis further confirmed the dramatic reduction in SLC2A2 protein in insulin-positive cells in these animals (Fig. 1D). Moreover, the ultrastructural analysis of β -cells by electron microscopy revealed an increased number of immature insulin secretory granules with low or intermediate electron density occupying most of the cytoplasm of Tg-IGF2 β -cells (Fig. 1E). Altogether, these results provide evidence of β -cell dysfunction and inadequate secretory response in Tg-IGF2 mice.

To gain insight into the IGF2-mediated molecular mechanisms that could account for β -cell dysfunction, we determined the expression of key genes involved in β -cell functionality. By qPCR, we observed that Insulin I mRNA levels were decreased in Tg-IGF2 compared with WT islets (Fig. 2A), and this agreed with the loss of the second phase of insulin release (Fig. 1A). The

expression of critical transcription factors for insulin production such as forebrain box A2 (*Foxa2*), neurogenic differentiation 1 (*Neurod1*), hepatic nuclear factor 4 α (*Hnf4 α*), v-maf musculoaponeurotic fibrosarcoma oncogene family (*Mafa*), and pancreatic and duodenal homeobox 1 (*Pdx1*) was also down-regulated (Fig. 2, B–F). Decreased production of MAFa and PDX1 was further confirmed by immunohistochemical analysis (Fig. 2, G and H). In addition, we performed microarray analysis on islets isolated from Tg-IGF2 and WT mice. After normalization and statistical filtering of the data, more than 300 genes with a 2-fold up- or down-regulation were identified, with the majority of them down-regulated and involved, according to gene ontology classification, in insulin secretion, development, cellular stress, immunity, transport, electron transport chain, and replication (Table 2). Noticeably, there was down-regulation of several genes characteristic of fully differentiated and functional β -cells, such as proprotein convertase subtilisin/kexin type 1 (*Pcsk1/PC1/3*), forkhead box A2

TABLE 2
Differentially expressed genes

Microarray expression analysis in islets from WT and Tg-IGF2 2-month-old mice. Classification of up- and down-regulated genes according to their biological function reported in the literature. RNA was isolated from three pools of islets from nine animals per group.

Gene description	Gene symbol	Fold change	p value	RefSeq
Development				
Myeloid ecotropic viral integration site-related gene 1	<i>Meis2</i>	0.67	0.03160	NM_010825
ISL1 transcription factor, LIM/homeodomain (islet 1)	<i>Isl1</i>	0.42	0.04980	NM_021459
Insulin promoter factor 1, homeodomain transcription factor	<i>Pdx1</i>	0.58	0.03740	NM_008814
Notch gene homolog 3 (<i>Drosophila</i>)	<i>Nocth3</i>	2.11	0.00484	NM_00876.2
Insulin secretion				
Activating transcription factor 2	<i>Atf2</i>	0.45	0.02090	NM_009715
Calcyphosphine 2	<i>Caps2</i>	0.35	0.03340	NM_178278
Dipeptidylpeptidase 4	<i>Dpp4</i>	0.60	0.01890	NM_010074
ERO1-like (<i>Saccharomyces cerevisiae</i>) /// hypothetical gene supported by AK009667; NM_01577	<i>Ero1l</i> /// <i>LOC434</i>	0.44	0.00139	NM_015774
Forkhead box A2	<i>Foxa 2</i>	0.58	0.04580	NM_010446
Gap junction membrane channel protein $\alpha 9$	<i>Gja9/Cx36</i>	0.38	0.03470	NM_010290
Potassium channel, subfamily K, member 16	<i>Kcnk16</i>	0.52	0.00696	NM_029006
Lipin 1	<i>Lpin1</i>	0.59	0.03970	NM_172950
Proprotein convertase subtilisin/kexin type 1	<i>Pcsk1/PC1/3</i>	0.46	0.01790	NM_13628
N-Ethylmaleimide-sensitive fusion protein	<i>Nsf</i>	0.70	0.03610	NM_008740
Rap guanine nucleotide exchange factor (GEF) 3	<i>Rapgef3</i>	2.15	0.02210	NM_144850
Synaptotagmin-like 4	<i>Syt4</i>	0.33	0.04260	NM_013757
Transthyretin	<i>Ttr</i>	0.40	0.02690	NM_013697
Urocortin 3	<i>Ucn3</i>	0.23	0.00894	NM_031250
Metal-response element-binding transcription factor 1	<i>Mtf1</i>	0.52	0.03510	NM_008636
Stress				
Autocrine motility factor receptor	<i>Amfr</i>	0.62	0.02980	NM_011787
Activating transcription factor 2	<i>Atf2</i>	0.45	0.02090	NM_009715
ATPase, Cu^{2+} transporting, α -polypeptide	<i>Atp7a</i>	0.73	0.03470	NM_009726
Der1-like domain family, member 2	<i>Der12</i>	0.53	0.01260	NM_033562
ERO1-like (<i>S. cerevisiae</i>) /// hypothetical gene supported by AK009667; NM_01577	<i>Ero1l</i>	0.44	0.00139	NM_015774
F-box and leucine-rich repeat protein 5	<i>Fbxl5</i>	0.65	0.03600	NM_178729
Forkhead box M1	<i>Foxm1</i>	1.33	0.03840	NM_027697
Heat shock protein 1	<i>Hspb1</i>	1.91	0.02100	NM_013560
Methyl CpG-binding protein 2	<i>Mecp2</i>	0.63	0.02480	NM_010788
Metal-response element-binding transcription factor 1	<i>Mtf1</i>	0.52	0.03510	NM_008636
Polo-like kinase 3 (<i>Drosophila</i>)	<i>Plk3</i>	0.42	0.00505	NM_013807
Autophagy 10-like (<i>S. cerevisiae</i>)	<i>Atg10</i>	0.64	0.04880	NM_025770
Immune response				
Alanyl-tRNA synthetase	<i>Aars</i>	0.68	0.04430	NM_146217
Glutamic acid decarboxylase 1	<i>Gad1</i>	0.21	0.01030	NM_008077
Cathepsin H	<i>Ctsh</i>	2.42	0.02940	NM_007801
Immunoglobulin heavy chain (J558 family)	<i>Igh-VJ558</i>	8.33	0.00888	NM_134051
Endothelial differentiation, sphingolipid G-protein-coupled receptor, 3	<i>Edg3</i>	2.96	0.03100	NM_010101
Histocompatibility 2, class II antigen A, α	<i>H2-Aa</i>	2.70	0.01940	NM_010378
Transport				
Solute carrier family 9 (sodium/hydrogen exchanger), isoform 3 regulator 2	<i>Slc9a3r2</i>	1.50	0.03590	NM_023055
Solute carrier family 35, member 3	<i>Slc35a3</i>	0.66	0.04800	NM_144902
Solute carrier family 2 (facilitated glucose transporter), member 2	<i>Slc2a2</i>	0.24	0.04170	NM_031197
solute carrier family 20, member 2	<i>Slc20a2</i>	0.60	0.04070	NM_011394
Solute carrier family 12, member 7	<i>Slc12a7</i>	0.57	0.03840	NM_011390
Solute carrier family 4, sodium bicarbonate cotransporter, member 7	<i>Slc4a7</i>	0.47	0.01890	NM_001033270
Solute carrier family 2 (facilitated glucose transporter), member 5	<i>Slc2a5</i>	0.61	0.04950	NM_0119741
Potassium large conductance calcium-activated channel, β member 2	<i>Kcnmb2</i>	0.50	0.00456	NM_028231
Activating transcription factor 2	<i>Atf2</i>	0.45	0.02090	NM_009715
Calcium channel, voltage-dependent, $\beta 2$ subunit	<i>Cacnb2</i>	0.68	0.02080	NM_023116
Transport chain				
ATPase, H^+ -transporting, V1 subunit B, isoform 2	<i>Atp6v1b2</i>	0.77	0.04870	NM_007509
ATPase, H^+ -transporting, V0 subunit C	<i>Atp6v0c</i>	0.61	0.04770	NM_009729
ATPase, aminophospholipid transporter (APLT), class I, type 8A, member 1	<i>Atp8a1</i>	0.62	0.04630	NM_009727
ATPase, Cu^{2+} -transporting, α -polypeptide	<i>Atp7a</i>	0.64	0.04200	NM_009726
ATPase, Ca^{2+} -sequestering	<i>Atp2c1</i>	0.55	0.03890	NM_175025
ATPase, Ca^{2+} -transporting, cardiac muscle, slow twitch 2	<i>Atp2a2</i>	0.49	0.03040	NM_009722
Cytochrome P450, family 4, subfamily v, polypeptide 3	<i>Cyp4v3</i>	0.53	0.03790	NM_133969
NADH dehydrogenase (ubiquinone) 1, subcomplex unknown, 2	<i>Ndufc2</i>	0.60	0.03220	NM_024220
Replication				
Cyclin D2	<i>Ccnd2</i>	0.49	0.04340	NM_009829
Prolactin receptor	<i>Prlr</i>	0.45	0.02650	NM_011169
B-cell translocation gene 1, anti-proliferative	<i>Btg1</i>	0.68	0.03340	NM_007569

(*Foxa2*), and lipin 1 (*lpin1*) involved in insulin secretion, insulin promoter factor 1, homeodomain transcription factor (*lpf1/Pdx1*), and solute carrier family 2, member 2 (*Slc2a2*). Altogether, these results support the notion that β -cells undergo a

process of dedifferentiation resulting in functional alterations due to the overexpression of IGF2.

IGF2 Overexpression Induces ER Stress in β -Cells—It has been reported that endoplasmic reticulum (ER) stress is involved in

IGF2 Overexpression in β -Cells




FIGURE 3. ER stress and autophagy in Tg-IGF2 islets. *A*, electron microscopy of β -cells in islets of 4-month-old WT and Tg-IGF2 mice. Rough endoplasmic reticulum showing a normal lamellar arrangement is observed in WT, whereas Tg-IGF2 presented altered ER lamellar arrangement and dilatations as indicated by *red arrows* (composite image). Original magnification $\times 100,000$. *B*, validation of differential gene expression of ER stress markers in WT (*white bars*) and Tg-IGF2 (*black bars*) islets from 4-month-old mice by qPCR. Data are expressed as mean \pm S.E. of at least three pools of islets from eight animals/group. *, $p < 0.05$; **, $p < 0.01$. *C*, electron microscopy of β -cells in islets of 4-month-old Tg-IGF2 mice. An autophagic body, a cytoplasmic inclusion surrounded by a double membrane containing part of the cytoplasm, is indicated by *red arrows* in an image corresponding to Tg-IGF2 β -cells ($\times 80,000$). *D*, immunohistochemical detection of insulin (*red*) and LGAL3/p62 (*green*) in pancreatic sections from 3-month-old mice. *Inset images* show in detail the positive p62 signal observed in β -cells of Tg-IGF2 islets. Scale bar, 8 μ m.

β -cell failure (31). Electron microscopy observation of Tg-IGF2 β -cells showed an expanded ER network (Fig. 3*A*, *arrows*), which is indicative of ER stress induced by the unfolded protein response (32). Accordingly, microarray analysis (Table 2) and qPCR (Fig. 3*B*) revealed down-regulation of the ATPase, Ca^{2+} -transporting cardiac muscle, slow twitch 2 (*Atp2a2/Serca2*) gene in Tg-IGF2 islets. The SERCA protein regulates Ca^{2+} flux in and out of the ER and, as a consequence, controls proper protein folding (33). Moreover, markers of ER stress such as spliced x-box-binding protein 1 (*Xbp1s*) and DNA damage-inducible transcript 3 (*Ddit3/Chop*) were also up-regulated in Tg-IGF2 (Fig. 3*B*).

Autophagy has also been associated with ER stress and plays an important role in diabetes pathogenesis (34). Electron

microscopy analysis revealed the presence in Tg-IGF2 β -cells of autophagic bodies or autophagosomes, identified as portions of the cytoplasm surrounded by a double membrane (Fig. 3*C*). An increase in the immunohistochemistry signal intensity of SQSTM1 (p62), a ubiquitin-binding protein also known as sequestosome 1 that co-localizes to autophagic bodies (35), was detected in most of the unstructured islets of Tg-IGF2 mice (Fig. 3*D*), confirming an increase in autophagy in Tg-IGF2 islets.

IGF2 Is Sufficient to Trigger β -Cell Dysfunction—To discard the possibility that the β -cell dysfunction observed in Tg-IGF2 mice was due to IGF2 overexpression from early embryonic development, islets isolated from adult WT animals were trans-




FIGURE 4. IGF2 effects on adult WT islets. *A*, expression of β -cell markers by qPCR analysis in WT islets transduced with null control vector (*Ad/Null*, white bars) or adenovirus expressing IGF2 (*Ad/IGF2*, black bars). *B*, expression of ER stress genes by qPCR in the same adenovirus-transduced islets. Three pools of islets from eight animals/group were used. *C*, gene expression analysis of β -cell markers by qPCR in WT islets incubated for 48 h with either vehicle (white bars) or 100 ng/ml recombinant IGF2 protein (black bars). Three pools of islets from eight animals/group were used. *D*, analysis by qPCR of *Foxa2*, *Slc2a2*/*Glut2*, and *Gck* genes in INS-1 cells treated for 48 h with vehicle (white bars), 100 ng/ml recombinant IGF2 protein (black bars), 200 nM of the PI3K inhibitor wortmannin (light gray bars), or both (dark gray bars). Results shown represent the data obtained for at least six wells/condition and from three independent experiments. Data are expressed as mean \pm S.E. *, $p < 0.05$; **, $p < 0.01$ versus vehicle-treated cells. #, $p < 0.05$, and ##, $p < 0.01$ versus recombinant IGF2-treated cells.

duced *in vitro* with adenoviral vectors encoding murine *Igf2* (*Ad/IGF2*) or with null vectors (*Ad/Null*) that served as controls. Islets transduced with *Ad/IGF2* vectors presented a similar phenotype to that observed in Tg-IGF2 islets, e.g. decreased *Slc2a2*, *Pdx1*, insulin I, and *Atp2a2*/*Serca2* expression and increased levels of *Ddit3*/*Chop*, although changes in *Serca2* and *Ddit3*/*Chop* did not reach statistical significance (Fig. 4, *A* and *B*). Thus, IGF2 overexpression in adult islets is sufficient to alter β -cell functionality by diminishing insulin production and inducing ER stress.

To study whether the phenotype of Tg-IGF2- and *Ad/IGF2*-transduced islets was due to the forced overexpression of IGF2 rather than a direct effect of IGF2 on β -cells, islets isolated from adult WT mice were also incubated for 48 h with recombinant IGF2 protein (rIGF2) to mimic the chronic exposure to IGF2 of Tg-IGF2 islets. Similarly to Tg-IGF2 and *Ad/IGF2*-transduced islets, rIGF2-treated islets presented a significant decrease in

Slc2a2 and insulin I expression and a nonstatistically significant down-regulation of *Gck*, *Pdx1*, and *Atp2a2*/*Serca2* (Fig. 4C). Similar results were obtained in rIGF2-treated INS-1 cells (data not shown).

Given that IGF2 binds with high affinity to the insulin (INSR), insulin-like growth factor 1 (IGF1R), and IGF2 receptors, we quantified the expression of these receptors in Tg-IGF2 islets. No differences were observed in the expression of *IGF2R*, *IGF1R*, or *INSR-B* (data not shown). However, a moderate (about 40%) statistically significant decrease in INSR-A expression was documented in Tg-IGF2 islets (WT 1.03 ± 0.11 versus Tg-IGF2 0.67 ± 0.05 , $p = 0.03$). Next, we analyzed the effects of IGF2 on gene expression by inhibiting both INSR and IGF1R downstream signaling pathways using wortmannin, a PI3K inhibitor (36). INS-1 cells cultured in the presence of rIGF2 showed decreased expression of markers of β -cell identity (Fig. 4D), which recapitulated the observations made in Tg-IGF2




FIGURE 5. **IGF2 overexpression leads to islet macrophage infiltration.** *A*, macrophage infiltration was determined by LGAL3/MAC-2 immunostaining on islet sections from 3-month-old mice. Original magnification $\times 10$ (scale bar, 100 μm) and insets $\times 40$ (scale bar, 50 μm). *B*, quantification of LGAL3/MAC-2-positive area. Data are expressed as mean \pm S.E. of percentage of LGAL3/MAC-2-positive islet areas/total islet areas. $n = 5$ animals/group. WT (white bars) and Tg-IGF2 (black bars). *C*, analysis of gene expression by qPCR of histocompatibility 2, class II antigen A α (*H2-Aa*), β -2 microglobulin (*B2m*) genes in WT (white bars), and Tg-IGF2 (black bars) islets from 4-month-old mice. Results are expressed as mean \pm S.E. of at least three pools of islets from eight animals/group. *, $p < 0.05$; **, $p < 0.01$.

islets (Figs. 1*B* and 2). The addition of wortmannin *per se* altered the expression of *Foxa2*, *Slc2a2*, and *Gck* (Fig. 4*D*). Despite this unexplained observation, we did document a significant inhibition of the effects of rIGF2 by wortmannin treatment (Fig. 4*D*), indicating that at least partially the IGF2 effects on β -cell expression were mediated by PI3K activation.

IGF2 Transgenic Islets Are More Susceptible to β -Cell Damage—Microarray analysis of Tg-IGF2 islets showed altered expression of several genes involved in the immune response, such as cathepsin H (*Ctsh*), immunoglobulin heavy chain (J558 family, *Igh-VJ558*), endothelial differentiation, sphingosine 1-phosphate receptor 3 (*Edg3*), and histocompatibility 2, class II antigen A, α (*H2-Aa*) (data not shown). Although CD4- and CD8- positive lymphocytes could not be detected in pancreata from Tg-IGF2 mice (Table 2), immunostaining against LGAL3/MAC-2 revealed the presence of macrophages infiltrating Tg-IGF2 islets (Fig. 5, *A* and *B*). qPCR analysis confirmed the increased expression of *H2-Aa* and β 2-microglobulin (*B2m*) (Fig. 5*C*), indicating an up-regulation of major histocompatibility

complex (MHC) class I and II antigens in Tg-IGF2 islets, likely due, at least in part, to macrophage infiltration. These results suggested that Tg-IGF2 β -cells could be more susceptible to immune cell-mediated damage.

To verify this hypothesis and to determine the degree of sensitivity to damage of their β -cells, Tg-IGF2 mice received five daily consecutive injections of very low doses of streptozotocin (STZ), 20 mg/kg bw, a dose too low to affect WT mice (29). In contrast to WT mice, which remained normoglycemic, Tg-IGF2 mice showed a progressive increase in blood glucose levels (Fig. 6*A*) with $\sim 45\%$ of transgenics becoming diabetic 6 months post-STZ administration (Fig. 6*B*). This increased incidence of diabetes was due to a drastic reduction in β -cell mass in STZ-treated Tg-IGF2 mice (Fig. 6, *C* and *D*) even though these mice have a pre-STZ β -cell mass 3-fold higher than that of WT mice (27). Thus, these results confirmed that local overexpression of IGF2 in β -cells increased susceptibility to injury caused by an external toxin, such as STZ.




FIGURE 6. Transgenic IGF2 mice are more sensitive to streptozotocin treatment. *A*, blood glucose levels were monitored in fed mice for 6 months following treatment with very low doses of STZ for 5 consecutive days (20 mg/kg bw). WT mice (white squares) and transgenic IGF2 mice (blue circles) ($n = 15$ animals/group). *B*, cumulative incidence of diabetes before and after STZ treatment. Mice were considered diabetic when two consecutive glucose measurements were >250 mg/dl. WT mice (white squares) and transgenic IGF2 mice (black squares) ($n = 15$ animals/group). *C*, insulin immunostaining of pancreatic sections 6 months after STZ treatment. Transgenic IGF2 mice clearly presented fewer insulin-positive cells. Representative images, original magnification $\times 10$ (scale bar, 100 μ m). *D*, β -cell mass was measured 6 months after STZ treatment in WT (white bars) and Tg-IGF2 (black bars) mice. Nine pancreatic sections from each individual mouse (four mice/group) were analyzed as described under "Experimental Procedures." Results are expressed as mean \pm S.E. #, $p < 0.05$ versus WT; **, $p < 0.01$ versus Tg-IGF2-STZ.

To further study the increased susceptibility of Tg-IGF2 β -cells to damage, Tg-IGF2 mice (C57Bl6/SJL background) were crossed with transgenic mice expressing human interferon β specifically in β -cells (Tg-IFN β , CD1 background). Tg-IFN β mice have islet lymphocytic infiltration and increased susceptibility to develop autoimmune diabetes with similar characteristics to humans type 1 diabetes (29). Although WT, Tg-IGF2, and Tg-IFN β mice (all in 50% C57Bl6/SJL-50% CD1 background) remained normoglycemic during the 12-month follow-up, double transgenic mice Tg-IFN β /IGF2 (50% C57Bl6/SJL-50% CD1 background) were highly hyperglycemic as early as 1 month of age (Fig. 7A), with 75% of them developing spontaneous diabetes during the first 2 months of life (Fig. 7B). At 3 month of age, WT and Tg-IFN β mice were normoinsulinemic and had normal β -cell mass, whereas Tg-IGF2 mice showed hyperinsulinemia and β -cell hyperplasia (Fig. 7, C and D). β -Cell hyperplasia in Tg-IGF2 mice was lower than what we previously described (Fig. 6D) (27). This was likely due to a variation in the genetic background from 100% C57Bl6/SJL in the original study, a background more prone to islet hyperplasia (37, 38), to 50% C57Bl6/SJL-50%CD1 in the Tg-IGF2 that were littermates of Tg-IFN β /IGF2. In agreement with the high inci-

dence of spontaneous diabetes, double Tg-IFN β /IGF2 mice had lower circulating levels of insulin and decreased β -cell mass (Fig. 7, C and D).

Discussion

Our group previously reported that local overexpression of IGF2 in β -cells predisposes to the development of T2D (27). Mice overexpressing IGF2 are mildly hyperglycemic, hyperinsulinemic, and insulin-resistant, and when subjected to a high-fat diet they develop overt diabetes (27). The increased β -cell mass and the high degree of islet disorganization present in Tg-IGF2 mice could account for the islet functional failure that characterizes these animals. The molecular bases underlying the observed phenotype, however, were not fully understood nor to what extent the overexpression of IGF2 during embryonic development contributed to the process. In this study, we demonstrate that β -cell-restricted IGF2 overexpression triggers β -cell dedifferentiation and dysfunction and increases their susceptibility to damage, highlighting the important role that this growth factor may have in the progression of T2D.

In our previous work, we reported that insulin expression in total pancreata and *in vitro* glucose-stimulated insulin secre-

IGF2 Overexpression in β -Cells




FIGURE 7. Double IFN β /IGF2 transgenic mice develop diabetes spontaneously. *A*, blood glucose levels were monitored in fed WT (white squares), transgenic IFN β (gray circles), transgenic IGF2 (black circles), and double transgenic IFN β /IGF2 mice (blue-white rhombus) mice from 0.5 to 12 months of age, $n = 40$ animals/group. *B*, cumulative incidence of diabetes in WT (white squares), transgenic IFN β (gray circles), transgenic IGF2 (black circles), and double transgenic IFN β /IGF2 (black rhombus) mice, $n = 40$ animals/group. Mice were considered diabetic when two consecutive glucose measurements were >250 mg/dl. *C*, serum insulin levels at 3 months of age in WT (white bars), Tg-IFN β (light gray bars), Tg-IGF2 (black bars), and double Tg-IGF2/IFN β (dark gray bars) mice. Results are expressed as mean \pm S.E. of 15 animals/group. *, $p < 0.05$ versus WT. *D*, β -cell mass was measured at 3 months of age in WT (white bars), Tg-IFN β (light gray bars), Tg-IGF2 (black bars), and double Tg-IFN β /IGF2 (dark gray bars) mice. Nine pancreatic sections from each individual mouse (four mice/group) were analyzed as described under "Experimental Procedures." Results are means \pm S.E. of six animals/group. *, $p < 0.05$; **, $p < 0.01$ versus WT.

tion by isolated islets were greater in Tg-IGF2 than in WT mice, which agreed with the hyperinsulinemia described in transgenic animals (27). For the *in vitro* experiments, an equal number of WT and Tg-IGF2 islets were used, but because transgenic islets are considerably larger, this study was not illustrative of the amount of insulin produced by each β -cell. In this work, we observed that Tg-IGF2 mice secreted less insulin in response to a glucose overload than WT mice. Other animal models with islet hyperplasia also display impaired insulin secretion, such as the IR/IRS-1 $^{-/-}$ and IRS1 $^{-/-}$ knock-out mice (39, 40), and this has been attributed to altered β -cell functionality. Noticeably, β -cell dysfunction and alterations in insulin secretion have been described to precede insulin resistance in human patients at risk of developing T2D (41, 42). Similarly, Tg-IGF2 mice show glucose intolerance from an early age (2 months) and develop insulin resistance when they are older than 4 months (27). Thus, β -cell failure would be a primary defect and not just a simple consequence of diabetes progression in this model. A clear reduction in the expression of the main glucose sensors of the β -cell, the glucose transporter GLUT2 and the glucose-phosphorylating enzyme glucokinase, was documented in islets

from Tg-IGF2 mice or in wild-type islets treated with recombinant IGF2 or with adenoviral vectors that mediate overexpression of IGF2 in β -cells. In addition, microarray profiling analysis of Tg-IGF2 islets revealed alterations in the expression of several genes involved in insulin secretion and cell transport, as well as of several transcription factors important for β -cell differentiation and functionality, such as *Pdx1*, *Isl1*, *MafA*, and *Beta2/Neurod1*. These changes could partially explain the clear β -cell dysfunction observed in our transgenic mice, mainly the reduced insulin synthesis and secretion, and the dedifferentiated phenotype of Tg-IGF2 β -cells. Other type 2 diabetic animal models, such as db/db mice and Zucker and Goto-Kakizaki rats, which have similar characteristics to Tg-IGF2 mice, *i.e.* islet hyperplasia, hyperinsulinemia, insulin resistance, and altered insulin secretion and response to glucose, also show a partial loss of β -cell phenotype (12, 43). Moreover, mice lacking FoxO1 specifically in β -cells have high blood glucose concentration associated with reduced β -cell mass that was attributed to β -cell dedifferentiation rather than to β -cell death (6). A recent study proposes that β -cell dedifferentiation, rather than apoptosis, is the main mechanism of loss of insulin-positive

cells in a mouse model of T2D and that intensive insulin therapy, through the reversal of hyperglycemia, leads to re-differentiation of β -cells to a mature phenotype (9). These observations suggest that β -cell dedifferentiation is a common process in T2D (6) and that at least some forms of diabetes mellitus may result from loss of β -cell identity (7, 8). Our study supports this notion by demonstrating that IGF2 alone is capable of inducing changes in the β -cell phenotype, increasing the susceptibility to develop diabetes. Of note, when Tg-IGF2 mice were treated with low nondiabetogenic STZ doses, ~45% of transgenic mice developed overt diabetes despite the fact that GLUT2, the transporter by which STZ enters the β -cell, was down-regulated.

Tg-IGF2 cells also presented other important cellular alterations. ER stress has been proposed as a mechanism to explain β -cell dysfunction and death in T2D (12, 44). Tg-IGF2 islets showed signs of ER stress. By electron microscopy, we observed the characteristic dilatations of the ER that accompany ER stress (31). In addition, the expression of markers of ER stress such as *Xbp1s* and *Ddit3/Chop* was increased, and the expression of *Serca2*, the main transporter of Ca^{2+} to the ER (45), was reduced. Some of these modifications in the expression of ER stress markers were also observed in wild-type islets treated with adenoviral vectors encoding *Igf2* or in wild-type islets treated with recombinant protein, indicating that IGF2 *per se* was sufficient to induce ER stress in β -cells.

On the other hand, Tg-IGF2 islets showed increased expression of genes involved in the immune response, such as MHC class I and II, and a considerable number of macrophages infiltrated transgenic islets. Similarly, islets from Goto-Kakizaki rats, a model of T2D, have macrophage infiltration and an increased number of MHC-II-positive cells (46, 47). Macrophage infiltration has also been observed in islets from human T2D subjects (41). The immunological alterations observed at the islet level could probably predispose IGF2 islets to be more susceptible to immune cell-mediated attack and β -cell damage. Indeed, double transgenic Tg-IFN β /IGF2 spontaneously developed overt diabetes at a very early age, without the need for administering STZ. These results indicate that islets overexpressing *Igf2* have increased susceptibility to β -cell death. In addition, increased macrophage infiltration and expression of MHC-II molecules could contribute to the increased sensitivity to STZ of Tg-IGF2 β -cells.

The reproducibility of the observations made in Tg-IGF2 islets or in wild-type islets treated with IGF2 recombinant protein or with *Igf2*-encoding adenoviral vectors supports the idea that most of the observations made were direct effects of IGF2. Therefore, these observations were not the result of the overexpression of the growth factor during fetal pancreatic development.

The binding of IGF2 to IGF-II/Man-6-P receptor, its own receptor, has no other biological function than clearing IGF2 from serum and targeting it for degradation in lysosomes (48), thus balancing IGF2 activity by controlling the extracellular levels of IGF2 (49). Our *in vitro* studies demonstrated that IGF2 effects on β -cells were at least partially mediated through IGF1R or INSR signaling, because inhibition of the common downstream effector PI3K reversed the down-regulation of

Glut2, *Foxa2*, and *Gck* expression. IGF2 can bind to INSR. In particular, isoform A (INSR-A) binds IGF2 with high affinity, and its activation by IGF2 elicits a unique signaling pathway different from that of insulin (50). Indeed, we observed a down-regulation of the INSR-A receptor, similar to what had previously been documented after hyperstimulation with insulin in various target tissues (51). Moreover, IGF2 can act through IGF1R and may even act through hybrid INSR-A/IGF1R receptors (52). Although we have observed detrimental effects of IGF2 on β -cell functionality, which resulted in increased sensitivity to β -cell damage, others have demonstrated that the interaction of IGF2 with the IGF1R is required for GLP-1-induced protection against β -cell apoptosis (53).

In summary, although there is still no direct evidence of the increase of IGF2 in T2D in humans, this study demonstrates that local overexpression of IGF2, specifically in β -cells, causes profound changes in β -cells that lead to their dedifferentiation, ER stress, and β -cell dysfunction. As a consequence of exposure to increased levels of IGF2, islets are predisposed to be more sensitive to immune- and nonimmune-mediated β -cell damage, triggering the onset of diabetes.

Acknowledgments—We thank Ivet Elias (Universitat Autònoma de Barcelona) for helpful discussions and Marta Moya (Universitat Autònoma de Barcelona) for technical support.

References

- Butler, P. C., Meier, J. J., Butler, A. E., and Bhushan, A. (2007) The replication of beta cells in normal physiology, in disease, and for therapy. *Nat. Clin. Pract. Endocrinol. Metab.* **3**, 758–768
- Chang-Chen, K. J., Mullur, R., Bernal-Mizrachi, E. (2008) β -Cell failure as a complication of diabetes. *Rev. Endocr. Metab. Disord.* **9**, 329–343
- Jonas, J.-C., Sharma, A., Hasenkamp, W., Ilkova, H., Patané, G., Laybutt, R., Bonner-Weir, S., and Weir, G. C. (1999) Chronic hyperglycemia triggers loss of pancreatic β cell differentiation in an animal model of diabetes. *J. Biol. Chem.* **274**, 14112–14121
- Weir, G. C., Aguayo-Mazzucato, C., Bonner-Weir, S. (2013) β -Cell dedifferentiation in diabetes is important, but what is it? *Islets* **5**, 233–237
- Weir, G. C., Marselli, L., Marchetti, P., Katsuta, H., Jung, M. H., Bonner-Weir, S. (2009) Towards a better understanding of the contributions of overwork and glucotoxicity to the beta-cell inadequacy of type 2 diabetes. *Diabetes Obes. Metab.* **11**, 82–90
- Talchai, C., Xuan, S., Lin, H. V., Sussel, L., Accili, D. (2012) Pancreatic β cell dedifferentiation as a mechanism of diabetic β cell failure. *Cell* **150**, 1223–1234
- Puri, S., Akiyama, H., Hebrok, M. (2013) VHL-mediated disruption of Sox9 activity compromises β -cell identity and results in diabetes mellitus. *Genes Dev.* **27**, 2563–2575
- Guo, S., Dai, C., Guo, M., Taylor, B., Harmon, J. S., Sander, M., Robertson, R. P., Powers, A. C., and Stein, R. (2013) Inactivation of specific β cell transcription factors in type 2 diabetes. *J. Clin. Invest.* **123**, 3305–3316
- Wang, Z., York, N. W., Nichols, C. G., Remedi, M. S. (2014) Pancreatic β cell dedifferentiation in diabetes and redifferentiation following insulin therapy. *Cell Metab.* **19**, 872–882
- Kitamura, T. (2013) The role of FOXO1 in β -cell failure and type 2 diabetes mellitus. *Nat. Rev. Endocrinol.* **9**, 615–623
- Iwawaki, T., and Oikawa, D. (2013) The role of the unfolded protein response in diabetes mellitus. *Semin. Immunopathol.* **35**, 333–350
- Chan, J. Y., Luzuriaga, J., Bensellam, M., Biden, T. J., Laybutt, D. R. (2013) Failure of the adaptive unfolded protein response in islets of obese mice is linked with abnormalities in β -cell gene expression and progression to diabetes. *Diabetes* **62**, 1557–1568

13. Tersey, S. A., Nishiki, Y., Templin, A. T., Cabrera, S. M., Stull, N. D., Colvin, S. C., Evans-Molina, C., Rickus, J. L., Maier, B., and Mirmira, R. G. (2012) Islet β -cell endoplasmic reticulum stress precedes the onset of type 1 diabetes in the nonobese diabetic mouse model. *Diabetes* **61**, 818–827
14. Engin, F., Yermalovich, A., Nguyen, T., Ngyuen, T., Hummasti, S., Fu, W., Eizirik, D. L., Mathis, D., and Hotamisligil, G. S. (2013) Restoration of the unfolded protein response in pancreatic β cells protects mice against type 1 diabetes. *Sci. Transl. Med.* **5**, 211ra156
15. Allagnat, F., Christulia, F., Ortis, F., Pirot, P., Lortz, S., Lenzen, S., Eizirik, D. L., and Cardozo, A. K. (2010) Sustained production of spliced X-box binding protein 1 (XBP1) induces pancreatic beta cell dysfunction and apoptosis. *Diabetologia* **53**, 1120–1130
16. LeRoith, D., Roberts, C. T., Jr. (2003) The insulin-like growth factor system and cancer. *Cancer Lett.* **195**, 127–137
17. Monk, D., Sanches, R., Arnaud, P., Apostolidou, S., Hills, F. A., Abu-Amero, S., Murrell, A., Friess, H., Reik, W., Stanier, P., Constância, M., and Moore, G. E. (2006) Imprinting of IGF2 P0 transcript and novel alternatively spliced INS-IGF2 isoforms show differences between mouse and human. *Hum. Mol. Genet.* **15**, 1259–1269
18. Nica, A. C., Ongen, H., Irminger, J.-C., Bosco, D., Berney, T., Antonarakis, S. E., Halban, P. A., and Dermitzakis, E. T. (2013) Cell-type, allelic, and genetic signatures in the human pancreatic beta cell transcriptome. *Genome Res.* **23**, 1554–1562
19. Awata, T., Kawasaki, E., Ikegami, H., Kobayashi, T., Maruyama, T., Nakanishi, K., Shimada, A., Iizuka, H., Kurihara, S., Osaki, M., Uga, M., Kawabata, Y., Tanaka, S., Kanazawa, Y., and Katayama, S. (2007) Insulin gene/IDDM2 locus in Japanese type 1 diabetes: contribution of class I alleles and influence of class I subdivision in susceptibility to type 1 diabetes. *J. Clin. Endocrinol. Metab.* **92**, 1791–1795
20. Pugliese, A., and Miceli, D. (2002) The insulin gene in diabetes. *Diabetes Metab. Res. Rev.* **18**, 13–25
21. Huxtable, S. J., Saker, P. J., Haddad, L., Walker, M., Frayling, T. M., Levy, J. C., Hitman, G. A., O'Rahilly, S., Hattersley, A. T., and McCarthy, M. I. (2000) Analysis of parent-offspring trios provides evidence for linkage and association between the insulin gene and type 2 diabetes mediated exclusively through paternally transmitted class III variable number tandem repeat alleles. *Diabetes* **49**, 126–130
22. Dai, N., Rapley, J., Angel, M., Yanik, M. F., Blower, M. D., and Avruch, J. (2011) mTOR phosphorylates IMP2 to promote IGF2 mRNA translation by internal ribosomal entry. *Genes Dev.* **25**, 1159–1172
23. Zeggini, E., Weedon, M. N., Lindgren, C. M., Frayling, T. M., Elliott, K. S., Lango, H., Timpson, N. J., Perry, J. R., Rayner, N. W., Freathy, R. M., Barrett, J. C., Shields, B., Morris, A. P., Ellard, S., Groves, C. J., et al. (2007) Replication of genome-wide association signals in UK samples reveals risk loci for type 2 diabetes. *Science* **316**, 1336–1341
24. Chon, S. J., Kim, S. Y., Cho, N. R., Min, D. L., Hwang, Y. J., and Mamura, M. (2013) Association of variants in PPAR γ 2, IGF2BP2, and KCNQ1 with a susceptibility to gestational diabetes mellitus in a Korean population. *Yonsei Med. J.* **54**, 352–357
25. Benrahma, H., Charoute, H., Lasram, K., Boulouiz, R., Atig, R. K., Fakiri, M., Rouba, H., Abdelhak, S., and Barakat, A. (2014) Association analysis of IGF2BP2, KCNJ11, and CDKAL1 polymorphisms with type 2 diabetes mellitus in a Moroccan population: a case-control study and meta-analysis. *Biochem. Genet.* **52**, 430–442
26. Höög, A., Sandberg-Nordqvist, A. C., Abdel-Halim, S. M., Carlsson-Skiriut, C., Guenifi, A., Tally, M., Ostenson, C. G., Falkmer, S., Sara, V. R., Efendić, S., Schalling, M., and Grimelius, L. (1996) Increased amounts of a high molecular weight insulin-like growth factor II (IGF-II) peptide and IGF-II messenger ribonucleic acid in pancreatic islets of diabetic Goto-Kakizaki rats. *Endocrinology* **137**, 2415–2423
27. Devedjian, J. C., George, M., Casellas, A., Pujol, A., Visa, J., Pelegrin, M., Gros, L., and Bosch, F. (2000) Transgenic mice overexpressing insulin-like growth factor-II in beta cells develop type 2 diabetes. *J. Clin. Invest.* **105**, 731–740
28. Pelegrin, M., Devedjian, J. C., Costa, C., Visa, J., Solanes, G., Pujol, A., Asins, G., Valera, A., and Bosch, F. (1998) Evidence from transgenic mice that interferon- β may be involved in the onset of diabetes mellitus. *J. Biol. Chem.* **273**, 12332–12340
29. Casellas, A., Salavert, A., Agudo, J., Ayuso, E., Jimenez, V., Moya, M., Muñoz, S., Franckhauser, S., and Bosch, F. (2006) Expression of IGF-I in pancreatic islets prevents lymphocytic infiltration and protects mice from type 1 diabetes. *Diabetes* **55**, 3246–3255
30. Pfaffl, M. W. (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* **29**, e45–e45
31. Sachdeva, M. M., Claiborn, K. C., Khoo, C., Yang, J., Groff, D. N., Mirmira, R. G., and Stoffers, D. A. (2009) Pdx1 (MODY4) regulates pancreatic beta cell susceptibility to ER stress. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 19090–19095
32. Back, S. H., and Kaufman, R. J. (2012) Endoplasmic reticulum stress and type 2 diabetes. *Annu. Rev. Biochem.* **81**, 767–793
33. Vangheluwe, P., Raeymaekers, L., Dode, L., Wuytack, F. (2005) Modulating sarco(endo)plasmic reticulum Ca²⁺-ATPase 2 (SERCA2) activity: cell biological implications. *Cell Calcium* **38**, 291–302
34. Martino, L., Masini, M., Novelli, M., Befly, P., Bugliani, M., Marselli, L., Masiello, P., Marchetti, P., and De Tata, V. (2012) Palmitate activates autophagy in INS-1E β -cells and in isolated rat and human pancreatic islets. *PLoS One* **7**, e36188
35. Wirawan, E., Vanden Berghe, T., Lippens, S., Agostinis, P., and Vandenaabee, P. (2012) Autophagy: for better or for worse. *Cell Res.* **22**, 43–61
36. Friedrichsen, B. N., Neubauer, N., Lee, Y. C., Gram, V. K., Blume, N., Petersen, J. S., Nielsen, J. H., and Møldrup, A. (2006) Stimulation of pancreatic beta-cell replication by incretins involves transcriptional induction of cyclin D1 via multiple signalling pathways. *J. Endocrinol.* **188**, 481–492
37. Swenne, I., and Andersson, A. (1984) Effect of genetic background on the capacity for islet cell replication in mice. *Diabetologia* **27**, 464–467
38. Kulkarni, R. N., Almind, K., Goren, H. J., Winnay, J. N., Ueki, K., Okada, T., and Kahn, C. R. (2003) Impact of genetic background on development of hyperinsulinemia and diabetes in insulin receptor/insulin receptor substrate-1 double heterozygous mice. *Diabetes* **52**, 1528–1534
39. Brüning, J. C., Winnay, J., Bonner-Weir, S., Taylor, S. I., Accili, D., and Kahn, C. R. (1997) Development of a novel polygenic model of NIDDM in mice heterozygous for IR and IRS-1 null alleles. *Cell* **88**, 561–572
40. Kulkarni, R. N. (2002) Receptors for insulin and insulin-like growth factor-1 and insulin receptor substrate-1 mediate pathways that regulate islet function. *Biochem. Soc Trans.* **30**, 317–322
41. Gerich, J. E. (2002) Is reduced first-phase insulin release the earliest detectable abnormality in individuals destined to develop type 2 diabetes? *Diabetes* **51**, S117–S121
42. Wajchenberg, B. L. (2007) Beta-cell failure in diabetes and preservation by clinical treatment. *Endocr. Rev.* **28**, 187–218
43. Ohneda, M., Johnson, J. H., Inman, L. R., Chen, L., Suzuki, K., Goto, Y., Alam, T., Ravazzola, M., Orci, L., and Unger, R. H. (1993) GLUT2 expression and function in beta-cells of GK rats with NIDDM. Dissociation between reductions in glucose transport and glucose-stimulated insulin secretion. *Diabetes* **42**, 1065–1072
44. Eizirik, D. L., Cardozo, A. K., and Cnop, M. (2008) The role for endoplasmic reticulum stress in diabetes mellitus. *Endocr. Rev.* **29**, 42–61
45. Tuusa, J. T., Markkanen, P. M., Apaja, P. M., Hakalahti, A. E., Petäjä-Repo, U. E. (2007) The endoplasmic reticulum Ca²⁺-pump SERCA2b interacts with G protein-coupled receptors and enhances their expression at the cell surface. *J. Mol. Biol.* **371**, 622–638
46. Inaba, W., Mizukami, H., Kamata, K., Takahashi, K., Tsuboi, K., Yagihashi, S. (2012) Effects of long-term treatment with the dipeptidyl peptidase-4 inhibitor vildagliptin on islet endocrine cells in nonobese type 2 diabetic Goto-Kakizaki rats. *Eur. J. Pharmacol.* **691**, 297–306
47. Ehses, J. A., Perren, A., Eppler, E., Ribaux, P., Pospisilik, J. A., Maor-Cahn, R., Gueripel, X., Ellingsgaard, H., Schneider, M. K., Biollaz, G., Fontana, A., Reinecke, M., Homo-Delarche, F., and Donath, M. Y. (2007) Increased number of islet-associated macrophages in type 2 diabetes. *Diabetes* **56**, 2356–2370
48. Nielsen, F. C. (1992) The molecular and cellular biology of insulin-like growth factor II. *Prog. Growth Factor Res.* **4**, 257–290
49. Forejt, J., and Gregorová, S. (1992) Genetic analysis of genomic imprinting: an Imprinter-1 gene controls inactivation of the paternal copy of the mouse Tme locus. *Cell* **70**, 443–450
50. Morcavallo, A., Gaspari, M., Pandini, G., Palummo, A., Cuda, G., Larsen,

- M. R., Vigneri, R., and Belfiore, A. (2011) Research resource: new and diverse substrates for the insulin receptor isoform A revealed by quantitative proteomics after stimulation with IGF-II or insulin. *Mol. Endocrinol.* **25**, 1456–1468
51. Okabayashi, Y., Maddux, B. A., McDonald, A. R., Logsdon, C. D., Williams, J. A., and Goldfine, I. D. (1989) Mechanisms of insulin-induced insulin-receptor down-regulation: decrease of receptor biosynthesis and mRNA levels. *Diabetes* **38**, 182–187
52. Djiogue, S., Nwabo Kamdje, A. H., Vecchio, L., Kipanyula, M. J., Farahna, M., Aldebasi, Y., and Seke Etet, P. F. (2013) Insulin resistance and cancer: the role of insulin and IGFs. *Endocr. Relat. Cancer* **20**, R1–R17
53. Cornu, M., Yang, J. Y., Jaccard, E., Poussin, C., Widmann, C., and Thorens, B. (2009) Glucagon-like peptide-1 protects beta-cells against apoptosis by increasing the activity of an IGF-2/IGF-1 receptor autocrine loop. *Diabetes* **58**, 1816–1825

Insulin-like Growth Factor 2 Overexpression Induces β -Cell Dysfunction and Increases Beta-cell Susceptibility to Damage

Alba Casellas, Cristina Mallol, Ariana Salavert, Veronica Jimenez, Miquel Garcia, Judith Agudo, Mercè Obach, Virginia Haurigot, Laia Vilà, Maria Molas, Ricardo Lage, Meritxell Morró, Estefania Casana, Jesús Ruberte and Fatima Bosch

J. Biol. Chem. 2015, 290:16772-16785.

doi: 10.1074/jbc.M115.642041 originally published online May 13, 2015

Access the most updated version of this article at doi: [10.1074/jbc.M115.642041](https://doi.org/10.1074/jbc.M115.642041)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 53 references, 23 of which can be accessed free at <http://www.jbc.org/content/290/27/16772.full.html#ref-list-1>