

FORCING AND ENTROPY OF STRIP PATTERNS OF QUASIPERIODIC SKEW PRODUCTS IN THE CYLINDER

LLUÍS ALSÈDÀ, FRANCESC MAÑOSAS, AND LEOPOLDO MORALES

ABSTRACT. We extend the results and techniques from [7] to study the combinatorial dynamics (*forcing*) and entropy of quasiperiodically forced skew-products on the cylinder. For these maps we prove that a cyclic permutation τ forces a cyclic permutation ν as interval patterns if and only if τ forces ν as cylinder patterns. This result gives as a corollary the Sharkovskii Theorem for quasiperiodically forced skew-products on the cylinder proved in [7].

Next, the notion of *s*-horseshoe is defined for quasiperiodically forced skew-products on the cylinder and it is proved, as in the interval case, that if a quasiperiodically forced skew-product on the cylinder has an *s*-horseshoe then its topological entropy is larger than or equals to $\log(s)$.

Finally, if a quasiperiodically forced skew-product on the cylinder has a periodic orbit with pattern τ , then $h(F) \geq h(f_\tau)$, where f_τ denotes the *connect-the-dots* interval map over a periodic orbit with pattern τ . This implies that if the period of τ is $2^n q$ with $n \geq 0$ and $q \geq 1$ odd, then $h(F) \geq \frac{\log(\lambda_q)}{2^n}$, where $\lambda_1 = 1$ and, for each $q \geq 3$, λ_q is the largest root of the polynomial $x^q - 2x^{q-2} - 1$. Moreover, for every $m = 2^n q$ with $n \geq 0$ and $q \geq 1$ odd, there exists a quasiperiodically forced skew-product on the cylinder F_m with a periodic orbit of period m such that $h(F_m) = \frac{\log(\lambda_q)}{2^n}$. This extends the analogous result for interval maps to quasiperiodically forced skew-products on the cylinder.

1. INTRODUCTION

In this paper we want to study the coexistence and implications between periodic objects of maps on the cylinder $\Omega = \mathbb{S}^1 \times \mathbb{I}$, of the form:

$$F: \begin{pmatrix} \theta \\ x \end{pmatrix} \longrightarrow \begin{pmatrix} R_\omega(\theta) \\ f(\theta, x) \end{pmatrix},$$

where $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z}$, $\mathbb{I} = [0, 1]$, $R_\omega(\theta) = \theta + \omega \pmod{1}$ with $\omega \in \mathbb{R} \setminus \mathbb{Q}$ and $f(\theta, x) = f_\theta(x)$ is continuous on both variables. To study this class of maps, in [7], were developed clever techniques that lead to a theorem of the Sharkovskii type for this class of maps and periodic orbits of appropriate objects.

We aim at extending these results and techniques to study the combinatorial dynamics (*forcing*) and entropy of the skew-products from the class $\mathcal{S}(\Omega)$ consisting on all maps of the above type.

As already remarked in [7], instead of \mathbb{S}^1 we could take any compact metric space Θ that admits a minimal homeomorphism $R: \Theta \longrightarrow \Theta$ such that R^ℓ is minimal for every $\ell > 1$. However, for simplicity and clarity we will remain in the class $\mathcal{S}(\Omega)$.

Date: November 11, 2014.

1991 Mathematics Subject Classification. Primary: 37C55, 34D08, 37C70.

Key words and phrases. Quasiperiodically forced systems on the cylinder, combinatorial dynamics, forcing entropy, irrational rotation.

The authors have been partially supported by MINECO grant numbers MTM2008-01486 and MTM2011-26995-C02-01.

Before stating the main results of this paper, we will recall the extension of Sharkovskii Theorem to $\mathcal{S}(\Omega)$ from [7], together with the necessary notation. We start by clarifying the notion of a periodic orbit for maps from $\mathcal{S}(\Omega)$. To this end we informally introduce some key notions that will be defined more precisely in Section 2.

Let X be a compact metric space. A subset $G \subset X$ is *residual* if it contains the intersection of a countable family of open dense subsets in X .

In what follows, $\pi: \Omega \rightarrow \mathbb{S}^1$ will denote the standard projection from Ω to the circle.

Instead of periodic points we use objects that project over the whole \mathbb{S}^1 , called *strips* in [7, Definition 3.9]. A *strip in Ω* is a closed set $B \subset \Omega$ such that $\pi(B) = \mathbb{S}^1$ (i.e., B projects on the whole \mathbb{S}^1) and $\pi^{-1}(\theta) \cap B$ is a closed interval (perhaps degenerate to a point) for every θ in a residual set of \mathbb{S}^1 .

Given two strips A and B , we will write $A < B$ and $A \leq B$ ([7, Definition 3.13]) if there exists a residual set $G \subset \mathbb{S}^1$, such that for every $(\theta, x) \in A \cap \pi^{-1}(G)$ and $(\theta, y) \in B \cap \pi^{-1}(G)$ it follows that $x < y$ and, respectively, $x \leq y$. We say that the strips A and B are *ordered*¹ (respectively *weakly ordered*) if either $A < B$ or $A > B$ (respectively $A \leq B$ or $A \geq B$).

Given $F \in \mathcal{S}(\Omega)$ and $n \in \mathbb{N}$, a strip $B \subset \Omega$ is called *n-periodic for F* ([7, Definition 3.15]), if $F^n(B) = B$ and the image sets $B, F(B), F^2(B), \dots, F^{n-1}(B)$ are pairwise disjoint and pairwise ordered.

To state the main theorem of [7] we need to recall the *Sharkovskii Ordering* ([9, 10]). The *Sharkovskii Ordering* is a linear ordering of \mathbb{N} defined as follows:

$$\begin{aligned} & 3_{\text{Sh}} > 5_{\text{Sh}} > 7_{\text{Sh}} > 9_{\text{Sh}} > \cdots_{\text{Sh}} > \\ & 2 \cdot 3_{\text{Sh}} > 2 \cdot 5_{\text{Sh}} > 2 \cdot 7_{\text{Sh}} > 2 \cdot 9_{\text{Sh}} > \cdots_{\text{Sh}} > \\ & 4 \cdot 3_{\text{Sh}} > 4 \cdot 5_{\text{Sh}} > 4 \cdot 7_{\text{Sh}} > 4 \cdot 9_{\text{Sh}} > \cdots_{\text{Sh}} > \\ & \quad \vdots \\ & 2^n \cdot 3_{\text{Sh}} > 2^n \cdot 5_{\text{Sh}} > 2^n \cdot 7_{\text{Sh}} > 2^n \cdot 9_{\text{Sh}} > \cdots_{\text{Sh}} > \\ & \quad \vdots \\ & \cdots_{\text{Sh}} > 2^n_{\text{Sh}} > \cdots_{\text{Sh}} > 16_{\text{Sh}} > 8_{\text{Sh}} > 4_{\text{Sh}} > 2_{\text{Sh}} > 1. \end{aligned}$$

In the ordering $_{\text{Sh}} \geq$ the least element is 1 and the largest one is 3. The supremum of the set $\{1, 2, 4, \dots, 2^n, \dots\}$ does not exist.

Sharkovskii Theorem for maps from $\mathcal{S}(\Omega)$ ([7]). *Assume that the map $F \in \mathcal{S}(\Omega)$ has a p -periodic strip. Then F has a q -periodic strip for every $q <_{\text{Sh}} p$.*

Our first main result (Theorem A) concerns the forcing relation. As we will see in detail, the *strips patterns* of periodic orbits of strips of maps from $\mathcal{S}(\Omega)$ can be formalized in a natural way as cyclic permutations, as in the case of the periodic patterns for interval maps. Our first main result states that a cyclic permutation τ forces a cyclic permutation ν as interval patterns if and only if τ forces ν as strips patterns.

Since the Sharkovskii Theorem in the interval follows from the forcing relation, a corollary of Theorem A is the Sharkovskii Theorem for maps from $\mathcal{S}(\Omega)$.

Next, an s -horseshoe for maps from $\mathcal{S}(\Omega)$ can be defined also in a natural way. Our second main result (Theorem B) states that if a map $F \in \mathcal{S}(\Omega)$ has an s -horseshoe then $h(F)$, the *topological entropy of F* , satisfies $h(F) \geq \log(s)$. This is a generalization of the well known result for the interval.

¹This notion will be defined with greater detail but equivalently in Definition 2.17. We are giving here this less technical definition just to simplify this general section.

The third main result of the paper (Theorem C) states that if a map $F \in \mathcal{S}(\Omega)$ has a periodic orbit of strips with strips pattern τ , then $h(F) \geq h(f_\tau)$, where f_τ denotes the *connect-the-dots* interval map over a periodic orbit with pattern τ . A corollary of this fact and the lower bounds of the topological entropy of interval maps from [3] is that, if the period of τ is $2^n q$ with $n \geq 0$ and $q \geq 1$ odd, then $h(F) \geq \frac{\log(\lambda_q)}{2^n}$, where $\lambda_1 = 1$ and, for each $q \geq 3$, λ_q is the largest root of the polynomial $x^q - 2x^{q-2} - 1$. Moreover, for every $m = 2^n q$ with $n \geq 0$ and $q \geq 1$ odd, there exists a quasiperiodically forced skew-product on the cylinder F_m with a periodic orbit of strips of period m such that $h(F_m) = \frac{\log(\lambda_q)}{2^n}$.

The paper is organized as follows. In Section 2 we introduce the notation and we state the results in detail and in Section 3 we prove Theorem A. Finally, in Section 4 we prove Theorems B and C.

2. DEFINITIONS AND STATEMENTS OF RESULTS

We start by recalling the notion of interval pattern and related results. Afterwards we will introduce the natural extension to the class $\mathcal{S}(\Omega)$ by defining the cylinder patterns.

In what follows we will denote the class of continuous maps from the interval \mathbb{I} to itself by $\mathcal{C}^0(\mathbb{I}, \mathbb{I})$.

2.1. Interval patterns. Given $f \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$, we say that $p \in \mathbb{I}$ is an n -periodic point of f if $f^n(p) = p$ and $f^j(p) \neq p$ for $j = 1, 2, \dots, n-1$. The set of points $\{p, f(p), f^2(p), \dots, f^{n-1}(p)\}$ will be called a *periodic orbit*. A periodic orbit $P = \{p_1, p_2, \dots, p_n\}$ is said to have the *spatial labelling* if $p_1 < p_2 < \dots < p_n$. In what follows, every periodic orbit will be assumed to have the spatial labelling unless otherwise stated.

Definition 2.1 (Interval pattern). Let $P = \{p_1 < p_2 < \dots < p_n\}$ be a periodic orbit of a map $f \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$ and let τ be a cyclic permutation over $\{1, 2, \dots, n\}$. The periodic orbit P is said to have the (*periodic*) *interval pattern* τ if and only if $f(p_i) = p_{\tau(i)}$ for $i = 1, 2, \dots, n$. The period of P , n , will also be called the *period* of τ . ▀

Remark 2.2. Every cyclic permutation can occur as interval pattern. ▀

To study the dynamics of functions from $\mathcal{C}^0(\mathbb{I}, \mathbb{I})$ we introduce the following ordering on the set of interval patterns.

Definition 2.3 (Forcing). Given two interval patterns τ and ν , we say that τ *forces* ν , as *interval patterns*, denoted by $\tau \implies_{\mathbb{I}} \nu$, if and only if every $f \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$ that has a periodic orbit with interval pattern τ also has a periodic orbit with interval pattern ν . By [1, Theorem 2.5], the relation $\implies_{\mathbb{I}}$ is a partial ordering. ▀

Next we define a *canonical map* for an interval pattern as follows.

Definition 2.4 (τ -linear map). Let $f \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$ and let $P = \{p_1, p_2, \dots, p_n\}$ be an n -periodic orbit of f with the spatial labelling $(p_1 < p_2 < \dots < p_n)$. We define the P -*linear map* f_P as the unique map in $\mathcal{C}^0(\mathbb{I}, \mathbb{I})$ such that $f_P(p_i) = f(p_i)$ for $i = 1, 2, \dots, n$, f_P is affine in each interval of the form $[p_i, p_{i+1}]$ for $i = 1, 2, \dots, n-1$, and f_P is constant on each of the two connected components of $\mathbb{I} \setminus [p_1, p_n]$. The map f_P is also called P -*connect-the-dots map*.

Observe that the map f_P is uniquely determined by P and $f|_P$.

Let τ be the pattern of the periodic orbit P . The map f_P will also be called a τ -*linear map* and denoted by f_τ . Then the maps f_τ are not unique but all maps $f_\tau|_{[\min P, \max P]}$ are topologically conjugate and, thus, they have the same topological entropy and periodic orbits. ▀

The next result is a useful characterization of the forcing relation of interval patterns in terms of the τ -linear maps.

Theorem 2.5 (Characterization of the forcing relation). *Let τ and ν be two interval patterns. Then, $\tau \Longrightarrow_{\mathbb{I}} \nu$ if and only if f_τ has a periodic orbit with pattern ν .*

2.2. Strips Theory. In this subsection we introduce a new (more restrictive) kind of strips with better properties and we study the basic properties that we will need throughout the paper. To introduce this new kind of strips we first need to introduce the notion of a *core* of a set.

Given a compact metric space (X, d) we denote the set of all closed (compact) subsets of X by 2^X , and we endow this space with the Hausdorff metric

$$\begin{aligned} H_d(B, C) &= \max\{\max_{b \in B} \min_{c \in C} d(c, b), \max_{c \in C} \min_{b \in B} d(c, b)\} \\ &= \max\{\max_{b \in B} d(b, C), \max_{c \in C} d(c, B)\}. \end{aligned}$$

It is well known that $(2^X, H_d)$ is compact. Also, given a set A we will denote the closure of A by \overline{A} .

Definition 2.6 ([7]). Let M be a subset of 2^Ω . We define the *core* of M , denoted M^c , as

$$\bigcap_{G \in \mathcal{G}(\mathbb{S}^1)} \overline{M \cap \pi^{-1}(G)},$$

where $\mathcal{G}(\mathbb{S}^1)$ denotes the set of all residual subsets of \mathbb{S}^1 . Observe that if M is compact, then $M^c \subset M$ and, $\pi(M) = \mathbb{S}^1$ implies $\pi(M^c) = \mathbb{S}^1$. \blacksquare

This definition of *core* is rather intricate. Below we settle an equivalent and more useful definition in the spirit of Lemma 3.2 and Remark 3.3 of [7]. The role of the residual of continuity in this equivalent definition is stated without proof in [7] and, hence, we include the proof for completeness.

Let $M \in 2^\Omega$ be such that $\pi(M) = \mathbb{S}^1$. We define the map $\phi_M : \mathbb{S}^1 \rightarrow 2^{\mathbb{I}}$ by $\phi_M(\theta) := M^\theta$, and $G_M := \{\theta \in \mathbb{S}^1 : \phi_M \text{ is continuous at } \theta\}$. It can be easily seen that ϕ_M is upper semicontinuous (i.e. for every $\theta \in \mathbb{S}^1$ and every open $U \subset \mathbb{I}$ such that $\phi_M(\theta) \subset U$, $\{z \in \mathbb{S}^1 : \phi_M(z) \subset U\}$ is open in \mathbb{S}^1). Hence, by [6, Theorem 7.10], the set G_M is residual. The set G_M will be called the *residual of continuity* of M .

Given $G \subset \mathbb{S}^1$ and a map $\varphi : G \rightarrow 2^{\mathbb{I}}$, $\text{Graph}(\varphi) := \{(\theta, \varphi(\theta)) : \theta \in G\} \subset \mathbb{S}^1 \times 2^{\mathbb{I}}$ denotes the *graph* of φ . By abuse of notation we will identify $\text{Graph}(\varphi)$ with the set $\bigcup_{\theta \in G} \{\theta\} \times \varphi(\theta)$. Hence, we will consider $\text{Graph}(\varphi)$ as a subset of Ω (or of $G \times \mathbb{I}$), and $\overline{\text{Graph}(\varphi)}$ is a compact subset of Ω .

Lemma 2.7. *Let M be a compact subset of Ω . Then,*

$$M^c = \overline{\text{Graph}(\phi_M|_G)} = \overline{M \cap \pi^{-1}(G)}$$

for every residual set $G \subset G_M$. Moreover, $M \cap \pi^{-1}(G) = M^c \cap \pi^{-1}(G)$ and $(M^c)^c = M^c$.

Proof. We start by proving the first statement of the lemma. Notice that if

$$(1) \quad \overline{M \cap \pi^{-1}(G_M)} \subset \overline{M \cap \pi^{-1}(H)} \quad \text{for every } H \in \mathcal{G},$$

then

$$\overline{M \cap \pi^{-1}(G)} \subset \overline{M \cap \pi^{-1}(G_M)} \subset M^c = \bigcap_{H \in \mathcal{G}} \overline{M \cap \pi^{-1}(H)} \subset \overline{M \cap \pi^{-1}(G)}.$$

Hence, we only have to prove (1).

Let $H \in \mathcal{G}$ and let $(\theta, x) \in M \cap \pi^{-1}(G_M)$ (i.e. $\theta \in G_M$ and $(\theta, x) \in M^\theta = \phi_M(\theta)$). Since H is residual, it is dense in \mathbb{S}^1 . Therefore, there exists a sequence $\{\theta_n\}_{n=1}^\infty \subset H$

converging to θ . Since $\theta \in G_M$, ϕ_M is continuous in θ . So, $\lim \phi_M(\theta_n) = \phi_M(\theta)$ and, for every $\varepsilon > 0$ exists $N \in \mathbb{N}$ such that $d((\theta, x), \phi_M(\theta_n)) \leq H_d(\phi_M(\theta), \phi_M(\theta_n)) < \varepsilon$ for every $n \geq N$. Since the sets $\phi_M(\theta_n)$ are compact, for every $n \in \mathbb{N}$, there exists $(\theta_n, x_n) \in \phi_M(\theta_n) \subset M \cap \pi^{-1}(H)$ such that $d((\theta, x), (\theta_n, x_n)) = d((\theta, x), \phi_M(\theta_n))$. Thus, $\lim(\theta_n, x_n) = (\theta, x)$ and, hence, $(\theta, x) \in \overline{M \cap \pi^{-1}(H)}$. This implies $M \cap \pi^{-1}(G_M) \subset \overline{M \cap \pi^{-1}(H)}$ which, in turn, implies (1).

By the first statement,

$$M \cap \pi^{-1}(G) \subset \overline{M \cap \pi^{-1}(G)} \cap \pi^{-1}(G) = M^c \cap \pi^{-1}(G) \subset M \cap \pi^{-1}(G).$$

Now, to end the proof of the lemma, take $\tilde{G} = G_M \cap G_{M^c}$, which is a residual set. By the part of the lemma already proven we have,

$$(M^c)^c = \overline{M^c \cap \pi^{-1}(\tilde{G})} = \overline{M \cap \pi^{-1}(\tilde{G})} = M^c.$$

□

Now we are ready to define the notion of band.

Definition 2.8 (Band). Every strip $A \subset \Omega$ such that $A^c = A$ will be called a *band*. □

Remark 2.9. In view of Lemma 2.7 a band could be equivalently defined as follows: A *band* is a set of the form $\text{Graph}(\varphi)$, where φ is a continuous map from a residual set of \mathbb{S}^1 to the connected elements (intervals) of $2^{\mathbb{I}}$. □

Given $F \in \mathcal{S}(\Omega)$, a strip A is *F*-invariant if $F(A) \subset A$ and *F*-strongly invariant if $F(A) = A$. As usual, the intersection of two *F*-invariant strips is either empty or an *F*-invariant strip. An invariant strip is called *minimal* if it does not have a strictly contained invariant strip.

Remark 2.10. From Corollary 3.5 and Lemmas 3.10 and 3.11 of [7] it follows that the bands in Ω have the following properties for every map from $\mathcal{S}(\Omega)$:

- (1) The image of a band is a band.
- (2) Every invariant strip contains an invariant minimal strip.
- (3) Every invariant minimal strip is a strongly invariant band.

□

Moreover, the Sharkovskii Theorem for maps from $\mathcal{S}(\Omega)$ is indeed,

Sharkovskii Theorem for maps from $\mathcal{S}(\Omega)$ ([7]). *Assume that $F \in \mathcal{S}(\Omega)$ has a p -periodic strip. Then F has a q -periodic band for every $q <_{\text{sh}} p$.*

Next we introduce a particular kind of bands that play a key role in this theory since they allow us to better study and work with the bands.

Given a set $A \subset \Omega$ and $\theta \in \Omega$ we will denote the set $A \cap \pi^{-1}(\theta)$ by A^θ .

Definition 2.11. A band A is called *solid* when A^θ is an interval for every $\theta \in \mathbb{S}^1$ and $\delta(A) := \inf\{\text{diam}(A^\theta) : \theta \in \mathbb{S}^1\} > 0$. Also, A is called *pinched* if A^θ is a point for each θ in a residual subset of \mathbb{S}^1 . □

Remark 2.12. From [7, Theorem 4.11] it follows that there are only two kind of strongly invariant bands: *solid* or *pinched*. □

Despite of the fact that the above notion of pinched band is completely natural, for several reasons that will become clear later (see also [2]) we prefer to view the pinched bands as *pseudo-curves* in the spirit of Remark 2.9:

Definition 2.13. Let G be a residual set of \mathbb{S}^1 and let $\varphi : G \rightarrow \mathbb{I}$ be a continuous map from G to \mathbb{I} . The set $\overline{\text{Graph}(\varphi)}$ will be called a *pseudo-curve*. □

The next remark summarizes the basic properties of the pseudo-curves.

Remark 2.14. The following properties of the pseudo-curves are easy to prove:

- (1) Every pseudo-curve is a band. In particular $\pi \left(\overline{\text{Graph}(\varphi)} \right) = \mathbb{S}^1$.
- (2) The image of a pseudo-curve is a pseudo-curve. Moreover, every invariant pseudo-curve is strongly invariant and minimal.

Now assume that $\overline{\text{Graph}(\varphi)}$ is a pseudo-curve where φ is a map from G to \mathbb{I} . Then,

- (3) $G_{\overline{\text{Graph}(\varphi)}} \supset G$ (see e.g. [8, Lemma 7.2]).
- (4) $\overline{\text{Graph}(\varphi)} \cap \pi^{-1}(G) = \text{Graph}(\varphi)$.

□

Next we want to define a partial ordering in the set of strips. We recall that a map g from \mathbb{S}^1 to \mathbb{I} is *lower semicontinuous* (respectively *upper semicontinuous*) at $\theta \in \mathbb{S}^1$ if for every $\lambda < g(\theta)$ (respectively $\lambda > g(\theta)$) there exists a neighbourhood V of θ such that $\lambda < g(V)$ (respectively $\lambda > g(V)$). When this condition holds at every point in \mathbb{S}^1 g is said to be *lower semicontinuous on \mathbb{S}^1* (respectively *upper semicontinuous on \mathbb{S}^1*).

Definition 2.15 ([7, Definition 4.1(a)]). Given $A \in 2^\Omega$ such that $\pi(A) = \mathbb{S}^1$ we define the functions

$$\begin{aligned} M_A(\theta) &:= \max\{x \in \mathbb{I} : (\theta, x) \in A\} \\ m_A(\theta) &:= \min\{x \in \mathbb{I} : (\theta, x) \in A\}. \end{aligned}$$

It can be seen that M_A is an upper semicontinuous function from \mathbb{S}^1 to \mathbb{I} and m_A is a lower semicontinuous function from \mathbb{S}^1 to \mathbb{I} . From [6, Theorem 7.10], each of the functions m_A and M_A is continuous on a residual set of \mathbb{S}^1 . We denote by G_{m_A} (respectively G_{M_A}) the residual set of continuity of m_A (respectively M_A). □

Remark 2.16. If A is a pseudo-curve, it follows from [8, Lemma 7.2] that $G_A = G_{M_A} = G_{m_A} = \{\theta \in \mathbb{S}^1 : M_A(\theta) = m_A(\theta)\}$ (that is, A is pinched in $G_A = G_{M_A} = G_{m_A}$) and, hence,

$$A = \overline{\text{Graph} \left(M_A|_{G_{M_A}} \right)} = \overline{\text{Graph} \left(m_A|_{G_{m_A}} \right)}.$$

□

Definition 2.17 ([7, Definition 3.13]). Given two strips A and B we write $A < B$ (respectively $A \leq B$) if there exists a residual set G in \mathbb{S}^1 such that $M_A(\theta) < m_B(\theta)$ (respectively $M_A(\theta) \leq m_B(\theta)$) for all $\theta \in G$. We say that two strips are *ordered* (respectively *weakly ordered*) if either $A < B$ or $A > B$ (respectively $A \leq B$ or $A \geq B$). □

Remark 2.18. It follows from the definition that two (weakly) ordered strips have pairwise disjoint interiors. □

The above ordering can be better formulated in terms of the *covers* of a strip.

Definition 2.19. Let $A \subset \Omega$ be a strip. We define the *top cover* of A as the pseudo-curve defined by $M_A|_{G_{M_A}}$:

$$A^+ := \overline{\text{Graph} \left(M_A|_{G_{M_A}} \right)},$$

and the *bottom cover* of A as the pseudo-curve defined by $m_A|_{G_{m_A}}$:

$$A^- := \overline{\text{Graph}\left(m_A|_{G_{m_A}}\right)}.$$

□

Remark 2.20. The sets A^+ and A^- are bands but in general do not coincide with $\overline{\text{Graph}(M_A)}$ and $\overline{\text{Graph}(m_A)}$ respectively. Moreover, if A is a pseudo-curve then, from Remark 2.16, $A^+ = A^- = A$. □

Remark 2.21. Let A and B be two strips. By Remark 2.16 we have, $A < B$ if and only if $A^+ < B^-$ and $A \leq B$ if and only if $A^+ \leq B^-$. □

To end this subsection we introduce the useful notion of *band between two pseudo-curves*. Although this definition is inspired in the definition of a *basic strip* from [7] (see Definition 3.5) we follow our approach based in pseudo-curves.

Definition 2.22. Let A and B be pseudo-curves such that $A < B$. We define the *band between A and B* as:

$$E_{AB} := \overline{\bigcup_{\theta \in G_{M_A} \cap G_{m_B}} \{\theta\} \times (M_A(\theta), m_B(\theta))}.$$

□

The properties of the set E_{AB} are summarized by:

Lemma 2.23. *Let A and B be pseudo-curves such that $A < B$. Then,*

- (a) $E_{AB}^- = A$ and $E_{AB}^+ = B$. Moreover, $(E_{AB})^\theta = \{\theta\} \times [M_A(\theta), m_B(\theta)]$ for every $\theta \in G_{M_A} \cap G_{m_B}$.
- (b) E_{AB} is a band.
- (c) $E_{AB} := \overline{\text{Int}(E_{AB})}$. In particular, E_{AB} has non-empty interior.

Proof. From the definition of E_{AB} it follows that

$$\text{Graph}\left(M_A|_{G_{M_A} \cap G_{m_B}}\right) \subset E_{AB}.$$

Thus,

$$A = A^c = \overline{\text{Graph}\left(M_A|_{G_{M_A} \cap G_{m_B}}\right)} \subset E_{AB}$$

by Remarks 2.14(1) and 2.16 and Lemma 2.7. Consequently, $m_{E_{AB}} \leq m_A$. Now we will prove that $m_{E_{AB}} \geq m_A$ and, hence, $m_{E_{AB}} = m_A$. To see this note that, for every $\theta \in \mathbb{S}^1$, there exists a sequence

$$\{(\theta_n, x_n)\}_{n \in \mathbb{N}} \subset \bigcup_{\theta \in G_{M_A} \cap G_{m_B}} \{\theta\} \times (M_A(\theta), m_B(\theta))$$

which converges to $(\theta, m_{E_{AB}}(\theta))$. Observe that $x_n \geq M_A(\theta_n) \geq m_A(\theta_n)$ for every n . Therefore, by Remark 2.16 $m_{E_{AB}}(\theta) = \lim_n x_n \geq \liminf_n m_A(\theta_n) \geq m_A(\theta)$.

Since $m_{E_{AB}} = m_A$, from Definition 2.19 and Remark 2.16 it follows that $E_{AB}^- = A$.

In a similar way we get that $M_{E_{AB}} = M_B$ and $E_{AB}^+ = B$.

Then, by the part already proven and Remark 2.16,

$$(2) \quad (E_{AB})^\theta = \{\theta\} \times [M_A(\theta), m_B(\theta)] \quad \text{for every } \theta \in G_{M_A} \cap G_{m_B}.$$

This ends the proof of (a).

Now we prove (b). From the previous statement it follows that E_{AB} is a strip. Hence, we have to show that $(E_{AB})^c = E_{AB}$ which, by Definition 2.6, it reduces to prove that $E_{AB} \subset (E_{AB})^c$. Moreover, it is enough to show that

$$(3) \quad E_{AB}^\theta \subset (E_{AB})^c \quad \text{for every } \theta \in G_{M_A} \cap G_{m_B}$$

because, by (2),

$$E_{AB} \subset \overline{\bigcup_{\theta \in G_{M_A} \cap G_{m_B}} E_{AB}^\theta} \subset \overline{(E_{AB})^c} = (E_{AB})^c.$$

To prove (3) observe that, since $G_{M_A} \cap G_{m_B} \cap G_{E_{AB}}$ is a residual set (contained in $G_{E_{AB}}$), from Lemma 2.7 we get

$$(4) \quad (E_{AB})^c = \overline{E_{AB} \cap \pi^{-1}(G_{M_A} \cap G_{m_B} \cap G_{E_{AB}})} = \overline{\bigcup_{\theta \in G_{M_A} \cap G_{m_B} \cap G_{E_{AB}}} E_{AB}^\theta}.$$

In particular,

$$\bigcup_{\theta \in G_{M_A} \cap G_{m_B} \cap G_{E_{AB}}} E_{AB}^\theta \subset (E_{AB})^c.$$

Fix $\theta \in (G_{M_A} \cap G_{m_B}) \setminus G_{E_{AB}}$. Since $G_{M_A} \cap G_{m_B} \cap G_{E_{AB}}$ is a residual set, there exists a sequence $\{\theta_n\}_{n=1}^\infty \subset G_{M_A} \cap G_{m_B} \cap G_{E_{AB}}$ whose limit is θ . The continuity of the functions M_A and m_B in $G_{M_A} \cap G_{m_B}$ implies that $\lim M_A(\theta_n) = M_A(\theta)$ and $\lim m_B(\theta_n) = m_B(\theta)$. Therefore, again by (2), every point of E_{AB}^θ is limit of points in $\{E_{AB}^{\theta_n}\}_{n=1}^\infty$. This implies that $E_{AB}^\theta \subset (E_{AB})^c$ by (4). This ends the proof of (b).

To prove (c) observe that $\overline{\text{Int}(E_{AB})} \subset E_{AB}$. So, it is enough to show that

$$\bigcup_{\theta \in G_{M_A} \cap G_{m_B}} \{\theta\} \times (M_A(\theta), m_B(\theta)) \subset \text{Int}(E_{AB}).$$

Take $(\theta, x) \in \{\theta\} \times (M_A(\theta), m_B(\theta))$ with $\theta \in G_{M_A} \cap G_{m_B}$. Since $x \neq M_A(\theta)$ and $x \neq m_B(\theta)$, there exists $\varepsilon > 0$ such that $x > M_A(\theta) + \varepsilon$ and $x < m_B(\theta) - \varepsilon$. On the other hand, the continuity of M_A and m_B on $G_{M_A} \cap G_{m_B}$ implies that there exist $\delta > 0$ such that $\theta' \in G_{M_A} \cap G_{m_B}$ and $|\theta - \theta'| < \delta$ implies $|M_A(\theta) - M_A(\theta')| < \varepsilon$ and $|m_B(\theta) - m_B(\theta')| < \varepsilon$. Now we define

$$r := \min\{\delta, |x - M_A(\theta) - \varepsilon|, |x - m_B(\theta) + \varepsilon|\} > 0.$$

Observe that, with this choice of r , $M_A(\theta) + \varepsilon \leq x - r < x + r \leq m_B(\theta) - \varepsilon$.

Let $U := \{(\theta', y) \in \Omega : |\theta - \theta'| < r \text{ and } |x - y| < r\}$ be an open neighbourhood of (θ, x) . We will prove that every $(\theta', y) \in U$ belongs to E_{AB} . If $\theta' \in G_{M_A} \cap G_{m_B}$, from the choice of δ and r , it follows that $(\theta', y) \in \{\theta'\} \times (x - r, x + r) \subset \{\theta'\} \times [M_A(\theta'), m_B(\theta')] \subset E_{AB}$. Now assume that $\theta' \notin G_{M_A} \cap G_{m_B}$ and consider a sequence $\{\theta_n\}_{n \in \mathbb{N}} \subset G_{M_A} \cap G_{m_B} \cap (\theta - r, \theta + r)$ converging to θ' . Clearly, $(\theta_n, y) \in U$ for every $n \in \mathbb{N}$ and, by the part already proven, $(\theta_n, y) \in E_{AB}$. Consequently, since E_{AB} is closed, $(\theta', y) = \lim(\theta_n, y) \in E_{AB}$. \square

2.3. Strip patterns. In this subsection we define the notion of strips pattern and forcing for maps from $\mathcal{S}(\Omega)$ along the lines of Subsection 2.1.

Definition 2.24 ([7, Definition 3.15]). Let $F \in \mathcal{S}(\Omega)$. We say that a strip $A \subseteq \Omega$ is a *p-periodic strip* if $F^p(A) = A$ and the strips $A, F(A), \dots, F^{p-1}(A)$ are pairwise disjoint and ordered. The set $\{A, F(A), \dots, F^{p-1}(A)\}$ is called an *n-periodic orbit of strips*.

By Remarks 2.10 and 2.12, it follows that we can restrict our attention to two kind of periodic orbit of bands: the solid ones and the pseudo-curves. \square

A periodic orbit of strips $\{B_1, B_2, \dots, B_p\}$ is said to have the *spatial labelling* if $B_1 < B_2 < \dots < B_p$. In what follows we will assume that every periodic orbit of strips has the spatial labelling.

Definition 2.25 (Strip pattern). Let $F \in \mathcal{S}(\Omega)$ and let $\mathcal{B} = \{B_1, B_2, \dots, B_n\}$ be a periodic orbit of strips. The *strips pattern* of \mathcal{B} is the permutation τ such that $F(B_i) = B_{\tau(i)}$ for every $i = 1, 2, \dots, n$.

When a map $F \in \mathcal{S}(\Omega)$ has a periodic orbit of strips with strips pattern τ we say that F *exhibits the pattern* τ . \square

Remark 2.26. Interval and strips patterns are formally the same algebraic objects; that is cyclic permutations. \square

Definition 2.27 (Forcing). Let τ and ν be strips patterns. We say that τ forces ν in Ω , denoted by $\tau \implies_{\Omega} \nu$, if and only if every map $F \in \mathcal{S}(\Omega)$ that exhibits the strips pattern τ also exhibits the quasiperiodic pattern ν . \square

The next theorem is the first main result of this paper. It characterizes the relation \implies_{Ω} by comparison with $\implies_{\mathbb{I}}$.

Theorem A. *Let τ and ν be patterns (both in \mathbb{I} and Ω). Then,*

$$\tau \implies_{\mathbb{I}} \nu \quad \text{if and only if} \quad \tau \implies_{\Omega} \nu.$$

The first important consequence of Theorem A is the next result which follows from the fact that the Sharkovskii theorem is a corollary of the forcing relation for interval maps.

Corollary 2.28. *The Sharkovskii Theorem for maps from $\mathcal{S}(\Omega)$ holds.*

Proof. Assume that $F \in \mathcal{S}(\Omega)$ exhibits a p -periodic strips pattern τ and let $q \in \mathbb{N}$ be such that $p_{\text{sh}} > q$. By [1, Corollary 2.7.4], $\tau \implies_{\mathbb{I}} \nu$ for some strips pattern ν of period q . Then, by Theorem A, $\tau \implies_{\Omega} \nu$ and, by definition, F also has a q -periodic orbit of strips (with strips pattern ν). Then the corollary follows from Remark 2.10(2,3). \square

Next we are going to study the relation between the forcing relation and the topological entropy of maps from $\mathcal{S}(\Omega)$. To this end we introduce the notion of horseshoe in $\mathcal{S}(\Omega)$.

Let $F \in \mathcal{S}(\Omega)$ and let A and B be bands in Ω . We say that A F -covers B if either $F(A^-) \leq B^-$ and $F(A^+) \geq B^+$, or $F(A^-) \geq B^+$ and $F(A^+) \leq B^-$.

Definition 2.29 (Horseshoe). An s -horseshoe for a map $F \in \mathcal{S}(\Omega)$ is a pair (J, \mathcal{D}) where J is a band and \mathcal{D} is a set of $s \geq 2$ pairwise weakly ordered bands, each of them with non-empty interior, such that L F -covers J for every $L \in \mathcal{D}$. Observe that, by Remark 2.18, the elements of \mathcal{D} have pairwise disjoint interiors. \square

The next theorem is the second main result of the paper. It relates the topological entropy of maps from $\mathcal{S}(\Omega)$ with horseshoes.

Theorem B. *Assume that $F \in \mathcal{S}(\Omega)$ has an s -horseshoe. Then*

$$h(F) \geq \log(s).$$

Next we want to introduce a class of maps that play the role of the connect-the-dots maps in the interval case and use them to study the topological entropy in relation with the periodic orbits of strips.

Definition 2.30 (Quasiperiodic τ -linear map). Given a strips pattern τ we define a quasiperiodic τ -linear map $F_\tau \in \mathcal{S}(\Omega)$ as:

$$F_\tau(\theta, x) := (R_\omega(\theta), f_\tau(x))$$

where R_ω is the irrational rotation by angle ω and f_τ is a τ -linear interval map (Definition 2.4 — recall that τ is also an interval pattern). \square

Remark 2.31. Since, by definition, f_τ has a periodic orbit with interval pattern τ , F_τ has a periodic orbit of bands (in fact curves which are horizontal circles) with strips pattern τ . \square

The next main result shows that the quasiperiodic τ -linear maps have minimal entropy among all maps from $\mathcal{S}(\Omega)$ which exhibit the strips pattern τ , again as in the interval case.

Theorem C. *Assume that $F \in \mathcal{S}(\Omega)$ exhibits the strips pattern τ . Then*

$$h(F) \geq h(F_\tau) = h(f_\tau).$$

Theorem C has an interesting consequence concerning the entropy of strips patterns that we define as follows.

Definition 2.32 (Entropy of strips patterns). Given a strips pattern τ we define the entropy of τ as

$$h(\tau) := \inf\{h(F) : F \in \mathcal{S}(\Omega) \text{ and } F \text{ exhibits the strips pattern } \tau\}.$$

\square

With this definition, in view of the Remark 2.31, Theorem C can be written as follows:

Theorem C. *Assume that $F \in \mathcal{S}(\Omega)$ exhibits the strips pattern τ . Then*

$$h(\tau) = h(F_\tau) = h(f_\tau).$$

By [1, Corollary 4.4.7] and [1, Lemma 4.4.11] we immediately get the following simple but important corollary of Theorem C which will allow us to obtain lower bounds of the topological entropy depending on the set of periods.

Corollary 2.33. *Assume that τ and ν are strips patterns such that $\tau \implies_\Omega \nu$. Then $h(\tau) \geq h(\nu)$.*

Corollary 2.34. *Assume that $F \in \mathcal{S}(\Omega)$ has a periodic orbit of strips of period $2^n q$ with $n \geq 0$ and $q \geq 1$ odd. Then,*

$$h(F) \geq \frac{\log(\lambda_q)}{2^n}$$

where $\lambda_1 = 1$ and, for each $q \geq 3$ odd, λ_q is the largest root of the polynomial $x^q - 2x^{q-2} - 1$. Moreover, for every $m = 2^n q$ with $n \geq 0$ and $q \geq 1$ odd, there exists a map $F_m \in \mathcal{S}(\Omega)$ with a periodic orbit of bands of period m such that $h(F_m) = \frac{\log(\lambda_q)}{2^n}$.

Proof. Let τ denote the strips pattern of a periodic orbit of strips of F of period $2^n q$. By Theorem C and [3] (see also Corollaries 4.4.7 and 4.4.18 of [1]) we get that

$$h(F) \geq h(f_\tau) \geq \frac{\log \lambda_q}{2^n}.$$

To prove the second statement we use [1, Theorem 4.4.17]: for every $m = 2^n q$ there exists a primary pattern ν_m of period m such that $h(f_{\nu_m}) = \frac{\log \lambda_q}{2^n}$. Then, from Theorem C, we can take $F_m = F_{\nu_m}$. \square

3. PROOF OF THEOREM A

To prove Theorem A we need some more notation and preliminary results.

An important tool in the study of patterns is the Markov graph. Signed Markov graphs are a specialization of Markov graphs. Next we define them and clarify the relation with our situation.

A *combinatorial (directed) signed graph* is defined as a pair $G = (V, \mathcal{A})$ where V is a finite set, called the *set of vertices*, and $\mathcal{A} \subset V \times V \times \{+, -\}$ is called the *set of signed arrows*. Given a *signed arrow* $\alpha = (I, J, s) \in \mathcal{A}$, I is the *beginning of* α , J is the *end of* α and s is the *sign of* α . Such an arrow α is denoted by $I \xrightarrow{s} J$.

3.1. Signed Markov graphs in the interval. We start by introducing the notion of *signed covering*. In what follows, $\text{Bd}(A)$ will denote the boundary of A .

Definition 3.1. Let $f \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$ and let $I, J \subset \mathbb{I}$ be two intervals. We say that I *positively F -covers* J , denoted by $I \xrightarrow{+} J$ (or $I \xrightarrow[f]{+} J$ if we need to specify the map), if $f(\min I) \leq \min J < \max J \leq f(\max I)$ and, analogously, we say that I *negatively F -covers* J , denoted by $I \xrightarrow{-} J$ (or $I \xrightarrow[f]{-} J$), if $f(\max I) \leq \min J < \max J \leq f(\min I)$. Observe that if $I \xrightarrow{s_1} J_1$ and $I \xrightarrow{s_2} J_2$ then $s_1 = s_2$.

We will write $I \xrightarrow{=s_1} J$ or $I \xrightarrow[f]{=s_1} J$ to denote that $f(I) = J$ and $I \xrightarrow[f]{s_1} J$ (in particular, $f(\text{Bd}(I)) = \text{Bd}(J)$). \blacksquare

We associate a signed graph to a periodic orbit of an interval map as follows.

Definition 3.2. Let $f \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$ and let P be a periodic orbit of f . A *P -basic interval* is the closure of a connected component of $[\min P, \max P] \setminus P$. The *P -signed Markov graph of f* is the combinatorial signed graph that has the set of all basic intervals as set of vertices V and the signed arrows in \mathcal{A} are the ones given by Definition 3.1. \blacksquare

Remark 3.3. Observe that the P -signed Markov graph of f depends only on $f|_P$ or more precisely on the pattern of P . It does not depend on the concrete choice of the points of P and on the graph of f outside P . Consequently, if P is a periodic orbit of $f \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$ and Q is a periodic orbit of $g \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$ with the same pattern then the P -signed Markov graph of f and the Q -signed Markov graph of g coincide. In particular, the P -signed Markov graph of f and the P -signed Markov graph of f_P coincide. \blacksquare

3.2. Signed Markov graphs in Ω . Now we also associate a signed graph to a periodic orbit of strips. We start by defining the notion of *signed covering* for bands. It is an improvement of the notion of F -covering introduced before.

Definition 3.4 (Signed covering [7, Definition 4.14]). Let $F \in \mathcal{S}(\Omega)$ and let A and B be bands in Ω . We say that A *positively F -covers* B , denoted by $A \xrightarrow{+} B$ (or $A \xrightarrow[F]{+} B$ if we need to specify the map), if² $F(A^-) \leq B^-$ and $F(A^+) \geq B^+$ and, analogously, we say that A *negatively F -covers* B , denoted by $A \xrightarrow{-} B$ (or $A \xrightarrow[F]{-} B$), if $F(A^-) \geq B^+$ and $F(A^+) \leq B^-$.

Observe that, as in the interval case (see Definition 3.1), if $A \xrightarrow{s_1} B_1$ and $A \xrightarrow{s_2} B_2$, then $s_1 = s_2$.

²Although these definitions are formally different from [7, Definition 4.14], they are equivalent by [7, Lemma 4.3(c,d)] and the definitions of the weak ordering of strips.

We will write $A \xrightarrow{s_1} B$ or $A \xrightarrow[F]{s_1} B$ to denote that $F(A) = B$ and $A \xrightarrow[F]{s_1} B$. \blacksquare

Next, by using the notion of band between two pseudo-curves, we will define the analogous of basic interval (*basic band*) and signed Markov graph for maps from $\mathcal{S}(\Omega)$.

Definition 3.5. Let $F \in \mathcal{S}(\Omega)$ and let $\mathcal{B} = \{B_1, B_2, \dots, B_n\}$ be a periodic orbit of strips of F with the spatial labelling (that is, $B_1 < B_2 < \dots < B_n$). For every $i = 1, 2, \dots, n-1$ the band (see Remark 2.21 and Lemma 2.23)

$$I_{B_i B_{i+1}} := E_{B_i^+ B_{i+1}^-} = \overline{\text{Int} \left(E_{B_i^+ B_{i+1}^-} \right)}$$

will be called a *basic band*. Observe that, from Lemma 2.23(a), $I_{B_i B_{i+1}}^- = B_i^+$ and $I_{B_i B_{i+1}}^+ = B_{i+1}^-$.

The \mathcal{B} -signed Markov graph of F is the combinatorial signed graph that has the set of all basic bands as set of vertices V and the signed arrows in \mathcal{A} are the ones given by Definition 3.4. \blacksquare

Clearly, all the basic bands are contained in $E_{B_1 B_n}$, $I_{B_i B_{i+1}} \leq I_{B_{i+1} B_{i+2}}$ for $i = 1, 2, \dots, n-2$ and if $I_{B_i B_{i+1}} \cap I_{B_j B_{j+1}} \neq \emptyset$ then $|i-j| = 1$.

Remark 3.6. As in the interval case (see Remark 3.3) the P -signed Markov graph of F is a pattern invariant. Moreover, if P is a periodic orbit of $F \in \mathcal{S}(\Omega)$ and Q is a periodic orbit of the interval map $f \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$ with the same pattern, then the P -signed Markov graph of F and the Q -signed Markov graph of f coincide. In particular, the P -signed Markov graph of F and the P -signed Markov graph of f_P coincide. \blacksquare

The following lemma summarizes the properties of basic bands and arrows. We will use it in the proof of Theorem A.

Lemma 3.7. *The following statements hold.*

(a) Let $F \in \mathcal{S}(\Omega)$ and let A and B be bands such that there is a signed arrow $A \xrightarrow{s} B$ from A to B in the signed Markov graph of F . Then,

(a.1) $F(A) \supset B$.

(a.2) $A \xrightarrow{s} D$ for every band $D \subset B$.

(a.3) There exists a band $C \subset A$ such that $C \xrightarrow{s} B$. Moreover, $F(C^+) \subset B^+$ and $F(C^-) \subset B^-$ if $s = +$, and $F(C^-) \subset B^+$ and $F(C^+) \subset B^-$ if $s = -$.

(a.4) Assume that $A \xrightarrow{s} \tilde{B}$ with $B \leq \tilde{B}$ and let C and \tilde{C} denote the bands given by (a.3) for B and \tilde{B} respectively. Then, $C \leq \tilde{C}$ if $s = +$, and $C \geq \tilde{C}$ if $s = -$.

(b) Let $F \in \mathcal{S}(\Omega)$ and let A be a band such that $A \xrightarrow{\pm} A$. Then there exists a band

$$A_\infty \subset A \text{ such that } A_\infty \xrightarrow{\pm} A_\infty.$$

Proof. Statement (a.1) is [7, Lemma 4.15] and (a.2) follows directly from the definitions. Statements (a.3,4) are [7, Lemma 4.19] while statement (b) is [7, Lemma 4.21]. \square

3.3. Loops of signed Markov graphs. Given a combinatorial signed Markov graph G , a sequence of arrows $\alpha = I_0 \xrightarrow{s_0} I_1 \xrightarrow{s_1} \dots \xrightarrow{s_{m-1}} I_{m-1}$ will be called a *path of length m* . The length of α will be denoted by $|\alpha|$. When a path begins

and ends in the same vertex (i.e. $I_{m-1} = I_0$) it will be called a *loop*. Observe that, then $I_1 \xrightarrow{s_1} I_2 \xrightarrow{s_2} \dots \xrightarrow{s_{m-2}} I_{m-1} \xrightarrow{s_m} I_1$ is also a loop in G . This loop is called a *shift* of α and denoted by $S(\alpha)$. For $n \geq 0$, we will denote by S^n the n -th iterate of the shift. That is,

$$S^n(\alpha) = I_{j_0} \xrightarrow{s_{j_0}} I_{j_1} \xrightarrow{s_{j_1}} I_{j_2} \xrightarrow{s_{j_2}} \dots \xrightarrow{s_{j_{m-2}}} I_{j_{m-1}},$$

where $j_r = r + n \pmod{m}$. Note that $S^{km}(\alpha) = \alpha$ for every $k \geq 0$.

Let $\alpha = I_0 \xrightarrow{s_0} I_1 \xrightarrow{s_1} \dots \xrightarrow{s_{m-2}} I_{m-1}$ and $\beta = J_0 \xrightarrow{r_0} J_1 \xrightarrow{r_1} \dots \xrightarrow{r_{l-2}} J_{l-1}$ be two paths such that the last vertex of α coincides with the first vertex of β (i.e. $I_{m-1} = J_0$). The path $I_0 \xrightarrow{s_0} I_1 \xrightarrow{s_1} \dots \xrightarrow{s_{m-1}} J_0 \xrightarrow{r_0} J_1 \xrightarrow{r_1} \dots \xrightarrow{r_{l-1}} J_{l-1}$ is the *concatenation* of α and β and is denoted by $\alpha\beta$. In this spirit, for every $n \geq 1$, α^n will denote the concatenation of α with himself n -times. the path α^n will be called the *n -repetition of α* . Also, α^∞ will denote the infinite path $\alpha\alpha\alpha\dots$.

A loop is called *simple* if it is not a repetition of a shorter loop. Observe that, in that case, the length of the shorter loop divides the length of the long one.

The next lemma translates the non-repetitiveness of a loop to conditions on its liftings. Its proof is folk knowledge.

Lemma 3.8. *Let α be a signed loop of length n in a combinatorial signed Markov graph G . If α is simple, $S^i(\alpha) \neq S^j(\alpha)$ for every $i \neq j$.*

Given a path $\alpha = I_0 \xrightarrow{s_0} I_1 \xrightarrow{s_1} \dots \xrightarrow{s_{m-1}} I_m$ we define the *sign* of α , denoted by $\text{Sign}(\alpha)$, as $\prod_{i=1}^m s_i$, where in this expression we use the obvious multiplication rules:

$$\begin{aligned} + \cdot + &= - \cdot - = +, \text{ and} \\ + \cdot - &= - \cdot + = -. \end{aligned}$$

Finally we introduce a (lexicographical) *ordering* in the set of paths of signed combinatorial graphs. To this end we start by introducing a linear ordering in the set of vertices. This ordering is arbitrary but fixed.

In the case of Markov graphs, the spatial labelling of orbits induces a natural ordering in the set of basic intervals or basic bands, which is the ordering that we are going to adopt. More precisely, if $P = \{p_0, p_1, \dots, p_{n-1}\}$ is a periodic orbit with the spatial labelling, then we endow the set of vertices (basic intervals) of the associated signed Markov graph with the following ordering:

$$[p_0, p_1] < [p_1, p_2] < \dots < [p_{n-2}, p_{n-1}].$$

Analogously, if $\mathcal{B} = \{B_0, B_1, \dots, B_{n-1}\}$ is a periodic orbit of strips with the spatial labelling, then we endow the set of vertices (basic intervals) of the associated signed Markov graph with the following ordering:

$$I_{B_0 B_1} < I_{B_1 B_2} < \dots < I_{B_{n-2} B_{n-1}}.$$

Then, the above ordering in the set of vertices naturally induces a *lexicographical ordering* in the set of paths of the signed combinatorial graph as follows. Let

$$\begin{aligned} \alpha &= I_0 \xrightarrow{s_0} I_1 \xrightarrow{s_1} \dots \xrightarrow{s_{n-1}} I_n \text{ and} \\ \beta &= J_0 \xrightarrow{r_0} J_1 \xrightarrow{r_1} \dots \xrightarrow{r_{m-1}} J_m \end{aligned}$$

be paths such that there exists $k \leq \min\{n, m\}$ with $I_k \neq J_k$ and $I_i = J_i$ for $i = 0, 1, \dots, k-1$ (recall that, by Definition 3.1, if $I_i = J_i$ then the signs s_i and r_i

of the corresponding arrows coincide). We write $\alpha < \beta$ if and only if

$$\begin{cases} I_k < J_k & \text{when } s = +, \text{ or} \\ I_k > J_k & \text{when } s = -, \end{cases}$$

where $s = \text{Sign} \left(I_0 \xrightarrow{s_0} I_1 \xrightarrow{s_1} \cdots I_{k-1} \xrightarrow{s_{k-1}} I_k \right) = s_0 s_1 \cdots s_{k-1}$.

Next we relate the loops in signed Markov graphs with periodic orbits.

Definition 3.9. Let $f \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$ and let p be a periodic point of f and let

$$\alpha = J_0 \xrightarrow{s_0} J_1 \xrightarrow{s_1} \cdots J_{n-1} \xrightarrow{s_{n-1}} J_0$$

be a loop in the P -signed Markov graph of f . We say that α and p are *associated* if p has period n and $f^i(p) \in J_i$ for every $i = 0, 1, \dots, n-1$. Observe that in such case $S^m(\alpha)$ and $f^m(p)$ are associated for all $m \geq 1$. \square

The next lemma relates the ordering of periodic points with the ordering of the associated loops. Its proof is a simple exercise.

Lemma 3.10. *Let $f \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$ and let f_P be a P -linear map, where P is a periodic orbit. Let x and y be two distinct periodic points of f_P associated respectively to two distinct loops α and β in the P -signed Markov graph of f_P . Then $x < y$ if and only if $\alpha < \beta$. Consequently, for every $n \geq 1$, $f^n(x) < f^n(y)$ if and only if $S^n(\alpha) < S^n(\beta)$.*

The next lemma is folk knowledge but we include the proof because we are not able to provide an explicit reference for it.

Lemma 3.11. *Let τ be a pattern and let $f_\tau = f_P$ be a P -linear map, where P is a periodic orbit of f_P of pattern τ . Assume that $\{q_0, q_1, q_2, \dots, q_m\}$ is a periodic orbit of f_τ with pattern $\nu \neq \tau$. Then there exists a unique loop α in the P -signed Markov graph of f_P associated to q_0 . Moreover, α is simple.*

Proof. The existence and unicity of the loop α follows from [1, Lemma 1.2.12]. We have to show that α is simple. Assume that α is the k repetition of a loop

$$\beta = J_0 \xrightarrow{s_0} J_1 \xrightarrow{s_1} \cdots J_{\ell-1} \xrightarrow{s_{\ell-1}} J_0$$

of length ℓ with $k \geq 2$ and $m = k\ell$. By [1, Lemma 1.2.6], there exist intervals $K_0 \subset J_0, K_1 \subset J_1, \dots, K_{\ell-1} \subset J_{\ell-1}$ such that $K_i \xrightarrow{s_i} K_{i+1}$ for $i = 0, 1, \dots, \ell-2$ and $K_{\ell-1} \xrightarrow{s_{\ell-1}} J_0$. Clearly, since f_P is P -linear, $f_P^\ell|_{K_0}$ is an affine map from K_0 onto J_0 . On the other hand, since q_0 is associated to $\alpha = \beta^k$ it follows that $f_P^i(q_0), f_P^{i+\ell}(q_0), \dots, f_P^{i+(k-1)\ell}(q_0) \in J_i$ for $i = 0, 1, \dots, \ell-1$ and, consequently, $q_0, f_P^\ell(q_0), \dots, f_P^{(k-1)\ell}(q_0) \in K_0$. Consequently, since $f_P^\ell \left(f_P^{(k-1)\ell}(q_0) \right) = f_P^m(q_0) = q_0$, it follows that $\{q_0, f_P^\ell(q_0), \dots, f_P^{(k-1)\ell}(q_0)\}$ is a periodic orbit $f_P^\ell|_{K_0}$ with period $k \geq 2$. The affinity of $f_P^\ell|_{K_0}$ implies that $f_P^\ell|_{K_0}$ is decreasing with slope -1 and $k = 2$. The fact that $f_P^\ell|_{K_0}(K_0) = J_0$ implies that $K_0 = J_0$ and the endpoints of J_0 are also a periodic orbit of $f_P^\ell|_{K_0}$ of period 2. In this situation P and $\{q_0, q_1, q_2, \dots, q_m\}$ both have the same period and pattern; a contradiction. \square

Now we want to extend the notion of associated periodic orbit and loop and Lemma 3.10 to periodic orbits of strips.

Definition 3.12. Let $F \in \mathcal{S}(\Omega)$ and let and let \mathcal{B} be a periodic orbit of strips of F . We say that a loop

$$\alpha = J_0 \xrightarrow{s_0} J_1 \xrightarrow{s_1} \cdots J_{n-1} \xrightarrow{s_{n-1}} J_0$$

in the \mathcal{B} -signed Markov graph of F and a strip A are *associated* if A is an n -periodic strip of F and $F^i(A) \in J_i$ for every $i = 0, 1, \dots, n-1$. Observe that in such case $S^m(\alpha)$ and $F^m(A)$ are associated for all $m \geq 1$. \blacksquare

The next lemma extends Lemma 1.2.7 and Corollary 1.2.8 of [1] to quasiperiodically forced skew products on the cylinder.

Lemma 3.13. *Let $F \in \mathcal{S}(\Omega)$ and let J_0, J_1, \dots, J_{n-1} be basic bands such that*

$$\alpha = J_0 \xrightarrow{s_0} J_1 \xrightarrow{s_1} \dots J_{n-1} \xrightarrow{s_{n-1}} J_0$$

is a simple loop in a signed Markov graph of F . Then there exists a periodic band $C \subset J_0$ associated to α (and hence of period n). Moreover, for every $i, j \in \{0, 1, \dots, n-1\}$, $F^i(C) < F^j(C)$ if and only if $S^i(\alpha) < S^j(\alpha)$.

Proof. Let A be a basic band and let $B_1 \leq B_2 \leq \dots \leq B_m$ be all basic bands F -covered by A . By Lemma 3.7(a.3,4) there exist bands $K(A, B_1) \leq K(A, B_2) \leq \dots \leq K(A, B_m)$ contained in A such that $K(A, B_i) \xrightarrow{s_A} B_i$ for $i = 1, 2, \dots, m$, where s_A denotes the sign of all arrows $A \xrightarrow{s_A} B_i$ (see Definition 3.4).

Now we recursively define a family of $2n$ bands in the following way. We set $K_{2n-1} := K(J_{n-1}, J_0) \subset J_{n-1}$ so that $K_{2n-1} \xrightarrow{s_{n-1}} J_0$.

Then, assume that $K_j \subset J_{j \pmod n}$ have already been defined for $j = i+1, i+2, \dots, 2n-1$ and $i \in \{0, 1, \dots, 2n-2\}$. Since $J_{\tilde{i}} \xrightarrow{s_{\tilde{i}}} J_{i+1 \pmod n}$ with $\tilde{i} = i \pmod n$, by Lemma 3.7(a.2,3), there exists a band $K_i \subset K(J_{\tilde{i}}, J_{i+1 \pmod n}) \subset J_{\tilde{i}}$ such that $K_i \xrightarrow{s_{\tilde{i}}} K_{i+1}$.

Now we claim that for every $i, j \in \{0, 1, \dots, n-1\}$ $S^i(\alpha) < S^j(\alpha)$ is equivalent to $K_i \leq K_j$. If $S^i(\alpha) \neq S^j(\alpha)$ there exists $k \in \{0, 1, \dots, n-1\}$ such that

$$\begin{aligned} S^i(\alpha) &= J_i \xrightarrow{s_i} J_{i+1} \xrightarrow{s_{i+1}} \dots J_{k+i-1} \xrightarrow{s_{k+i-1}} J_{k+i} \xrightarrow{s_{k+i}} J_{k+i+1} \dots \text{ and} \\ S^j(\alpha) &= J_i \xrightarrow{s_i} J_{i+1} \xrightarrow{s_{i+1}} \dots J_{k+i-1} \xrightarrow{s_{k+i-1}} J_{k+j} \xrightarrow{s_{k+j}} J_{k+j+1} \dots \end{aligned}$$

with $J_{k+i \pmod n} \neq J_{k+j \pmod n}$ (where every sub-index in the above paths must be read modulo n). By construction, $K_{k+i} \subset J_{k+i \pmod n}$ and $K_{k+j} \subset J_{k+j \pmod n}$. Hence, $K_{k+i} \leq K_{k+j}$ if and only if $J_{k+i \pmod n} < J_{k+j \pmod n}$. By definition

$$K_{k+i-1} \xrightarrow{s_{k+i-1} \pmod n} K_{k+i} \quad \text{and} \quad K_{k+i-1} \subset K(J_{k+i-1 \pmod n}, J_{k+i \pmod n}),$$

and

$$K_{k+j-1} \xrightarrow{s_{k+j-1} \pmod n} K_{k+j} \quad \text{and} \quad K_{k+j-1} \subset K(J_{k+i-1 \pmod n}, J_{k+j \pmod n}).$$

Thus, $K_{k+i-1} \leq K_{k+j-1}$ if and only if $K_{k+i} \leq K_{k+j}$ and $s_{k+i-1 \pmod n} = +$. So, $K_{k+i-1} \leq K_{k+j-1}$ if and only if $J_{k+i \pmod n} < J_{k+j \pmod n}$ and $s_{k+i-1 \pmod n} = +$. By iterating this argument $k-1$ times backwards we get that $K_i \leq K_j$ if and only if $J_{k+i \pmod n} < J_{k+j \pmod n}$ and

$$\text{Sign} \left(J_i \xrightarrow{s_i} J_{i+1} \xrightarrow{s_{i+1}} \dots J_{k+i-1} \xrightarrow{s_{k+i-1}} J_{k+i} \right) = s_i s_{i+1} \dots s_{k+i-1} = +$$

(where every sub-index in the above formula is modulo n). This ends the proof of the claim.

Observe that, since $K_n \subset J_0$, from the construction of the sets K_n we get that $K_{n-1} \subset K_{2n-1}, K_{n-2} \subset K_{2n-2}, \dots, K_0 \subset K_n$ and $K_0 \xrightarrow[\frac{\text{Sign}(\alpha)}{F^n}]{} K_n$. Then, by Lemma 3.7(a.2,b) there exists a band $C \subset K_0 \subset J_0$ such that $C \xrightarrow[\frac{\text{Sign}(\alpha)}{F^n}]{} C$ and $F^i(C) \subset K_i \subset J_i$ for $i = 0, 1, \dots, n-1$.

Since C is a periodic strip, $F^i(C)$ and $F^j(C)$ are either disjoint or equal. Hence, by the claim, $F^i(C) < F^j(C)$ if and only if $S^i(\alpha) < S^j(\alpha)$. Now, Lemma 3.8 tells us that $S^i(\alpha) \neq S^j(\alpha)$ whenever $i \neq j$. Consequently, $F^i(C) \neq F^j(C)$ whenever $i \neq j$ and C has period n . This ends the proof of the lemma. \square

Remark 3.14. From the construction in the above proof it follows that if $F \in \mathcal{S}(\Omega)$ and

$$\alpha = J_0 \xrightarrow{s_0} J_1 \xrightarrow{s_1} \cdots J_{n-1} \xrightarrow{s_{n-1}} J_0$$

is a loop in the a signed Markov graph of F by basic bands, then there exist bands $K_0 = K_0(\alpha) \subset J_0$, $K_1 \subset J_1$, \dots , $K_{n-1} \subset J_{n-1}$ such that $K_i \xrightarrow{s_i} K_{i+1}$ for $i = 0, 1, \dots, n-2$ and $K_{n-1} \xrightarrow{s_{n-1}} J_0$. In particular, $K_0 \xrightarrow{\frac{\text{Sign}(\alpha)}{F^n}} J_0$. Moreover, if β is another loop such that $\alpha^\infty \neq \beta^\infty$, then $K_0(\alpha)$ and $K_0(\beta)$ have pairwise disjoint interiors. \blacksquare

3.4. Proof of Theorem A. We start this subsection with a lemma that studies the periodic orbits of the uncoupled quasiperiodically forced skew-products on the cylinder (in particular for the maps F_τ).

Lemma 3.15. *Let $f \in \mathcal{C}^0(\mathbb{I}, \mathbb{I})$ and let F be a map from $\mathcal{S}(\Omega)$ such that $F(\theta, x) = (R_\omega(\theta), f(x))$. Then, the following statements hold.*

- (a) *Assume that $P = \{p_1, p_2, \dots, p_n\}$ is a periodic orbit of f with pattern τ . Then $\mathbb{S}^1 \times P$ is a periodic orbit of F with pattern τ .*
- (b) *If B is a periodic orbit of strips of F with pattern τ then there exists a periodic orbit P of f with pattern τ such that $\mathbb{S}^1 \times P$ is a periodic orbit of F with pattern τ and $\mathbb{S}^1 \times P \subset B$. In particular, every cyclic permutation is a pattern of a function of $F \in \mathcal{S}(\Omega)$.*

Proof. The first statement follows directly from the definition of a pattern. Now we prove (b). Let $B = \{B_1, B_2, \dots, B_n\}$ be periodic orbit of strips of F with pattern τ (that is, $F(B_i) = B_{\tau(i)}$ for $i = 1, 2, \dots, n$). Since $F = (R_\omega, f)$ it follows that $F^k = (R_\omega^k, f^k)$ for every $k \in \mathbb{N}$ (so the iterates of F are also uncoupled quasiperiodically forced skew-products). So, since $F^n(B_i) = B_i$ for every i , it follows that the strips B_i are horizontal. That is, for every i there exists a closed interval $J_i \subset \mathbb{I}$ such that $B_i = \mathbb{S}^1 \times J_i$. Moreover, since the strips are pairwise disjoint, so are the intervals J_i . Clearly, $f(J_i) = J_{\tau(i)}$ for every i and, hence, $f^n(J_1) = J_1$. So, there exists a point $p_1 \in J_1$ such that $f^n(p_1) = p_1$ and $f^k(p_1) \in f^k(J_1) = J_{\tau^k(1)}$ for $k \geq 0$. Since the intervals J_i are pairwise disjoint, the set $P = \{p_1, f(p_1), \dots, f^{n-1}(p_1)\}$ is a periodic orbit of f of period n such that $\mathbb{S}^1 \times P \subset B$. Moreover, if we set $f^k(p_1) = p_{\tau^k(1)}$ for $k = 1, 2, \dots, n-1$, then P has the spatial labelling and it follows that the pattern of P is τ . \square

Proof of Theorem A. First we prove that $\tau \implies_\Omega \nu$ implies $\tau \implies_{\mathbb{I}} \nu$. The assumption $\tau \implies_\Omega \nu$ implies that every map $F \in \mathcal{S}(\Omega)$ that exhibits the strips pattern τ also exhibits the strips pattern ν . In particular, the map F_τ has a periodic orbit of strips with pattern ν . By Lemma 3.15, f_τ has a periodic orbit with pattern ν . Therefore, $\tau \implies_{\mathbb{I}} \nu$ by the characterization of the forcing relation in the interval (Theorem 2.5).

Now we prove that $\tau \implies_{\mathbb{I}} \nu$ implies $\tau \implies_\Omega \nu$. Clearly, we may assume that $\nu \neq \tau$. We have to show that every $F \in \mathcal{S}(\Omega)$ that has a periodic orbit of strips $B = \{B_0, B_1, \dots, B_{n-1}\}$ with strips pattern τ also has a periodic orbit of strips with strips pattern ν .

We consider the map $f_\tau = f_P$ where P is a periodic orbit with pattern τ . By Theorem 2.5, f_τ has periodic orbit $Q = \{q_0, q_1, \dots, q_{n-1}\}$ with pattern ν . Since Q has the spatial labelling, $q_0 = \min Q$,

Since $\nu \neq \tau$, by Lemma 3.11, there exists a simple loop

$$\alpha = I_0 \xrightarrow{s_0} I_1 \xrightarrow{s_1} \cdots \xrightarrow{s_{n-2}} I_{n-1} \xrightarrow{s_{n-1}} I_0$$

in the P -signed Markov graph of f_τ associated to q_0 . Moreover, by Definition 3.9,

$$\begin{aligned} q_0 &\sim I_0 \xrightarrow{s_0} I_1 \xrightarrow{s_1} \cdots \xrightarrow{s_{n-2}} I_{n-1} \xrightarrow{s_{n-1}} I_0 \\ f_\tau(q_0) &\sim I_1 \xrightarrow{s_1} I_2 \xrightarrow{s_2} \cdots \xrightarrow{s_{n-1}} I_0 \xrightarrow{s_0} I_1 \\ f_\tau^2(q_0) &\sim I_2 \xrightarrow{s_2} I_3 \xrightarrow{s_3} \cdots \xrightarrow{s_{n-1}} I_0 \xrightarrow{s_0} I_1 \xrightarrow{s_1} I_2 \\ &\vdots \\ f_\tau^{n-1}(q_0) &\sim I_{n-1} \xrightarrow{s_{n-1}} I_0 \xrightarrow{s_0} I_1 \xrightarrow{s_1} I_2 \cdots \xrightarrow{s_{n-2}} I_{n-1}, \end{aligned}$$

where the symbol \sim means ‘‘associated with’’. By Remark 3.6 (see also Remark 3.3), the above loop α also exists in the B -signed Markov graph of F by replacing the basic intervals $I_i = [q_i, q_{i+1}]$ by the basic bands $I_{B_i B_{i+1}}$:

$$\alpha = I_{B_0 B_1} \xrightarrow{s_0} I_{B_1 B_2} \xrightarrow{s_1} \cdots \xrightarrow{s_{n-2}} I_{B_{n-2} B_{n-1}} \xrightarrow{s_{n-1}} I_{B_0 B_1}.$$

By Lemma 3.13 and Definition 3.12, F has a periodic band Q_0 associated to α (and hence of period n), and

$$\begin{aligned} Q_0 &\sim I_{B_0 B_1} \xrightarrow{s_0} I_{B_1 B_2} \xrightarrow{s_1} \cdots \xrightarrow{s_{n-2}} I_{B_{n-2} B_{n-1}} \xrightarrow{s_{n-1}} I_{B_0 B_1} \\ F(Q_0) &\sim I_{B_1 B_2} \xrightarrow{s_1} I_{B_2 B_3} \xrightarrow{s_2} \cdots \xrightarrow{s_{n-2}} I_{B_{n-2} B_{n-1}} \xrightarrow{s_{n-1}} I_{B_0 B_1} \xrightarrow{s_0} I_{B_1 B_2} \\ F^2(Q_0) &\sim I_{B_2 B_3} \xrightarrow{s_2} \cdots \xrightarrow{s_{n-2}} I_{B_{n-2} B_{n-1}} \xrightarrow{s_{n-1}} I_{B_0 B_1} \xrightarrow{s_0} I_{B_1 B_2} \xrightarrow{s_1} I_{B_2 B_3} \\ &\vdots \\ F^{n-1}(Q_0) &\sim I_{B_{n-2} B_{n-1}} \xrightarrow{s_{n-1}} I_{B_0 B_1} \xrightarrow{s_0} I_{B_1 B_2} \xrightarrow{s_1} \cdots \xrightarrow{s_{n-2}} I_{B_{n-2} B_{n-1}}. \end{aligned}$$

By Lemmas 3.10 and 3.8, the order of the points $f_\tau^i(q_0)$ induces an order on the shifts of the loop $S^i(\alpha)$, with the usual lexicographical ordering and, by Lemma 3.13, the order of the shifts $S^i(\alpha)$ induces the same order on the bands $F^i(Q_0)$. Thus, for every $i, j \in \{0, 1, \dots, n-1\}$, $i \neq j$, $F^i(Q_0) < F^j(Q_0)$ if and only if $f_\tau^i(q_0) < f_\tau^j(q_0)$. So, $\{Q_0, F(Q_0), F^2(Q_0), \dots, F^{n-1}(Q_0)\}$ and $\{q_0, q_1, \dots, q_{n-1}\}$ have the same pattern ν . This concludes the proof. \square

4. PROOF OF THEOREMS B AND C

We start by proving Theorem B.

The next technical lemma is inspired in [1, Lemma 4.3.1].

Lemma 4.1. *Let $F \in \mathcal{S}(\Omega)$ and let (J, \mathcal{D}) be an s -horseshoe of F . Then, there exists \mathcal{D}_n , a set of s^n pairwise weakly ordered bands contained in J , each of them with non-empty interior, such that (J, \mathcal{D}_n) is a s^n -horseshoe for F^n .*

Proof. We use induction. For $n = 1$ there is nothing to prove.

Suppose that the induction hypothesis holds for some n and let $D \in \mathcal{D}$ and $C \in \mathcal{D}_n$. Since $C \subset J$ has non-empty interior and $D \xrightarrow{\pm} J$, by Lemma 3.7(a.2,3), there exists a band $B(D, C) \subset D$ with non-empty interior such that $B(D, C) \xrightarrow{\pm} C$. Moreover, given $C' \in \mathcal{D}_n$ with $C' \neq C$, $B(D, C)$ and $B(D, C')$ can be chosen to be weakly ordered because C and C' are weakly ordered by assumption. Since, $C \in \mathcal{D}_n$, $B(D, C) \xrightarrow[F^{n+1}]{\pm} J$. Thus, the family

$$\mathcal{D}_{n+1} = \{B(D, C) : D \in \mathcal{D} \text{ and } C \in \mathcal{D}_n\}$$

consists of s^{n+1} pairwise weakly ordered bands contained in J , each of them with non-empty interior, such that $B(D, C)$ F^{n+1} -covers J . Consequently, (J, \mathcal{D}_{n+1}) is an s^{n+1} -horseshoe for F^{n+1} . \square

Proof of Theorem B. Fix $n > 0$. By Lemma 4.1, F^n has a s^n -horseshoe (J, \mathcal{D}) . Remove the smallest and the biggest band of \mathcal{D} and call K the smallest band that contains the remaining elements of \mathcal{D} . Clearly, K is contained in the interior of J . Thus, by Lemma 3.7(a.2,3), each element D of \mathcal{D} contains in its interior a band $A(D)$ such that $A(D) \xrightarrow[F^n]{\pm} K$. Then there exists an open cover \mathcal{B} of the strip J (formed by open sets B such B^θ is an open interval for every $\theta \in \mathbb{S}^1$), such that for each $D \in \mathcal{D}|_K$, the band $A(D)$ intersects only one element $B(D)$ of \mathcal{B} (then it has to be contained in it) and if $D, D' \in \mathcal{D}|_K$ with $D \neq D'$ then $B(D) \neq B(D')$. For $D_0, D_1, \dots, D_{k-1} \in \mathcal{D}|_K$ the set $\cap_{i=0}^{k-1} F^{-n}(A(D_i))$ is non-empty and intersects only one element of $\mathcal{B}_{F^n}^k$, namely $\cap_{i=0}^{k-1} F^{-n}(B(D_i))$. Therefore the sets $\cap_{i=0}^{k-1} F^{-n}(A(D_i))$ are different for different sequences $(D_0, D_1, \dots, D_{k-1})$, and thus

$$\mathcal{N}(\mathcal{B}_{F^n}^k) \geq (\text{Card } \mathcal{D} - 2)^k,$$

where $\mathcal{N}(\mathcal{B}_{F^n}^k)$ is defined as in [1, Section 4.1]. Hence,

$$h(F) = \frac{1}{n} h(F^n) \geq \frac{1}{n} h(F^n, \mathcal{B}) \geq \frac{1}{n} \log(\text{Card}(\mathcal{D}) - 2) = \frac{1}{n} \log(s^n - 2).$$

Since n is arbitrary, we obtain $h(F) \geq \log(s)$. \square

Now we aim at proving Theorem C. To this end we have to introduce some more notation and preliminary results concerning the *topological entropy*.

Given a map $f \in \mathcal{S}(\Omega)$, $h(F|_{I_\theta})$ is defined for every $I_\theta := \{\theta\} \times \mathbb{I}$ (despite of the fact that it is not F -invariant) by using the Bowen definition of the topological entropy (c.f. [5, 4]). Moreover, the Bowen Formula gives

$$\max\{h(R), h_{\text{fib}}(F)\} \leq h(F) \leq h(R) + h_{\text{fib}}(F)$$

where

$$h_{\text{fib}}(F) = \sup_{\theta \in \mathbb{S}^1} h(F|_{I_\theta}).$$

Since $h(R) = 0$, it follows that $h(F) = h_{\text{fib}}(F)$.

In the particular case of the uncoupled maps $F_\tau = (R, f_\tau)$ we easily get the following result:

Lemma 4.2. *Let τ be a pattern (both in \mathbb{I} and Ω). Then $h(F_\tau|_{I_\theta}) = h(f_\tau)$ for every $\theta \in \mathbb{S}^1$. Consequently,*

$$h(F_\tau) = h_{\text{fib}}(F_\tau) = h(f_\tau).$$

Given a signed Markov graph G with vertices I_1, I_2, \dots, I_n we associate to it a $n \times n$ transition matrix $T_G = (t_{ij})$ by setting $t_{ij} = 1$ if and only if there is a signed arrow from the vertex I_i to the vertex I_j in G . Otherwise, t_{ij} is set to 0.

The spectral radius of a matrix T , denoted by $\rho(T)$, is equal to the maximum of the absolute values of the eigenvalues of T .

Lemma 4.3. *Let P be a periodic orbit of strips of $F \in \mathcal{S}(\Omega)$ and let T be the transition matrix of the P -signed Markov graph of F . Then*

$$h(F) \geq \max\{0, \log(\rho(T))\}.$$

Proof. If $\rho(T) \leq 1$ then there is nothing to prove. So, we assume that $\rho(T) > 1$. Let J be the i -th P -basic band and let s be the i -th entry of the diagonal of T^n . By [1, Lemma 4.4.1] there are s loops of length n in the P -signed Markov graph

of F beginning and ending at J . Hence, if $s \geq 2$, F^n has an s -horseshoe (J, \mathcal{D}) by Remark 3.14. By Theorem B, $h(F) = \frac{1}{n}h(F^n) \geq \frac{1}{n}\log(s)$.

If there are k basic bands, the trace of T^n is not larger than k times the maximal entry on the diagonal of T^n . Hence, $h(F) \geq \frac{1}{n}\log\left(\frac{1}{k}\text{tr}(T^n)\right)$. Therefore, by [1, Lemma 4.4.2],

$$h(F) \geq \limsup_{n \rightarrow \infty} \frac{1}{n} \log\left(\frac{1}{k} \text{tr}(T^n)\right) = \limsup_{n \rightarrow \infty} \frac{1}{n} \log(\text{tr}(T^n)) = \log(\rho(T)).$$

□

Proof of Theorem C. Let P be a periodic orbit of strips with pattern τ and let T be the transition matrix of the P -signed Markov graph of F . Let $f_\tau = f_Q$ be a Q -linear map in $\mathcal{C}^0(\mathbb{I}, \mathbb{I})$, where Q is a periodic orbit of f_Q with pattern τ . In view of Remark 3.6 (see also Remark 3.3), T is also the transition matrix of the Q -signed Markov graph of f_τ . Consequently, by [1, Theorem 4.4.5], $h(f_\tau) = \max\{0, \log(\rho(T))\}$. By Lemmas 4.3 and 4.2,

$$h(F) \geq \max\{0, \log(\rho(T))\} = h(f_\tau) = h(F_\tau).$$

□

REFERENCES

- [1] Lluís Alsedà, Jaume Llibre, and Michał Misiurewicz. *Combinatorial dynamics and entropy in dimension one*, volume 5 of *Advanced Series in Nonlinear Dynamics*. World Scientific Publishing Co. Inc., River Edge, NJ, second edition, 2000.
- [2] Lluís Alsedà, Francesc Mañosas, and Leopoldo Morales. A skew-product map on the cylinder without fixed-curves. preprint, 2014.
- [3] Louis Block, John Guckenheimer, Michał Misiurewicz, and Lai Sang Young. Periodic points and topological entropy of one-dimensional maps. In *Global theory of dynamical systems (Proc. Internat. Conf., Northwestern Univ., Evanston, Ill., 1979)*, volume 819 of *Lecture Notes in Math.*, pages 18–34. Springer, Berlin, 1980.
- [4] R. Bowen. Erratum to “Entropy for group endomorphisms and homogeneous spaces”. *Trans. Amer. Math. Soc.*, 181:509–510, 1973.
- [5] Rufus Bowen. Entropy for group endomorphisms and homogeneous spaces. *Trans. Amer. Math. Soc.*, 153:401–414, 1971.
- [6] Gustave Choquet. *Topology*. Translated from the French by Amiel Feinstein. Pure and Applied Mathematics, Vol. XIX. Academic Press, New York-London, 1966.
- [7] Roberta Fabbri, Tobias Jäger, Russel Johnson, and Gerhard Keller. A Sharkovskii-type theorem for minimally forced interval maps. *Topol. Methods Nonlinear Anal.*, 26(1):163–188, 2005.
- [8] Sam B. Nadler, Jr. *Continuum theory*, volume 158 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, 1992. An introduction.
- [9] O. M. Šarkovskii. Co-existence of cycles of a continuous mapping of the line into itself. *Ukrain. Mat. Ž.*, 16:61–71, 1964.
- [10] A. N. Sharkovskii. Coexistence of cycles of a continuous map of the line into itself. In *Thirty years after Sharkovskii’s theorem: new perspectives (Murcia, 1994)*, volume 8 of *World Sci. Ser. Nonlinear Sci. Ser. B Spec. Theme Issues Proc.*, pages 1–11. World Sci. Publ., River Edge, NJ, 1995. Translated by J. Tolosa, Reprint of the paper reviewed in MR1361914 (96j:58058).

DEPARTAMENT DE MATEMÀTIQUES, EDIFICI CC, UNIVERSITAT AUTÒNOMA DE BARCELONA,
08913 Cerdanyola del Vallès, BARCELONA, SPAIN
E-mail address: alseda@mat.uab.cat

DEPARTAMENT DE MATEMÀTIQUES, EDIFICI CC, UNIVERSITAT AUTÒNOMA DE BARCELONA,
08913 Cerdanyola del Vallès, BARCELONA, SPAIN
E-mail address: manyosas@mat.uab.cat

DEPARTAMENT DE MATEMÀTIQUES, EDIFICI CC, UNIVERSITAT AUTÒNOMA DE BARCELONA,
08913 Cerdanyola del Vallès, BARCELONA, SPAIN
E-mail address: mleo@mat.uab.cat