

This is the submitted version of the article: Cerdà, AP. et al. *Long term enhanced solid-state fermentation: inoculation strategies for amylase production from soy and bread wastes by Thermomyces sp. In a sequential batch operation* in Journal of environmental chemical engineering (Ed. Elsevier), vol. 4, issue 2 (June 2016), p. 2394-2401.

DOI 10.1016/j.jece,2016.04.022

© 2018. This manuscript version is made available under the Creative Commons CC BY NC ND license



1 **Abstract**

2 The effect of two thermophilic amylase producing strains was evaluated using different  
3 mixtures of soy and bread wastes. *Thermomyces* sp. was found to be better inoculum than  
4 *Geobacillus* sp. for a soy and bread waste mixture (90:10 w/w respectively) producing a  
5 maximum enzyme activity of  $39.9 \cdot 10^3$  U g<sup>-1</sup> dry substrate. Three strategies (a, b, c) were  
6 evaluated for solid-state fermentation (SSF) operation in sequential batches. Fermented  
7 solids from each batch were used to inoculate the following batch: a) solids at the moment of  
8 maximum biological activity; b) final solids (end of the process, maximum amylase  
9 production); c) final solids after enzymatic extraction. The evaluated strategies led to an  
10 increase in amylase production of 50, 500 and 98% for each strategy, respectively. This  
11 indicates the suitability of fermented solids to act as inoculum and the enhancement of  
12 amylase production compared to traditional batches. As one of the main challenges of SSF is  
13 the maintenance of a productive process along time, these results confirm SSF as an  
14 excellent option to produce amylases from organic wastes.

15

16 **Keywords:** solid-state fermentation (SSF); *Thermomyces*; inoculation strategies; amylase;  
17 sequential batch; waste valorisation.

18

19

## 1 **1. Introduction**

2 In the last decade solid-state fermentation (SSF) has received increased attention from  
3 researchers as a bio-based process. This technology allows for the valorisation of wastes for  
4 the production of valuable commodities such as enzymes [1-3].

5 SSF presents some advantages in comparison with conventional fermentations such as higher  
6 productivity or reduced energy requirements, and low wastewater output [3] However,  
7 important drawbacks are described concerning the SSF scale up largely due to heat transfer,  
8 sterilization costs and culture heterogeneity [4,5]. Abraham et al. [6] proposed to perform  
9 SSF as a thermophilic batch process under near-adiabatic conditions, similar to the  
10 composting process which is a robust, well known, easily scalable and low-cost process. This  
11 self-heating process does not involve heating/cooling costs since the temperature evolve due  
12 to the heat released in the biodegradation process. Based on this proposal, El-Bakry et al. [7]  
13 proved that thermophilic strains inoculated on non-sterile wastes were able to grow and  
14 compete with autochthonous microbiota and significantly increase the protease production.  
15 Still the setting up of a long term operation in continuous or semi-continuous regime remains  
16 a challenge and very few attempts have been reported [8-10].

17 Enzymatic products such as amylases have great potential in different biotechnological  
18 applications such as food, fermentation, detergent, textile and paper industries. Some  
19 thermophilic amylase-producers microorganisms have been found during the highest bio-  
20 oxidative activity stage during composting processes [11-13]. In addition, it has been well  
21 established that these microorganisms are able to produce substantial amounts of amylase in  
22 SSF systems [2,14-16].

23 Several attempts for the optimization of enzymatic production have been reported, such as  
24 nutritional supplementation for microorganism growth [16-18], use of engineered strains  
25 [16,19] or operational strategies such as batch or fed-batch [9,20,21]. However, most SSF for

1 the production of enzymatic compounds have been tested at lab scale and under sterile  
2 substrate conditions [10].

3 The first aim of this work was to develop a SSF process for the production of amylases using  
4 a thermophilic strain growing on non-sterile wastes in the lab scale (500-mL reactors) and  
5 bring it to the bench scale (10-L reactors), based on the proposal of Abraham et al. [6] and  
6 El-Bakry et al. [7]. Second goal was to perform a first approach to the assessment of the  
7 effect of inoculation strategies in sequential batch operation in order to develop a long term  
8 operation process in a solid-state fermentation configuration at the bench scale. SSF of  
9 organic wastes operated in sequential batches was evaluated for the production of amylases  
10 using thermophilic and amylase producer microorganisms (*Thermomyces* sp. and  
11 *Geobacillus* sp.).

12

13 **2. Materials and methods**

14

15 *2.1 Microorganisms*

16

17 Thermophilic and amylase producers *Geobacillus* sp. ATCC-31198 and *Thermomyces*  
18 sp. ATCC-200065 were incubated in Petri dishes on supplier's recommended agar media:  
19 ATCC® Media 1207 and ATCC® Media 350 respectively.

20 Both strains were grown in a liquid medium containing the same formulation than that used  
21 for Petri dishes. Final pH was adjusted to  $7.0 \pm 0.2$  and 100 mL were incubated in a 500 mL  
22 Erlenmeyer flask at 150 rpm and 55°C for 16h. Final culture broth was used as inoculum in  
23 all lab scale fermentations and in the propagation reactor for sequential batch experiments in  
24 a 10% w/w ratio.

25

## 1 2.2 Wastes

2

3 Soy fibre residues were provided by Soy Nature (Barcelona, Spain). Bread waste (expired  
4 pre-cooked baguettes) was procured from a local market. Both wastes were obtained in a  
5 sufficient amount to do the entire experiments with the same material. Materials were used  
6 as received in 500 mL reactors and in the scale-up experiments. In the case of the inoculation  
7 strategies experiments, aliquots were frozen separately prior to their use. Characterization of  
8 both substrates is presented in Table 1. Moisture was initially adjusted to 70% using tap  
9 water in all the experiments.

10

## 11 2.3 Selection of inoculum and substrates ratio at lab scale

12

13 Solid state fermentations (lab scale) were carried out at 55°C for 7 days in 500 mL  
14 Erlenmeyer flasks with 100 g of wet substrate and at fixed airflow of 20 mL min<sup>-1</sup> by means  
15 of a mass flow meter. Air was humidified prior entering the reactors. 10 g of wood sticks  
16 were added to the substrate to provide porosity. Oxygen content was measured continuously  
17 in the reactors gas output. Specific oxygen uptake rate (sOUR) was calculated with equation  
18 1.

19

$$20 \quad sOUR = F \cdot (0.209 - y_{O_2}) \cdot \frac{P \cdot 32 \cdot 60 \cdot 10^3}{R \cdot T \cdot DW \cdot 10^3} \quad \text{Equation (1)}$$

21

22 where: sOUR is the Oxygen Uptake Rate (mg O<sub>2</sub> g<sup>-1</sup> DM h<sup>-1</sup>); *F*, airflow into the reactor (mL  
23 min<sup>-1</sup>); *y*<sub>O<sub>2</sub></sub>, is the oxygen molar fraction in the exhaust gases (mol O<sub>2</sub> mol<sup>-1</sup>); *P*, pressure of  
24 the system assumed constant at 101325 Pa; 32, oxygen molecular weight (g O<sub>2</sub> mol O<sub>2</sub><sup>-1</sup>); 60,  
25 conversion factor from minute to hour; 10<sup>3</sup>, conversion factor from mL to L; *R*, ideal gas

1 constant ( $8310 \text{ Pa L K}^{-1} \text{ mol}^{-1}$ );  $T$ , temperature at which  $F$  is measured (K);  $DW$ , initial dry  
2 weight of solids in the reactor (g);  $10^3$ , conversion factor from g to mg.

3 Samples were taken daily for amylase activity determination. Firstly, *Geobacillus* sp. and  
4 *Thermomyces* sp. were evaluated in the solid-state fermentation of soy and bread wastes for  
5 the production of amylases and the best inoculum was selected for further experiments.  
6 Secondly, mixtures of different soy:bread ratios (0:100, 10:90, 50:50, 90:10 and 100:0  
7 (w/w)) were evaluated and the best mixture selected for following experiments in reactors.  
8 Experiments were carried out in triplicates.

9

#### 10 *2.4 Bench scale reactors operation*

11

12 Experiments were performed in 4.5 L cylindrical reactors previously described by Abraham  
13 et al. [6] with 1.2 kg of wet substrate. Reactors were thermally isolated to operate under near-  
14 adiabatic conditions and with on-line temperature and oxygen monitoring. Continuous  
15 aeration was provided to the reactor by means of a mass flow meter, starting with a rate of  
16  $100 \text{ mL min}^{-1}$  which was manually increased (maximum air flow  $1000 \text{ mL min}^{-1}$ ) according  
17 to the oxygen content in the reactor ensuring aerobic conditions. The oxygen concentration  
18 in the exhaust gases was measured by means of an oxygen sensor (Crowcon's Xgard, United  
19 Kingdom). sOUR is the specific oxygen uptake rate and was calculated from airflow and  
20 oxygen content and expressed as mg of oxygen consumed per gram of initial dry mass in the  
21 reactor and per hour (Equation 1). Total oxygen consumed was calculated as the area below  
22 the sOUR curve by the trapezoidal rule and expressed as mg of oxygen consumed per gram  
23 of initial dry mass. Temperature was monitored by means of a Pt-100 probe located at the  
24 central point of the reactor.

1 All fermentations were carried out by using the inoculum and soy:bread mixture previously  
2 selected at the lab scale. Wood chips were added as bulking agent in 1:1 (v/v) ratio, in order  
3 to provide porosity to the mixture and improve oxygen transfer in the reactors.

4

#### 5 *2.4.1 Scaling-up of amylase production*

6 The scaling-up effect on the process defined at lab scale was assessed by monitoring two 4.5  
7 L reactors for 14 days in order to have a complete profile of all relevant parameters of  
8 fermentation. One reactor was operating without inoculation as control. A second reactor  
9 was inoculated directly with liquid *Thermomyces* sp. culture. Amylase activity was  
10 determined at days 0, 4, 7, 10 and 14 of operation.

11

#### 12 *2.4.2 Inoculation strategies for operation in sequential batches*

13 One first batch was performed as a propagation reactor and it was inoculated directly with  
14 *Thermomyces* sp. liquid culture in a 10% w/w ratio according to El-Bakry et al. [7]. Samples  
15 were taken from this reactor in two different moments to act as inoculum for the following  
16 batches. Three sequential batches (SB) were performed and each batch was inoculated with  
17 solids from the previous batch. Inoculation size was 10% w/w (fermented solids / fresh  
18 solids in the reactor) in all cases. The three strategies assayed are described below:

19 a) Strategy MOUR: Using as inoculum the fermented solids obtained at the moment of  
20 maximum sOUR, that is, solids with *Thermomyces* sp. at maximum growth rate.

21 b) Strategy MAA: Using as inoculum the fermented solids obtained at the end of the process,  
22 that is, solids obtained at the moment of maximum amylase activity.

23 c) Strategy MAAE: Using as inoculum the solids obtained after amylase recovery when its  
24 maximum activity was reached at the end of the process.

25



1 *2.5 Enzyme extraction and amylase activity assay*

2

3 Fermented solids were mixed thoroughly on a magnetic stirrer during 30 min with 0.1M  
4 phosphate buffer, at a pH of 6.8 and ratio of 1:5 (w:v) respectively. The extract was  
5 separated by centrifugation at 10000 rpm during 10 min followed by a filtration with a  
6 0.45 µm filter. The supernatant was used for the amylase activity determination assay.

7 The reaction mixture consisted of 0.2 mL enzyme extract, 0.25 mL of 0.1% soluble starch  
8 solution and 0.5 mL of phosphate buffer (0.1M, pH 6.8). The reaction was stopped by adding  
9 0.25 mL of 0.1N HCl, followed by the addition of 0.25 mL of I/KI solution (2% KI in 0.2%  
10 I) [22]. The colour intensity developed was measured in a spectrophotometer at 690 nm. One  
11 unit of amylase activity (U) is defined as the amount of enzyme able to generate a reduction  
12 of 0.01% in the blue colour intensity of starch in iodine solution per minute. In this work,  
13 amylase activity is reported as U per gram of dry substrate, considering soy and/or bread  
14 wastes as substrate, without considering the wood sticks or the wood chips. All  
15 measurements of amylase activity were performed in triplicates.

16

17 *2.6 Sampling and analytical methods*

18

19 Sampling was performed on a composite sample of the reactor, after homogenizing all the  
20 reactor material.

21 Moisture content, total and volatile solids, pH and EC were determined according to the  
22 standard procedures of The US Department of Agriculture and The US Composting Council  
23 [23]. Soluble starch content was determined by a commercial assay kit (catalogue item  
24 STA20) purchased from Sigma-Aldrich (St. Louis, USA).

25

### 1 3. Results and Discussion

2

#### 3 3.1 Selection of the inoculum and mixtures at lab scale

4

5 Fig. 1 shows sOUR and amylase activity profiles for experiments in 500 mL reactors for  
6 inoculum selection using soy fibre (Fig. 1a, 1b) and bread waste (Fig. 1c, 1d) as substrate. As  
7 observed in Fig. 1a and 1c, higher sOUR values were obtained in reactors inoculated with  
8 *Thermomyces* sp. and *Geobacillus* sp. strains than in non-inoculated reactors, when using soy  
9 fibre and bread waste as sole substrate. This increase was more obvious in the case of soy  
10 fibre, which presented a higher sOUR than bread waste confirming a higher biodegradability.  
11 sOUR values were maximum at around 24h in all the cases and slowly decreased after that.  
12 Similar results were obtained for amylase activity: inoculation showed a positive effect on  
13 amylase production for both substrates, and higher enzymatic yields were obtained for soy  
14 fibre than for bread wastes. Maximum activity was obtained 48 h after reaching the  
15 maximum value of sOUR and slightly decreased after that. For both substrates, reactors  
16 inoculated with *Thermomyces* sp. presented higher values of sOUR and amylase activity than  
17 reactors with *Geobacillus* sp. Best process performance was obtained using soy fibre and  
18 *Thermomyces* sp. with a maximum sOUR of  $11.8 \text{ mgO}_2 \text{ g}^{-1}\text{DM h}^{-1}$  and a maximum amylase  
19 production of  $34 \cdot 10^3 \text{ U g}^{-1}\text{DS}$ . Fungi were expected to be better inoculum due to their ability  
20 to produce all required enzymes to fully degrade polymeric compounds and also to be able to  
21 survive in extreme conditions [24]. Also, fungi play a key role in solid fermentations such as  
22 composting processes, where *Thermomyces* sp. has been detected during thermophilic and  
23 highest amylolytic activity production stage [12,25]. These authors reported the  
24 predominance of *Thermomyces* sp. in temperatures ranging 50-60°C and alkaline pH, which  
25 is also the range of its optimum growth regarding temperature and pH.

1 In consequence, *Thermomyces* sp. was selected as the best strain. Published literature [26]  
2 reports the synergistic effect of different wastes used as substrates for amylase production by  
3 SSF. Accordingly, further experiments were performed with different mixtures of soy and  
4 bread wastes to assess this potential synergistic effect. Table 2 summarizes maximum sOUR  
5 and amylase activity for all evaluated mixtures using different ratios of soy:bread. It was  
6 found that sOUR and amylase production were improved at high soy fibre content due to its  
7 higher content and availability of easily biodegradable organic matter. Also the mineral  
8 content of soy fibre is higher than the bread waste content, especially in Mg, P, and K which  
9 are required for the growth of amylase producing microorganisms [27,28]. Starch contained  
10 on bread most probably acted as inducer for amylase production as reported by some  
11 researchers [18,21] and nutrients present on soy provided proper conditions for the  
12 proliferation of microorganisms. It has been reported that several cations present in great  
13 amounts on soy fibre such as  $\text{Ca}^{2+}$ , acted as stabilizing agent or co-factors for amylases  
14 [16,18], generating an amylase activity increase, which is in accordance with obtained  
15 results.

16 All the mixtures showed a maximum amylase production on day 4 of process. Maximum  
17 amylase activity was obtained at a 90:10 soy:bread ratio, with a value over  $39.9 \cdot 10^3 \text{ U g}^{-1}\text{DS}$   
18 and maximum sOUR of  $11.5 \text{ mgO}_2 \text{ g}^{-1}\text{DM h}^{-1}$ . Lowest amylase production was obtained  
19 with bread waste as sole substrate, with a maximum activity of  $10.9 \cdot 10^3 \text{ U g}^{-1}\text{DS}$ . As it is  
20 possible to observe in all enzymatic profiles, amylase production is not directly related to  
21 maximum biological activity or growth. This fact was expected as amylase production is  
22 regulated by starch inducible system and repressed in presence of glucose or mannose  
23 isomers [29]. Maximum amylase was observed 48h after maximum sOUR as also reported  
24 by Kunamneni et al. [15] and Petrova et al. [16].

25

### 1 3.2 Scale up effect on amylase production

2

3 The selected process (substrates mixture 90:10 w/w and *Thermomyces* sp) was evaluated in  
4 4.5 L reactors and results are presented in Fig. 2. Maximum sOUR and temperature were  
5 obtained at day 2 of operation in both control and inoculated reactors. As observed at lab  
6 scale, maximum biological activity did not match with amylase maximum production, which  
7 was obtained 48h after the maximum sOUR (day 4) in both cases. There was a clear effect of  
8 inoculation in terms of biological activity and amylase activity production [7]. *Thermomyces*  
9 sp. inoculated reactor presented a temperature of 65°C and a maximum sOUR of 11.1 mgO<sub>2</sub>  
10 g<sup>-1</sup>DM h<sup>-1</sup> which represented an 8 and 47% increase respectively in comparison with non-  
11 inoculated reactors. Also, using *Thermomyces* sp. as inoculum presented a positive effect on  
12 amylase activity with a 36% increment in comparison with control reactor, reaching a  
13 maximum of 41.03·10<sup>3</sup> U g<sup>-1</sup>DS.

14 Similar results were obtained at 4.5 L and 500 mL scales for maximum sOUR and amylase  
15 activity. This confirms the easy scalability of the self-heating SSF and its suitability to work  
16 with thermophilic strains with non-sterile substrates.

17

### 18 3.3 Inoculation strategies on amylase production by sequential batches

19

20 Process profile for the propagation reactor is presented in Fig. 3a, 4a and 5a. A lag phase of  
21 nearly 2 days was observed. This was due to the use of previously frozen substrates. As  
22 reported by Pognani et al. [30], freezing for less than one year does not affect the  
23 biodegradable organic matter content or the biological activity measured as sOUR, but a lag  
24 phase may appear. Thermophilic conditions were reached after that and remained nearly for  
25 48h, achieving maximum sOUR of 23 mgO<sub>2</sub> g<sup>-1</sup>DM h<sup>-1</sup> after sampling the reactor. Sampling

1 sOUR was  $17 \text{ mgO}_2 \text{ g}^{-1}\text{DM h}^{-1}$  and obtained amylase activity of  $31.7 \cdot 10^3 \text{ U g}^{-1}\text{DS}$ . Total  
2 oxygen consumed was  $725 \text{ mgO}_2 \text{ g}^{-1}\text{DM}$  in 5 days. At the end of the fermentation (48h after  
3 maximum sOUR) amylase activity was 15% higher with a value of  $36.6 \cdot 10^3 \text{ U g}^{-1}\text{DS}$ .

4

### 5 *3.3.1 Strategy MOUR*

6 Fig. 3 presents the process profiles for the propagation reactor plus the three sequential  
7 batches (SB1, SB2 and SB3) performed with inoculation using solids obtained at the moment  
8 of maximum sOUR (gap in temperature curve of SB2 was due to data acquisition failure). In  
9 all four cases, maximum sOUR was achieved after sampling, which could be of importance  
10 considering the quality of the inoculums for the sequential batches [7].

11 The three sequential batches presented maximum temperatures ranging between  $53\text{-}61^\circ\text{C}$   
12 similar to the propagation reactor. As for sOUR profile, maximum values were obtained also  
13 after sampling, reaching values of  $21, 23, 19$  and  $19 \text{ mgO}_2 \text{ g}^{-1}\text{DM h}^{-1}$  for the propagation  
14 reactor and the SB 1, 2 and 3 respectively (Fig. 3).

15 Amylase activity obtained during maximum sOUR were  $31.7, 21.9, 7.1$  and  $43.0 \cdot 10^3 \text{ U g}^{-1}$   
16  $\text{DS}$  and during maximum amylase activity  $36.6, 39.1, 47.2$  and  $29.3 \cdot 10^3 \text{ U g}^{-1}\text{DS}$  for the  
17 propagation reactor and the SB 1, 2 and 3 respectively.

18 In general terms, sOUR increase and lag phase reduction was only obtained during SB1.  
19 Latter inoculations resulted in lower or equal sOUR, obtained only after homogenization of  
20 reactors in the sampling process. In addition, each sequential inoculation produced an  
21 increase of 0.8, 0.9 and 15.1% in total oxygen consumption in comparison with propagation  
22 reactor.

23 Inoculation with solids from maximum sOUR period from the propagation reactor was  
24 expected to provide a reduction in lag phase and an increase in sOUR. This is based on the  
25 fact that fungi from the class Eurotiomycetes, such as *Thermomyces* sp. have been found

1 predominant in thermophilic stage during composting processes [11,12,31], indicating that a  
2 suitable environment for inoculated *Thermomyces* sp. is provided. This fact was observed  
3 through all sequential batches, where a slight increase in sOUR and total oxygen  
4 consumption was obtained.

5

### 6 3.3.2 Strategies MAA and MAAE

7 Fig. 4 presents process profiles for the propagation reactor and the SB 1, 2 and 3 for MAA  
8 strategy. Temperature profile was similar in all four processes, reaching values near 60°C.  
9 Maximum sOUR values were in the range of 21-25 mgO<sub>2</sub> g<sup>-1</sup>DM h<sup>-1</sup> except for the first  
10 sequential batch (Fig. 4b). However, total oxygen consumption presented a sustained  
11 increase of 1.9, 54.2 and 76.6% in the SB 1, 2 and 3 respectively in comparison with the  
12 propagation reactor. Also lag phase was gradually reduced in each SB, and sOUR peaked in  
13 the mesophilic phase in SB 2 and 3 with values 8-9 mgO<sub>2</sub> g<sup>-1</sup>DM h<sup>-1</sup>.

14 During thermophilic period (temperature >45°C), high amylase activity was produced in all  
15 batches. Highest amylase production of 228.9·10<sup>3</sup> U g<sup>-1</sup>DS was obtained at the end of the  
16 MAA-SB2, which represents a 500% increase in comparison with the propagation reactor. In  
17 the MAA-SB3, amylase activity was 156.9·10<sup>3</sup> U g<sup>-1</sup>DS, 329% higher than the propagation  
18 reactor.

19 Fig. 5 shows the process profile for the propagation reactor and the three sequential batches,  
20 using final solids after the extraction of soluble and enzymatic components as inoculums.  
21 This solid material would be the actual final output of the process after amylase recovery as  
22 targeted product.

23 Maximum sOUR values in the three sequential batches were lower than in the propagation  
24 reactor. A 16.6 % decrease in the total oxygen consumption on the first sequential batch SB1  
25 was obtained but increased by 1.1 and 23.2 % in SB2 and SB3. Thermophilic phase was

1 longer in the three SB than in the propagation reactor. Also amylase activity was higher in  
2 the sequential batches than in propagation reactor, with an increase of 27, 98, 56% in SB 1, 2  
3 and 3 respectively. Maximum activity of the sequential experiments was obtained at the end  
4 of batch 2, achieving a value of  $72.4 \cdot 10^3 \text{ U g}^{-1}\text{DS}$ .

5 Regarding MAA and MAAE strategies, temperature profiles and total oxygen consumption  
6 presented a different trend than the propagation reactor. Temperature is known as a crucial  
7 parameter to describe microbial activity during aerobic fermentation of solid heterogeneous  
8 material. It has been stated that inoculation with different microorganisms to a solid state  
9 fermentation could lead to a rapid rate of temperature elevation and for a more prolonged  
10 time of high-temperature process [32].

11 In MAA strategy, inoculation with fermented solid inoculum with high enzymatic and  
12 soluble content in sequential batches may have generated a fast hydrolysis of available  
13 substrate for the consumption of all present microorganisms. On the other hand, an  
14 extraction of soluble and enzymatic compounds in the inoculums could have generated a  
15 reduction in sOUR values and in the oxygen consumption rate, due to a lack of hydrolytic  
16 capacity in comparison with propagation reactor and the other evaluated batches with  
17 different inoculation strategies.

18 A reduction in amylase activity in the SB3 in both MAA and MAAE strategies can be  
19 attributed to several factors such as the possible repression on enzymatic synthesis [14,18],  
20 presence of hydrolytic enzymes such as proteases [6], to a depletion of the amorphous form  
21 of the solid substrate [21] or even more, to a change in the production profile of amylases.

22 It is a fact that sugars and low polymerization degree polysaccharides are normally fast  
23 consumed by microflora present during fermentation. It is reasonable to assume that with  
24 high production of amylase activity, high amounts of sugars are released which could cause  
25 catabolic repression of amylase synthesis. On the other hand, researchers have presented that

1 catabolic repression at high reducing sugars content occurs even in metabolically engineered  
2 de-repressed strains [19].

3 The lack of literature about amylase production by inoculation strategies in sequential solid  
4 state fermentations and the many different methods of amylase activity determination made  
5 difficult to compare the obtained results with other reported researches. However, other  
6 enzymes have been produced in fed batch strategy [9], with results found in accordance to  
7 this study. In any case it is proved that the operation strategy is a key factor for the  
8 implementation of solid state fermentation at the commercial level. Further research shall  
9 focus on determining the number of sequential batches that can be performed following this  
10 strategy. Nevertheless, both MAA and MAAE strategies allowed a remarkable enhancement  
11 on amylase activity. In our opinion, both strategies are suitable for industrial operation: in  
12 strategy MAA, 10% of the solids would be diverted to inoculate the following batch prior to  
13 enzyme extraction; in strategy MAAE, 10% of the solids after extraction would be recycled  
14 to inoculate the next batch. To finally decide which the best strategy is, it would be necessary  
15 to perform a complete economic and environmental assessment.

16

#### 17 **4. Conclusions**

18

19 Operation of SSF of soy and bread wastes in sequential batches has been proven as a suitable  
20 strategy for amylase production with *Thermomyces* sp. Additionally, it opens a new  
21 valorisation alternative for these wastes. A SSF process was operated for 19 days and the  
22 enhancement of amylase production was accomplished. In terms of amylase yield, the most  
23 suitable inoculation strategy was using fermented solids at the end of the batch to inoculate  
24 the following batch, obtaining a 500% increase in productivity and eliminating the need of  
25 producing fresh inoculum for each batch. Nonetheless this, the final strategy selection should



1 consider economic and environmental aspects. The development of these operational  
2 strategies could benefit process economics and provides the possibility of a continuous  
3 operation for amylases and potentially for other enzymes.

4

#### 5 **Acknowledgments**

6 This research has been funded by the Spanish Ministerio de Economía y Competitividad  
7 (project CTM2015-69513-R). Alejandra Cerda thanks to Universitat Autònoma de Barcelona  
8 for her pre-doctoral grant. Mamdouh El-Bakry was recipient of a postdoctoral fellowship  
9 from Universitat Autònoma de Barcelona.

10

## 1 **References**

- 2 [1] El-Bakry, M., Abraham, J., Cerda, A., Barrena, R., Ponsá, S., Gea, T., Sánchez, A. 2015.  
3 From Wastes to High Value Added Products: Novel Aspects of SSF in the Production of  
4 Enzymes. *Crit. Rev. Env. Sci. Technol.*, 45, 1999-2042.
- 5 [2] Saxena, R., Singh, R. 2011. Amylase production by solid-state fermentation of agro-  
6 industrial wastes using *Bacillus* sp. *Braz. J. Microbiol.*, 42, 1334-1342.
- 7 [3] Pitol, L.O., Biz, A., Mallmann, E., Krieger, N., Mitchell, D.A. 2016. Production of  
8 pectinases by solid-state fermentation in a pilot-scale packed-bed bioreactor. *Chemical Eng.*  
9 *J.*, 283, 1009-1018.
- 10 [4] Mitchell, D., Berovič, M., Krieger, N. 2006. Solid-State Fermentation Bioreactor  
11 Fundamentals: Introduction and Overview. in: *Solid-State Fermentation Bioreactors*, (Eds.)  
12 D. Mitchell, M. Berovič, N. Krieger, Springer Berlin Heidelberg, pp. 1-12.
- 13 [5] Casciadori, F.P., Bück, A., Thoméo, J.C., Tsotsas, E. 2016. Two-phase and two-  
14 dimensional model describing heat and water transfer during solid-state fermentation within a  
15 packed-bed bioreactor. *Chemical Eng. J.*, 287, 103-116.
- 16 [6] Abraham, J., Gea, T., Sánchez, A. 2013. Potential of the solid-state fermentation of soy  
17 fibre residues by native microbial populations for bench-scale alkaline protease production.  
18 *Biochem. Eng. J.*, 74, 15-19.
- 19 [7] El-Bakry, M., Gea, T., Sánchez, A. 2016. Inoculation effect of thermophilic  
20 microorganisms on protease production through solid-state fermentation under non-sterile  
21 conditions at lab and bench scale (SSF). *Bioprocess Biosyst. Eng.*, DOI: 10.1007/s00449-  
22 016-1540-x.
- 23 [8] Alkasrawi, M., Jrai, A.A., Al-Muhtaseb, A.H. 2013. Simultaneous saccharification and  
24 fermentation process for ethanol production from steam-pretreated softwood: Recirculation  
25 of condensate streams. *Chemical Eng. J.*, 225, 574-579.

- 1 [9] Cheirsilp, B., Kitcha, S. 2015. Solid state fermentation by cellulolytic oleaginous fungi  
2 for direct conversion of lignocellulosic biomass into lipids: Fed-batch and repeated-batch  
3 fermentations. *Ind. Crops Prod.*, 66, 73-80.
- 4 [10] Farinas, C.S. 2015. Developments in solid-state fermentation for the production of  
5 biomass-degrading enzymes for the bioenergy sector. *Renew. Sustain. Energy Rev.*, 52, 179–  
6 188
- 7 [11] López-González, J.A., Suárez-Estrella, F., Vargas-García, M.C., López, M.J., Jurado,  
8 M.M., Moreno, J. 2015. Dynamics of bacterial microbiota during lignocellulosic waste  
9 composting: Studies upon its structure, functionality and biodiversity. *Bioresour. Technol.*,  
10 175, 406-416.
- 11 [12] López-González, J.A., Vargas-García, M.d.C., López, M.J., Suárez-Estrella, F., Jurado,  
12 M.d.M., Moreno, J. 2015. Biodiversity and succession of mycobiota associated to agricultural  
13 lignocellulosic waste-based composting. *Bioresour. Technol.*, 187, 305-313.
- 14 [13] Zeng, G., Yu, M., Chen, Y., Huang, D., Zhang, J., Huang, H., Jiang, R., Yu, Z. 2010.  
15 Effects of inoculation with *Phanerochaete chrysosporium* at various time points on enzyme  
16 activities during agricultural waste composting. *Bioresour. Technol.*, 101, 222-227.
- 17 [14] Babu, K.R., Satyanarayana, T. 1995.  $\alpha$ -Amylase production by thermophilic *Bacillus*  
18 *coagulans* in solid state fermentation. *Process Biochem.*, 30, 305-309.
- 19 [15] Kunamneni, A., Permaul, K., Singh, S. 2005. Amylase production in solid state  
20 fermentation by the thermophilic fungus *Thermomyces lanuginosus*. *J. Biosci, Bioeng.*, 100,  
21 168-171.
- 22 [16] Petrova, S.D., Ilieva, S.Z., Bakalova, N.G., Atev, A.P., Bhat, M.K., Kolev, D.N. 2000.  
23 Production and characterization of extracellular  $\alpha$ -amylases from the thermophilic fungus  
24 *Thermomyces lanuginosus* (wild and mutant strains). *Biotechnol. Letters*, 22, 1619-1624.

- 1 [17] Hashemi, M., Mousavi, S.M., Razavi, S.H., Shojaosadati, S.A. 2013. Comparison of  
2 submerged and solid state fermentation systems effects on the catalytic activity of *Bacillus*  
3 sp. KR-8104  $\alpha$ -amylase at different pH and temperatures. *Ind. Crops Prod.*, 43, 661-667.
- 4 [18] Mukherjee, A.K., Borah, M., Rai, S.K. 2009. To study the influence of different  
5 components of fermentable substrates on induction of extracellular  $\alpha$ -amylase synthesis by  
6 *Bacillus subtilis* DM-03 in solid-state fermentation and exploration of feasibility for inclusion  
7 of  $\alpha$ -amylase in laundry detergent formulations. *Biochem. Eng. J.*, 43, 149-156.
- 8 [19] Rajagopalan, G., Krishnan, C. 2010. Hyper-production of alpha-amylase from agro-  
9 residual medium with high-glucose in SSF using catabolite derepressed *Bacillus subtilis*  
10 KCC103. *J. Basic Microbiol.*, 50, 336-43.
- 11 [20] Astolfi, V., Joris, J., Verlindo, R., Oliveira, J.V., Maugeri, F., Mazutti, M.A., de  
12 Oliveira, D., Treichel, H. 2011. Operation of a fixed-bed bioreactor in batch and fed-batch  
13 modes for production of inulinase by solid-state fermentation. *Biochem. Eng. J.*, 58–59, 39-  
14 49.
- 15 [21] Tao, S., Beihui, L., Peng, L., Deming, L., Zuohu, L. 1998. New solid-state fermentation  
16 process for repeated batch production of fibrinolytic enzyme by *Fusarium oxysporum*.  
17 *Process Biochem.*, 33, 419-422.
- 18 [22] Swain, M.R., Kar, S., Padmaja, G., Ray, R.C. 2006. Partial characterization and  
19 optimization of production of extracellular  $\alpha$ -amylase from *Bacillus subtilis* isolated from  
20 culturable cow dung microflora. *Polish J. Microbiol.*, 55, 289-296.
- 21 [23] The US Department of Agriculture and The US Composting Council, Test methods for  
22 the examination of composting and compost, Edaphos International, Houston, 2001
- 23 [24] De Gannes, V., Eudoxie, G., Hickey, W.J. 2013. Insights into fungal communities in  
24 composts revealed by 454-pyrosequencing: implications for human health and safety.  
25 *Frontiers Microbiol.*, 4, 164.

- 1 [25] Zhang, X., Zhong, Y., Yang, S., Zhang, W., Xu, M., Ma, A., Zhuang, G., Chen, G., Liu,  
2 W. 2014. Diversity and dynamics of the microbial community on decomposing wheat straw  
3 during mushroom compost production. *Bioresour. Technol.*, 170, 183-195.
- 4 [26] De Castro, R.J.S., Sato, H.H. 2013. Synergistic effects of agroindustrial wastes on  
5 simultaneous production of protease and  $\alpha$ -amylase under solid state fermentation using a  
6 simplex centroid mixture design. *Ind. Crops Prod.*, 49, 813-821.
- 7 [27] Kim, H.W., Lee, Y.L., Kim, Y.H. 2015. Efficacy of pectin and insoluble fiber extracted  
8 from soy hulls as a functional non-meat ingredient. *Food Sci. Technol.*, 64(2), 1071-1077.
- 9 [28] N.B. Tuncel, N. Yılmaz, H. Kocabıyık, A. Uygur. 2014. The effect of infrared stabilized  
10 rice bran substitution on B vitamins, minerals and phytic acid content of pan breads: Part II.  
11 *J. Cereal Sci.*, 59(2), 162–166.
- 12 [29] Song, M.W. 1990. Regulation of Alpha-amylase Synthesis in *Bacillus Subtilis*.  
13 University of Wisconsin-Madison.
- 14 [30] Pognani, M., Barrena, R., Font, X., Sánchez, A. 2012. Effect of freezing on the  
15 conservation of the biological activity of organic solid wastes. *Bioresour. Technol.*, 104, 832-  
16 836.
- 17 [31] Ugwuanyi, J.O., Harvey, L.M., McNeil, B. 2004. Development of thermophilic  
18 populations, amylase and cellulase enzyme activities during thermophilic aerobic digestion of  
19 model agricultural waste slurry. *Process Biochem.*, 39, 1661-1669.
- 20 [32] Jurado, M.M., Suárez-Estrella, F., López, M.J., Vargas-García, M.C., López-González,  
21 J.A., Moreno, J. 2015. Enhanced turnover of organic matter fractions by microbial  
22 stimulation during lignocellulosic waste composting. *Bioresour. Technol.*, 186, 15-24.
- 23  
24  
25

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17

Table 1. Characterization of raw materials used on SSF (average  $\pm$  standard deviation)

Parameter*	Soy fibre	Bread waste
Dry mass (%)	17.7 $\pm$ 1.3	62.8 $\pm$ 0.8
Organic matter (%)	96.8 $\pm$ 1.3	96.95 $\pm$ 0.2
C (%)	67.5 $\pm$ 1.8	48.1 $\pm$ 2.1
N (%)	4.4 $\pm$ 0.7	2.5 $\pm$ 0.2
pH	7.0 $\pm$ 0.7	6.1 $\pm$ 0.1
EC	0.9 $\pm$ 0.3	3.6 $\pm$ 0.6
Starch (%)	3.1 $\pm$ 0.1	75.7 $\pm$ 0.1

\* all percentages in dry basis except for dry mass, in wet basis.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

Table 2. Maximum sOUR and amylase activity obtained in SSF using different soy:bread ratios

soy:bread (%w/w)	Maximum sOUR (mgO <sub>2</sub> g <sup>-1</sup> DM h <sup>-1</sup> )	Maximum amylase activity (10 <sup>3</sup> U g <sup>-1</sup> DS)
0:100	3.4	10.9 ± 0.5
10:90	4.1	14.7 ± 0.7
50:50	5.8	18.9 ± 0.9
90:10	11.5	39.9 ± 0.9
100:00	11	34.6 ± 1.2

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

**Figure list**

Figure 1. Solid-state fermentation at 500 mL scale using *Thermomyces* sp. and *Geobacillus* sp. as inoculums. Profiles of sOUR and amylase activity are presented using soy fibre a) and b) and bread wastes c) and d) respectively.

Figure 2. Solid-state fermentation at 4.5 L scale using soy:bread 90:10 ratio as substrate. Profiles of sOUR, temperature and amylase activity are presented using a) no inoculation, b) *Thermomyces* sp. inoculation.

Figure 3. MOUR strategy in 4.5 L reactors using soy:bread 90:10 ratio as substrate. Profiles of sOUR, temperature and amylase activity are presented for the propagation reactor and the three sequential batches MOUR-SB1, MOUR-SB2 and MOUR-SB3.

Figure 4. MAA strategy in 4.5 L reactors using soy:bread 90:10 ratio as substrate. Profiles of sOUR, temperature and amylase activity are presented for the propagation reactor and three sequential batches MAA-SB1, MAA-SB2 and MAA-SB3.

Figure 5. MAAE strategy in 4.5 L reactors using soy:bread 90:10 ratio as substrate. Profiles of sOUR, temperature and amylase activity are presented for the propagation reactor and three sequential batches MAAE-SB1, MAAE-SB2 and MAAE-SB3.






Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.