




Reliability and validity of the Positive Mental Health Questionnaire in a sample of Spanish university students

Journal:	<i>Journal of Psychiatric and Mental Health Nursing</i>
Manuscript ID	JPM-16-0166.R2
Manuscript Type:	Original Article
Keywords:	Mental Health Promotion, Scales and Assessment, Public Health, Health Promotion



1
2
3 ***Reliability and validity of the Positive Mental Health Questionnaire in a***
4 ***sample of Spanish university students***
5
6
7
8
9

10
11 **Accessible summary**
12

13
14 - **What is known on the subject**
15

16
17 In general, the current studies of positive mental health use questionnaires or parts
18 thereof. However, while these questionnaires evaluate aspects of positive mental
19 health, they fail to measure the construct itself.
20
21
22

23
24
25
26 - **What this paper adds to existing knowledge**
27

28
29 The widespread use and the lack of specific questionnaires for evaluating the
30 positive mental health construct justify the need to measure the robustness of the
31 Positive Mental Health Questionnaire. Also six factors are proposed to measure
32 positive mental health.
33
34
35
36

37
38 - **What are the implications for practice**
39

40
41 The availability of a good questionnaire to measure positive mental health in
42 university students is useful for promoting not only mental health but also to
43 strengthen the curricula of future professionals.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Introduction

Nursing has a relevant role in managing mental health. It is important to identify and thereafter enhance positive aspects of mental health among university nursing students.

Aim

The aim of the present study was to analyze the psychometric properties of the Positive Mental Health Questionnaire (PMHQ) in terms of reliability and validity using confirmatory factor analysis in a sample of university students.

Method

A cross-sectional study was carried out in a sample of 1091 students at 4 nursing schools in Catalonia, Spain. The reliability of the PMHQ was measured by means of Cronbach's alpha coefficient, and the test-retest stability was measured with the intraclass correlation coefficient (ICC). Confirmatory factor analysis was used to determine the validity of the factorial structure.

Results

Cronbach's alpha coefficient was satisfactory (>0.70) for four of the six subscales or dimensions and ranged from 0.54 to 0.79. ICC analysis was satisfactory for the six subscales or dimensions. The hypothesis was confirmed in the analysis of the correlations between subclasses and the overall scale, with the strongest correlations being found between the majority of the subscales and the overall scale. Confirmatory factor analysis showed that the model proposed for the factors fit the data satisfactorily.

Discussion

This scale is a valid and reliable instrument for evaluating positive mental health in university students.

Implications for Practice

A good questionnaire to measure positive mental health in university students is useful for promoting not only mental health but also to strengthen the curricula of future professionals.

Keywords: Mental Health Promotion; Health Promotion; Public Health; Scales and Assessment

Relevance Statement: The availability of a good questionnaire to measure positive mental health in university students is useful for promoting not only mental health but also to strengthen the curricula of future professionals.

In addition, an instrument such as this would be very useful for nurses to assess positive aspects of mental health of patients and develop care plans to reinforce positive mental health (Orem & Vardiman, 1995).

In psychiatry the construct of positive mental health is useful for the creation of programmes for the promotion of mental health and the prevention of mental illness as well as evaluating their effectiveness. This construct would also be of use in the approach to perspectives on the current understanding of therapeutic activity that includes aspects of resiliency and recovery.

BACKGROUND

In 2001 the World Health Organization (WHO) provided a specific definition of mental health that emphasized the positive perspective: [mental health is a] "*state of well-being in which the individual realizes his skills, copes with the normal stresses of life, can work productively and fruitfully, and is able to make a contribution to his community.*" This definition has strongly reinforced the need to fully accept the original ideas put forward at the outset of the 20th century that saw mental health as something

1
2
3 more than the mere absence of illness. Indeed, this position has led to the formulation
4
5 of explanatory models of health, and by extension mental health, as a continuum, with
6
7 the promotion of mental health as a primary objective (Barry, 2009; Herman, Saxena &
8
9 Moodie, 2005; Jané-Llopis, 2007; World Health Organization, 2015). It has also
10
11 generated models with dimensions and separate but inter-related axes, based on
12
13 positive concepts such as happiness, flourishing, satisfaction with life, positive affect,
14
15 and meaning of life (Huppert & So, 2013; Kashdan, Biswas-Diener & King, 2009;
16
17 Keyes, 2002; Lamers, 2012; Peterson, Park & Seligman, 2005; Westerhof & Keyes,
18
19 2010).

20
21
22
23 A strong positive current on mental health has emerged from psychology emphasizing
24
25 the need to exploit and develop the strengths of the human being (Bolier, Haverman,
26
27 Westerhof, Riper, Smit & Bohlmeijer, 2013; Kobau, Seligman, Peterson, Diener, Zack,
28
29 Chapman & Thompson, 2011; Peterson & Seligman, 2004). As early as 1994, Bowling
30
31 stated that although a high percentage of the population suffered or was in danger of
32
33 suffering mental illness (up to 20% according to some reports), it was also necessary to
34
35 attend the healthy population to prevent the appearance of mental health problems and
36
37 to enhance positive mental health (Bowling, 1994). Along parallel lines, from the
38
39 perspective of mental health professionals, positive mental health would allow
40
41 professionals to focus on the healthy aspects of the individual in terms of resilience
42
43 (Vilete, Figueira, Andreoli, Ribeiro, Quintana, Mari & Coutinho, 2014) and/or recovery
44
45 (Provencher & Keyes, 2011).

46
47
48
49 In order to be able to develop interventions aimed at promoting positive mental health
50
51 conceptual models are first needed to guide the actions to be taken. Furthermore, we
52
53 also need instruments that can measure levels of positive mental health both before (to
54
55 carry out health diagnosis) and after (to evaluate results in health) the intervention.
56
57
58
59
60

1
2
3 Following extensive review of the literature Lluch found that the work of Jahoda (1958)
4
5 on conceptual aspects of positive mental health had not been implemented. In his
6
7 doctoral thesis, Lluch made three consecutive studies and structured positive mental
8
9 health in a multifactor model made up of 6 factors to explain that positive mental health
10
11 involves both theoretical and practical aspects which address concepts independently
12
13 or with relationships between two or three concepts. For example, personal satisfaction
14
15 with autonomy or self-control (Barry, Clarke, Jenkins & Patel, 2013; Hofmann,
16
17 Luhmann, Fisher, Vohs & Baumeister, 2014; Min, Lee & Lee, 2013; Orpana, Vachon,
18
19 Dykxhoorn, McRae & Jayaraman, 2016). In addition, there are multicomponent
20
21 intervention programs in mental health that use techniques such as problem solving
22
23 and social skills combined (Forsman, Nordmyr & Wahlbeck, 2011; van der Stouwe,
24
25 Asscher, Hoeve, Van der Laan & Stams, 2016). Indeed, no previous study has
26
27 integrated the six factors within a conceptual framework to provide a model category
28
29 highlighting the need to further explore its structure and configuration for future
30
31 confirmation as a model.
32

33
34
35
36 The study on Positive Mental Health (PMH) performed by Luch in 1999 had two
37
38 complementary objectives: to define a conceptual model of positive mental health and
39
40 to construct an instrument to make the conceptual model operative to evaluate positive
41
42 mental health. The result of this effort was the creation of the Multifactor Model of
43
44 Positive Mental Health (MM-PMH). This model is made up of 6 factors that define the
45
46 construct of Positive Mental Health (Lluch, 1999, 2003): Personal Satisfaction (F1),
47
48 Prosocial Attitude (F2), Self-control (F3), Autonomy (f4), Problem-Solving -and Self-
49
50 actualization (F5) and Interpersonal Relationship Skills (F6) (Table 1).
51

52
53
54
55 On the basis of MM-PMH the Positive Mental Health Questionnaire (PMHQ) was
56
57 created to make the model operative and to measure positive mental health. The
58
59
60

1
2
3 PMHQ questionnaire comprises 39 items which are unevenly distributed across the six
4
5 factors that define the construct. The distribution of the items of each factor is
6
7 described in Methods. Initial validation of the PMHQ was carried out in a sample of
8
9 students from the nursing school of the University of Barcelona (Lluch, 2003). Gold
10
11 standard assessments were followed to establish the reliability and validity of the new
12
13 instrument (Nunnally & Bernstein, 1994). In the exploratory factor analysis the six
14
15 factors extracted accounted for 46.8% of the total variance of the questionnaire. In the
16
17 resulting factor matrix, the weights of each item with respect to the factor extracted
18
19 were above 0.40 in all cases (Hair, J. F., Anderson, R. E., Tatham, R. L., & Black, W.
20
21 C, 1998). Overall, the psychometric results were favourable, but the sample size of this
22
23 initial study was small and confirmatory factor analysis was not performed.
24

25
26 The PMHQ has been used by various research teams, especially in Spain, Portugal,
27
28 and Latin America (Amar, Palacio, Llinas, Puerta, Sierra, Pérez & Velásquez, 2008;
29
30 Lluch-Canut, Puig-Llobet, Sánchez-Ortega, Roldán-Merino, Ferré-Grau & Positive
31
32 Mental Health Research Group, 2013; Sequeira, Carvalho, Sampaio, Sá, Lluch &
33
34 Roldán, 2014). However, very few studies have analysed its psychometric properties
35
36 (Lluch et al. 2013; Sequeira et al., 2014) and to our knowledge, this is the first
37
38 confirmatory factor analysis of the questionnaire.
39

40
41 The widespread use and the lack of specific questionnaires for evaluating the positive
42
43 mental health construct justify the need to measure the robustness of the Positive
44
45 Mental Health Questionnaire, as has been pointed out by the EUROHIS group
46
47 (Meltzer, 2003). The only specific instrument for evaluating positive mental health as a
48
49 construct was described in a recent study carried out in an Asian population on the
50
51 relevance of religious and spiritual practices in mental health (Vaingankar,
52
53 Subramaniam, Abdin, Picco, Chua, Eng, Sambavisam, Shafie, Zhang & Chong, 2014).
54
55 In general, the current studies of positive mental health use questionnaires such as the
56
57 Warwick-Edinburgh Mental Well-being Scale-WEMWBS (Tennant, Hiller, Fishwick,
58
59
60

1
2
3 Platt, Joseph, Weich, Parkinson, Secker & Stewart-Brown, 2007), the Sense of
4
5 Coherence Scale-SOC (Antonovsky, 1993), the Mental Health Continuum-Short Form-
6
7 MHC-SF (Keyes, 2002), the WHO-five Well-being Index (Bech, 2004) or the
8
9 Affectometer 2 (Kamman & Flett, 1983) that evaluate aspects of mental well being, but
10
11 they fail to measure the construct “positive mental health” itself (Dreger, Buck & Bolte,
12
13 2014; Lamers, 2012). According to Cronbach & Meehl (1955), the main difficulty of a
14
15 construct is that it is *“a concept for which there is not a single observable referent,*
16
17 *which cannot be directly observed, and for which there exist multiple referents, but*
18
19 *none all-inclusive”*.

20
21
22 The question is, however, how many factors or criteria are required to define mental
23
24 health? Researchers in Canada have used the definition of mental health formulated by
25
26 the PHAC (Public Health Agency of Canada) to operationalize the positive mental
27
28 health construct into five components (Canadian Institute for Health Information, 2009,
29
30 2011). In our work, we took the conceptualization of positive mental health described
31
32 by Jahoda (1958) as the initial reference point followed by the formulation of the MM-
33
34 PMH, which as noted above, defines the construct in terms of 6 factors.

35
36
37
38 Nursing professionals need validated instruments that may be used in daily practice.
39
40 Thus, the model of positive mental health as expressed in this study may be of great
41
42 importance for both the prevention and promotion of mental health.

43
44
45
46 The aim of this study was to analyse the psychometric properties of the PMHQ in terms
47
48 of reliability and validity by means of confirmatory factor analysis in a sample of
49
50 university students.

51 52 53 54 **METHODS**

55 56 57 58 59 60 **DESIGN**

1
2
3 We performed a cross-sectional study to validate the psychometric properties of the
4
5 PMHQ in a sample of 1091 nursing students at 4 university nursing schools in
6
7 Catalonia, Spain.
8
9

10 11 **PARTICIPANTS AND SETTING**

12
13 Data were collected from first-year nursing students during the years 2012-2014. The
14
15 inclusion criteria were: registration in the program and voluntary participation in the
16
17 study. The 4 university schools of nursing included: Bellvitge and Sant Joan de Dèu,
18
19 both affiliated with the University of Barcelona, and Blanquerna and Rovira i Virgili, the
20
21 latter two being private institutions. The total student population during the study period
22
23 was of approximately 1500 students.
24
25

26 27 **INSTRUMENTS**

28
29 The following demographic data of the participants were collected: age, sex, nursing
30
31 school, physical and mental problems in the preceding month, and visits to a physician,
32
33 psychologist, or psychiatrist in the preceding month.
34

35
36 The PMHQ (Positive Mental Health Questionnaire), Lluch (1999, 2003): this
37
38 questionnaire comprises 39 items which are unevenly distributed across the six factors
39
40 that define the construct: F1. Personal Satisfaction (8 items), F2. Prosocial Attitude (5
41
42 items), F3. Self-control (5 items), F4. Autonomy (5 items), F5. Problem-solving and
43
44 Self-actualization (9 items), and F6. Interpersonal Relationship Skills (7 items). The
45
46 items are expressed as positive or negative statements which are responded to using a
47
48 scale ranging from 1 to 4, according to how frequently they occur: always or almost
49
50 always, quite often, sometimes, never or rarely. The distribution of the 39 items of the
51
52 PMHQ among the six factors is shown in Table 1. The PMHQ was constructed in
53
54 Spanish and has been translated and adapted to Portuguese (Sequeira, Carvalho,
55
56 Sampaio, Sá, Lluch & Roldán, 2014).
57
58
59
60

ETHICAL CONSIDERATIONS

This study was approved by the Ethics and Research Committee of the University of Barcelona. All participants were informed of the purpose of the study and provided oral informed consent to participate. The study was anonymous and confidential.

DATA COLLECTION PROCEDURES

The questionnaire was administered on one day at each school to all students attending classes on that day and who voluntarily agreed to participate.

Four weeks later, the PMHQ was again administered to the same 380 participants at the Bellvitge School of Nursing (University of Barcelona) during the academic year 2013-2104 to determine test-retest reliability.

STATISTICAL ANALYSIS

Data analysis was performed using SPSS for Windows 17.0 (SPSS, Chicago, IL, USA).

One-sample Kolmogorov-Smirnov Z tests were used to assess normality, and descriptive statistics were used to summarize the scale.

Item analyses included calculation of item means, standard deviations, percentage ceiling and floor effects, and corrected item-total correlation. Ceiling and floor effects are the percentage of people with the highest score and with the lower score respectively. The item-total correlation is the correlation between the score of one item and the sum of the scores of the remaining items. Internal consistency reliability was evaluated using Cronbach's alpha coefficient for the total and each of the subscales (Nunnally & Bernstein, 1994).

A coefficient alpha value of 0.70 (Nunnally, J. & Bernstein, 1994) or above was considered acceptable for this scale. Test-retest reliability was examined within a 4-week time frame using the intraclass correlation coefficient (ICC) criteria (two-factor and mixed effects model).

1
2
3
4
5 A correlation analysis was performed using the Spearman correlation between the
6
7 scales in PMHQ to evaluate convergent and discriminant validity based on the
8
9 hypothesis that the correlation between each subscale and the overall scale should be
10
11 higher than the correlations among the subscales (Fayers & Machin, 2000). The
12
13 discriminant validity was also evaluated by comparing the means of the positive mental
14
15 health questionnaire in the two groups of students according to whether they had
16
17 reported having a mental health problem and whether they had visited a
18
19 psychologist/psychiatrist on at least one occasion. The Student's t test was used for
20
21 this analysis.
22

23 The construct validity of the PMHQ was determined using confirmatory factor analysis
24
25 (CFA). CFA models were estimated using structural equation modeling (EQS 6.1 for
26
27 Windows, Multivariate Software, Inc., Encino, CA, USA). The generalized least squares
28
29 parameter estimation method of a polychoric correlation matrix was used. This method
30
31 has the same properties as the maximum likelihood method, although with less
32
33 stringent criteria of normality, and it is mainly used for measuring ordinal items (Batista-
34
35 Foguet, Coenders & Alonso, 2004). Model fit was determined with several methods
36
37 since several authors have suggested using a number of indicators to determine the fit
38
39 of the models (Bollen & Long, 1993). The statistics provided were the χ^2 test, the ratio
40
41 between χ^2 and the degrees of freedom ($\chi^2/d.f.$), the root mean square error of
42
43 approximation (RMSEA), the standardised root mean square residual (SRMR), the
44
45 Bentler-Bonett normed fit index (BBNFI), and the Bentler-Bonett (1980) non-normed fit
46
47 index (BBNNFI). The BBNFI, and BBNNFI indexes can vary from 0 (poor fit) to 1
48
49 (perfect fit). The RMSEA reflects the extent to which the model approximates a
50
51 reasonable fit, and values close to or below 0.07 are generally recommended (Steiger,
52
53 2007). Using a two-index presentation strategy, Hu and Bentler (1999) proposed a
54
55 cutoff of 0.09 for SRMR and 0.06 for RMSEA (values close to or below). According to
56
57 Hinkin (1995) $\chi^2/d.f.$ may be used when there is a ratio of 5:1 or less.
58
59
60

1
2
3 Statistical significance was set at a p value 0.05.
4
5

6 7 **RESULTS**

8
9 The questionnaires were completed by 1091 participants. The sociodemographic data
10 and clinical characteristics of the participants are shown in Table 2. The average age of
11 the participants was 21 years and most were female (87.5%). The majority of students
12 stated that they had no physical or mental health problems and also declared that they
13 had not seen a doctor, psychologist, or psychiatrist in the previous month.
14
15
16
17
18
19
20
21
22
23
24
25
26

27 **ITEM ANALYSIS**

28
29 The mean item value ranged from 2.47 (item 18) to 3.79 (item 3), and the standard
30 deviation ranged from 0.51 (item 3) to 0.90 (item 35). The item with the greatest ceiling
31 effect was item 3 (*I find it particularly difficult to listen to people telling me their*
32 *problems*) (82.7%), and the items with the greatest floor effect were item 10 (*I worry a*
33 *lot about what others think of me*) (12.6%), item 18 (*I consider myself to be a good*
34 *psychologist*) (12.6%), and item 19 (*It troubles me when people criticize me*) (13.0%)
35
36
37
38
39
40
41
42
43 (Table 3).
44

45 **RELIABILITY**

46
47 Cronbach's alpha coefficient was satisfactory (>0.70) for 4 of the 6 subscales or
48 dimensions and ranged from 0.54 to 0.79 (Table 4).
49

50
51 The internal consistency of the total PMHQ yielded an alpha of 0.89, indicating good
52 internal consistency for the 39-item PMHQ.
53
54
55
56
57
58
59
60

1
2
3 Most of the 39-item PMHQ had item-total correlations >0.20 . Only two items had
4 corrected item-total scale correlation coefficients <0.20 (items 1 and 3). These items
5 were: *I find it especially difficult to accept others when their attitudes are different from*
6 *mine*(item 1) and *I find it particularly difficult to listen to people telling me their problems*
7 (item 3) (Table 3).
8
9
10
11
12

13
14
15 ICC analysis demonstrated that the 4-week test-retest reliability was 0.92 (95%
16 confidence interval 0.91–0.93, $n = 380$) and was satisfactory for the 6 subscales or
17 dimensions (Table 4).
18
19
20
21
22

23 24 **CONVERGENT AND DISCRIMINANT VALIDITY**

25
26 The hypothesis was confirmed in the analysis of the correlations between the
27 subclasses and the overall scale, with the strongest correlations being found between
28 the majority of the subscales and the overall scale. Subscale F5 (Problem-solving and
29 Self-Actualization) most strongly correlated with the overall scale ($\rho=0.80$), while
30 subscale F2 (Prosocial Attitude) showed the weakest correlation with the overall scale
31 ($\rho=0.42$).
32
33
34
35
36
37

38 The strongest correlation among the subscales was that of F3 (Self-control) and F5
39 (Problem-solving and Self-Actualization) ($\rho=0.50$), while the weakest was between
40 F2 (Prosocial Attitude) and F4 (Autonomy) ($\rho=0.06$) (Table 5). All of the correlations
41 were significant ($p<0.001$). The mean total scores of the scale were greater in students
42 declaring not having had mental health problems ($t(28.52) = 3.90, p=0.001$) and in
43 those stating that they had not visited a psychologist/psychiatrist ($t(157.20) = 3.45,$
44 $p=0.001$).
45
46
47
48
49
50
51
52
53
54

55 **CONSTRUCT VALIDITY**

56
57
58
59
60

1
2
3 The result of the χ^2 test was significant ($\chi^2 = 2930.07$; $p < 0.001$), indicating that the
4 hypothesis of a perfect fit model should be rejected. It was considered that other
5 statistics were needed to assess the model due to the sensitivity of the χ^2 test to
6 sample size. The value for RMSEA was 0.05 and the SRMR was 0.05 which are below
7 the recommended critical limits of 0.06 and 0.09, respectively. The $\chi^2/d.f.$ ratio was
8 equal to 4.26, which is within the acceptable value for this ratio of up to a maximum of
9 5. The value for BBNFI was 0.77, being 0.81 for BBNNFI. According to the results
10 presented above, the model proposed for the factors satisfactorily fit the data.
11
12
13
14
15
16
17
18
19

20
21 The parameters estimated by the model were all significantly different from zero
22 (Figure 1). No items had loads under 0.30 in the factor analysis (Kline, 2011).
23
24
25
26

27 DISCUSSION

28
29
30 The results obtained using several different analyses have shown that the PMHQ has
31 good psychometric properties. The scale was developed with the aim of evaluating the
32 positive mental health construct considering 6 dimensions or factors for the definition of
33 the construct: Personal Satisfaction (F1), Prosocial Attitude (F2), Self-control (F3),
34 Autonomy (F4), Problem-solving and Self-actualization (F5), and Interpersonal
35 Relationship Skills (F6).
36
37
38
39
40
41
42

43 In relation to the analysis of the items, it was of note that item 3 presented an elevated
44 ceiling effect (82.7 %) which may simply be due to nursing students having a greater
45 predisposition to listen to people with problems. Regarding reliability, 4 of the 6 factors
46 proposed (F1, F3, F4, F5 and F6) scored good levels of reliability in terms of internal
47 consistency. In the present study Factor F2 (Prosocial Attitude) scored the lowest
48 Cronbach's alpha coefficient. This result is similar to that of other studies in which the
49 results ranged from a maximum score of 0.60 in the study by Lluch et al. (2013) in a
50
51
52
53
54
55
56
57
58
59
60

1
2
3 sample of people with chronic physical health disorders to a score of 0.51 reported by
4
5 Sequeira et al. (2014) in a sample of Portuguese university students.
6

7
8 In regard to the stability over time of the PMHQ, our test-retest correlations were high
9
10 as in other studies, ranging from 0.98 in the study by Sequeira et al. (2014) to 0.92 for
11
12 the overall scale in our study. In previous studies including analysis of stability over
13
14 time, test-retest evaluation was performed within time periods ranging from 30 days, as
15
16 in the present study, to 60 days as in the study by Sequeira et al. (2014).
17

18
19
20 In view of all these results we can therefore conclude that the PMHQ possesses a
21
22 good level of reliability for measuring the construct of positive mental health.
23

24
25
26 In the present study, the results obtained from analysis of the PMHQ items were higher
27
28 (range of 2.47 for item 18, to 3.79 for item 3) than those obtained in a previous study
29
30 (Albacar, Lleixá, Lluçh, Sequeira, Carvalho & Roldán, 2015), (range of 1.22 for item 23,
31
32 to 2.31 for item 18). These results may be due to our population being made up of
33
34 university students while that of Albacar et al. (2015) included care-givers of patients
35
36 diagnosed with schizophrenia whose response may have been influenced by tiredness.
37
38 Furthermore, the range of the standard deviation in the present study was slightly
39
40 narrower than that of Albacar et al. (2015), ranging from 0.51 (item 3) to 0.90 (item 35)
41
42 in our study and from 0.53 (item 23) and 1.13 (item 35) in the latter. In both studies the
43
44 item showing the greatest deviation was number 35, *'I am able to say no when I want*
45
46 *to.'*
47

48
49 On comparing the correlation among factors and subscales of the PMHQ of the present
50
51 study with that of the studies by Lluçh (1999, 2003), Lluçh et al. (2013), and Sequeira
52
53 et al. (2014), the scores were similar for the three studies, and in all three cases the six
54
55 factors or subscales showed statistically significant correlations below 0.70.
56
57
58
59
60

1
2
3 Since the present study is the first confirmatory factor analysis of the PMHQ our results
4 cannot be compared with others. As shown in results, the model proposed for the
5 factors satisfactorily fit the data and the parameters estimated by the model were all
6 significantly different from zero. No items had loads under 0.30 in the factor analysis.
7
8 The results obtained are in agreement with the initial factorial data for PMHQ obtained
9 in the exploratory factor analysis by Lluch (1999, 2003). All of the items on the
10 questionnaire scored factor loads greater than 0.30 (ranging from 0.42 to 0.78). Similar
11 results were obtained in the exploratory factor analysis carried out by Sequeira et al.
12 (2014) in which the item saturation ranged from 0.38 to 0.84. The results of the
13 confirmatory factor analysis of the present study confirm the distribution of the items in
14 the factors as well as the multifactorial structure of the PMHQ according to the model
15 proposed in support of the instrument.
16
17

18
19 On comparing our results with those using the positive mental health instrument of
20 Vaingarker et al (2014) the psychometric reliability was similar to that of the PMHQ,
21 with Cronbach's alpha values ranging between 0.73 and 0.91, and intraclass
22 correlation coefficients (ICC) ranging between 0.74 and 0.96. Similarly, the load factors
23 of the instrument and the fit indexes were also similar to those obtained with the
24 PMHQ, ranging from 0.208 to 0.882. The correlations among subscales of the PMHQ
25 were smaller (range 0.07 to 0.51) than those of Vaingarkar et al. (2014) (range 0.25 to
26 0.81) indicating that the factors or subscales of the PMHQ have good levels of
27 discrimination. In both studies the correlations among subscales or factors were
28 statistically significant.
29
30

31
32 Finally, it was of note that the reliability and validity scores for the PMHQ questionnaire
33 were similar to the psychometric characteristics of other scales which also evaluate
34 constructs or dimensions of a positive nature and are widely used in the field of mental
35 health. These scales include the General Well-Being Schedule, GWBS (Dupuy, 1984),
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the Warwick-Edinburgh Mental Well-being Scale-WEMWBS (Tennant, Hiller, Fishwick,
4 Platt, Joseph, Weich, Parkinson, Secker & Stewart-Brown, 2007), the Sense of
5 Coherence Scale-SOC (Antonovsky, 1993), the Mental Health Continuum-Short Form-
6 MHC-SF (Keyes, 2002), the WHO-five Well-being Index (Bech, 2004), and the
7 Affectometer 2 (Kamman & Flett, 1983), among others.
8
9
10
11
12

13
14
15 However, these scales do not assess the construct "positive mental health". Moreover
16 as stated by Lehtinen, Sholman & Kovess-Masfety (2005), "*Happiness or life*
17 *satisfaction are necessarily not the same as positive mental health, although they can*
18 *be seen as essential components of the construct. More research on the epidemiology*
19 *of positive mental health is evidently needed*".
20
21
22
23
24
25
26

27 Conceptual approaches and instruments are needed to encourage the development of
28 the construct "positive mental health". Moreover, the PMHQ questionnaire and its
29 supporting conceptual model provide a new approach to the present international
30 evidence to advance in the study of the positive aspects of mental health.
31
32
33
34
35
36

37 Limitations

38
39 Nevertheless, the present study has some limitations. First, the questionnaire was
40 performed in a sample of students who voluntarily participated and thus, may not be
41 representative of the target student population. In addition, the students who accepted
42 to participate may have had better positive mental health the day the questionnaire was
43 given, thereby overestimating the results. The response rate was 73 % (1091/1500)
44 and, therefore, the influence of this bias would be minimal.
45
46
47
48
49
50
51

52
53 Secondly, in relation to the homogeneity of the sample, the results should be
54 considered with caution. We agree with Bech (2012) that the validation of measuring
55 instruments is an ongoing process that requires very large samples. Moreover,
56
57
58
59
60

1
2
3 evaluation of the construct further enhances the complexity of the validation of
4
5 measurement and thus, additional studies with more heterogeneous samples should
6
7 be performed. In this study 88.2% of the sample was made up of women with possible
8
9 differences in the perception of positive mental health compared to men.
10

11
12
13 On the other hand, it should be taken into account that the test-retest was undertaken
14
15 in only one teaching centre. However, the sociodemographic characteristics of the
16
17 students from all the centres were similar and should not bias the results.
18
19

20
21 Finally, it would also be important to evaluate the predictive validity with longitudinal
22
23 studies to analyse the correlation of this questionnaire with future measures of mental
24
25 health.
26
27

28 29 30 31 **CONCLUSIONS**

32
33 The PMHQ has demonstrated good measuring properties both at the overall level and
34
35 for the 6 factors with which it is constituted. Its internal consistency and stability over
36
37 time were favourable for five of the six factors that comprise the questionnaire: (F1)
38
39 Personal Satisfaction, (F2) Prosocial Attitude, (F3) Self-control, (F4) Autonomy, (F5)
40
41 Problem-solving and Self-actualization, and (F6) Interpersonal Relationship Skills.
42
43 Factor 2 (Prosocial Attitude) showed more moderate results in terms of reliability and
44
45 should be further evaluated in future studies. As to validity, the results of the
46
47 confirmatory factor analysis support the multifactor model of positive mental health that
48
49 underpins the questionnaire, yielding favourable data both for the 6 factors that make
50
51 up the PMHQ and for the 39 questionnaire items distributed among the 6 factors.
52
53

54 55 56 **RELEVANCE FOR CLINICAL PRACTICE**

1
2
3 The availability of a good questionnaire to measure positive mental health in university
4 students is useful for promoting not only mental health but also to strengthen the
5 curricula of future professionals.
6
7

8
9
10 In addition, an instrument such as this would be very useful for nurses to assess
11 positive aspects of mental health of patients and develop care plans to reinforce
12 positive mental health (Orem & Vardiman, 1995).
13

14
15
16 In psychiatry the construct of positive mental health is useful for the creation of
17 programmes for the promotion of mental health and the prevention of mental illness as
18 well as evaluating their effectiveness. This construct would also be of use in the
19 approach to perspectives on the current understanding of therapeutic activity that
20 includes aspects of resiliency and recovery.
21
22
23
24
25
26
27
28
29
30
31

32 REFERENCES:

- 33
34 Albacar-Ribóo, N., Lleixá-Fortuño, M., Lluch Canut, MT., Sequeira, C., Carvalho, JC. &
35 Roldán-Merino, J. (2015). Psychometric properties of the self-administered
36 version of the Self-Care Requisites Scale among carers for the mentally ill.
37 *Revista Portuguesa de Enfermagem de Saúde Mental* **13**, 53-60.
38
39
40
41 Amar, J., Palacio, J., Llinas, H., et al. (2008). Calidad de vida y salud mental positiva
42 en menores trabajadores de Tolviejo (Quality of life and positive mental health
43 in a youngworker of Tolviejo). *Suma Psicológica* **15** (2), 385-403.
44
45
46
47 Antonovsky, A. (1993). The structure and properties of the Sense of Coherence Scale.
48 *Soc Sci Med* **36** (6), 725-733.
49
50
51
52 Barry, M.M. (2009). Addressing the determinants of positive mental health: concepts,
53 evidence and practice. *International Journal of Mental Health Promotion* **11** (3),
54 4-17.
55
56
57
58
59
60

- 1
2
3 Barry, M.M., Clarke, A. M., Jenkins, R. & Patel, V. (2013). A systematic review of the
4 effectiveness of mental health promotion interventions for young people in low
5 and middle income countries. *BMC Public Health* **13**, 835.
6
7
8
9 Batista-Foguet, J.M., Coenders, G. & Alonso, J. (2004). Confirmatory Factor Analysis.
10 Its utility for Validating Health-Related Questionnaires. *Med Clin* **122**(1), 21-27.
11
12
13 Bech, P. (2004). Measuring the dimensions of psychological general well-being by the
14 WHO-5. *QoLNews* **32**, 15–6.
15
16
17 Bech, P. (2012). *Clinical Psychometrics*. Southern Gate, Chichester. West Sussex
18 (UK): John Wiley & Sons, Ltd.
19
20
21 Bentler, PM. & Bonett, DG. (1980). Significance tests and goodness-of-fit in the
22 analysis of covariance structures. *Psychological Bulletin* **88**, 588-600.
23
24
25 Bolier, L., Haverman, M., Westerhof, G.J., Riper, H., Smit, F. & Bohlmeijer, E. (2013).
26 Positive psychology interventions: a meta-analysis of randomized controlled
27 studies. *BMC Public Health* **13**, 119.
28
29
30
31 Bollen, KA. & Long, JS. (1993). *Testing Structural Equation Models*. Newbury Park,
32 CA: Sage.
33
34
35 Bowling, A. (1994). *Measuring Health. A Review of Quality of Life Measurement Scales*.
36 Open University Press.
37
38
39 Canadian Institute for Health Information. (2009). *Improving the Health of Canadians:
40 Exploring Positive Mental Health*. Ottawa.
41
42
43 Canadian Institute for Health Information. (2011). *Recognizing and Exploring Positive
44 Mental Health-Policy Dialogue: Synthesis and Analysis*. Ottawa.
45
46
47
48 Cronbach, LJ. & Meehl, PE. (1955). Construct validity in psychological tests.
49 *Psychological Bulletin* **52**, 281-302.
50
51
52 Dreger, S., Buck, C. & Bolte, G. (2014). Material, psychosocial and sociodemographic
53 determinants are associated with positive mental health in Europe: a cross-
54 sectional study. *BMJ Open*. doi: 10.1136/bmjopen-2014-005095.
55
56
57
58
59
60

- 1
2
3 Dupuy, HJ. (1984). Assessment of quality of life in clinical trials of cardiovascular
4 therapies. In: NK. Wenger., ME. Mattson., CD. Furberg., & J. Elison (Eds). *The*
5 *Psychological Well-Being (PGWB) Index*. (pp.184-188). Nueva York: Le Jacq
6 publishing.
7
8
9
10 Fayers, PM. & Machin, D. (2000). *Quality of life: Assessment, analysis and*
11 *interpretation*. London: John Wiley & Sons.
12
13
14 Forsman, A.K., Nordmyr., J. & Wahlbeck, K. (2011). Psychosocial interventions for the
15 promotion of mental health and the prevention of depression among older
16 adults *Health Promot. Int* **26** (1): 85-107. doi: 10.1093/heapro/dar074
17
18
19
20 Hair, J. F., Anderson, R. E., Tatham, R. L., & Black, W. C. (1998). *Multivariate data*
21 *analysis, 5th*. NY: Prentice Hall International.
22
23
24
25 Herman, H., Saxena, S. & Moodie, R. (2005). *Promoting mental health: concepts,*
26 *emerging evidence, practice. A report of the World Health Organization,*
27 *Department of Mental Health and Substance Abuse in collaboration with the*
28 *Victorian Health Promotion Foundation and The University of Melbourne.*
29 Geneva: World Health Organization.
30
31
32
33 Hinkin, TR. (1995). A review of scale development practices in the study of
34 organizations. *Journal of Management*, 21(5), 967-988.
35
36
37
38 Hofmann, W., Luhmann, M., Fisher, R.R., Vohs, K.D. & Baumeister, R.F. (2014). Yes,
39 but are they happy?. Effects of trait self-control on affective well-being and life
40 satisfaction. *Personality* **82**(4), 265-277.
41
42
43
44 Hu, L.T. & Bentler, P.M. (1999). Cutoff Criteria for Fit Indexes in Covariance Structure
45 Analysis: Conventional Criteria Versus New Alternatives. *Structural Equation*
46 *Modeling* **6**(1),1-55.
47
48
49 Huppert, FA. & So, TTC. (2013). Flourishing across Europe. Application of a new
50 conceptual framework for defining well-being. *Soc Indic Res* **110**, 837-861.
51
52
53
54 Jahoda, M. (1958). *Current concepts of positive mental health*. New York: Basic Books.
55
56
57 Jané-Llopis, E. (2007). Mental health promotion: concepts and strategies for reaching
58 the population. *Health Promotion Journal of Australia* **18**,191-197.
59
60

- 1
2
3 Kamman, R. & Flett, R. (1983). Affectometer 2: A scale to measure current level of
4 general happiness. *Australian Journal of Psychology* **35**, 259-265.
5
6
7 Kashdan, TB., Biswas-Diener, R., & King, LA. (2009). Two traditions of happiness
8 research, not two distinct types of happiness. *The Journal of Positive*
9 *Psychology* **4**, 208-211.
10
11
12 Keyes, CLM. (2002). The mental health continuum. From languishing to flourishing in
13 life. *Journal of Health and Social Behavior* **43**, 207-222.
14
15
16
17 Kline, R. B. (2011). *Principles and practice of structural equation modeling*. New York:
18 The Guilford Press.
19
20
21 Kobau, R., Seligman, M.E.P., Peterson, C., et al. (2011). Mental Health Promotion in
22 Public Health: Perspectives and Strategies From Positive Psychology.
23 *American Journal of Public Health* **101** (8),1-9.
24
25
26
27 Lamers, SMA. (2012). *Positive mental health: Measurement, relevance and*
28 *implications*. PhD Thesis. University of Twente: Enschede the Netherlands.
29
30
31 Lehtinen, V., Sholman, B. & Kovess-Masfety, V. (2005). Level of positive mental health
32 in the European Union: Results from the Eurobarometer 2002 survey. *Clin Pract*
33 *Epidemiol Ment Health* **1**(9).
34
35
36
37 Lluch, MT. (1999). *Construcción de una escala para evaluar la salud mental positiva*.
38 PhD Thesis. University of Barcelona: Faculty of Psychology.
39
40
41 Lluch, M.T. (2003) Construcción y análisis psicométrico de un cuestionario para
42 evaluar la salud mental positiva. *Psicología Conductual* **11**(1),61-78.
43
44
45 Lluch-Canut, T., Puig-Llobet, M., Sánchez-Ortega, A., Roldán-Merino, J., Ferré-
46 Grau, C. & Positive Mental Health Research Group (2013). Assessing positive
47 mental health in people with chronic physical health problems: correlations with
48 socio-demographic variables and physical health status. *BMC Public Health* **13**,
49 928.
50
51
52
53
54 Meltzer, H. (2003). *Development of common instruments for mental health*. In: A.
55 Norsikov & C. Gudex C (Eds), *EUROHIS: Developing Common Instruments for*
56 *Health Surveys* (pp. 35-60). Amsterdam (The Netherlands): IOS Press.
57
58
59
60

- 1
2
3 Min, J-A., Lee, Ch-U. & Lee, Ch (2013). Mental Health Promotion and Illness
4 Prevention: A Challenge for Psychiatrists. *Psychiatry Investig* **10**(4),307–316.
5
6
7 Nunnally, J. & Bernstein, I. (1994). Psychometric Theory, 3rd edn. New York: McGraw-
8 Hill.
9
10
11 Orem, D.E. & Vardiman, E.M. (1995). Orem's nursing theory and positive mental
12 health: practical considerations. *Nursing Science Quarterly* **8**(4), 165-173.
13
14 Orpana, H., Vachon, J., Dykxhoorn, J., McRae, L. & Jayaraman, G. (2016). Monitoring
15 positive mental health and its determinants in Canada: the development of the
16 Positive Mental Health Surveillance Indicator Framework. Health Promotion and
17 Chronic Disease Prevention in Canada Research, *Policy and Practice* **36**(1),1-10.
18
19
20
21
22 Peterson, C., Park, N. & Seligman, M. (2005). Orientations to happiness and life
23 satisfaction: The full life versus the empty life. *Journal of Happiness Studies* **6**,
24 25-41.
25
26
27
28 Peterson C, Seligman MEP. (2004). Character strengths and virtues. A handbook and
29 classification. New York: Oxford University Press.
30
31
32 Provencher, HL. & Keyes, CL. (2011). Mental health recovery: bridging mental illness
33 with positive mental health. *Journal of Public Mental Health* **10** (1), 57-69.
34
35
36 Sequeira, C., Carvalho, JC., Sampaio, F., Sá, L., Lluch, T. & Roldán, J. (2014).
37 Evaluation of the psychometric properties of the Positive Mental Health
38 Questionnaire in Portuguese higher education students. *Rev Portuguesa de*
39 *Enfermagem de Saúde Mental* **11**, 45-53.
40
41
42
43 Steiger, J.H. (2007). Understanding the limitations of global fit assessment in structural
44 equation modelling. *Personality and Individual Differences* **42** (5), 893-98.
45
46
47
48 Tennant, R., Hiller, L. Fishwick, R., et al. (2007). The Warwick-Edinburgh Mental Well-
49 being Scale (WEMWBS): development and UK validation. *Health Qual Life*
50 *Outcome* **5**, 36.
51
52
53 Vaingankar, JA., Subramaniam, M., Abidin, E., et al. (2014). Development, validity and
54 reliability of the short multidimensional positive mental health instrument.
55 *Quality of Life Research* **23**(5), 1459-1477.
56
57
58
59
60

1
2
3 Van der Stouwe, T., Asscher, J.J., Hoeve, M., Van der Laan, P.H. & Stams, J.M.
4 (2016). Social skills training for juvenile delinquents: post-treatment changes.
5 *Journal of Experimental Criminology* 1-22. doi:10.1007/s11292-016-9262-2
6
7

8
9
10 Vilete, L., Figueira, I., Andreoli, S.B., et al. (2014). Resilience to trauma in the two
11 largest cities of Brazil: a cross-sectional study. *BMC Psychiatry* **14**, 255.
12

13
14 Westerhof, GJ. & Keyes., CLM. (2010). Mental illness and mental health: two continua
15 model across the lifespan. *J Adult Dev* **17**, 110-119.
16

17
18 World Health Organization (2015). *The European Mental Health Action Plan 2013-*
19 *2020*. Copenhagen: WHO Regional Office for Europe.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Multifactor Model of Positive Mental Health (Lluch, 1999) and Distribution of items of each factor

Positive Mental Health factors	Definition	Items
F1: Personal Satisfaction	<ul style="list-style-type: none"> - Self-concept/Self-esteem - Satisfaction with personal life - Optimistic outlook on the future 	4, 6, 7, 12, 14, 31, 38, 39
F2: Prosocial Attitude	<ul style="list-style-type: none"> - Active predisposition towards society - Altruistic social attitude; attitude of helping/supporting others - Acceptance of others and of differential social characteristics 	1, 3, 23, 25, 37
F3: Self-control	<ul style="list-style-type: none"> - Ability to cope with stress/situations of conflict - Emotional balance/emotional control - Tolerance of frustration, anxiety and stress 	2, 5, 21, 22, 26
F4: Autonomy	<ul style="list-style-type: none"> - Able to have one's own standards - Independence - Self-regulation of one's behavior - Sense of personal security/self-confidence 	10, 13, 19, 33, 34
F5: Problem-solving and Self-actualization	<ul style="list-style-type: none"> - Analytical capacity - Ability to make decisions - Flexibility/ability to adapt to change - Attitude of continuous growth and personal development 	15, 16, 17, 27, 28, 29, 32, 35, 36
F6: Interpersonal Relationship Skills	<ul style="list-style-type: none"> - Ability to establish interpersonal relationships - Empathy/ability to understand the feelings of others - Ability to give emotional support - Ability to establish and maintain close interpersonal relationships 	8, 9, 11, 18, 20, 24, 30

Table 2
Sociodemographic and clinical characteristics of the study sample (n=1091)

	n	%
Mean age (years) min. 18 max 56	21.37 (Standard Deviation: 4.8)	
Sex		
Male	129	11.8
Female	962	88.2
School / Program		
Bellvitge - University of Barcelona	821	75.2
University Rovira i Virgili	74	6.8
Sant Joan de Dèu - University of Barcelona	124	11.4
Blanquerna	72	6.6
Physical health problems		
yes	97	8.9
no	601	86.1
Mental health problems		
yes	28	4.0
no	669	96.0
Visits to doctor		
yes	234	33.6
no	463	66.4
Visits to psychologist / psychiatrist		
yes	125	17.9
no	573	82.1

Table 3 Descriptive statistics of the items of the scale (PMHQ)
Item Mean, Standard Deviations, % Ceiling, % Floor, and Corrected Item – total Correlation

Summary of the contents of the items	Mean	Standard Deviation	% Floor	% Ceiling	Corrected Item – Total Correlation
Item 1: I find it especially difficult to accept others when their attitudes are different from mine.	3.26	.68	1.8	38.0	.198
Item 2: Problems often cause me to feel blocked.	2.86	.74	4.9	16.7	.480
Item 3: I find it particularly difficult to listen to people telling me their problems.	3.79	.51	0.7	82.7	.153
Item 4: I like myself as I am.	2.86	.82	4.1	24.3	.492
Item 5: I am able to control myself when I feel negative emotions.	2.54	.80	7.6	11.9	.397
Item 6: I feel like I am about to explode.	3.08	.65	2.0	23.9	.363
Item 7: I find life to be boring and monotonous.	3.49	.64	1.1	55.7	.417
Item 8: I find it particularly difficult to provide emotional support to others.	3.45	.70	1.9	55.8	.308
Item 9: I find it hard to establish deep and satisfying interpersonal relationships with some people.	3.32	.75	2.6	47.2	.308
Item 10: I worry a lot about what others think of me.	2.63	.86	12.6	13.2	.384
Item 11: I feel that I have a strong ability to put myself in the shoes of others and to understand their responses.	3.11	.77	1.6	34.4	.366
Item 12: I see the future with pessimism.	3.39	.68	1.7	49.2	.442
Item 13: The opinions of others have a strong influence on me when I have to make decisions.	2.91	.76	4.9	19.8	.422
Item 14: I see myself as less important than those around me.	3.44	.75	2.7	57.5	.500
Item 15: I am able to make decisions on my own.	3.44	.72	1.3	56.6	.520
Item 16: I try to look for the positive side when bad things happen to me.	2.90	.88	5.1	29.5	.497
Item 17: I try to improve myself as a person.	3.49	.69	0.7	59.2	.388
Item 18: I consider myself to be a good psychologist.	2.47	.86	12.6	12.5	.216
Item 19: It troubles me when people criticize me.	2.68	.89	13.0	16.4	.372
Item 20: I think that I am a sociable person.	3.29	.76	1.8	45.9	.412
Item 21: I am able to control myself when I have negative thoughts.	2.74	.80	3.8	18.9	.433
Item 22: I am able to maintain a high level of self-control in conflictive situations in my life.	2.75	.75	2.8	15.9	.504
Item 23: I feel that I am someone to be trusted.	3.67	.54	0.3	70.6	.332
Item 24: I find it particularly hard to understand the feelings of others.	3.53	.61	0.9	58.5	.267
Item 25: I consider the needs of others.	3.15	.72	0.9	34.3	.232
Item 26: When I experience unpleasant external pressure I am able to maintain my personal balance.	2.56	.77	7.1	10.6	.481
Item 27: When there are changes in my surroundings I try to adapt to them.	3.33	.69	0.4	46.0	.520
Item 28: In the face of a problem I am able to ask for information.	3.04	.80	2.6	31.9	.363
Item 29: I find changes in my daily routine to be stimulating.	2.86	.78	2.5	22.2	.334
Item 30: I find it hard to relate openly with my teachers/bosses.	3.20	.79	4.2	39.2	.336
Item 31: I feel inept and useless.	3.71	.54	0.3	75.1	.529
Item 32: I try to develop my abilities to the maximum.	3.27	.70	1.0	41.2	.503
Item 33: I find it hard to hold my own opinions.	3.59	.64	1.3	67.0	.395
Item 34: When I have to make big decisions I feel very unsure of myself.	2.75	.88	11.9	17.9	.497
Item 35: I am able to say no when I want to.	3.08	.90	4.1	40.5	.332
Item 36: When I am faced with a problem I try to find possible solutions.	3.48	.64	0.5	55.6	.538
Item 37: I like to help others.	3.67	.57	0.3	71.7	.206
Item 38: I feel unsatisfied with myself.	3.22	.78	4.5	39.7	.434
Item 39: I feel unsatisfied with the way I look.	3.06	.81	5.2	31.0	.409

Table 4

Cronbach's alpha coefficient and intraclass correlation coefficient test-retest

	Cronbach's α	Intraclass Correlation Coefficient	Confidence Interval 95%
Personal satisfaction	0.794	0.890	0.865-0.910
Prosocial attitude	0.543	0.760	0.707-0.804
Self-control	0.772	0.767	0.715-0.810
Autonomy	0.752	0.890	0.865-0.910
Problem-solving	0.778	0.874	0.846-0.897
Interpersonal Relationship Skills	0.636	0.840	0.804-0.869
TOTAL	0.890	0.925	0.907-0.939


Table 5.
Correlations among the subscales of the PMHQ Scale.

	Personal satisfaction	Prosocial attitude	Self-control	Autonomy	Problem solving	Interpersonal Relationship Skills
Personal satisfaction	1					
Prosocial attitude	0.15	1				
Self-control	0.45	0.18	1			
Autonomy	0.44	0.06	0.34	1		
Problem solving	0.46	0.34	0.50	0.39	1	
Interpersonal Relationship Skills	0.34	0.34	0.28	0.25	0.41	1
TOTAL Positive Mental Health	0.74	0.42	0.68	0.62	0.80	0.62

n = 1091; All correlation coefficients are significant at $P < 0.001$

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: Factor loadings derived from the LS estimation (least squares) Confirmatory Factor Analysis (λ_{ij})



99x133mm (300 x 300 DPI)