

WILEY

Online Proofing System Instructions

The Wiley Online Proofing System allows proof reviewers to review PDF proofs, mark corrections, respond to queries, upload replacement figures, and submit these changes directly from the locally saved PDF proof.

1. For the best experience reviewing your proof in the Wiley Online Proofing System ensure you are connected to the internet. This will allow the PDF proof to connect to the central Wiley Online Proofing System server. If you are connected to the Wiley Online Proofing System server you should see a green check mark icon above in the yellow banner.



Connected



Disconnected

2. Please review the article proof on the following pages and mark any corrections, changes, and query responses using the Annotation Tools outlined on the next 2 pages.



3. Save your proof corrections by clicking the “Publish Comments” button in the yellow banner above. Corrections don’t have to be marked in one sitting. You can publish comments and log back in at a later time to add and publish more comments before you click the “Complete Proof Review” button below.



4. If you need to supply additional or replacement files bigger than 5 Megabytes (MB) do not attach them directly to the PDF Proof, please click the “Upload Files” button to upload files:

5. When your proof review is complete and all corrections have been published to the server by clicking the “Publish Comments” button, please click the “Complete Proof Review” button below:

IMPORTANT: Did you reply to all queries listed on the Author Query Form appearing before your proof?

IMPORTANT: Did you click the “Publish Comments” button to save all your corrections? Any unpublished comments will be lost.


IMPORTANT: Once you click “Complete Proof Review” you will not be able to add or publish additional corrections.

USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION

Required software to e-Annotate PDFs: **Adobe Acrobat Professional** or **Adobe Reader** (version 11 or above). (Note that this document uses screenshots from **Adobe Reader DC**.)
 The latest version of Acrobat Reader can be downloaded for free at: <http://get.adobe.com/reader/>

Once you have Acrobat Reader open on your computer, click on the **Comment** tab (right-hand panel or under the Tools menu).

This will open up a ribbon panel at the top of the document. Using a tool will place a comment in the right-hand panel. The tools you will use for annotating your proof are shown below:



1. Replace (Ins) Tool – for replacing text.

 Strikes a line through text and opens up a text box where replacement text can be entered.

How to use it:

- Highlight a word or sentence.
- Click on .
- Type the replacement text into the blue box that appears.

Experimental data if available. For ORFs to be had to meet all of the following criteria:



2. Strikethrough (Del) Tool – for deleting text.

 Strikes a red line through text that is to be deleted.

How to use it:

- Highlight a word or sentence.
- Click on .
- The text will be struck out in red.

Experimental data if available. For ORFs to be had to meet all of the following criteria:

1. Small size (35–250 amino acids).
2. Absence of similarity to known proteins.
3. Absence of functional data which could not be the real overlapping gene.
4. Greater than 25% overlap at the N-terminus terminus with another coding feature; over both ends; or ORF containing a tRNA.

3. Commenting Tool – for highlighting a section to be changed to bold or italic or for general comments.

 Use these 2 tools to highlight the text where a comment is then made.

How to use it:

- Click on .
- Click and drag over the text you need to highlight for the comment you will add.
- Click on .
- Click close to the text you just highlighted.
- Type any instructions regarding the text to be altered into the box that appears.

informal invariance: [1] or A: Math. Gen., Vol. 12, N



lified theory for a matrix. 'ol. 8, 1984, pp. 305–323. d manuscript, 1984. ching fractions for D0 → K+K relation in D0 decays' Phys


4. Insert Tool – for inserting missing text at specific points in the text.

 Marks an insertion point in the text and opens up a text box where comments can be entered.

How to use it:

- Click on .
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the box that appears.

Meiosis has a central role in the sexual reproduction of nearly all eukaryotes. Saccharom: analysis of meiosis, esp by a simple change of n conveniently monitored cells. Sporulation of Sac cell, the a/a cell, and is of a fermentable carbon sporulation and are refe [2b]. Transcription of meiosis, in S. cerevisiae activator, IME1 (inducer of the gene RME1 funct Rme1p to exert repress of GAL1 gene expression) and HGR1 are required [1, 2, 3, 4]. These ge



5. Attach File Tool – for inserting large amounts of text or replacement figures.

 Inserts an icon linking to the attached file in the appropriate place in the text.

How to use it:

- Click on  .
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.

The attachment appears in the right-hand panel.

chondrial preparator
ative damage injury
re extent of membra
l, malondialdehyde (TBARS) formation.
used by high perform


6. Add stamp Tool – for approving a proof if no corrections are required.

 Inserts a selected stamp onto an appropriate place in the proof.

How to use it:

- Click on  .
- Select the stamp you want to use. (The **Approved** stamp is usually available directly in the menu that appears. Others are shown under *Dynamic, Sign Here, Standard Business*).
- Fill in any details and then click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).

of the business cycle, starting with the
on perfect competition, constant ret
roduction. In this environment goods
extra... market...
he...
etermined by the model. The New-Key
otaki (1987), has introduced produc
general equilibrium models with nomin
ed and...
Most of this...
APPROVED




7. Drawing Markups Tools – for drawing shapes, lines, and freeform annotations on proofs and commenting on these marks.


Allows shapes, lines, and freeform annotations to be drawn on proofs and for comments to be made on these marks.

How to use it:

- Click on one of the shapes in the **Drawing Markups** section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, right-click on shape and select **Open Pop-up Note**.
- Type any text in the red box that appears.



For further information on how to annotate proofs, click on the **Help** menu to reveal a list of further options:



Author Query Form

Journal: ANDR

Article: 12504

Dear Author,

During the copyediting of your manuscript the following queries arose.

Please refer to the query reference callout numbers in the page proofs and respond to each by marking the necessary comments using the PDF annotation tools.

Please remember illegible or unclear comments and corrections may delay publication.

Many thanks for your assistance.

Query reference	Query	Remarks
1	AUTHOR: Please check and approve the edits made in the article running title.	APROBADO Por 1001869 fecha 16:24 , 14/05/2018
2	AUTHOR: Please verify that the linked ORCID identifiers are correct for each author.	APROBADO Por 1001869 fecha 16:24 , 14/05/2018
3	AUTHOR: Please confirm that given names (red) and surnames/family names (green) have been identified correctly.	APROBADO Por 1001869 fecha 16:24 , 14/05/2018
4	AUTHOR: Please check that authors and their affiliations are correct.	APROBADO Por 1001869 fecha 16:25 , 14/05/2018
5	AUTHOR: Please give address information for 'Sigma-Aldrich': town, state (if applicable), and country.	CORREGIDO 16:25 , 14/05/2018
6	AUTHOR: Bradford, 1970 has been changed to Bradford, 1976 so that this citation matches the Reference List. Please confirm that this is correct.	CORREGIDO 16:25 , 14/05/2018
7	AUTHOR: Ramió-Lluch <i>et al.</i>, 2001 has been changed to Ramió-Lluch <i>et al.</i>, 2011 so that this citation matches the Reference List. Please confirm that this is correct.	APROBADO Por 1001869 fecha 16:26 , 14/05/2018
8	AUTHOR: Please suggest whether the term "olygomycin A" can be changed as "oligomycin A" in the sentence "In addition to this. . .of their lifespan."	CORREGIDO 16:26 , 14/05/2018
9	AUTHOR: Bromfield et al. (2014) has not been cited in the text. Please indicate where it should be cited; or delete from the Reference List.	CORREGIDO 16:27 , 14/05/2018

Funding Info Query Form

Please confirm that the funding sponsor list below was correctly extracted from your article: that it includes all funders and that the text has been matched to the correct FundRef Registry organization names. If a name was not found in the FundRef registry, it may not be the canonical name form, it may be a program name rather than an organization name, or it may be an organization not yet included in FundRef Registry. If you know of another name form or a parent organization name for a “not found” item on this list below, please share that information.

FundRef name	FundRef Organization Name
Ministry of Economy and Competitiveness, Spain	

spermatozoa has been clearly associated with circadian modulation in seasonal breeders, such as the sheep (Casao *et al.*, 2010), its relationship with the circadian rhythm in non-seasonal breeders, such as the pig, seems to be dismissed (González-Arto *et al.*, 2016).

Focusing on the effects upon sperm function, melatonin seems to have a vital regulatory role for sperm capacitation in the sheep and water buffalo (Casao *et al.*, 2009; Ashrafi *et al.*, 2013). Furthermore, melatonin has been shown to improve motility and other sperm functional parameters in human, ram, equine and boar spermatozoa (reviewed in Cebrián-Pérez *et al.*, 2014). The mechanism through which melatonin exerts these effects has been suggested to be linked with a reduction in oxidative stress by scavenging intracellular free radicals (Reiter *et al.*, 2000). In agreement with this hypothesis, *in vitro* treatment of spermatozoa with melatonin decreases intracellular levels of reactive oxygen (ROS) and nitrogen species (RNS; Rao & Gangadharan, 2008; Du Plessis *et al.*, 2010; Jang *et al.*, 2010; Najafi *et al.*, 2018), membrane lipid peroxidation (Gadella *et al.*, 2008; Du Plessis *et al.*, 2010; Da Silva *et al.*, 2011), apoptosis markers (Casao *et al.*, 2010; Da Silva *et al.*, 2011; Espino *et al.*, 2011; Najafi *et al.*, 2018) and DNA fragmentation (Sarabia *et al.*, 2009). In spite of all these data, little is known about whether melatonin could exert any effect on sperm function through a mechanism independent from its antioxidant properties. This is especially relevant for some species such as the pig, in which ROS levels produced by their spermatozoa in response to cryopreservation are marginal when compared to other species, such as the horse and cattle (Bilodeau *et al.*, 2000; Guthrie & Welch, 2006; Yeste *et al.*, 2013, 2015a,b). Thus, ROS production or accumulation seems to play a minor role to explain specific events, such as boar sperm cryodamage (Yeste *et al.*, 2013, 2015b). Likewise, changes in ROS levels are also low during the achievement of *in vitro* boar sperm capacitation (IVC), which again suggests they play a marginal role (Awda *et al.*, 2009). Taking all these data into account, we can hypothesise that melatonin could affect boar sperm function through mechanisms other than its ability to modulate intracellular ROS levels.

This study sought to determine the influence of melatonin on the achievement of IVC and subsequent progesterone-induced *in vitro* acrosome exocytosis (IVAE) of boar spermatozoa, as well as on their ability to adhere and further penetrate *in vitro* matured oocytes. With this purpose, two experiments were devised. In the first one, boar spermatozoa were subjected to IVC/IVAE in the presence of increasing concentrations of melatonin, added either before or after 4 h of incubation under capacitating conditions. Several parameters related to the achievement of IVC and IVAE were evaluated. In the second experiment, *in vitro* fertilisation (IVF) was conducted with spermatozoa previously capacitated with 1 μM melatonin. The sperm ability to bind the oocyte ZP and penetrate *in vitro* matured oocytes was assessed.

MATERIALS AND METHODS

Seminal samples

A total of 57 ejaculates collected from 32 healthy Pietrain boars aged between two and three years were used. These animals were housed in climate-controlled commercial farms (Servicios Genéticos Porcinos, S.L., Roda de Ter, Spain), fed with a commercial adjusted diet and provided with water *ad libitum*. Boar

housing fulfilled with the welfare standards established by European regulations on livestock species, specifically, on pig farms. Furthermore, and despite not being required as we did not manipulate any boar and only worked with seminal doses provided by the commercial farm, the experimental protocol was approved by the Ethics Committee of our institution (Bioethics Commission, Autonomous University of Barcelona; Bellaterra, Cerdanyola del Vallès, Spain). In all the cases, samples came from sperm-rich fractions that were obtained through manual collection with the conventional hand-gloved method. Upon collection, samples were immediately diluted with a commercial extender (Androstar Plus[®]; Minitub Ibérica SL, Tarragona, Spain) to a final sperm concentration of 2×10^7 spermatozoa/mL and cooled down to 16 °C. Diluted semen was packaged in 90-mL commercial AI doses and transported in an insulated container at 16 °C for approximately 45 min, which was the time required to arrive to our laboratory.

In vitro capacitation and progesterone-induced acrosome exocytosis

As aforementioned, two experiments were set. The first experiment was subdivided into two parts. The first one aimed at testing how melatonin affected the achievement of IVC (i.e. addition at 0 h). The second part sought to determine the impact of melatonin upon triggering IVAE in fully *in vitro* capacitated boar spermatozoa (i.e. addition at 4 h). With this purpose, different concentrations of melatonin were added after 4 h of incubation in CM, as this time period has previously been reported to induce IVC in boar spermatozoa (Ramió *et al.*, 2008). In this case, the addition of melatonin was performed together with that of progesterone, which was used to induce the acrosome reaction (Jimenez *et al.*, 2003; Wu *et al.*, 2006). Regardless of when melatonin was added (i.e. either at 0 h or at 4 h), five different treatments were assayed: a positive control (C+), which consisted of spermatozoa incubated in CM containing bicarbonate and bovine serum albumin (BSA); three treatments with different concentrations of melatonin (0.5, 1, 5 μM) in CM; and a negative control (C-), which consisted of spermatozoa incubated in non-capacitating medium without bicarbonate or BSA (NCM). As aforementioned, the tested concentrations of melatonin were within the physiological range of the genital tract, as described in previous works (Luboshitzky *et al.*, 2002; Casao *et al.*, 2009, 2010; Ashrafi *et al.*, 2013; Cebrián-Pérez *et al.*, 2014; Pérez-Patiño *et al.*, 2016).

For both parts, 50 mL of a given semen sample was split into five aliquots of 10 mL each. Aliquots were centrifuged at 600 g and 16 °C for 10 min and resuspended either with NCM (C-), CM (C+) or melatonin treatments (i.e. CM supplemented with melatonin at final concentrations of 0.5, 1 or 5 μM). In all cases, final sperm concentration was adjusted to 2×10^7 sperm/mL. The composition of NCM was 20 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (Hepes) buffer (pH = 7.4), 112 mM NaCl, 3.1 mM KCl, 5 mM glucose, 21.7 mM sodium L-lactate, 1 mM sodium pyruvate, 0.3 mM NaHPO₄, 0.4 mM MgSO₄ and 4.5 mM CaCl₂ (osmolarity: 287 mOsm/kg \pm 6 mOsm/kg). Capacitation medium (CM) consisted of NCM supplemented with 37.6 mM NaHCO₃ and 5 mg/mL BSA (pH adjusted to 7.4; osmolarity: 304 mOsm/kg \pm 5 mOsm/kg).

Aliquots were incubated at 38.5 °C and 5% CO₂ in humidified air for 4 h either with or without melatonin, as described in

Ramió *et al.* (2008). Samples were taken at 0 h and 4 h of incubation for analysis of sperm parameters. After 4 h of incubation, spermatozoa were subjected to progesterone-induced in vitro acrosome exocytosis (IVAE) through adding 10 mg/mL progesterone. After thoroughly mixing, sperm samples were incubated at 38.5 °C and 5% CO₂ in humidified air for a further hour. Separate aliquots were taken at 1, 5 and 60 min after the addition of progesterone. In the case of the second part, melatonin was directly added to the corresponding tube at final concentrations of 0.5, 1 or 5 µM after 4 h of starting the experiment rather than at 0 h.

At each relevant time point (i.e. 0, 4 h, and after 1, 5 and 60 min of progesterone addition), the achievement of both IVC and IVAE was evaluated on the basis of the following parameters: sperm motility, agglutination, viability, acrosome exocytosis, membrane lipid disorder and tyrosine phosphorylation of P32 protein, as a specific capacitation marker of boar spermatozoa (Bravo *et al.*, 2005). Furthermore, intracellular ROS levels, free cysteine residues in sperm head and tail extracts and lectin distribution over sperm membrane were also evaluated as parameters that could be related to the effects of melatonin on IVC/IVAE.

Unless stated otherwise, all fluorochromes and lectins were purchased from Molecular Probes® (Invitrogen, ThermoFisher; Eugene, OR, USA) and diluted with dimethyl sulfoxide (DMSO; Sigma-Aldrich); Saint Louis, MO, USA

CORREGIDO
16:16, 14/05/2018

Evaluation of sperm motility and agglutination

Sperm motility and agglutination were evaluated by utilising a commercial, computer-assisted sperm analysis (CASA) system (Integrated Sperm Analysis System V1.0; Proiser, Valencia, Spain). This system is based on the analysis of 25 consecutive digitalised photographic images obtained from a single field at a magnification of 100× (Olympus BX41 microscope; Olympus 10 × 0.30 PLAN objective lens, negative phase contrast; Olympus Europa GmbH, Hamburg, Germany). These 25 consecutive photographs were taken at a time lapse of 1 sec, which implies that an image was captured every 40 ms. Five to six separate fields were taken for each replicate, and five replicates were evaluated per sample and treatment. Prior to evaluation with CASA, a 5 µL drop was placed onto a warmed Makler chamber (Sefi Medical Instruments, Haifa, Israel). In the case of samples evaluated at 0 h, they were previously warmed at 38 °C for 15 min in a water bath. Recorded sperm motility and kinematic parameters were those described in Ramió *et al.* (2008). Settings for the CASA system were as follows: area of particles: 10–80 µm²; curvilinear velocity (VCL): 1–500 µm/sec; mean velocity (VAP): 1–500 µm/sec; linearity coefficient (LIN): 10–98%; straightness coefficient (STR): 10–98%; mean amplitude of lateral head displacement (ALH): 0–100 µm; and beat cross-frequency (BCF): 0–100 Hz. A spermatozoon was considered to be motile when its VAP was higher than 10 µm/sec.

Furthermore, the number of spermatozoa included in agglutination complexes and the percentage of agglutinated spermatozoa that showed apparent tail movement were determined by evaluation of each consecutive photograph obtained from CASA analyses. Specifically, we determined the number of sperm heads agglutinated divided by the total number of sperm heads,

and the number of beating tails observed in each agglutination complex divided by the total number of tails.

Flow cytometry analyses

Sperm viability, acrosome exocytosis, membrane lipid disorder and intracellular ROS levels were evaluated by flow cytometry. Information about flow cytometry analyses is given according to the recommendations of the International Society for Advancement of Cytometry (ISAC; Lee *et al.*, 2008). Prior to evaluation, sperm concentration was adjusted to 1 × 10⁶ spermatozoa/mL in a volume of 0.5 mL. Thereafter, spermatozoa were stained with the appropriate combinations of fluorochromes, following the protocols described below.

Samples were evaluated through a Cell Laboratory QuantaSC cytometre (Beckman Coulter, Fullerton, CA, USA) and were excited using single-line visible light from an argon laser (wavelength: 488 nm; power: 22 mW). Sheath flow rate was set at 4.17 µL/min, and electronic volume (EV; equivalent to forward scatter) and side scatter (SS) were recorded for each event. Calibration of the equipment was periodically performed using 10-µm Flow-Check fluorospheres (Beckman Coulter). Three optical filters with the following characteristics were used: FL1 (green fluorescence): Dichroic/Splitter, DRLP: 550 nm, band-pass filter: 525 nm, detection width 505–545 nm; FL2 (orange fluorescence): DRLP: 600 nm, BP filter: 575 nm, detection width: 560–590 nm; and FL3 (red fluorescence): long pass filter: 670 nm. Signals were logarithmically amplified, and photomultiplier (PMT) settings were adjusted to each particular staining method; compensation was used to minimise spillover of the fluorescence into a different channel. The analyser threshold was adjusted on the EV channel to exclude subcellular debris (particle diameter < 7 µm) and cell aggregates (particle diameter > 12 µm) and sperm-specific events were positively gated on the basis of EV/SS distributions. Three independent replicates were examined, and 10,000 events were evaluated per replicate. Information on the events was collected as list-mode data files (LMD), and data were analysed through Cell Lab QuantaSC MPL Analysis Software (version 1.0; Beckman Coulter).

In all assessments except SYBR14/PI, data obtained from flow cytometry experiments were corrected according to the procedure described in Petrunkina *et al.* (2010).

Evaluation of sperm viability

Sperm viability was assessed using the LIVE/DEAD® Sperm Viability Kit (SYBR14/PI) following the protocol set in Garner & Johnson (1995). With this purpose, sperm samples were incubated with SYBR14 (final concentration = 100 nM) at 38 °C for 10 min and then with propidium iodide (PI; final concentration = 10 µM) at the same temperature for 5 min. Fluorescence emitted by SYBR14 was measured through FL1, whereas that emitted by PI was detected through FL3. Three sperm populations were identified as follows: (i) viable green-stained spermatozoa (SYBR14⁺/PI⁻); (ii) non-viable, red-stained spermatozoa (SYBR14⁻/PI⁺); and (iii) non-viable spermatozoa that were stained both in green and in red (SYBR14⁺/PI⁺). Non-sperm particles (debris) were found in the SYBR14⁻/PI⁻ quadrant. Single-stained samples were used for setting PMT voltages of EV, FL1

and FL3, and for compensation of SYBR14 spillover into the FL3 channel (2.45%).

Evaluation of acrosome exocytosis

True acrosome exocytosis was determined through costaining of spermatozoa with *Arachis hypogaea* agglutinin (PNA) conjugated with fluorescein isothiocyanate (FITC-PNA) and ethidium homodimer (3,8-diamino-5-ethyl-6-phenylphenanthridinium bromide; EthD-1). This protocol was originally described by Cooper & Yeung (1998) and has been adapted to boar spermatozoa in our laboratory. In brief, samples were incubated with EthD-1 (final concentration: 2.5 $\mu\text{g}/\text{mL}$) at 38 °C for 5 min in the dark. Following this step, samples were centrifuged at 2000 *g* for 30 sec and then resuspended with PBS containing 4 mg/mL bovine serum albumin (BSA) to remove the free dye. Thereafter, samples were again centrifuged at the aforementioned conditions and then fixed and permeabilised by adding 100 μL of ice-cold methanol (100%) for 30 sec. Methanol was removed by centrifugation at 2000 *g* for 30 sec, and pellets were resuspended with 250 μL PBS. Following this step, 0.8 μL PNA-FITC (final concentration: 2.5 μM) was added, and samples were incubated at 15 °C in the dark for 15 min. Next, samples were washed twice with PBS at 2000 *g* for 30 sec and finally resuspended in PBS.

Following staining, samples were evaluated with the flow cytometry and the following four sperm populations were identified: (i) viable spermatozoa with an intact acrosome membrane (PNA-FITC⁺/EthD-1⁻); (ii) viable spermatozoa with a non-intact acrosome membrane (PNA-FITC⁻/EthD-1⁻); (iii) non-viable spermatozoa with an intact acrosome membrane (PNA-FITC⁺/EthD-1⁺); and (iv) non-viable spermatozoa with a non-intact acrosome membrane (PNA-FITC⁻/EthD-1⁺). Fluorescence of EthD-1 was detected through FL3, whereas that of PNA-FITC was detected through FL1.

Evaluation of sperm membrane lipid disorder

Lipid disorder of boar sperm membrane was evaluated by costaining with merocyanine-540 (M540) and YO-PRO-1, following a procedure modified from Rathi *et al.* (2001). Briefly, spermatozoa were stained with M540 (final concentration: 2.6 μM) and YO-PRO-1 (final concentration: 25 nM) and incubated at 38 °C for 10 min in the dark. Red fluorescence from M540 was collected through FL3, and green fluorescence from YO-PRO-1 was collected through FL1. The following four sperm populations were distinguished: (i) viable spermatozoa with low membrane lipid disorder (M540⁻/YO-PRO-1⁻); (ii) viable spermatozoa with high membrane lipid disorder (M540⁺/YO-PRO-1⁻); (iii) non-viable spermatozoa with low membrane lipid disorder (M540⁻/YO-PRO-1⁺); and (iv) non-viable spermatozoa with high membrane lipid disorder (M540⁺/YO-PRO-1⁺). In this test, data were not compensated.

Evaluation of intracellular levels of superoxides and peroxides

Intracellular superoxide ($\text{O}_2^{\bullet-}$) and peroxide (H_2O_2) levels were determined using two different oxidation-sensitive fluorescent probes: hydroethidine (HE) and 2',7'-dichlorodihydrofluorescein diacetate (H_2DCFDA). Following a procedure modified from Guthrie & Welch (2006), a simultaneous differentiation of viable from non-viable spermatozoa was performed using either YO-PRO-1 or PI.

In the case of superoxides, samples were stained with HE (final concentration: 4 μM) and YO-PRO-1 (final concentration: 25 nM) and incubated at 15 °C for 40 min in the dark [17]. Hydroethidine is freely permeable to cells and is oxidised by $\text{O}_2^{\bullet-}$ to ethidium (E^+) and other products. Fluorescence of ethidium (E^+) was detected through FL3, and that of YO-PRO-1 was collected through FL1. Viable spermatozoa with high intracellular superoxide levels were positive for ethidium and negative for YO-PRO-1 ($\text{E}^+/\text{YO-PRO-1}^-$).

With regard to peroxides, spermatozoa were stained with H_2DCFDA at a final concentration of 200 μM and PI at a final concentration of 10 μM , and incubated at 15 °C for 60 min in the dark. H_2DCFDA is a cell-permeable, non-fluorescent probe that is intracellularly de-esterified and converted into highly fluorescent, 2',7'-dichlorofluorescein (DCF^+) upon oxidation (Guthrie & Welch, 2006). This DCF^+ fluorescence was collected through FL1, whereas PI fluorescence was detected through FL3. Data were not compensated, and viable spermatozoa with high intracellular peroxide levels were positive for DCF and negative for PI (DCF^+/PI^-).

In both cases, unstained and single-stained samples were used for setting EV, FL1 and FL3 PMT voltages and data were not compensated.

Immunoblotting

Aliquots of 1 mL corresponding to each experimental point were centrifuged at 1000 *g* and 15 °C for 30 sec, and pellets were stored at -80 °C until the beginning of the assay. Pellets were resuspended and sonicated in 300 μL ice-cold lysis buffer containing 50 mM Tris-HCl, 1 mM EDTA, 10 mM EGTA, 25 mM dithiothreitol, 1.5% (v:v) Triton[®] X-100, 1 mM PMSF, 10 $\mu\text{g}/\text{mL}$ leupeptin, 1 mM orthovanadate and 1 mM benzamidine (pH = 7.4). After 30 min on ice, the homogenised suspensions were centrifuged at 600 *g* and 4 °C for 20 min, and total protein content in supernatants was calculated through the Bradford method (Bradford, 1976) using **APROBADO** kit (Bio-Rad Laboratories). Afterwards, samples **Por 1001869 fecha 16:14, 14/05/2018** (1 : 5; v:v) containing 250 mM Tris-HCl (pH = 6.8), 50 mM dithiothreitol, 10% (w:v) SDS, 0.5% (v:v) bromophenol blue and 50% (v:v) glycerol and stored at -20 °C until gel electrophoresis (SDS-PAGE; Laemmli, 1970).

Prepared samples were loaded onto 0.75-mm gels containing 10% acrylamide (w:v). After running the gels at constant voltage (180 V), proteins were transferred onto Immun-Blot[®] low-fluorescence polyvinylidene fluoride (PVDF) membranes (Bio-Rad) using the Trans-Blot[®] Turbo Transfer System with Trans-Blot[®] Turbo Midi Transfer Packs (Bio-Rad). Membranes were subsequently immersed for 60 min into blocking solution, consisting of Tris-buffered saline solution added with 5% (w:v) BSA and 0.1% (v:v) Tween-20. Thereafter, membranes were incubated with a mouse monoclonal PY20 antiphosphotyrosine antibody (ref. P4110; Sigma-Aldrich; dilution factor: 1 : 1000 (v:v) in blocking solution) at 4 °C for 8 h. Membranes were washed three times with blocking solution (5 min per wash) and then incubated at 15 °C for 1 h with a horseradish peroxidase (HRP)-conjugated, polyclonal rabbit anti-mouse antibody (Dako; Glostrup, Denmark) at a dilution of 1 : 5000 (v:v) in blocking solution. After washing membranes with blocking solution for six times (5 min per wash), membranes were incubated with chemiluminescent HRP substrate (ImmunoCruz Western Blotting Luminol

Reagent; Santa Cruz Biotechnology®, Dallas, Texas, USA) at 15 °C for 2 min, following manufacturer's instructions. Revealed images were analysed through IMAGEJ ver. 1.49 (National Institute of Health, Bethesda, MD, USA), and the intensity/densitometry of each band was quantified. Following this, membranes were stripped and then incubated with a mouse monoclonal anti- β -tubulin (ref. T5201; Sigma-Aldrich; 1 : 1000 (v:v) in blocking solution) and the same secondary HRP antibody. Images were also analysed through ImageJ. A total of seven semen samples were used for Western blot assays.

Evaluation of free cysteine residues in spermatozoa and tail extracts

Determination of free cysteine radicals in sperm head and tail extracts as an indirect measure of disrupted disulphide bridges within proteins was carried out following the protocol described in Flores *et al.* (2011). Briefly, samples were centrifuged at 600 g and 16 °C for 10 min and then resuspended in an ice-cold lysis buffer made up as follows: 50 mM Tris buffer, 150 mM NaCl, 1% (v:v) Nonidet, 0.5% (w:v) sodium deoxycholate, 1 mM benzamide, 10 μ g/mL leupeptin, 0.5 mM phenylmethylsulfonyl fluoride (PMSF) and 1 mM Na₂VO₄ (pH adjusted to 7.4). Samples were homogenised through sonication (12 pulses; Ikasonic U50 sonicator; Ika Labor Technik, Staufen, Germany), and obtained homogenates were centrifuged at 850 g and 4 °C for 20 min. After this centrifugation step, the supernatant mainly contained the sperm tails, whereas the pellet mainly contained the sperm heads. This was confirmed by separate evaluations through phase-contrast microscopy of both fractions, the pellets being previously resuspended with 300 μ L Tris buffer. Indeed, percentages of tails in supernatants and heads in pellets were found to be above 90%, respectively (data not shown).

Levels of free cysteine radicals in both fractions (i.e. supernatants and Tris-solubilised pellets) were determined using the 2,2'-dithiodipyridine technique (2,2'-dipyridyl disulphide; Sigma-Aldrich) as described in Brocklehurst *et al.* (1979). Briefly, 10 μ L of the supernatant or resuspended pellet was added with 990 μ L of an aqueous solution containing 0.4 mM 2,2'-dithiodipyridine. Standard curves were generated with 10- μ L aliquots containing different concentrations of cysteine (Sigma-Aldrich; from 0.1 to 5 mM), which were also added with 990 μ L of 0.4 mM 2,2'-dithiodipyridine. Samples were incubated at 37 °C for 60 min, and levels of free cysteine radicals were subsequently determined through spectrophotometry at a wavelength of 340 nm. The results obtained were normalised through a parallel determination of the total protein content by the Bradford method (Bradford, 1976), using a commercial kit (Quick Start Bradford Protein Assay; Bio-Rad, Hercules, CA, USA). Five replicates per sample and treatment were evaluated, and the corresponding mean \pm standard error of the mean (SEM) was calculated.

Distribution of lectins over sperm membrane

Four FITC-conjugated lectins were used as follows: *Triticum vulgare* agglutinin (WGA), *Solanum lycopersicum* lectin (STL), *Pisum Sativum* agglutinin (PSA) and *Arachis Hypogaea* agglutinin (PNA). Semen samples were centrifuged at 1000 g and 15 °C for 30 sec, and the resultant pellets were resuspended with 400 μ L PBS containing 4% (w:v) paraformaldehyde. Fixation was conducted at 4 °C in the dark for 2 h. Samples were

subsequently spread onto poly-lysine (1% poly-lysine solution in water; Sigma-Aldrich)-coated microscope slides and then left to dry. Samples were permeabilised by incubation with 0.3% (v:v) Triton® X-100 in PBS (pH = 7.4) at 15 °C for 10 min. Next, slides were washed three times with PBS and then blocked through incubation with PBS containing 0.1% (v:v) Tween-20 and 5% (w:v) BSA at 15 °C for 60 min. After blocking, samples were incubated at 15 °C in a humid chamber for 1 h with the corresponding lectin at the following dilutions in PBS: 1 : 200 (w:v) for WGA and PSA, 1 : 300 (w:v) for PNA and 1 : 50 (w:v) for STL. Slides were further washed three times with PBS (5 min each wash) and then mounted with antifading medium Vectashield H-1000 (Vector Laboratories, Burlingame CA, USA). After being covered by coverslips, slides were compressed to eliminate any excess of liquid. Coverslips were finally sealed with colourless nail polish, and slides were stored at 4 °C in the dark until observation. Negative control experiments were performed omitting the lectin.

Samples were observed using a confocal laser scanning microscope (Leica TCS 4D; Leica Lasertechnik, Heidelberg, Germany) at 63 \times magnification. The light source was an argon/krypton laser. Successive confocal slices of images (image thickness: 0.5 μ m) were integrated to create three-dimensional images that were saved in TIFF format. Each lectin generated distinct staining patterns that were examined in non-capacitated, capacitated and acrosome-exocytosed spermatozoa.

In vitro oocyte–sperm co-incubation and evaluation of sperm adhesiveness and penetration ability

As previously mentioned, the current work was divided into two experiments. In the second experiment, and following the results obtained in the first one, one melatonin treatment (1 μ M) was compared with the control. The sperm ability to bind the ZP and to penetrate in vitro matured oocytes was evaluated after previous incubation with 1 μ M melatonin, following a modified protocol from Castillo-Martín *et al.* (2014).

Ovaries were obtained from a local slaughterhouse and were brought to the laboratory in a 0.9% (w:v) NaCl solution containing 100 μ g kanamycin sulphate per mL previously warmed at 37 °C. Oocyte–cumulus cell complexes (COCs) were collected from follicles of 3–6 mm diameter and only those showing at least two layers of cumulus cells and a homogeneous cytoplasm were selected. COCs were washed twice with DPBS supplemented with 4 mg/mL polyvinyl alcohol (PVA) and then with maturation medium, previously equilibrated at 38.5 °C and 5% CO₂ in humidified air for at least 3 h. Groups of 50 oocytes were cultured in 500 μ L maturation medium for 22 h at 38.5 °C and 5% CO₂ in humidified air. Thereafter, oocytes were transferred to fresh maturation medium without hormones or dibutyl cAMP, and cultured for further 22 h. The maturation medium was NCSU-37 (Petters & Wells, 1993) supplemented with 0.57 mM cysteine, 1 mM dibutyl cAMP, 5 μ g/mL insulin, 50 μ M β -mercaptoethanol, 10 IU/mL equine chorionic gonadotrophin (Folligon, Intervet International BV, Boxmeer), 10 IU/mL human chorionic gonadotrophin (Veterin Corion, Divasa Farmavic, Barcelona, Spain) and 10% (v:v) pig follicular fluid.

After maturation, oocytes were mechanically stripped of cumulus cells by gentle aspiration with a pipette. Denuded oocytes were washed with TALP medium, and groups of 25

oocytes were then transferred to each well of four-well Nunc multidishes (Nunc; Roskilde, Denmark) containing 250 μ L TALP medium, previously equilibrated at 38.5 °C under 5% CO₂ in humidified air. The composition of TALP medium was as follows: 114.06 mM NaCl, 3.2 mM KCl, 8 mM calcium lactate·5H₂O, 0.5 mM MgCl₂·6H₂O, 0.35 mM NaH₂PO₄, 25.07 mM NaHCO₃, 10 mL/L sodium lactate, 5 mM glucose, 2 mM caffeine, 1 g/L PVA, and 0.17 mM kanamycin sulphate supplemented with 3 mg/mL fatty acid-free BSA (FAF-BSA) and 1.1 mM sodium pyruvate (Rath *et al.*, 1999).

Two hundred fifty microlitres of sperm suspensions from each treatment group was added to the fertilisation wells at a final concentration of 5×10^4 sperm/mL. Those spermatozoa had previously been incubated in CM at 38.5 °C and 5% CO₂ in humidified air for 4 h, either in the presence or absence of 1 μ M melatonin. Specifically, three treatments were set as follows: (i) control, which consisted of oocytes incubated with spermatozoa previously incubated in CM without melatonin; (ii) experimental treatment 1, which consisted of oocytes co-incubated with spermatozoa previously in vitro capacitated with CM added with 1 μ M melatonin at 38.5 °C and 5% CO₂ in humidified air for 4 h; and (iii) experimental treatment 2, which consisted of oocytes added with both spermatozoa (previously incubated in CM without melatonin) and melatonin to a final concentration of 1 μ M. In all treatments, co-incubation of spermatozoa with in vitro matured oocytes was performed at 38.5 °C and 5% CO₂ in humidified air for 1 h. Free, non-attached spermatozoa were removed by washing oocyte–sperm complexes with TALP medium, and 500 μ L fresh TALP medium was subsequently added. Oocyte–sperm complexes were incubated at 38.5 °C and 5% CO₂ in humidified air for further 7 h and subsequently prepared for nuclear staining. Following this, oocytes were gently aspirated with a pipette, washed with TALP medium and subsequently transferred to a new well containing 500 μ L TALP. Oocyte–sperm complexes were maintained in this medium at 38.5 °C and 5% CO₂ in humidified air for further 7 h and then collected to perform the following nuclear staining protocol.

Oocyte–sperm complexes were washed with warmed PBS and then fixed with 4% (w:v) paraformaldehyde in PBS at 38.5 °C for 30 min. After fixation, complexes were washed twice with PBS and subsequently stained with 1% (v:v) Hoechst[®] 33342 in PBS at 15 °C for 25 min. Oocyte–sperm complexes were then washed two times with PBS, mounted on glass slides and examined under a TCS 4D laser confocal scanning microscope (Leica Lasertechnik) at 63 \times magnification. The following parameters were evaluated: (i) spermatozoa bound to the ZP: number of nuclear spermatozoa attached to the ZP; (ii) total penetration rate: number of oocytes that showed evident signs of sperm penetration divided by the total number of sperm–oocytes complexes; lack of penetration consisted of sperm–oocytes complexes that showed a unique nucleus with or without apparent polar bodies; (iii) percentage of monospermy: number of oocytes showing the presence of two pronuclei, or one sperm head inside the oocyte with or without signs of decondensation, divided by the total number of sperm–oocytes complexes; and (iv): percentage of polyspermy: number of oocytes showing more than two pronuclei, or more than one sperm head inside the oocyte with or without signs of decondensation, divided by the total number of sperm–oocytes complexes.

Statistical analyses

Statistical analyses were performed using a statistical package (IBM SPSS for Windows version 21.0, IBM Corp; Chicago, IL, USA). Data are presented as mean \pm standard error of the mean (SEM), and the level of significance was set at $p \leq 0.05$.

In the case of experiment 1, data were first tested for normality and homogeneity of variances through Shapiro–Wilk and Levene tests, respectively. When required, data (x) were transformed through arcsine square root ($\arcsin\sqrt{x}$) before a general mixed model (i.e. with repeated measures) was run. In this model, the intersubject factor was the treatment (i.e. composition of capacitation media), and the intrasubject factor was the incubation time (i.e. 0 h, 4 h, 4 h + 1 min, 4 h + 5 min, 4 h + 60 min). In all cases, each sperm functional parameter was the dependent variable, and pairwise comparisons were made with Sidak post hoc test. When no transformation remedied the normality, nonparametric procedures were conducted with raw data. Friedman's test and the Wilcoxon matched-pairs test were performed as nonparametric alternatives to repeated measures ANOVA.

With regard to experiment 2, the number of spermatozoa attached to oocyte ZP was checked for normality and homogeneity of variances as previously described, and compared through one-way ANOVA followed by post hoc Sidak's test. For the analysis of monospermy/polyspermy, a chi-square test (χ^2) was used.

RESULTS

Effects of melatonin on viability, acrosome exocytosis and capacitation-like changes in sperm membrane

As shown in Figure S1A, incubation of boar spermatozoa in CM for 4 h reduced their viability, which went from 80.4% \pm 3.7% at 0 h to 67.9% \pm 2.8% after 4 h of incubation. This decline was maintained after the addition of progesterone. While the addition of melatonin to CM at 0 h or 4 h did not significantly modify the observed drop in sperm viability, the extent of that decrease was higher when spermatozoa were incubated in NCM (Figure S1A,B).

Percentages of true acrosome exocytosis (PNA-FITC⁻/EthD-1⁻) were very low in cells incubated in CM during 4 h. The addition of progesterone at 4 h induced an increase in this percentage, which reached maximal values after 60 min of that addition (67.4% \pm 2.3%; Figure S2A,B). This increase was not observed when spermatozoa were incubated in NCM. The addition of melatonin either at 0 h or at 4 h did not modify the pattern observed in spermatozoa incubated in CM (Figure S2A,B).

Incubation of boar spermatozoa in CM significantly ($p < 0.05$) increased the percentage of viable spermatozoa with high membrane lipid disorder (from 9.4% \pm 2.6% at 0 h to 45.7% \pm 4.6% at 4 h; Figure S3A,B). The subsequent addition of progesterone was associated with a progressive decrease in this percentage, which reached values of 34.6% \pm 3.0% after 60 min of progesterone addition. The addition of melatonin either at 0 h or at 4 h did not change the dynamics observed in spermatozoa incubated in CM (i.e. positive control; Figure S3A,B).

Effects of melatonin on P32 tyrosine phosphorylation levels

As expected, incubation of boar spermatozoa in CM for 4 h induced a noticeable increase in tyrosine phosphorylation (pTyr)


levels of P32 protein (from 100.0 arbitrary units at 0 h of incubation to 231.7 ± 14.3 arbitrary units after 4 h), which was roughly maintained after progesterone addition (Figures S4 and S5). Addition of melatonin at 0 h did not significantly modify that pattern (Figure S4). The addition of progesterone after 4 h of incubation in CM did not have any prominent effect on pTyr-P32 levels. Only the treatment containing melatonin at $5 \mu\text{M}$ showed a slight decrease in pTyr-P32 values when compared to incubation in CM (60 min after progesterone addition; melatonin at $5 \mu\text{M}$: 203.5 ± 7.4 arbitrary units vs. CM: 228.2 ± 7.6 arbitrary units; see Figure S5). Finally, the addition of $5 \mu\text{M}$ melatonin at 4 h was found to decrease the intensity of tyrosine phosphorylation in P32 band after 5 min and 60 min of progesterone addition (60 min after progesterone addition; melatonin at $5 \mu\text{M}$: 206.8 ± 6.5 arbitrary units vs. CM: 226.1 ± 6.9 arbitrary units; see Figure S5).

Effects of melatonin on sperm motility

Total motility of spermatozoa incubated in CM significantly ($p < 0.05$) decreased throughout the experiment, reaching minimal values of $27.0\% \pm 2.5\%$ after 60 min of the addition of progesterone (Fig. 1). Incubation of spermatozoa in NCM led to even worse motility values, with complete immobilisation at the end of the experiment. The addition of melatonin at 0 h induced an immediate decrease in total motility, which was more apparent at the highest melatonin concentration (0 h: $47.2\% \pm 3.0\%$ in melatonin at $5 \mu\text{M}$ vs. $64.0\% \pm 3.9\%$ in CM; Fig. 1A). This adverse effect on sperm motility was observed throughout all the incubation period. Melatonin also decreased sperm motility when added together with progesterone at 4 h, the treatments containing melatonin at 1 and $5 \mu\text{M}$ showing values near to complete immobility after 60 min of progesterone addition (Fig. 1B).

Regarding kinetic parameters, spermatozoa incubated in CM for 4 h showed significant ($p < 0.05$) increases in several parameters, including VCL, VAP and ALH (as an example, VAP at 0 h of incubation in CM: $65.7 \mu\text{m}/\text{sec} \pm 2.1 \mu\text{m}/\text{sec}$ vs. VAP after 4 h of incubation in CM: $72.9 \mu\text{m}/\text{sec} \pm 2.9 \mu\text{m}/\text{sec}$; Table 1). However, the addition of melatonin at 0 h significantly ($p < 0.05$) decreased VAP values (after 4 h of incubation; $57.9 \mu\text{m}/\text{sec} \pm 1.7 \mu\text{m}/\text{sec}$ in the treatment containing $0.5 \mu\text{M}$ melatonin vs. $72.9 \mu\text{m}/\text{sec} \pm 2.9 \mu\text{m}/\text{sec}$ in CM; Table 1). When melatonin was added together with progesterone at 4 h, there was an immediate decrease in VAP (1 min after progesterone addition: $40.9 \mu\text{m}/\text{sec} \pm 2.0 \mu\text{m}/\text{sec}$ in the treatment containing $1 \mu\text{M}$ melatonin vs. $59.8 \mu\text{m}/\text{sec} \pm 2.7 \mu\text{m}/\text{sec}$ in CM; Table 1), LIN (1 min after progesterone addition: $25.7\% \pm 1.6\%$ in the treatment containing $1 \mu\text{M}$ melatonin vs. $39.9\% \pm 1.7\%$ in CM; Table 2) and STR (1 min after progesterone addition: $58.1\% \pm 1.8\%$ in the treatment containing $1 \mu\text{M}$ melatonin vs. $67.7\% \pm 2.8\%$ in CM; Table 2). Melatonin-induced decreases of both VAP and STR were recovered after 5 min and 60 min of the addition of progesterone and melatonin at 0.5 and $1 \mu\text{M}$ (VAP after 60 min of the addition of progesterone and melatonin at $1 \mu\text{M}$: $50.1 \mu\text{m}/\text{sec} \pm 3.1 \mu\text{m}/\text{sec}$ vs. $56.8 \mu\text{m}/\text{sec} \pm 3.3 \mu\text{m}/\text{sec}$ in CM; STR after 60 min of the addition of progesterone and melatonin at $1 \mu\text{M}$: $77.1\% \pm 4.4\%$ vs. $76.2\% \pm 3.3\%$ in CM; Tables 1 and 2). In contrast, LIN was only recovered after 60 min of the addition of progesterone and melatonin at the same concentrations (LIN after 60 min of the addition of progesterone and melatonin at $1 \mu\text{M}$: $32.7\% \pm 1.5\%$ vs. $37.1\% \pm 2.0\%$ in CM; Table 2).

Figure 1 Effects of melatonin on total motility of boar spermatozoa subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis. (A): Melatonin added at 0 h. (B): Melatonin added together with progesterone at 4 h. White bars: spermatozoa incubated in NCM medium (C−). Light grey bars: spermatozoa incubated in CM medium (C+). Medium grey bars: spermatozoa incubated in CM added with $0.5 \mu\text{M}$ melatonin. Dark green bars: spermatozoa incubated in CM added with $1 \mu\text{M}$ melatonin. Black bars: spermatozoa incubated in CM added with $5 \mu\text{M}$ melatonin. Asterisks indicate significant ($p < 0.05$) differences between a given treatment and C+ samples. Figure shows means \pm SEM for seven separate experiments.



Effects of melatonin on sperm agglutination

Incubation of spermatozoa in CM increased their degree of agglutination, which was $60.9\% \pm 7.5\%$ at 4 h (Fig. 2). Agglutinations were of medium size (Figure S6D,E), and about 45% of agglutinated spermatozoa showed appreciable tail beating at 4 h of incubation (Fig. 3). Although the percentage of agglutinated spermatozoa did not vary after the addition of progesterone (Fig. 2, Figure S6F), the percentage of agglutinated spermatozoa with appreciable tail beating showed a transient increase upon progesterone addition and then started to decrease, reaching a value of $17.3\% \pm 2.6\%$ at the end of the experiment (Fig. 3). In contrast to the aforementioned, spermatozoa incubated in NCM did show low percentages of agglutinated spermatozoa (Fig. 2, Figure S6A–C). The addition of melatonin at 0 h induced an immediate and

Table 1 Effects of melatonin on curvilinear velocity (VCL) and average path velocity (VAP) of boar spermatozoa subjected to in vitro capacitation and subsequent, progesterone-induced in vitro acrosome exocytosis

Incubation time	0 h	4 h	1 min	5 min	60 min
VCL ($\mu\text{m}/\text{sec}$)					
C-	77.0 \pm 2.9 ^{a*}	39.0 \pm 1.1 ^{b*}	39.1 \pm 0.9 ^{b*}	38.5 \pm 1.2 ^{c*}	0 ^{d*}
C+	65.7 \pm 2.1 ^a	72.9 \pm 2.9 ^b	74.1 \pm 3.2 ^b	81.4 \pm 3.7 ^b	81.6 \pm 4.3 ^b
0.5 μM melatonin	67.5 \pm 2.3 ^a	55.6 \pm 1.5 ^{b*}	57.9 \pm 1.7 ^{b*}	57.0 \pm 2.2 ^{b*}	44.5 \pm 1.2 ^{c*}
1 μM melatonin	56.2 \pm 1.8 ^{a*}	60.7 \pm 2.5 ^{a*}	59.1 \pm 2.6 ^{a*}	61.2 \pm 3.5 ^{a*}	41.2 \pm 2.8 ^{b*}
5 μM melatonin	62.5 \pm 2.5 ^a	0 ^{b*}	0 ^{b*}	0 ^{b*}	0 ^{b*}
0.5 μM melatonin+PG	65.7 \pm 2.1 ^a	72.9 \pm 2.9 ^b	68.2 \pm 2.7 ^a	90.8 \pm 4.7 ^c	76.5 \pm 3.9 ^b
1 μM melatonin+PG	65.7 \pm 2.1 ^a	72.9 \pm 2.9 ^b	67.7 \pm 2.6 ^a	89.9 \pm 4.5 ^c	76.8 \pm 4.2 ^b
5 μM melatonin+PG	65.7 \pm 2.1 ^a	72.9 \pm 2.9 ^b	71.4 \pm 3.1 ^{ab}	81.1 \pm 4.3 ^c	66.8 \pm 4.0 ^{a*}
VAP ($\mu\text{m}/\text{sec}$)					
C-	35.5 \pm 1.4 ^{a*}	25.1 \pm 1.1 ^{b*}	27.5 \pm 1.6 ^{b*}	33.6 \pm 0.9 ^{b*}	0 ^{c*}
C+	46.0 \pm 2.4 ^a	62.6 \pm 3.2 ^b	59.8 \pm 2.7 ^b	56.7 \pm 2.8 ^b	56.8 \pm 3.3 ^b
0.5 μM melatonin	48.1 \pm 2.0 ^a	44.6 \pm 2.2 ^{a*}	47.0 \pm 2.7 ^{a*}	62.6 \pm 3.7 ^b	85.6 \pm 5.4 ^{c*}
1 μM melatonin	39.8 \pm 1.4 ^{a*}	37.4 \pm 1.2 ^{a*}	47.2 \pm 2.6 ^{b*}	67.2 \pm 3.6 ^{c*}	92.4 \pm 6.1 ^{d*}
5 μM melatonin	39.8 \pm 1.6 ^{a*}	0 ^{b*}	0 ^{b*}	0 ^{b*}	0 ^{b*}
0.5 μM melatonin+PG	46.0 \pm 2.4 ^a	62.6 \pm 3.2 ^b	40.8 \pm 2.0 ^{a*}	52.2 \pm 4.1 ^a	49.1 \pm 3.0 ^a
1 μM melatonin+PG	46.0 \pm 2.4 ^a	62.6 \pm 3.2 ^b	40.9 \pm 2.0 ^{a*}	51.9 \pm 3.9 ^a	50.1 \pm 3.1 ^a
5 μM melatonin+PG	46.0 \pm 2.4 ^a	62.6 \pm 3.2 ^b	38.8 \pm 1.8 ^{c*}	35.4 \pm 3.1 ^{c*}	39.5 \pm 1.7 ^{c*}

Spermatozoa were subjected to IVC and further IVAE as described in the Materials and Methods section. Determination of motion parameters through CASA and statistical analyses has been also described in the Material and Methods section. Spermatozoa were incubated in a non-capacitating medium (NCM, C-) or in capacitating medium without (CM, C+) or with melatonin at final concentrations of 0.5 μM (0.5 μM melatonin), 1 μM (1 μM melatonin) and 5 μM (5 μM melatonin). After 4 h of incubation, progesterone (PG) was added. Simultaneously, three more aliquots were incubated in capacitating medium and, after 4 h of incubation, were added with progesterone and 0.5 μM melatonin (0.5 μM melatonin+PG), progesterone with 1 μM melatonin (1 μM melatonin+PG) and progesterone with 5 μM (5 μM melatonin+PG). In all cases, spermatozoa were subsequently incubated, and aliquots were taken after 1, 5 and 60 min of progesterone addition. Different superscript letters (a-d) indicate significant differences ($p < 0.05$) between columns within a given row. Asterisks indicate significant differences ($p < 0.05$) when compared with C+ (CM) at the same time point. Results are shown as means \pm SEM for seven separate experiments.

Table 2 Effects of melatonin on linearity (LIN) and straightness (STR) coefficients of boar spermatozoa subjected to in vitro capacitation and subsequent, progesterone-induced in vitro acrosome exocytosis

Incubation time	0 h	4 h	1 min	5 min	60 min
LIN (%)					
C-	23.6 \pm 1.1 ^{a*}	35.1 \pm 1.3 ^{b*}	36.3 \pm 1.6 ^b	33.6 \pm 1.5 ^{b*}	0 ^{c*}
C+	36.2 \pm 1.4 ^a	44.8 \pm 2.4 ^b	39.9 \pm 1.7 ^a	47.9 \pm 2.3 ^b	37.1 \pm 2.0 ^a
0.5 μM melatonin	42.0 \pm 1.9 ^a	34.8 \pm 1.8 ^{b*}	33.5 \pm 1.6 ^b	43.2 \pm 2.2 ^a	33.2 \pm 2.1 ^b
1 μM melatonin	36.2 \pm 1.7 ^a	30.8 \pm 2.1 ^{a*}	33.6 \pm 1.4 ^a	44.9 \pm 2.4 ^b	35.3 \pm 1.8 ^a
5 μM melatonin	39.8 \pm 2.2 ^a	0 ^{b*}	0 ^{b*}	0 ^{b*}	0 ^{b*}
0.5 μM melatonin+PG	36.2 \pm 1.4 ^a	44.8 \pm 2.4 ^b	26.8 \pm 1.2 ^{c*}	36.5 \pm 1.8 ^{a*}	32.7 \pm 1.5 ^a
1 μM melatonin+PG	36.2 \pm 1.4 ^a	44.8 \pm 2.4 ^b	25.7 \pm 1.6 ^{c*}	36.3 \pm 1.9 ^{a*}	33.2 \pm 1.8 ^a
5 μM melatonin+PG	36.2 \pm 1.4 ^a	44.8 \pm 2.4 ^b	26.0 \pm 1.3 ^{c*}	30.1 \pm 1.9 ^{c*}	28.2 \pm 1.2 ^{c*}
STR (%)					
C-	55.8 \pm 2.3 ^{a*}	48.5 \pm 2.4 ^{b*}	44.6 \pm 2.0 ^{b*}	43.6 \pm 1.9 ^{b*}	0 ^{c*}
C+	63.3 \pm 2.4 ^a	70.9 \pm 3.0 ^b	67.7 \pm 2.8 ^{ab}	71.0 \pm 2.9 ^b	76.2 \pm 3.3 ^b
0.5 μM melatonin	64.7 \pm 2.3 ^a	83.9 \pm 3.4 ^{b*}	67.1 \pm 3.0 ^a	69.0 \pm 3.6 ^a	81.2 \pm 4.5 ^b
1 μM melatonin	62.1 \pm 2.4 ^a	52.1 \pm 1.5 ^{b*}	68.6 \pm 2.5 ^{ab}	79.1 \pm 3.6 ^c	73.9 \pm 3.1 ^b
5 μM melatonin	63.1 \pm 2.4 ^a	0 ^{b*}	0 ^{b*}	0 ^{b*}	0 ^{b*}
0.5 μM melatonin+PG	63.3 \pm 2.4 ^a	70.9 \pm 3.0 ^b	60.6 \pm 2.5 ^a	76.1 \pm 3.7 ^b	77.1 \pm 4.4 ^b
1 μM melatonin+PG	63.3 \pm 2.4 ^a	70.9 \pm 3.0 ^b	58.1 \pm 1.8 ^a	70.5 \pm 2.6 ^b	76.0 \pm 4.5 ^b
5 μM melatonin+PG	63.3 \pm 2.4 ^a	70.9 \pm 3.0 ^b	55.6 \pm 1.6 ^{b*}	64.6 \pm 1.9 ^{a*}	64.8 \pm 2.9 ^{a*}

Spermatozoa were subjected to IVC and further IVAE as described in the Material and Methods section. Determination of motion parameters through CASA and statistical analyses has been also described in the Material and Methods section. Spermatozoa were incubated in a non-capacitating medium (NCM, C-) or in capacitating medium without (CM, C+) or with melatonin at final concentrations of 0.5 μM (0.5 μM Melatonin), 1 μM (1 μM melatonin) and 5 μM (5 μM Melatonin). After 4 h of incubation, progesterone (PG) was added. Simultaneously, three more aliquots were incubated in capacitating medium and, after 4 h of incubation, were added with progesterone and 0.5 μM melatonin (0.5 μM melatonin+PG), progesterone with 1 μM melatonin (1 μM melatonin+PG) and progesterone with 5 μM (5 μM melatonin+PG). In all cases, spermatozoa were subsequently incubated, and aliquots were taken after 1, 5 and 60 min of progesterone addition. Different superscript letters (a-c) indicate significant differences ($p < 0.05$) between columns within a given row. Asterisks indicate significant differences ($p < 0.05$) when compared with C+ (CM) at the same time point. Results are shown as means \pm SEM for seven separate experiments.

significant ($p < 0.05$) increase in the percentage of agglutinated spermatozoa (5 μM melatonin: 56.2% \pm 6.4% vs. CM: 26.1% \pm 3.8%; Fig. 2A and Figure S6G). This increase continued and reached values of about 80–85% at 4 h, when more than a hundred spermatozoa were observed in a single

agglutination (Figure S6H). Similar results were found throughout the remaining experimental period (Fig. 2A, Figure S6I).


Regarding the percentage of agglutinated spermatozoa with appreciable tail beating, melatonin induced a significant

($p < 0.05$) decrease in all the tested concentrations, reaching minimal values at 4 h (melatonin μM : $12.8\% \pm 1.9\%$ vs. CM: $43.2\% \pm 3.2\%$; Fig. 3A). After progesterone addition, a similar decreasing pattern was observed. Finally, the addition of both 1 and 5 μM melatonin at 4 h did counteract the decreases in the percentages of agglutination and of agglutinated spermatozoa with tail beating observed in control samples 60 min after progesterone addition (Fig. 3; Figure S6J).

Effects of melatonin on intracellular ROS levels

Incubation of boar spermatozoa in CM induced a slight, but significant ($p < 0.05$) increase in the percentage of viable spermatozoa with high intracellular H_2O_2 levels, which went from $1.6\% \pm 0.2\%$ at 0 h to $5.8\% \pm 1.3\%$ at 4 h (Fig. 5A,B). This was in

Figure 2 Effects of melatonin on the percentage of agglutinated cells of boar spermatozoa subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis. (A): Melatonin added at 0 h. (B): Melatonin added together with progesterone at 4 h. White bars: spermatozoa incubated in NCM medium (C-). Light grey bars: spermatozoa incubated in CM medium (C+). Medium grey bars: spermatozoa incubated in CM added with 0.5 μM melatonin. Dark green bars: spermatozoa incubated in CM added with 1 μM melatonin. Black bars: spermatozoa incubated in CM added with 5 μM melatonin. Asterisks indicate significant ($p < 0.05$) differences between a given treatment and C+ samples. Figure shows means \pm SEM for seven separate experiments.



contrast with sperm cells incubated in NCM in which the extent of that increase was higher ($9.4\% \pm 2.5\%$ at 4 h; Fig. 4). The subsequent addition of progesterone did not significantly modify the percentage of high- H_2O_2 cells in spermatozoa incubated in CM, whereas those incubated in NCM showed a slight and gradual increase, reaching values of $13.6\% \pm 2.9\%$ after 60 min of progesterone addition (Fig. 4).

The addition of melatonin at 0 h did not significantly affect the pattern observed in CM, except in the case of melatonin 5 μM , where there was a significant ($p < 0.05$) decrease in the percentage of cells with high H_2O_2 levels after 1 min of progesterone addition that was not further recovered (Fig. 4A). The addition of melatonin at 1 or 5 μM at 4 h showed a significant ($p < 0.05$) decrease in this percentage at 1 min post-progesterone addition (Fig. 4B).

Percentages of viable spermatozoa with high intracellular O_2^{\bullet} levels slightly decreased throughout incubation time and

Figure 3 Effects of melatonin on the percentage of agglutinated cells with beating tails of boar spermatozoa subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis. (A): Melatonin added at 0 h. (B): Melatonin added together with progesterone at 4 h. White bars: spermatozoa incubated in NCM medium (C-). Light grey bars: spermatozoa incubated in CM medium (C+). Medium grey bars: spermatozoa incubated in CM added with 0.5 μM melatonin. Dark green bars: spermatozoa incubated in CM added with 1 μM melatonin. Black bars: spermatozoa incubated in CM added with 5 μM melatonin. Asterisks indicate significant ($p < 0.05$) differences between a given treatment and C+ samples. Figure shows means \pm SEM for seven separate experiments.




Figure 4 Effects of melatonin on the percentage of viable spermatozoa with high intracellular peroxide levels of boar spermatozoa subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis. (A): Melatonin added at 0 h. (B): Melatonin added together with progesterone at 4 h. White bars: spermatozoa incubated in NCM medium (C-). Light grey bars: spermatozoa incubated in CM medium (C+). Medium grey bars: spermatozoa incubated in CM added with 0.5 μM melatonin. Dark green bars: spermatozoa incubated in CM added with 1 μM melatonin. Black bars: spermatozoa incubated in CM added with 5 μM melatonin. Asterisks indicate significant ($p < 0.05$) differences between a given treatment and C+ samples. Figure shows means \pm SEM for seven separate experiments.





Figure 5 Effects of melatonin on the percentage of viable spermatozoa with high intracellular superoxide levels of boar spermatozoa subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis. (A): Melatonin added at 0 h. (B): Melatonin added together with progesterone at 4 h. White bars: spermatozoa incubated in NCM medium (C-). Light grey bars: spermatozoa incubated in CM medium (C+). Medium grey bars: spermatozoa incubated in CM added with 0.5 μM melatonin. Dark green bars: spermatozoa incubated in CM added with 1 μM melatonin. Black bars: spermatozoa incubated in CM added with 5 μM melatonin. Asterisks indicate significant ($p < 0.05$) differences between a given treatment and C+ samples. Figure shows means \pm SEM for seven separate experiments.



went from $1.4\% \pm 0.2\%$ at 0 h to $6.2\% \pm 1.7\%$ at 4 h (Fig. 5A,B). Subsequent addition of progesterone did not have a remarkable effect on this parameter, and a slight increase was seen after 60 min of progesterone addition ($9.0\% \pm 2.8\%$; Fig. 5A,B). In the case of incubation in NCM, the values were significantly higher, reaching values of $19.8\% \pm 4.1\%$ after 60 min of progesterone addition. The addition of melatonin at any of the tested concentrations either at 0 h or at 4 h did not differ from spermatozoa incubated in CM (Fig. 5A,B).

Effects of melatonin on the free cysteine residues in both head and tail sperm extracts

Incubation of boar spermatozoa in CM induced a progressive increase in the free cysteine levels from head extracts which

went from $3.9 \text{ nmol/g protein} \pm 0.3 \text{ nmol/g protein}$ at 0 h to $17.2 \text{ nmol/g protein} \pm 2.3 \text{ nmol/g protein}$ at 4 h (Fig. 6A,B). This increase was not observed in sperm cells incubated in NCM. Subsequent addition of progesterone to spermatozoa incubated in CM did not increase these levels, and there was a slight gradual decrease with values of $11.3 \text{ nmol/g protein} \pm 1.7 \text{ nmol/g protein}$ after 60 min of progesterone addition (Fig. 6A,B).

Addition of melatonin at 0 h had a dramatic effect on free cysteine levels from head extracts, as almost abolished the increase observed in CM (Fig. 6A). This effect was maintained after the addition of progesterone. When melatonin was added at 4 h, a similar decreasing effect on free cysteine levels of sperm head

extracts was immediately observed (i.e. 1 min after melatonin and progesterone addition), especially at the highest concentrations.

Free cysteine levels of sperm tail extracts also increased in spermatozoa incubated in CM. Values went from 3.7 nmol/g protein \pm 0.6 nmol/g protein at 0 h to 9.3 nmol/g protein \pm 1.9 nmol/g protein at 4 h (Fig. 7A,B). These values were roughly maintained after the addition of progesterone. The addition of melatonin at 0 h almost abolished that increase at any concentration tested (Fig. 7A). On the contrary, the addition of melatonin at 4 h had no clear effect on this parameter before 60 min after progesterone addition, when free cysteine levels of sperm tail extracts were found to increase in a melatonin





dose-dependent manner (5 μ M: 12.8 nmol/g protein \pm 2.4 nmol/g protein vs. CM: 8.9 nmol/g protein \pm 1.7 nmol/g; Fig. 7A).

Effects of melatonin on distribution of lectins over sperm membrane

At the beginning of incubation in CM, WGA signal was located at the sperm head and the whole tail, although the maximal intensity of the signal was observed at the acrosomal edge (Figure S7). Spermatozoa incubated in NCM showed a similar staining, but the acrosomal signal was much less intense. After 4 h of incubation in CM, the intensity of the acrosome-located signal increased and uniformly distributed throughout the entire acrosome (Figure S7). These changes were not detected in

Figure 6 Effects of melatonin on the head intracellular free cysteine radicals levels of boar spermatozoa subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis. (A): Melatonin added at 0 h. (B): Melatonin added together with progesterone at 4 h. White bars: spermatozoa incubated in NCM medium (C-). Light grey bars: spermatozoa incubated in CM medium (C+). Medium grey bars: spermatozoa incubated in CM added with 0.5 μ M melatonin. Dark green bars: spermatozoa incubated in CM added with 1 μ M melatonin. Black bars: spermatozoa incubated in CM added with 5 μ M melatonin. Asterisks indicate significant ($p < 0.05$) differences between a given treatment and C+ samples. Figure shows means \pm SEM for seven separate experiments.

Figure 7 Effects of melatonin on the tail intracellular free cysteine radical levels of boar spermatozoa subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis. (A): Melatonin added at 0 h. (B): Melatonin added together with progesterone at 4 h. White bars: spermatozoa incubated in NCM medium (C-). Light grey bars: spermatozoa incubated in CM medium (C+). Medium grey bars: spermatozoa incubated in CM added with 0.5 μ M melatonin. Dark green bars: spermatozoa incubated in CM added with 1 μ M melatonin. Black bars: spermatozoa incubated in CM added with 5 μ M melatonin. Asterisks indicate significant ($p < 0.05$) differences between a given treatment and C+ samples. Figure shows means \pm SEM for seven separate experiments.



spermatozoa incubated in NCM for 4 h. The subsequent addition of progesterone to CM induced further modifications in the acrosome signal of WGA. In effect, although small changes were observed after 1 min of adding progesterone, with some sperm cells losing the inner lectin signal and others showing an irregular acrosome marking, the progesterone-induced changes were much more apparent after 5 min, when a high number of sperm cells showed a diffuse inner or irregular acrosome signal (Figure S7). These patterns were also observed after 60 min of progesterone addition. Addition of melatonin at 0 h induced changes in the WGA-staining pattern. At 4 h, most of the spermatozoa incubated with melatonin showed much more intense acrosome signal than control spermatozoa (CM), but there were also sperm cells that showed no lectin signal in the post-acrosomal area and even spermatozoa with no signal over the head (Figure S7). The addition of progesterone to spermatozoa incubated for 4 h in treatments containing melatonin did not induce immediate changes in WGA distribution. However, the acrosome staining in spermatozoa incubated with melatonin was more intense than that observed in control spermatozoa incubated in CM after 60 min of incubation (Figure S7). In contrast, the addition of 1 μM melatonin at 4 h did not change WGA distribution when compared to control spermatozoa incubated in CM (Figure S7).

Regarding STL, it was mainly found in sperm head and midpiece (in some cells, STL was observed in the entire tail) at 0 h, with a more intense signal in the acrosome area (Figure S8). Following incubation in CM for 4 h, two different STL-staining patterns were observed. Whereas one pattern consisted of an intense and uniform signal throughout the entire acrosome area, STL staining in the other was mainly restricted to the acrosomal edge (Figure S8). In both patterns, STL signal in the midpiece was much decreased or totally absent. The addition of progesterone at 4 h decreased the intensity of acrosome signal in practically all sperm cells (which was already apparent after 1 min of progesterone addition) but increased that of the tail (Figure S8). Addition of 1 μM melatonin at 0 h exhibited similar STL-staining patterns to the control (Figure S8) and there was a clear loss of STL signal in the acrosome following progesterone addition, the STL-staining being restricted to the acrosome edge. While the addition of 1 μM melatonin at 4 h did not clearly affect the STL pattern observed in spermatozoa incubated in CM, a high proportion of spermatozoa showed a clearly intense acrosome signal after 1 min of the addition of melatonin and progesterone, which differed from spermatozoa incubated in CM (Figure S8).

At 0 h, PSA staining was observed in the whole cell (spermatozoa incubated in CM), although the most intense marking was detected in the entire acrosomal area. In the case of spermatozoa incubated in NCM, the acrosomal signal was much less intense and restricted to the acrosome edge (Figure S9). Incubation in CM for 4 h induced an increase in the PSA-staining of the sperm head and tail, despite the post-acrosomal region being devoid of PSA signal. Subsequent addition of progesterone induced a rapid loss of PSA signal in the acrosome, which was evident in a high percentage of spermatozoa after 1 min of the addition of the hormone (Figure S9). In spite of this, there were other spermatozoa that showed an intense acrosome signal. The addition of 1 μM melatonin either at 0 h or at 4 h did not have a clear effect on PSA localisation during IVC and IVAE (Figure S9).

Regarding the localisation of PNA, it was exclusively found at the whole acrosome surface and there were no changes after 4 h of incubation either in CM or in NCM (Figure S10). As expected, the addition of progesterone at 4 h increased the proportion of spermatozoa with less intensity of PNA marking and faint staining restricted to the acrosome edge (Figure S10). These changes were very rapid, as they were already observed after 1 min of progesterone addition. Addition of melatonin either at 0 h or at 4 h showed no differences when compared to spermatozoa incubated in CM (Figure S10).

Effects of melatonin on the ability of in vitro capacitated boar spermatozoa to adhere and penetrate pig oocytes

As shown in Table 4, the number of spermatozoa attached to the ZP (78.4 ± 1.8), the total penetration rate (90.6%) and the percentage of monospermy (69.8%) were higher in the control group (CM) than in the other treatments. Previous incubation of spermatozoa with 1 μM melatonin significantly ($p < 0.05$) decreased the number of spermatozoa adhered to the ZP (68.2 ± 2.7 vs. 78.4 ± 1.8 in CM). Furthermore, incubation with melatonin at 1 μM significantly ($p < 0.05$) decreased the proportion of polyspermic oocytes (14.5% vs. 20.8% in CM). The addition of 1 μM melatonin after in vitro capacitation for 4 h had no effect on polyspermy, but increased the number of spermatozoa attached to the ZP (88.9 ± 1.7 vs. 78.4 ± 1.8 in CM; Table 4).

DISCUSSION

The results shown herein suggest that one of the most important effects of melatonin during the achievement of boar spermatozoa IVC is the increase in cell adhesiveness. The increase in sperm adhesiveness would influence important aspects of boar sperm capacitation such as sperm motility and sperm-zona pellucida interaction through the activation of sperm agglutination, as the results suggested (Fig. 2 and Tables 1–3). Additionally, melatonin did not affect several of the most important capacitation markers, such as membrane lipid disorder, tyrosine phosphorylation levels of P32 and the ability to reach acrosome exocytosis after progesterone stimulation. This lack of additional effects could suggest that the melatonin action of IVC is mainly focused on motility and/or sperm adhesiveness. Regarding the relationship between sperm motility and agglutination, it is worth noting that IVC in species such as the monkey, cattle, sheep and pig leads to an increase in the percentage of agglutinated spermatozoa (Boatman & Bavister, 1984; Ehrenwald *et al.*, 1990; Funahashi & Day, 1993; Lefebvre & Suarez, 1996; Leahy *et al.*, 2016). In fact, sperm agglutination does not only result from IVC, but also from other factors, such as the presence of antibodies (Yakirevich & Naota, 1999) or in response to cell degeneration (Harayama *et al.*, 1998). Furthermore, several components of capacitation media, such as heparin in cattle, BSA in horse, and bicarbonate and calcium in pigs, monkeys and cattle increase sperm agglutination (Lindahl & Sjöblom, 1981; Boatman & Bavister, 1984; Ehrenwald *et al.*, 1990; Funahashi & Day, 1993; Lefebvre & Suarez, 1996; Harayama *et al.*, 1998; Harayama & Kato, 2002). Therefore, in our experimental conditions, melatonin could have enhanced the agglutination-promoting effect of CM components. This hypothesis could explain why the addition of melatonin at concentrations lower than those assayed in this work (100 pM) does not agglutinate ram spermatozoa (Casao *et al.*, 2009). Based on CM composition, one could

1 suggest that bicarbonate is the agglutinating factor whose action
 2 is potentiated by the addition of melatonin. This hypothesis is
 3 based on previous works from our laboratory, in which IVC of
 4 boar spermatozoa was achieved in a medium without bicarbon-
 5 ate and not much agglutination was observed (Ramió *et al.*,
 6 2008; Ramió-Lluch *et al.*, 2011; Ramió-Lluch *et al.*, 2014). While
 7 the exact mechanism through which
 8 sperm agglutination, our results sug-
 9 gest it is unlikely to be related to changes in the glycolalix com-
 10 position of membrane surface, as lectin-binding assays did not
 11 show apparent changes (Figures S7–S10). Therefore, we could
 12 propose two explanations. The first one would be related to a
 13 cAMP-mediated mechanism via the activation of the bicarbon-
 14 ate-sensitive adenylyl cyclase and PKA (Harayama & Kato,
 15 2002). Another possible mechanism would involve the mainte-
 16 nance of disulphide bonds, as melatonin induced a clear
 17 decrease in the intracellular free cysteine levels of boar sperma-
 18 tozoa. In this respect, it is worth noting that penicillamine has a
 19 potent action against agglutination in ram spermatozoa sub-
 20 jected to IVC (Leahy *et al.*, 2016), and one of the mechanisms
 21 through which it exerts that effect is linked to its direct action
 22 on disulphide radicals, which are converted into sulfhydryl
 23 groups (Talevi *et al.*, 2007; Gualtieri *et al.*, 2009). These data
 24 would be in concordance with the results obtained in this study,
 25 suggesting that the effects of melatonin on sperm agglutination
 26 and free cysteine levels could be linked. However, more work is
 27 needed to further elucidate this point.

28 The results observed following sperm–oocyte co-incubation
 29 could also be a consequence of melatonin action on sperm
 30 agglutination rather than on other capacitation-related changes
 31 such as the ability to trigger acrosome exocytosis following the
 32 appropriate stimuli, namely progesterone. Thus, the reduction
 33 in the number of spermatozoa adhered to the ZP when they were
 34 previously capacitated in the presence of melatonin could be a
 35 consequence of an increased degree of sperm agglutination,
 36 thereby lowering the number of free spermatozoa able to adhere
 37 the ZP. Furthermore, as shown in Table 4, the addition of mela-
 38 tonin after IVC significantly increased the number of spermato-
 39 zoa bound to the ZP. The combined analysis of these results
 40 suggests that the melatonin-induced agglutinating effect in IVC
 41 conditions could be linked to an unspecific, increased adherence
 42 ability of boar spermatozoa to other sperm cells or the oocyte. In
 43 turn, the observed decrease in polyspermy in spermatozoa
 44 capacitated in the presence of melatonin could result from the
 45 decrease in the number of adhered spermatozoa.

46 Regarding the effects of melatonin on the proportions of boar
 47 spermatozoa with high intracellular ROS levels, results displayed
 48 in Figs 4 and 5 indicate that the antioxidant action of melatonin
 49 does not seem to play a prominent role in the effects observed
 50 during IVC and IVAE and, specifically, in the increase in sperm
 51 agglutination. While this conclusion could be surprising at first
 52 glance, one should note that boar spermatozoa are characterised
 53 by a very low ROS production rate, even when subjected to treat-
 54 ments that, such as freeze-thawing, induce the generation of
 55 high intracellular ROS in other species (Bilodeau *et al.*, 2000;
 56 Guthrie & Welch, 2006; Yeste *et al.*, 2013, 2015a). This feature is
 57 different from other species, such as the horse, in which ROS
 58 production is more intense (see Gibb & Aitken, 2016; as a
 59 review). As a consequence, in species such as equine, bovine
 60 and human, melatonin has a clear antioxidant effect and affects

membrane lipid peroxidation and ROS levels (Gadella *et al.*,
 2008; Rao & Gangadharan, 2008; Du Plessis *et al.*, 2010; Jang
et al., 2010; Da Silva *et al.*, 2011; Najafi *et al.*, 2018). In fact, the
 antioxidant effect of melatonin in species such as the horse is
 not only related to a direct action on the oxidative potential but
 also to intracellular Na⁺ concentrations, which also affect the
 overall redox status (Ortega Ferrusola *et al.*, 2017). Taking all of
 these data into consideration, an expected antioxidant effect of
 melatonin on boar spermatozoa should be very subtle, if detect-
 able. A possible reason to explain why boar spermatozoa do not
 accumulate high ROS levels, and thus why melatonin would
 exert a slight action on this parameter, could be linked to a spe-
 cies-specific mitochondria function. Previous reports have sug-
 gested that, despite being important to maintain crucial sperm
 functions, such as motility, mitochondrial-produced energy is
 low in boar spermatozoa (Rodríguez-Gil & Bonet, 2016). Indeed,
 O₂ consumption rate and intracellular ATP levels have been
 reported to be low during IVC and subsequent IVAE (Ramió-
 Lluch *et al.*, 2014). In addition to this, induction of IVC in the
 presence of oligomycin A, a specific inhibitor of the mito-
 chondrial ATP synthase, does not decrease either O₂ consumption
 rates or ATP levels (Ramió-Lluch *et al.*, 2014), which suggests
 that boar sperm mitochondria are in an uncoupled status
 throughout most of their lifespan. A consequence of this uncou-
 pled status is the low rate of ROS production due to substimula-
 tion of the electronic chain, which is the most important ROS-
 synthesising point (Rodríguez-Gil & Bonet, 2016). Interestingly,
 progesterone-induced acrosome exocytosis is concomitant with
 a sudden and intense peak in both O₂ consumption rate and
 intracellular ATP levels (Balis *et al.*, 1999; Gualtieri *et al.*, 2009),
 which suggests that mitochondria are coupled at this moment.
 Following this rationale, mitochondrial coupling upon proges-
 terone addition would be associated with a transient increase in
 ROS generation, especially that of peroxides (Tait & Green,
 2012). This hypothesis matches with our observations, as the
 addition of melatonin together with that of progesterone was
 found to decrease the proportions of viable spermatozoa with
 high peroxide levels (see Fig. 4A). However, the fact that such a
 decrease did not modify the percentage of acrosome-exocytosed
 spermatozoa induced by progesterone suggests that peroxide
 levels do not play a vital role for acrosome reaction in boar sper-
 matozoa, at least under our in vitro conditions.

As indicated when discussing the effects on sperm agglutina-
 tion and adhesiveness, melatonin decreased the intracellular
 free cysteine levels in both sperm head and tail extracts (see
 Figs 6 and 7). Free cysteine levels are an indirect marker of the
 number of disrupted disulphide bonds, as despite not all free
 cysteine radicals resulting from the breakage of disulphide
 bonds, a significant percentage has this origin (Yeste *et al.*, 2013,
 2014). Our results also indicated that the melatonin-induced
 decrease in free cysteine residues of sperm head has no impact
 on achieving the IVC and subsequent progesterone-induced
 IVAE. Another issue is the observed increase in free cysteine resi-
 dues in sperm tail extracts during IVC and the melatonin-coun-
 teracting effect of that increase. Disulphide bonds are important
 for the maintenance of a proper sperm flagellum structure (Ijiri
et al., 2014). Specifically, disulphide bonds are crucial for a pro-
 tein associated with the outer dense fibre 1 (ODF1; Cabrilla-
et al., 2011). Thus, one could suggest that changes in free cys-
 teine residues of sperm tail would affect the structure of the

APROBADO not determined
 Por 1001869 fecha 16:15, 14/05/2018

CORREGIDO
 16:22, 14/05/2018

Table 3 Effects of melatonin on mean amplitude of lateral head displacement (ALH) and frequency of head displacement (BCF) of boar spermatozoa subjected to *in vitro* capacitation and subsequent, progesterone-induced *in vitro* acrosome exocytosis

Incubation time	0 h	4 h	1 min	5 min	60 min
ALH (μm)					
C-	2.56 \pm 0.05 ^{a*}	2.57 \pm 0.09 ^{a*}	2.18 \pm 0.11 ^{b*}	2.14 \pm 0.10 ^{b*}	0 ^{c*}
C+	3.80 \pm 0.12 ^a	4.13 \pm 0.13 ^b	4.93 \pm 0.15 ^c	4.64 \pm 0.15 ^c	4.00 \pm 0.11 ^b
0.5 μM melatonin	3.31 \pm 0.09 ^{a*}	2.88 \pm 0.08 ^{b*}	4.10 \pm 0.12 ^{c*}	3.63 \pm 0.12 ^{a,c*}	5.13 \pm 0.21 ^{d*}
1 μM melatonin	3.50 \pm 0.14 ^a	2.72 \pm 0.09 ^{b*}	3.68 \pm 0.08 ^{a*}	3.49 \pm 0.11 ^{a*}	3.85 \pm 0.10 ^a
5 μM Melatonin	3.89 \pm 0.13 ^a	0 ^{b*}	0 ^{b*}	0 ^{b*}	0 ^{b*}
0.5 μM melatonin+PG	3.80 \pm 0.12 ^a	4.13 \pm 0.13 ^b	4.82 \pm 0.20 ^b	5.09 \pm 0.20 ^b	3.81 \pm 0.14 ^a
1 μM melatonin+PG	3.80 \pm 0.12 ^a	4.13 \pm 0.13 ^b	5.38 \pm 0.25 ^c	4.96 \pm 0.19 ^c	3.09 \pm 0.11 ^{d*}
5 μM melatonin+PG	3.80 \pm 0.12 ^a	4.13 \pm 0.13 ^b	5.18 \pm 0.20 ^c	4.93 \pm 0.18 ^c	2.99 \pm 0.06 ^{d*}
BCF (Hz)					
C-	2.92 \pm 0.04 ^{a*}	3.97 \pm 0.26 ^{b*}	3.88 \pm 0.21 ^{b*}	4.01 \pm 0.24 ^{b*}	0 ^{c*}
C+	6.68 \pm 0.17 ^a	6.54 \pm 0.19 ^a	7.18 \pm 0.18 ^b	6.28 \pm 0.21 ^a	6.03 \pm 0.19 ^c
0.5 μM melatonin	6.73 \pm 0.15 ^a	6.70 \pm 0.16 ^a	7.20 \pm 0.12 ^b	6.97 \pm 0.24 ^{b*}	6.85 \pm 0.21 ^{ab*}
1 μM melatonin	6.16 \pm 0.11 ^{a*}	5.95 \pm 0.11 ^{a*}	6.76 \pm 0.20 ^{b*}	6.90 \pm 0.24 ^{b*}	7.20 \pm 0.27 ^{c*}
5 μM melatonin	6.05 \pm 0.10 ^{a*}	0 ^{b*}	0 ^{b*}	0 ^{b*}	0 ^{b*}
0.5 μM melatonin+PG	6.68 \pm 0.17 ^a	6.54 \pm 0.19 ^a	6.86 \pm 0.21 ^a	5.46 \pm 0.09 ^{b*}	4.37 \pm 0.06 ^{c*}
1 μM melatonin+PG	6.68 \pm 0.17 ^a	6.54 \pm 0.19 ^a	7.48 \pm 0.29 ^b	4.98 \pm 0.09 ^{c*}	3.91 \pm 0.08 ^{d*}
5 μM melatonin+PG	6.68 \pm 0.17 ^a	6.54 \pm 0.19 ^{ab}	6.44 \pm 0.17 ^{b*}	4.92 \pm 0.08 ^{c*}	3.85 \pm 0.07 ^{d*}

Spermatozoa were subjected to IVC and further IVAE as described in the Material and Methods section. Determination of motion parameters through CASA and statistical analyses has been also described in the Material and Methods section. Spermatozoa were incubated in a non-capacitating medium (NCM, C-) or in capacitating medium without (CM, C+) or with melatonin at final concentrations of 0.5 μM (0.5 μM melatonin), 1 μM (1 μM melatonin) and 5 μM (5 μM melatonin). After 4 h of incubation, progesterone (PG) was added. Simultaneously, three more aliquots were incubated in capacitating medium and, after 4 h of incubation, were added with progesterone and 0.5 μM melatonin (0.5 μM melatonin+PG), progesterone with 1 μM melatonin (1 μM melatonin+PG) and progesterone with 5 μM (5 μM melatonin+PG). In all cases, spermatozoa were subsequently incubated, and aliquots were taken after 1, 5 and 60 min of progesterone addition. Different superscript letters (a-c) indicate significant differences ($p < 0.05$) between columns within a given row. Asterisks indicate significant differences ($p < 0.05$) when compared with C+ (CM) at the same time point. Results are shown as means \pm SEM for seven separate experiments.

Table 4 Effects of melatonin on the adherence and penetration abilities of boar spermatozoa subjected to co-incubation with *in vitro* matured porcine oocytes

	Control ($n = 53$)	Capacitation with 1 μM melatonin ($n = 55$)	Co-incubation with 1 μM melatonin ($n = 57$)
Adhered spermatozoa/oocyte	78.4 \pm 1.8 ^a	68.2 \pm 2.7 ^b	88.9 \pm 1.7 ^c
Total penetration rate (%)	90.6 ^a	89.1 ^a	93.0 ^a
Monospermy (%)	69.8 ^a	74.5 ^a	70.2 ^a
Polyspermy (%)	20.8 ^a	14.5 ^b	22.8 ^a

Control: Spermatozoa subjected to a previous standard *in vitro* capacitation procedure through incubation at 38.5 $^{\circ}\text{C}$ and 5% CO_2 for 4 h. Capacitation with 1 μM melatonin: Spermatozoa subjected to *in vitro* capacitation in a medium added with 1 μM melatonin. Co-incubation with 1 μM melatonin: Spermatozoa subjected to a previous standard *in vitro* capacitation procedure for 4 h. Melatonin at 1 μM was added when spermatozoa and oocytes were co-incubated. Different superscript letters between columns within a given row indicate significant ($p < 0.05$) differences between groups.

flagellum which could, in turn, induce subtle changes in sperm motion. Related to this, cleavage of disulphide bonds in mouse hexokinase-I isozyme 1 (HK1) is related to the initiation of sperm motility (Nakamura *et al.*, 2008). Thus, it could be hypothesised that changes in the number of disrupted disulphide bonds along the flagellum, specifically in flagellum-bond-related proteins such as sperm hexokinase-1 (Cabrillana *et al.*, 2011, 2016), could also be related to the changes in motion parameters observed during IVC. Herein, melatonin at 0.5 and 1 μM was found to abolish the IVC-linked changes in kinetic parameters such as VCL, VAP and ALH (see Tables 1-3). While this effect could also be due to other mechanisms, such as sperm agglutination, the possibility that melatonin affects sperm motility through regulating the number of disrupted disulphide bonds should not be dismissed.

It is worth noting that the addition of melatonin together with progesterone after 4 h of IVC has almost no effects on subsequent acrosome exocytosis or on the sperm ability to adhere and penetrate oocytes *in vitro*. Taking into account that the main difference of spermatozoa before and after their incubation for 4 h in CM is the achievement of a feasible capacitation

status, these results suggest that the observed effects of melatonin on parameters such as the adhesiveness and free cysteine levels of the tail are linked to the precise sperm status. Thus, and as indicated by membrane lipid disorder, sperm motility and sensitivity to progesterone, capacitated differs from uncapacitated spermatozoa in their response to melatonin. At this moment, it is not possible to ascertain which the basis for these differences in melatonin action is. Further research focusing on the function of specific melatonin receptors MT1 and MT2, which are present in boar spermatozoa (González-Arto *et al.*, 2016), is warranted.

In conclusion, melatonin modulates the achievement of IVC and subsequent progesterone-induced IVAE in boar spermatozoa via mechanism/s involved in the control of sperm motion through changes in the number of tail disulphide bridges, adhesiveness and further oocyte penetration ability. In addition, prevention of the IVC-induced increase in the disulphide bonds of sperm head proteins mediated by melatonin could also be relevant. Remarkably, melatonin effects on IVC/IVAE in boar spermatozoa do not seem to be related to a direct action on intracellular ROS levels, thus opening up alternative, perhaps



receptor-mediated, pathways to explain the effects of this hormone upon sperm capacitation.

ACKNOWLEDGEMENTS

We wish to thank Dr. Manel Garcia (Serveis Científico-Tècnics, University of Barcelona, Spain) for his very useful technical advice in confocal microscopy analyses. The authors also thank Frigorífics Costa Brava, S.A. (Riudellots de la Selva, Spain) for donating the ovaries used in IVF experiments.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported herein.

AUTHORS' CONTRIBUTIONS

M.R. and R.B. carried out the majority of the experimental work, collaborated in the design of the experiments and wrote the manuscript. A.Pl., M.S., A.Pe. and T.R. collaborated with M.R. and R.B. in performing laboratory work. J.M.F.N. conducted confocal analysis of lectin location. T.M.B., A.C. and J.A.C.P. collaborated in designing the experiments and critically revised the manuscript. S.B. and M.C.M. helped conduct flow cytometry analyses and IVF experiments. As joint senior author, M.Y. was involved in flow cytometry analyses and IVF experiments, designed the experiments, analysed the data, wrote the manuscript and gave his final approval. J.E.R.G. designed the experiments, analysed the data, wrote the manuscript and gave his final approval.

FUNDING

This work was supported by Grants AGL2013-43328-P and RYC-2014-15581 (Ministry of Economy and Competitiveness, Spain).

REFERENCES

Ashrafi I, Kohram H & Ardabili FF. (2013) Antioxidative effects of melatonin on kinetics, microscopic and oxidative parameters of cryopreserved bull spermatozoa. *Anim Reprod Sci* 139, 25–30.

Awda BJ, Mackenzie-Bell M & Buhr MM. (2009) Reactive oxygen species and boar sperm function. *Biol Reprod* 81, 553–561.

Balis UJ, Behnia K, Dwarakanath B & Bathia SN. (1999) Oxygen consumption characteristics of porcine hepatocytes. *Metab Eng* 1, 49–62.

Bilodeau JF, Chatterjee S, Sirard NA & Gagnon C. (2000) Levels of antioxidant defenses are decreased in bovine spermatozoa after a cycle of freezing and thawing. *Mol Reprod Dev* 55, 282–288.

Boatman DE & Bavister BD. (1984) Stimulation of rhesus monkey sperm capacitation by cyclic nucleotide mediators. *J Reprod Fertil* 71, 357–366.

Bradford MM. (1976) Rapid and sensitive method for quantitation of microgram quantities of protein utilizing principle of protein-dye binding. *Anal Biochem* 72, 248–254.

Bravo MM, Aparicio IM, García-Herreros M, Gil MC, Peña FJ & García-Marín LJ. (2005) Changes in tyrosine phosphorylation associated with true capacitation and capacitation-like state in boar spermatozoa. *Mol Reprod Dev* 71, 88–96.

Brocklehurst K, Stuchbury T & Malthouse JPG. (1979) Reactivities of neutral and cationic forms of 2,2'-dipyridyl disulfide towards thiolate anions - detection of differences between the active-centers of actinidin, papain and ficin by a 3-protonic-state reactivity probe. *Biochem J* 183, 233–238.

Bromfield EG, Aitken RJ, Gibb Z, Lambourne SR & Nixon B. (2014) Capacitation in the presence of methyl- β -cyclodextrin results in enhanced zona pellucida-binding ability of stallion spermatozoa. *Reproduction* 147, 153–166.

Cabrillana ME, Monclus MA, Saez Lancellotti TE, Boarelli PV, Clementi MA, Vincenti AE, Yunes RFM & Fornés MW. (2011) Characterization of flagellar cysteine-rich sperm proteins involved in motility, by the combination of cellular fractionation, fluorescence detection, and mass spectrometry analysis. *Cytoskeleton* 68, 491–500.

Cabrillana ME, Uribe P, Villegas JV, Álvarez J, Sánchez R & Fornés MW. (2016) Thiol oxidation by nitrosative stress: cellular localization in human spermatozoa. *Syst Biol Reprod Med* 62, 325–334.

Casao A, Mendoza N, Pérez-Pé R, Grasa P, Abecia JA, Forcada F, Cebrián-Pérez JA & Muñio-Blanco T. (2009) Melatonin prevents capacitation and apoptotic-like changes of ram spermatozoa and increases fertility rate. *J Pineal Res* 48, 39–46.

Casao A, Cebrián I, Asumpção ME, Pérez-Pé R, Abecia JA, Forcada F, Cebrián-Pérez JA & Muñio-Blanco T. (2010) Seasonal variations of melatonin in ram seminal plasma are correlated to those of testosterone and antioxidant enzymes. *Reprod Biol Endocrinol* 8, 59.

Castillo-Martín M, Bonet S, Morató R & Yeste M. (2014) Comparative effects of adding β -mercaptoethanol or L-ascorbic acid to culture or vitrification-warming media on IVF porcine embryos. *Reprod Fertil Dev* 26, 875–882.

Cebrián-Pérez JA, Casao A, Gonzalez-Arto M, dos Santos Hamilton TR, Pérez-Pé R & Muñio-Blanco T. (2014) Melatonin in sperm biology: breaking paradigms. *Reprod Domest Anim* 49, 1–21.

Cooper TG & Yeung CH. (1998) A flow cytometric technique using peanut agglutinin for evaluating acrosomal loss from human spermatozoa. *J Androl* 19, 542–550.

Da Silva CM, Macías-García B, Miró-Morán A, González-Fernández L, Morillo-Rodríguez A, Ortega-Ferrusola C, Gallardo-Bolaños JM, Stilwell G, Tapia JA & Peña FJ. (2011) Melatonin reduces lipid peroxidation and apoptotic-like changes in stallion spermatozoa. *J Pineal Res* 51, 172–179.

Du Plessis SS, Hagenaar K & Lampiao F. (2010) The in vitro effects of melatonin on human sperm function and its scavenging activities on NO and ROS. *Andrologia* 42, 112–116.

Ehrenwald E, Foote RH & Parks JE. (1990) Bovine oviductal fluid components and their potential role in sperm cholesterol efflux. *Mol Reprod Develop* 25, 195–204.

Espino J, Ortiz Á, Bejarano I, Lozano GM, Monllor F, García JF, Rodríguez AB & Pariente JA. (2011) Melatonin protects human spermatozoa from apoptosis via melatonin receptor- and extracellular signal-regulated kinase-mediated pathways. *Fertil Steril* 95, 2290–2296.

Flores E, Ramio-Lluch L, Bucci D, Fernandez-Novell JM, Peña A & Rodríguez-Gil JE. (2011) Freezing-thawing induces alterations in histone H1-DNA binding and the breaking of protein-DNA disulfide bonds in boar sperm. *Theriogenology* 76, 1450–1464.

Funahashi H & Day BN. (1993) Effects of follicular fluid at fertilization in vitro on sperm penetration in pig oocytes. *J Reprod Fertil* 99, 97–103.

Gadella BM, Tsai PS, Boerke A & Brewis IA. (2008) Sperm head membrane reorganisation during capacitation. *Int J Dev Biol* 52, 473–480.

Garner DL & Johnson LA. (1995) Viability assessment of mammalian sperm using sybr-14 and propidium iodide. *Biol Reprod* 53, 276–284.

Gibb Z & Aitken RJ. (2016) The impact of sperm metabolism during in vitro storage: the stallion as a model. *Biomed Res Int* 2016, 1–8.

González-Arto M, Carrillo AV, Martínez-Pastor F, Fernández-Alegre E, Roca J, Miró J, Rigau T, Rodríguez-Gil JE, Pérez-Pé R, Muñio-Blanco T, Cebrián-Pérez JA & Casao A. (2016) Melatonin receptors MT1 and MT2 are expressed in spermatozoa from several seasonal and nonseasonal breeder species. *Theriogenology* 86, 1958–1968.

- Gualtieri R, Mollo V, Duma G & Talevi R. (2009) Redox control of surface protein sulphhydryls in bovine spermatozoa reversibly modulates sperm adhesion to the oviductal epithelium and capacitation. *Reproduction* 138, 33–43.
- Guthrie HD & Welch GR. (2006) Determination of intracellular reactive oxygen species and high mitochondrial membrane potential in Percoll-treated viable boar sperm using fluorescence-activated flow cytometry. *J Anim Sci* 84, 2089–2100.
- Harayama H & Kato S. (2002) Relationship between bicarbonate and cyclic nucleotide in the promoting effects on head-to-head agglutination in boar spermatozoa. *Asian J Androl* 4, 87–96.
- Harayama H, Miyake M, Shidara O, Iwamoto E & Kato S. (1998) Effects of calcium and bicarbonate on head-to-head agglutination in ejaculated boar spermatozoa. *Reprod Fertil Dev* 10, 445–450.
- Ijiri TW, Vadnais ML, Huang AP, Lin AM, Levin LR, Buck J & Gerton GL. (2014) Thiol changes during epididymal maturation: a link to flagellar angulation in mouse spermatozoa? *Andrology* 2, 65–75.
- Jang HY, Kim YH, Kim BW, Park IC, Cheong HT, Kim JT, Park CK, Kong HS, Lee HK & Yang BK. (2010) Ameliorative effects of melatonin against hydrogen peroxide induced oxidative stress on boar sperm characteristics and subsequent in vitro embryo development. *Reprod Domest Anim* 45, 943–950.
- Jimenez I, Gonzalez-Marquez H, Ortiz R, Herrera JA, Garcia A, Betancourt M & Fierro R. (2003) Changes in the distribution of lectin receptors during capacitation and acrosome reaction in boar spermatozoa. *Theriogenology* 59, 1171–1180.
- Laemmli UK. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680–685.
- Leahy T, Rickard JP, Aitken RJ & de Graaf SP. (2016) Penicillamine prevents ram sperm agglutination in media that support capacitation. *Reproduction* 151, 167–177.
- Lee JA, Spidlen J, Boyce K, Cai J, Crosbie N, Dalphin M, Furlong J, Gasparetto M, Goldberg M, Goralczyk EM, Hyun B, Jansen K, Kollmann T, Kong M, Leif R, McWeeney S, Moloshok TD, Moore W, Nolan G, Nolan J, Nikolich-Zugich J, Parrish D, Purcell B, Qian Y, Selvaraj B, Smith C, Tchuvatkina O, Wertheimer A, Wilkinson P, Wilson C, Wood J, Zigon R, International Society for Advancement of Cytometry Data Standards Task Force, Scheuermann RH & Brinkman RR. (2008) MIFlowCyt: the minimum information about a flow cytometry experiment. *Cytometry* 73A, 926–930.
- Lefebvre R & Suarez SS. (1996) Effect of capacitation on bull sperm binding to homologous oviductal epithelium. *Biol Reprod* 54, 575–582.
- Lindahl PE & Sjöblom P. (1981) On mechanisms of head-to-head association in bovine spermatozoa. *Biol Reprod* 25, 29–43.
- Luboshitzky R, Shen-Orr Z & Herer P. (2002) Seminal plasma melatonin and gonadal steroids concentrations in normal men. *Arch Androl* 48, 225–232.
- Najafi A, Adutwum E, Yari A, Salehi E, Mikaeili S, Dashtestani F, Abolhassani F, Rashki L, Shiasi S & Asadi E. (2018) Melatonin affects membrane integrity, intracellular reactive oxygen species, caspase3 activity and AKT phosphorylation in frozen thawed human sperm. *Cell Tissue Res* 372, 149–159.
- Nakamura N, Shibata H, O'Brien DA, Mori C & Eddy EM. (2008) Spermatogenic cell-specific type 1 hexokinase is the predominant hexokinase in sperm. *Mol Reprod Dev* 75, 632–640.
- Ortega Ferrusola C, Anel-López L, Ortiz Rodríguez JM, Martín Muñoz P, Álvarez M, de Paz P, Masot J, Redondo E, Balao da Silva C, Morrell JM, Rodríguez-Martínez H, Tapia JA, Gil MC, Anel L & Peña FJ. (2017) Stallion spermatozoa surviving freezing and thawing experience membrane depolarization and increased intracellular Na⁺. *Andrology* 5, 1174–1182.
- Pérez-Patiño C, Barranco I, Casao A, Cebrián-Pérez JA, Muñio-Blanco T, Martínez EA & Roca J. (2016) The melatonin concentration in boar seminal plasma: a predictive in vivo fertility marker? *Anim Reprod Sci* 169, 131.
- Petrunkina AM, Waberski D, Bollwein H & Sieme H. (2010) Identifying non-sperm particles during flow cytometric physiological assessment: a simple approach. *Theriogenology* 73, 995–1000.
- Petters RM & Wells KD. (1993) Culture of pig embryos. *J Reprod Fertil* 48, 61–73.
- Ramió L, Rivera MM, Ramírez A, Concha II, Peña A, Rigau T & Rodríguez-Gil JE. (2008) Dynamics of motile-sperm subpopulation structure in boar ejaculates subjected to “in vitro” capacitation and further “in vitro” acrosome reaction. *Theriogenology* 69, 501–512.
- Ramió-Lluch L, Fernández-Novell JM, Peña A, Colás C, Cebrián-Pérez JA, Muñio-Blanco T, Ramírez A, Concha II, Rigau T & Rodríguez-Gil JE. (2011) In vitro capacitation and acrosome reaction are concomitant with specific changes in mitochondrial activity in boar sperm: evidence for a nucleated mitochondrial activation and for the existence of a capacitation-sensitive subpopulational structure. *Reprod Domest Anim* 46, 664–673.
- Ramió-Lluch L, Yeste M, Fernández-Novell JM, Estrada E, Rocha L, Cebrián-Pérez JA, Muñio-Blanco T, Concha II, Ramírez A & Rodríguez-Gil JE. (2014) Oligomycin A-induced inhibition of mitochondrial ATP-synthase activity suppresses boar sperm motility and in vitro capacitation achievement without modifying overall sperm energy levels. *Reprod Fertil Dev* 26, 883–897.
- Rao MV & Gangadharan B. (2008) Antioxidative potential of melatonin against mercury induced intoxication in spermatozoa in vitro. *Toxicol In Vitro* 22, 935–942.
- Rath D, Long CR, Dobrinsky JR, Welch GR, Schreier LL & Johnson LA. (1999) In vitro production of sexed embryos for gender preselection: high-speed sorting of X-chromosome-bearing sperm to produce pigs after embryo transfer. *J Anim Sci* 77, 3346–3352.
- Rathi R, Colenbrander B, Bevers MM & Gadella BM. (2001) Evaluation of in vitro capacitation of stallion spermatozoa. *Biol Reprod* 65, 462–470.
- Reiter RJ, Tan DX, Osuna C & Gitto E. (2000) Actions of melatonin in the reduction of oxidative stress - A review. *J Biomed Sci* 7, 444–458.
- Reiter RJ, Tan DX, Manchester LC, Paredes SD, Mayo JC & Sainz RM. (2009) Melatonin and reproduction revisited. *Biol Reprod* 81, 445–456.
- Rodríguez-Gil JE & Bonet S. (2016) Current knowledge on boar sperm metabolism: comparison with other mammalian species. *Theriogenology* 85, 4–11.
- Sarabia L, Maurer I & Bustos-Obregon E. (2009) Melatonin prevents damage elicited by the organophosphorous pesticide diazinon on mouse sperm DNA. *Ecotoxicol Environ Saf* 72, 663–668.
- Tait SWG & Green DR. (2012) Mitochondria and cell signalling. *J Cell Sci* 125, 807–815.
- Talevi R, Zagami M, Castaldo M & Gualtieri R. (2007) Redox regulation of sperm surface thiols modulates adhesion to the fallopian tube epithelium. *Biol Reprod* 76, 728–735.
- Wu JT, Chiang KC & Cheng FP. (2006) Expression of progesterone receptor(s) during capacitation and incidence of acrosome reaction induced by progesterone and zona proteins in boar spermatozoa. *Anim Reprod Sci* 93, 34–45.
- Yakirevich EMD & Naota Y. (1999) Characterization of a potent sperm-agglutinating monoclonal antibody and its cognate antigens. *Fertil Steril* 71, 502–510.
- Yeste M, Flores E, Estrada E, Bonet S, Rigau T & Rodríguez-Gil JE. (2013) Reduced glutathione and procaine hydrochloride protect the nucleoprotein structure of boar spermatozoa during freeze-thawing by stabilising disulfide bonds. *Reprod Fertil Dev* 25, 1036–1050.
- Yeste M, Estrada E, Pinart E, Bonet S, Miró J & Rodríguez-Gil JE. (2014) The improving effect of reduced glutathione on boar sperm cryotolerance is related with the intrinsic ejaculate freezability. *Cryobiology* 68, 251–261.
- Yeste M, Estrada E, Rocha LG, Marín H, Rodríguez-Gil JE & Miró J. (2015a) Cryotolerance of stallion sperm is related to ROS production

and mitochondrial membrane potential rather than to the integrity of sperm nucleus. *Andrology* 3, 395–407.

Yeste M, Fernández-Novell JM, Ramió-Lluch L, Estrada E, Rocha LG, Cebrián-Pérez JA, Muiño-Blanco T, Concha II, Ramírez A & Rodríguez-Gil JE. (2015b) Intracellular calcium movements of boar spermatozoa during in vitro capacitation and subsequent acrosome exocytosis follow a multiple-storage place, extracellular calcium-dependent model. *Andrology* 3, 729–747.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information section at the end of the article.

Figure S1 Effects of melatonin on the percentage of viability of boar spermatozoa subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis.

Figure S2 Effects of melatonin on the percentage of true acrosome exocytosis of boar spermatozoa subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis.

Figure S3 Effects of melatonin on the percentage of cells with capacitation-like membrane lipid disorder of boar spermatozoa subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis.

Figure S4 Effects of melatonin added at 0 h on tyrosine phosphorylation levels of P32 protein in boar sperm subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis.

Figure S5 Effects of the addition of melatonin to CM at 4 h on tyrosine phosphorylation levels of the P32 protein in boar spermatozoa subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis.

Figure S6 Images showing the effect of 1 μM melatonin on the formation of cell agglutinations in boar sperm subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis.

Figure S7 Effects of melatonin (1 μM) on the distribution of WGA lectin in boar sperm subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis.

Figure S8 Effects of 1 μM melatonin on the distribution of STL lectin in boar sperm subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis.

Figure S9 Effects of 1 μM melatonin on the distribution of PSA lectin in boar sperm subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis.

Figure S10 Effects of 1 μM melatonin on the distribution of PNA lectin in boar sperm subjected to in vitro capacitation and subsequent progesterone-induced acrosome exocytosis.

Figure S11 Examples of sperm-oocyte complexes considered as monospermic and polyspermic.