

This is a pre-copyedited, author-produced version of an article accepted for publication in Biology of reproduction (Ed. Oxford University Press) following peer review.

The version of record:

Gòdia, M. et al *A history of why fathers' RNA matters* in Biology of reproduction, vol. 99, issue 1 (July 2018), p. 147-159

Is available online at: DOI 10.1093/biolre/loy007

**Title:**

A History of Why Fathers' RNA Matters

**Running title:**

Population, roles, and plasticity of sperm RNAs

**Summary sentence:**

A historical perspective of the discovery and delivery of sperm RNAs that initiate development is presented. How these RNAs provide a record of, and components essential to fertility, embryo development, and modulating offspring's phenotype are discussed.

**Keywords:**

sperm RNA, recurrent miscarriage, transgenerational epigenetics, allostatic load

**Authors and affiliations:**

Marta Gòdia<sup>1</sup>, Grace Swanson<sup>2,3</sup>, Stephen A. Krawetz<sup>2, 3, 4\*</sup>

1. Animal Genomics Group, Center for Research in Agricultural Genomics (CRAG) (CSIC-IRTA-UAB-UB), Cerdanyola del Vallès (Barcelona), Catalonia, Spain.
2. Department of Obstetrics and Gynecology, Wayne State University, Detroit, Michigan.
3. Center for Molecular Medicine and Genetics, Wayne State University, Detroit, Michigan.

4. C.S. Mott Center for Human Growth and Development, Wayne State University, Detroit, Michigan.

\*Corresponding Author. Tel: +1 313 577 6770; Fax: +1 313 577 8554; C.S. Mott Center for Human Growth and Development, Wayne State University School of Medicine, 275 E. Hancock, Detroit, MI, 48201

Email: steve@compbio.med.wayne.edu

***Grant support:***

M.G. was funded with a FPI Ph.D. grant from the Spanish Ministry of Education (BES-2014-070560) and a Short-Stay fellowship from the Spanish Ministry of Education (EEBB-I-2017-12229) at S. A. K. laboratory. G.S. is supported in part by a Postdoctoral Recruiting Fellowship from Wayne State University. Support from the Charlotte B. Failing Fetal Therapy and Diagnosis endowed Professorship (along with the EMD Serono 2016 Grant for Fertility Innovation) to S.A.K. is gratefully acknowledged.

**Abbreviations**

*ACTB* *Actin beta*

*AGO* *Argonaute*

*BMI* *Body mass index*

*CARM1* *Histone-arginine methyltransferase CARM1*

*CD45* *Leukocyte common antigen*

circRNA circular RNA

DGCR8 DiGeorge Syndrome Critical Region Gene 8

Dicer Endoribonuclease Dicer

*DNMT1* DNA methyltransferase 1

*DMNT3* DNA methyltransferase 3

*E2R* Ubiquitin Conjugating Enzyme E2 R2

EGA Embryonic genome activation

EGFP Enhanced green fluorescent protein

FISH Fluoresce *in-situ* hybridization

FPKM Fragments Per Kilobase of transcript per Million

*GAS5* Growth arrest specific 5

H3K27 Histone 3 lysine 27

H3K4 Histone 3 lysine 4

*Hdac11* Histone deacetylase 11

HPA Hypothalamic-pituitary-adrenal

*IAP* Intracisternal A Protein

ISH *In Situ* Hybridization

*ITGB* Integrin subunit beta 1

*Kit* KIT proto-oncogene receptor tyrosine kinase

*LGC1* Low glutelin content 1

LINE Long interspersed nuclear element

LINE1 Long interspersed element 1

*Lnc2* lncRNA transcript 2

*Lnc3* lncRNA transcript 3

lncRNA long non-coding RNA

LTR Long terminal repeat

MII egg Meiosis II egg

$m^5C$  5-methylcytidine

$m^2G$  N<sup>2</sup>-methylguanosine

MAR Matrix attachment region

MERVL Mouse Endogenous retroElement

*MILI* Piwi like RNA-mediated gene silencing 2

*MIWI* Piwi like RNA-mediated gene silencing 1

*MIWI2* Piwi like RNA-mediated gene silencing 4

miRNA microRNA

mRNA messenger RNA

*MYC* MYC Proto-Oncogene, BHLH Transcription Factor

NGS Next Generation Sequencing

PARN poly(A)-specific ribonuclease

piRNA Piwi-interacting RNAs

Piwi *P* element-induced *wimpy* testes

PLCZ Phospholipase C zeta

PNLDC1 PARN like, ribonuclease domain containing 1

POU5F1 POU class 5 homeobox 1

*PRM1* Protamine 1

*PRM2* Protamine 2

*PTPRC* Protein tyrosine phosphatase, receptor type, C

*RASGRP1* RAS guanyl releasing protein 1

RISC RNA-induced silencing complex

RM Recurrent miscarriage

RNU RNA U small nuclear

RPKM Reads Per Kilobase of transcript per Million

rRNA ribosomal RNA

SAGE Serial analysis of gene expression

SINE Short interspersed nuclear element

siRNA small interfering RNA

sncRNA small non-coding RNA

sno75 piwi RNA 75

SOX2 Sex-determining region Y box 2

RE RNA element

TCEP Tris (2-carboxyethyl) phosphine hydrochloride

TE Transposable element

*TNP2* Nuclear transition protein 2

tRNA transfer RNA

*TRAIL* Tumor necrosis factor (TNF) superfamily member 10

3' UTR 3' untranslated region

**Abstract:**

Having been debated for many years, the presence and role of spermatozoal RNAs is resolving, and their contribution to development is now appreciated. Data from different species continues to show that sperm contain a complex suite of coding and non-coding RNAs that play a role in an individual's life course. Mature sperm RNAs provide a retrospective of spermatogenesis, with their presence and abundance reflecting sperm maturation, fertility potential, and the paternal contribution to the developmental path the offspring may follow.

Sperm RNAs delivered upon fertilization provide some to the initial contacts with the oocyte, directly confronting the maternal with the paternal contribution as a prelude to genome consolidation. Following syngamy, early embryo development may in part be modulated by

paternal RNAs that can include epididymal passengers. This provides a direct path to relay an experience and then initialize a paternal response to the environment to the oocyte and beyond. Their epigenetic impact is likely felt prior to embryonic genome activation when the population of sperm delivered transcripts markedly changes. Here, we review the insights gained from sperm RNAs over the years, the subtypes, and the caveats of the RNAs described. We discuss the role of sperm RNAs in fertilization and embryo development, and their possible mechanism(s) influencing the offspring's phenotype. Approaches to meet the future challenges as the study of sperm RNAs continues, will include, among others, elucidating the potential mechanisms underlying how paternal allostatic load, the constant adaptation of health to external conditions, may be relayed by sperm RNAs to affect a child's health.

### ***Introduction***

Although spermatozoa were thought for many years to merely contribute their half of the genome to the offspring, it is now appreciated that sperm delivers its entire structure, from which selective components are used to build a healthy child [1]. These include, phospholipase C zeta (PLC $\zeta$ ) and other factors that yield a pulsatile  $\text{Ca}^{2+}$  response, and in humans, the sperm centriole organizing center [reviewed in 2]. In contrast, certain structures including the mitochondria are ubiquitinated and targeted for degradation [reviewed in 2]. The suite of RNAs that a sperm carries is also part of the package. The RNAs can reflect the fidelity of spermatogenesis that impacts embryo development [3, 4], that, in turn, may affect offspring phenotype [5-8]. This has prompted the field to begin to dissect their role in the fetal origins of adult disease, a concept laid out in the Barker hypothesis [9] that continues to be tested as our environment changes. Health is maintained and reflects a biological endpoint termed allostatic load [10], that is both directed by, and responds to, past and present events [11, 12]. A system overload, even by one too many inputs, leaves the potential for adverse health effects to arise.

The relationship between the allostatic load and offspring health has recently been implicated in epidemiological studies and recapitulated in mouse models [13-15], with sperm RNAs as potential messengers between generations.

In this review we will provide a historical perspective of sperm RNAs, and discuss the characteristics of this unique population and the caveats of studying sperm RNAs. Their contribution to the oocyte and their role in embryo development is considered. We conclude with a discussion of how sperm RNAs may respond to their environment epigenetically relaying the father's experience to the offspring.

### ***History of sperm RNAs***

Sperm RNAs had been considered an artifact and if present, as having no role in fertilization and obviously not embryo development. The hypothesis was founded upon the extrusion of RNAs from the cell as part of the residual body during spermatogenesis and their lack of intact ribosomal RNA (rRNAs) [16]. This led to the conclusion that if any RNA remained, it was residual, possessing no function in the absence of a complete transcript. This has not stood the test of time. With marked technological advances that the field has experienced over the years, the presence and some of the roles of sperm RNAs (Figure 1) are now established with possibly others on the horizon.

Early studies suggested that spermatozoal RNAs were present in the epididymis and ductus deferens from mouse sperm, but were dismissed as reflective of mitochondrial contamination [17]. It was not until the late 1980's, that the presence of sperm RNAs was beginning to be considered by various independent techniques e.g., using immunogold staining in ferns [18], and finally in humans and rats by RNase colloidal gold. This provided an estimate of 70 and 100 fg RNA per cell respectively, with the majority primarily localized within the nucleus [19]. RT-

PCR and *In Situ* Hybridization (ISH) reconfirmed these observations in human sperm with the detection of *MYC* [20]. However, the field was still not convinced attributing these observations to spurious hybridization or priming events. Several independent studies unknown to each other were undertaken in the 1990s that addressed the issue of the presence of RNAs in sperm using various techniques. These included RT-PCR, ISH and RT-qPCR, following the exclusion of samples with genomic DNA or contamination by somatic cell RNAs. Detection of *PRM2*, *E2R*, and *ACTB* by RT-PCR [21], *ITGB1* via PCR [22], and *PRM1*, *PRM2*, *TNP2* in humans [23] and mice [24] by ISH, as well as *LGC1* by ISH and RT-PCR in lily plants was observed [25]. It was not until 1999, when Miller, Krawetz and colleagues attempted to characterize human sperm RNA transcripts at the sequence level [26]. A group of translationally quiescent RNAs, the majority of which were derived from repetitive elements with the near or complete absence of 28S and 18S rRNAs were revealed [26]. This was a curious observation in a transcriptionally inactive sperm cell [16, 27]. Translation of nuclear encoded RNAs by mitochondrial ribosomes in mature sperm during sperm capacitation has been proposed [28, 29], but these observations still await independent confirmation.

Microarray analysis that followed provided the first characterization of sperm mRNAs from healthy human donors [30]. A set of transcripts shared between sperm samples functionally enriched for spermatogenesis and early development were resolved, suggesting that the retention and presence of certain RNAs was not stochastic [30]. In maize (*Zea mays*), a set of sperm RNA transcripts were identified by cDNA sequencing with some derived by selective partitioning [31]. Delivery of human sperm RNAs to the oocyte was validated in the hamster penetration assay opening the door to a role in early embryonic development, and the male assuming a greater role in the birth of a healthy child [1]. Antisense and micro RNAs (miRNAs) were also identified amongst this population, and thus inferred to be delivered to the oocyte

upon fertilization. This cemented the foundation for the beginnings of mechanistic proposals regulating parental gene activity through targeting paternal or maternal RNAs, thereby regulating early genomic events in the embryo [32, 33]. Characterization of the human sperm transcriptome by Serial Analysis of Gene Expression (SAGE), i.e., tag sequencing [34], identified a series of highly abundant transcripts in fertile donors, enriched for roles related to spermatogenesis, sperm function, fertility, and conception [35], in accordance with previous microarray studies [30]. This further encouraged interest to assign potential roles for sperm RNAs. Microarray analysis was beginning to lay a path towards developing diagnostic strategies to understand male infertility [3, 4, 36].

Advances continued over the years and in 2005, shortly after the identification of miRNAs in human sperm by microarrays [33], miRNAs were observed in mice using a microarray and RT-qPCR approach [37]. Other small non-coding (snc) RNAs like Piwi-interacting RNAs (piRNAs) were subsequently detected in mouse testis by pyrosequencing [38] but left their description in sperm wanting. The low abundance of paternal miRNAs in zygotes led to the assumption they possessed a limited role, if any, in fertilization [37]. Yet, Rassoulzadegan and colleagues provided the first study that suggested that miRNAs could influence offspring phenotype, a phenomenon known as paramutation [39]. As with most studies in this field, this initial foray into sperm RNA mediated transgenerational epigenetics was controversial and met with skepticism by others, but this concept has also withstood the test of time.

In recent years, several compelling independent studies have corroborated the initial findings of Rassoulzadegan and colleagues that epigenetic modifications can be transmitted from father to offspring via paternal RNAs [reviewed in 13, 40, 41, 42]. Various components or stressors including mental and physical stress, induced by the physical environment, toxins, or diet that together constitute the allostatic load experienced by the father have been individually well-

studied in mice and rats [5, 40, 43]. Both miRNAs [7] and piRNAs [6] have been implicated in paternal stress studies and have provided a framework. Similarly, paternal diet can coincide with alterations in the abundance of miRNAs and transfer RNAs (tRNAs) leading to transgenerational effects whether they are subject to a high-fat-high-sugar diet [13], a high fat diet [5], or a low protein diet [8]. Still, the results of these studies, especially in humans require further study (see: “RNAs and Transgenerational Epigenetic Inheritance”).

Next Generation Sequencing (NGS) provided a boost to RNA-seq studies early in the first decade of the millennium. The first sncRNA-seq study performed in human sperm showed the presence of several sncRNAs including miRNAs, piRNAs, and repeat-associated small RNAs [44]. Spurred on by this work, sncRNA-seq in mouse sperm followed, that also reported a highly enriched fraction of tRNAs [45]. Several sperm RNA-seq studies have now characterized the population of transcripts found in sperm including human [46], mouse [47], and agricultural livestock species such as bovine [48, 49] and stallion [50]. Moreover, an in-depth study in mature human sperm has shown that the ribosomal RNAs that remain, while abundant, are fragmented [51]. The continually decreasing cost of sequencing has enabled the RNA-seq approach to become widely adopted and is now a cornerstone of medicine and agricultural research [52, 53] and is becoming the diagnostic standard. Nevertheless, sperm NGS techniques present several intricacies in comparison to somatic or other germinal cells. Hence, experimental design and analyses are crucial for reliable interpretation that yield meaningful results.

### ***Methods, techniques and caveats when considering sperm RNA***

The unique characteristics of sperm cells compared to somatic cells necessitates careful consideration as a new study is conceived or secondary analysis of a publicly available dataset

is undertaken. The low concentration of sperm RNAs within each cell must be considered along with the technology that has been employed. Human sperm carries approximately 50 fg of long RNA (> 200 nt) and 0.3 fg of sncRNAs (< 200 nt) per sperm cell, which is very similar to early estimates [19]. However, this is dwarfed in comparison to somatic cells which contain 10 pg of long RNA and 1-3 pg of sncRNAs [54-56], and necessitates the use of highly purified samples to eliminate maturing cells, somatic cells, and genomic DNA that can obscure the results.

Purification protocols have been developed to separate mature spermatozoa from seminal plasma, as well as immature sperm cells, leukocytes, epithelial cells, and bacteria. Three different approaches have generally been employed. The first, swim-up or sperm migration, in which 0.5 – 1 ml of semen is placed in a 45° angled centrifuge tube under a medium salt solution and incubated at 37°C for 60 minutes [57]. The sperm cells swim out of the semen to the medium, where they are aspirated using a sterile pipette. This approach selects motile sperm, however, the number of spermatozoa recovered can be low [57]. The second, density gradient centrifugation, uses an isotonic salt solution with saline-coated silica particles, such as PureSperm® (Nidacon), to separate spermatozoa according to density [58]. In most mammalian studies, the starting concentration used is 50% [54]. As with the swim-up method, this approach favors motile and morphologically normal spermatozoa [58], but the number of spermatozoa recovered is significantly higher by this method [59]. The third method employs a somatic cell lysis buffer that typically contains 0.1% SDS and 0.5% Triton (X-100). This effectively lyses somatic cells, and leaves the sperm head intact [30, 60]. While this method is effective in eliminating somatic cells, this treatment has been proven to compromise the midpiece, and can solubilize sperm-membrane structures, with preferential loss of mitochondrial RNAs [30, 58, 60]. Although a higher recovery rate was observed, RNA yield was significantly lower than density gradient purification [58]. Optical microscopy to visually confirm the lack of immature sperm, somatic or bacterial cells is typically used as an initial screen.

Species specific protocol optimization of sperm RNA extraction has proven useful once RNase free reagents have been confirmed as RNase free. Extraction methods generally include a homogenization and cell disruption using a chaotropic and reducing agent, e.g., TRIzol<sup>TM</sup> and TCEP (Tris (2-carboxyethyl) phosphine hydrochloride), typically followed by commercial RNA isolation protocols [58, 61, 62]. A final DNase treatment step is requisite for eliminating residual genomic DNA. Several strategies to confirm spermatozoal recovery have been employed. For example, following sperm RNA extraction, the absence of intact rRNAs 18S and 28S as assessed by a Bioanalyzer trace would be consistent with their targeted removal during spermatogenesis [51]. The presence of transcripts from *CD45/PTPRC* (expressed in most somatic cells) is typically used as a somatic marker. RT-PCR and/or RT-qPCR quantitation can simultaneously assess the presence of genomic DNA when intron spanning primers are used. The ultimate test for sperm RNA purity is by comparative sequence analysis [63].

The construction of high throughput sperm RNA sequencing libraries is confounded by several factors including selective fragmentation of coding transcripts [46] as well as residual fragmented rRNAs [51] and low total RNA yield [54]. While typical RNA-seq library construction protocols require 1 µg of total RNA, sperm only yields between 30 - 80 ng of total RNA per million cells [54-56]. In most tissue samples, poly(A<sup>+</sup>) selection has been used to selectively enrich the population of mRNAs as compared to mitochondrial and rRNAs. RNA-seq library construction using poly(A<sup>+</sup>) selection can result in 3' bias due to the exclusion of all short polyadenylated tailed transcripts, those lacking this modification, and nonpolyadenylated transcripts or the majority of the other segments from fragmented transcripts [55]. A random amplification strategy can, somewhat correct for potential 3' bias. Ribosomal RNA depletion or primer design can also be used to reduce the number of sequencing reads assigned to this biotype. Different commercial library preparation kits and methods have been tested in various

species, including human [64], bovine [48], and porcine (Gòdia M, Quoos Mayer F, Nafissi J, Catelló A, Rodríguez-Gil JE, Sánchez A, and Clop A, unpublished results), resulting in different outcomes with preferred protocols being developed. When comparing two studies, it is important to remember that the data obtained will vary in terms of the library preparation protocol [64]. In addition, quantification of sperm RNAs from sequencing data is dependent on the approach used to analyze the data and this can vary widely. Bioinformatics analysis can follow two different paths. The most commonly used method is “directed analysis”. After read mapping, bioinformatic analyses are used to quantify the abundance of each of the annotated genes from the genome studied. This can be refined to quantify RNA levels for each of the annotated transcripts, enabling the detection of alternatively spliced isoforms, but this still remains a challenge when multiple isoforms are simultaneously expressed. Transcript quantitation is measured by FPKM (Fragments Per Kilobase of transcript per Million), when a paired-end sequencing approach is employed or RPKM (Reads Per Kilobase of transcript per Million), when single-end sequenced. These measurements consider the total number of reads mapped to the transcript after adjusting for the full length of the gene/transcript and applying a library size (sequencing depth) correction. This analytical strategy has been widely used in sperm RNA profiling as it provides a basic way forward to direct transcript comparison with other tissues and/or between species [46-50, 64]. However, as substantial portion of the sperm transcripts are fragmented [46], all exons may not be represented in an equivalent manner and depending on the question posed, this might obscure the results. “Discovery” strategies that utilize a biologically meaningful minimal unit of detection can provide an unbiased alternative. This approach considers sperm RNAs as a collection of sperm RNA elements (REs) [sperm REs: 53]. REs are short sized sequences, formed by a set number of reads and joined as a contig, independent of their annotation. Resultant REs are considered independent of each other and are annotated by genomic location as exonic, intronic, intergenic, close to exonic

regions, or novel as existing within an unannotated region (Figure 2). This approach permits a comprehensive examination [53] and addresses low quality genome annotations in some species.

### ***The population of Sperm RNAs***

Mature spermatozoal RNAs represent a wide range of coding and non-coding RNAs, including long and short non-coding RNAs. This rich number and classes of RNAs are implicated in different roles, including spermatogenesis, regulation, fertilization, embryo development, and offspring phenotype [reviewed in 53, 55, 65]. Early microarray studies showed that a vast majority of transcripts were shared among healthy sperm donors, enriched in sequences reminiscent of past function [30], suggesting that their distribution among sperm was not stochastic. Human sperm RNA-seq enabled a more complete understanding of the unique features of the sperm transcript population [46]. Over 22,000 transcripts were detected with approximately 700 of which were moderate to high abundance [46]. These RNAs present a complex heterogeneous profile of intact and variously fragmented transcripts [46]. As shown in human [46] and other species including mice [47], horse [50], and bovine [48, 49], a substantial proportion of RNAs are fragmented. Nevertheless, like sncRNAs, they might also function in sperm maturation and the initial events as part of and following fertilization. Integrative analysis between human RNA-seq datasets of testes, mature sperm, and seminal fluid has provided additional insight that highlights the importance of precise transcriptional regulation during spermatogenesis, to maintain the integrity of gamete generation and maturation [66]. The presence or absence of shared RNAs and corresponding proteins between the different tissue types, is indicative of cross-talk between the cellular and extracellular environment from the prostate, epididymis, and seminal vesicles [66].

### **Long non-coding RNAs**

Long non-coding RNAs (lncRNAs) can modulate both transcriptional and posttranscriptional processes [reviewed in 55], while often serving a nuclear structural role [67]. Approximately 1/3 of sperm RNAs are confined within the bounds of the perinuclear theca encapsulated nucleus [52]. The complex packaging of the paternal genome by protamines alongside nucleosome-bound DNA form a unique chromatin structure. In part, chromatin remodeling has been attributed to lncRNAs that directly complex with chromatin [41] as a function of its affinity to the DNA or DNA-binding proteins. Interaction with target sites through long-range chromatin interactions exemplify a means of regulation often mediated by their 3D organization [reviewed in 68, 69]. Interestingly this structure appears in close association with a repertoire of RNAs [67], positioned by chromosomal regions attached to the nuclear matrix by the Matrix Attachment Regions (MARs) [70].

Several classes of lncRNAs have been observed in human [46, 53] and mouse [71] sperm. This includes members of the RNA U Small Nuclear (RNU) family. These components of the sperm spliceosome are intact and very abundant [46], perhaps suggestive of an early post fertilization role. Mouse sperm and testes are rich in lncRNAs of which *Lnc2* and *Lnc3* are the most abundant as evidenced by Fluoresce *In-Situ* Hybridization (FISH), RT-PCR, and RT-qPCR [71]. Interestingly they appear enriched in sperm when compared with testes [71]. lncRNA profiles in both testes and sperm can be affected by environmental exposures, e.g., cadmium, that impacts spermatogenesis and male fertility [72]. This is consistent with the view that lncRNAs modulate target genes that play a regulatory role in spermatogenesis, fertility, and embryo development [71-74].

Other types of non-coding RNAs have been described in germinal cells. These include the intronic retained elements and the recently discovered circular RNAs (circRNAs). Intronic

retained elements up to full-length introns have been reported in sperm, with more than 200 distinct Res of this class identified [46]. They appear more abundant in sperm than in testes [reviewed in 55]. Although suggested as retained in mature sperm it has yet to be resolved as to how they evade degradation [reviewed in 55]. circRNAs are stable 3' and 5' covalent linked non-coding RNAs formed from the same transcribed segment that are inherently resistant to exonuclease degradation [75] thus evading degradation. The function of circRNAs is origin dependent. Exonic circRNAs are primarily located in the cytoplasm, thought to act as "sponges", and counteract mRNA repression by miRNAs. Nuclear localized intronic circRNAs and exon-intron circRNAs, are thought to regulate their parental genes through direct cis/trans interaction [reviewed in 75, 76]. They have recently been detected in testes and seminal plasma suggestive of a role in gamete generation [77]. Gene Ontology of testes circRNAs shows enrichment for genes related to spermatogenesis, sperm motility, and fertilization [77]. Their stability and presence in seminal plasma may provide a source of infertility biomarkers. Some of the sperm RNA elements described in previous studies [46, 53, 56] warrant further consideration as they do possess characteristic circRNA signatures.

### ***Small non-coding RNAs***

Small non-coding RNAs encompass a variety of different RNA types that play crucial roles in the maintenance and function of the germline genome [reviewed in 42, 55, 78]. Major functions may be classified by RNA type and include regulation of gene expression by miRNAs [44], defense against transposable, repetitive elements or viruses by small interfering RNAs (siRNAs) [79] or piRNAs [80], and protein synthesis and signal modulation by tRNAs and their fragments [5, 45]. Together they can play a role in genome structure and integrity. Characterization of the population of sncRNAs from human and mouse has revealed their diversity, and provided a

glimpse into their roles in spermatogenesis, early embryo development, and how they may modify the genome to heritably affect offspring [44, 45].

### ***miRNAs***

The most well characterized sncRNAs are miRNAs, contributing to approximately 7% of the total sncRNAs to the fertile human sperm profile [44]. They play a crucial role in spermatogenesis and fertility, and have been reported to modulate expression during the different stages of sperm maturation [reviewed in 81]. miRNAs are an integral part of the RNA-induced silencing complex (RISC) and together with the AGO proteins generally target the 3' UTRs of mRNAs through sequence complementary to repress mRNA expression through degradation or activation [reviewed in 55]. On one hand, miR-140, miR-21, miR-152, and miR-148a have been shown to repress expression of RNAs encoding epigenetic modifiers, e.g., including *DMNT3* and *RASGRP1*. This is consistent with their absence in mature sperm [46]. On the other hand, *DNMT1* transcripts are present in sperm [46], concordant with their epigenetic role suggesting they escape this form of repression. Specific paternal origin miRNAs such as miR-34c have also been described [44]. Attempts to study its role have led to different conclusions. While *in vitro* they appear required for the first cleavage following fertilization [82], *in vivo*, it was not essential for fertilization or embryo development but was crucial for spermatogenesis, as its absence disrupted spermatogenesis leading to murine infertility [83].

### ***siRNAs***

Small interfering RNAs, also known as RNA-mediated interference RNAs, are active mediators of transcriptional and post-transcriptional gene-silencing [reviewed in 84]. As with miRNAs, siRNAs act in concert with RISC and AGO proteins to target complementary RNAs for translational inhibition or degradation [reviewed in 84]. siRNAs are primarily known to function in the host defense against transposable elements (TE) and RNA viruses, and aid in the

maintenance of heterochromatic DNA [reviewed in 84]. As shown in plants, somatic cell TE-derived siRNAs migrate into sperm cells contributing to TE silencing prior to fertilization [reviewed in 85]. Preliminary work on the role of siRNAs in fertilization using Dicer knockout mice, presented an altered profile of siRNAs in spermatozoa (and miRNAs), where sperm microinjection in wild-type oocytes resulted in embryos with reduced developmental potential [79]. Other independent studies have attempted to discern the role between siRNAs and miRNAs in spermatogenesis. While miRNAs require DICER and DGCR8 proteins for maturation, siRNAs only require DICER [reviewed in 86]. Results have shown that a conditional *Dicer* knockout presents severe sperm morphological defects as compared to the conditional *Dgcr8* mice mutant, implicating siRNAs in mammalian spermatogenesis [86, 87].

### ***piRNAs and Transposable Elements***

Piwi interacting RNAs are specialized RNAs that interact with Piwi (*P* element-induced *wimpy testes*) proteins, a gonad type of AGO proteins, which mediate RNA silencing of transposable elements [reviewed in 88, 89]. Murine Piwi proteins MIWI, MILI, and MIWI2, are essential to spermatogenesis and their absence is associated with male infertility [reviewed in 78, 88]. In humans, ~17% of the sncRNAs correspond to piRNAs [44]. Their presence still remains controversial although piRNAs are now beginning to be considered by others [90]. Altered levels of PIWI-Like 1 and 2 mRNAs have now been detected in men with decreased sperm count and motility by RT-qPCR [91]. The PIWI-Like 1 RNA remains essentially intact in mature spermatozoa as determined by RNA-Seq [53, data not shown]. Given these independent observations, their presence will likely become accepted.

In murine spermatocytes and spermatids, the vast majority of piRNAs map to specific genomic regions [reviewed in 78], and ~17% map to repeat sequences such as DNA transposons, Short Interspersed Nuclear Elements (SINEs), and Long Interspersed Nuclear Elements (LINEs) [reviewed in 89]. This is not surprising since TEs constitute a large fraction of the eukaryotic genome. SINES, LINEs, and Long Terminal Repeats (LTRs) are the most abundant [44]. LINE1 is the most common and well-studied retrotransposon in germ cells [reviewed in 88]. piRNA biogenesis is regulated by the PARN family proteins [92]. The role of one of its members, PNLDC1 has been examined *in vivo* [93]. In the corresponding mouse knockout model, LINE1 retrotransposon silencing was disrupted leading to aberrant piRNA biogenesis and spermatogenesis [93]. Studies in mutant Piwi proteins have shed light into their roles in spermatogenesis. *Miwi2* knockout mice showed spermatogenic arrest, increased LINE1 retrotransposon expression, and loss of methylation in testes, suggesting possible regulatory roles of Piwi proteins and piRNAs in methylation [80]. Moreover, *Mili* mice knockout mutants exhibited loss of DNA methylation of the LINE1 element and an increased expression of both LINE1 and *IAP* [94]. Interestingly, the piRNAs produced from lncRNAs have the capability to act as histone modifiers. For example, the piRNA sno75, derived from the lncRNA GAS5, has been shown to increase transcription of *TRA1L* by guiding H3K4 methylation and H3K27 demethylation [95]. Transposable elements can be regulated by DNA methylation [reviewed in 96], which may also be guided and modulated by RNAs [reviewed in 68]. This DNA methylation is maintained by DNMT1 [reviewed in 96], whose transcript is present in sperm [44]. It has also been suggested that piRNAs play a central role in the confrontation and consolidation when the sperm and oocyte meet to initially assess genetic compatibility through their interaction with repetitive elements [reviewed in 55] (see: “Contributions from sperm to the oocyte and early development”).

### ***tRNAs***

Transfer RNAs and their fragments are some of the more abundant sperm sncRNAs [reviewed in 97]. Their abundance is directly linked to the general translational needs of a given cell type rather than to a specific gene, reflective of their metabolic state. The original description of tRNAs in mouse sperm showed an enrichment of 5' end of tRNAs [45]. While generally scarce in testis, their abundance increases as the maturing sperm exits from testis as it passes through the caput, corpus and cauda in [8]. This has supported the view that the majority of tRNA fragments in mature spermatozoa arise from trafficking via epididymosomes from the epididymis to the sperm [8], with some up-regulated in mice fed a high fat diet [5].

In addition to fragmentation that yields 5' enrichment of mouse sperm tRNA fragments [45], modifications have also been reported [5]. These include the incorporation of 5-methylcytidine ( $m^5C$ ) in response to high or low fat diets [5] which may act to increase their stability. This chemical post-transcriptional modification may provide a mechanism to signal current metabolic state reflecting a change in environment [97]. It has also been argued that tRNAs fragments may act to repress embryonic Mouse Endogenous Retroelements (MERVLs). This is particularly attractive since some are in close proximity to genes expressed during the early stages of embryo development. It has thus been proposed that they regulate a significant proportion of the transcriptome in development [reviewed in 98], perhaps affecting placental size by altering metabolic pathways [8].

### ***Contributions from sperm to the oocyte and early development***

Upon fertilization the spermatozoon and oocyte confront one another for the first time towards consolidation [2, 99]. During this initial event when the ooplasm is exposed to the paternal

contribution it is also potentially exposed to a host of retroviruses and retrotransposons that have the potential to alter the zygotic genome [99]. They can insert within active regions of the genome as it is rearranged and/or affect the expression of neighboring genes by functioning as an enhancer, altering splicing or polyadenylation [100]. LINE1 transcription affects embryo chromatin accessibility in mice, and may play a role in coordinating the activity of multiple genes throughout the genome adding to genomic stability for the embryo [98]. Yet, in contrast to MERVLs, sperm LINE1 elements do not contain genes implicated in embryonic genome activation (EGA) [98]. By a mechanism of RNA activity through maternal and paternal sncRNAs, such as miRNAs and piRNAs [2, 44, 55], parasite RNAs are degraded, leading to a consolidated state of genetic compatibility [99]. Embryo development goes forward if these checkpoints are passed. While the potential contribution of repetitive elements in early development remains unclear, embryonic arrest at the 2 or 4-cell stage in mice coincides with the disruption of LINE1-encoded reverse transcriptase [reviewed in 101].

In humans, the transition to an embryo requires approximately 3 days and comprises the migration and fusion of the germ cell pronuclei and several cleavage divisions [reviewed in 102]. The human embryo is relatively transcriptionally silent until day 3. Having reached the 4-8 cell stage the major wave of human embryonic genome activation occurs [reviewed in 102, 103]. Following EGA, the embryo continues development, including implantation of the blastocyst into the uterine wall at day 7 [reviewed in 102]. Although humans and mice morphologically share embryo similarities, key differences necessitate that cross-species extrapolations be considered with caution. In mouse embryos, EGA occurs 26 – 29 h post-fertilization at or around the first cellular division so the transcriptionally silent period is substantially reduced [reviewed in 102]. Similarly, the timing of compaction is shorter in mice, and human embryos hold at least one extra round of cell division before implantation [reviewed in 102]. It is not surprising that marked

differences between gene expression, genome instability, and epigenetic modifications between human and mouse embryos are apparent [reviewed in 102]. While the quantity of paternal RNAs delivered by sperm to the oocyte may seem small compared to the oocyte, it is sufficient to play a role in transgenerational inheritance altering offspring phenotype [99, 104].

The role of sperm RNAs in EGA in mice has recently been examined *in silico* providing key elements for testing [105]. This provided a RNA-seq survey of sperm, MII eggs, and zygotes, supporting a potential role of paternal RNA or proteins to interact with maternal cofactors contributing to EGA [105]. A sperm RNA corresponding to an intragenic LTR of *Hdac11*, present in sperm but absent in MII eggs and zygotes, suggested that the paternally derived *Hdac11* LTR or others might complement maternal cofactors or pathways and participate together in EGA [105], not unlike the well-known oocyte activator factor PLCZ [106].

The first days of embryo development are critical as specification begins. Perturbation may resolve as the habitual first trimester pregnancy loss, which occurs in 15 - 25% of pregnancies [107]. Although most of the efforts to understand possible causes have focused on the mother reflecting the direct relationship with the fetus, the possible role of the male contribution has only begun to be discussed [108]. The factors from which sperm may contribute to miscarriage have been the focus of various studies. The influence of DNA fragmentation, aneuploidy, and integrity, as well as sperm morphology, have been considered to be correlated with Recurrent Miscarriage (RM), characterized as two or more consecutive failed pregnancies [107], yet, results remain controversial and inconsistent [109-112]. A recent study has focused on the possible role of sperm RNAs in RM etiology [108]. The ratio of protamine RNAs (*PRM1/PRM2*) was significantly different between spermatozoa when RM couples were compared to healthy donors suggesting that abnormal protamine packaging can negatively affect embryo development [108]. Whether this reflects histone retention and/or abnormal epigenetic marks

[108, 113] remains to be resolved. Interestingly, alterations in the protamine transcript ratio have also been associated with reduced sperm quality, including low concentration, reduced motility, abnormalities, and increased DNA fragmentation [114].

### ***RNAs and Transgenerational Epigenetic Inheritance***

The study of Rassoulzadegan and colleagues was the first report providing evidence of mammalian RNA-mediated epigenetic inheritance. Here, it was observed that wild type mouse offspring of *Kit* mutant parents, exhibited white patches characteristic of the *Kit* mutant phenotype coupled with altered RNA levels of *Kit* in sperm [39]. In the testis, a relatively dispersed population of RNAs was observed, including the identification of abnormal short mRNAs, derived from the wild type allele responsible for the *Kit* mutant phenotype. These abundant RNAs were only found in the mature sperm of heterozygotic offspring [39], suggesting that the increase was responsible for the disruption of wild type phenotype. To test this hypothesis, two miRNAs thought to target *Kit* mRNA, miR-221 and miR-222, were injected into one-cell embryos. The resultant mice exhibited the same *Kit* mutant phenotype, leading to the conclusion that the presence of certain miRNAs early in embryogenesis will result in a stable, heritable change in gene expression and correspondingly, phenotype [39]. This study also highlighted miRNAs as possible modifiers of the epigenome [39], that perhaps directed in some manner DNA methylation, the most widely recognized mechanism of epigenetic heritability [reviewed in 40].

Several sncRNAs can regulate DNA methylation and histone modifications [reviewed in 68] and, as above miRNAs, are known to influence DNA methyltransferases. CARM1, is an embryonic stem cell pluripotency factor involved in the H3 promoter methylation of two transcription factors,

POU5F1 and SOX2, providing an active chromatin mark upon induction [reviewed in 55]. miR-181c, is known to target *CARM1*, and its immature form, pri-miR-181c, is abundant in human sperm [46]. Upon delivery to the oocyte, pri-miR-181c is processed to its mature form that is coupled with a 70% decrease in these and 27 other *CARM1* associated target genes by the morula stage [46]. Recent results from a paternal chronic stress study identified that zygotic microinjection of 9 abundant miRNAs resulted in targeted degradation of maternal mRNA transcripts, including genes involved in chromatin remodeling [7]. These results infer a role for sperm miRNAs in ensuring or modifying cell specification.

Both multigenerational and transgenerational epigenetic inheritance imposed upon future generations have important ramifications for medicine and agriculture. The concept of fetal origins of adult health and disease [9] has given rise to numerous studies that illustrate the ability for the effects of an exacerbating input environment (allostatic overload [10]) in the father, to be inherited by later generations. While, male mediated multigenerational inheritance only passes from the affected father to child in a transient manner, transgenerational inheritance extends, affecting the father's grandchildren and/or beyond. Unlike multigenerational transmission, transgenerational inheritance can represent a permanent change to the gene pool [115]. Mechanistically, these responses to environmental factors, including stress, toxins, and diet may initially be mediated by epigenetic effectors, including RNAs. The ability of nucleic acids, including sncRNAs, to distribute throughout the body encapsulated in vesicles, provides the opportunity to influence sperm [reviewed in 115] that is normally protected from external effects by the blood-testes barrier. Data has also suggested that reverse transcription, by the LINE1 retrotransposon reverse transcriptase, might be a route for passing non-Mendelian traits to the offspring [101, 116]. The transmission of information from somatic to germ cells by exogenous RNA was suggested from a series of experiments in which mice xenografted with

human A-375 melanoma cells stably expressed EGFP that was then distributed through the bloodstream. Interestingly, transcripts were also detected in mouse spermatozoa, with exosomes as the suggested mechanism of transport [117]. In this paradigm, spermatozoa would take up exogenous DNA or transcribe exogenous RNA for delivery to the oocyte at fertilization [94]. This sperm mediated reverse transcriptase may act to expand the exogenous DNA or RNAs once in the sperm [116] providing a means to directly impact early embryo development.

Several compelling studies have exhibited the capacity for the environment experienced by the father to influence the offspring without the offspring ever being in contact with the environment [reviewed in 40, 42]. For example, both chronic and acute stress models have been used to explore how the paternal environment affects progeny. For example, chronic stress mouse models have identified increased levels of a group of sperm miRNAs [7] that act to specifically mark the response. Offspring developed a depressed hypothalamic-pituitary-adrenal (HPA) response in the absence of their father or shared paternal environment. This depressed HPA response was modulated by altered expression in the paraventricular nucleus of the hypothalamus [7]. Microinjection of the 9 miRNAs into the wild type oocyte resulted in zygotes expressing the same altered pattern of gene expression [7]. An acute stress mouse model resulted in depressive-like behaviors, exhibiting altered levels of both sperm miRNAs and piRNAs compared to controls [6]. Male offspring presented a similar depressive-like behavior as the father, in addition to altered levels of miRNAs within different brain regions and in serum. Interestingly, male offspring also exhibited altered glucose and insulin metabolic traits [6]. These results suggested that behavioral phenotypes could be transmitted down the male line, in the absence of paternal rearing, using a RNA mechanism [6, 7].

The impact of paternal diet has also drawn attention. For example, male mice subjected to a high-fat-high-sugar diet presented upregulated levels of miR-19b [13]. When miR-19b was microinjected into zygotes, offspring presented both obesity and reduced glucose tolerance, which persisted through several generations [13]. A low-protein diet in mice has been found to alter levels of sperm miRNAs, piRNAs, and several tRNAs [8]. In mice, alterations in miRNA levels as well as 5-methylcytidine ( $m^5C$ ) and  $N^2$ -methylguanosine ( $m^2G$ ) modified tRNAs in sperm have also been observed in response to a high-fat diet [5]. Microinjection of total sperm RNA from high-fat diet mice at fertilization resulted in offspring exhibiting impaired glucose tolerance, although insulin sensitivity was not observed [5]. This inferred that while modified sperm miRNA and tRNA may result in inherited metabolic traits, other layers of regulation may be involved in the inheritance of metabolic diseases such as diabetes. In humans, epidemiological studies on male-line transmission as exemplified in those of a Swedish Överkalix cohort [15, 118] span at least three generations. The impact of a grandfathers access to excess food during his childhood correlated with a higher incidence of both diabetes related mortality and mortality risk ratios in his grandsons [15, 118], while fathers with low food access as a child correlated with a decreased mortality by cardiovascular disease in their sons [118]. These results suggested a role of paternal influence in transgenerational responses in humans. As measured by BMI, male obesity has been shown to alter the human sperm methylome, with its reversal following bariatric surgery [119]. While alterations of sperm ncRNAs were also identified [119], their association was not well defined. Nevertheless, these results identify potential alterations in epigenetic effectors that may be transferred to the oocyte at fertilization, to influence embryo development and potentially be inherited by offspring.

### ***Challenges and Questions that Remain***

The low viability of parthenogenic mice [120] from two maternal genomes, sheds light on the importance of sperm beyond providing genomic information for successful zygote and embryo formation. The role of sperm RNAs is indicative of the intricate regulatory mechanisms of spermatogenesis, fertilization, and embryogenesis. Different classes of sperm RNAs have been found to act with different *modus operandi* to maintain the integrity of the genome, regulate gene expression, and chromatin state (Figure 3).

Currently, 10 - 15% of reproductive aged couples are affected by infertility [reviewed in 55]. Existing diagnostics rely on observable semen parameters [reviewed in 65] leaving many cases of male infertility unexplained. Differential RNA profiles exhibited by infertile men have been identified [reviewed in 4, 53, 55], and altered presence and/or abundance of RNAs are being used as molecular biomarkers to address reproductive health concerns. This includes the 648 sperm REs that appear essential for natural conception [53]. circRNAs which can be stable at room temperature for several hours [reviewed in 75] may offer yet another avenue. Factors such as advanced paternal age and increased BMI have been amplified in recent years [reviewed in 121, 122], and are increasingly linked to reproductive success. Standard semen parameters decline with advanced age [123] and increased BMI [124]. Similarly, both paternal age and BMI impact the potential for successful live birth [125, 126]. It is likely that these factors, among others, correlate with reproductive potential through the alteration of sperm RNAs. While diet modulates RNA abundance in sperm [5], these alterations have not been assessed in terms of reproductive potential, although work on this subject should be forthcoming.

Sperm RNAs used as biomarkers are also being developed in livestock, both as a model system and for enhanced breeding. In agriculture, identification of genetic markers of sperm quality, fertility, and those minimizing frozen-thawed cryo-damage are presently being pursued to optimize animal selection [50, 127-129]. Bovine is a well-studied agricultural species due to

its high economic impact. The corresponding sperm microarray studies have focused on sperm quality [127], fertility [130], and cryo-damage [131], while NGS technology is beginning to be applied to sperm motility [52]. Other agricultural species including stallion [50], porcine [128], and chicken [129] are still focused on the fundamental profiling of sperm RNAs. As the cost of sequencing decreases their potential impact for agriculture, and novel studies continues to grow.

Research focusing on the concept of health maintenance and lifestyle continues to move forward. Given the comparatively long-generation times between humans as compared to mice [132], the majority of human studies addressing epigenetic inheritance are epidemiologically based [15, 118]. Their significance is highlighted by the observation that access to food during a males' slow growth period correlates with the risk of diabetes and cardiovascular-related mortality, as well as over-all mortality risk, in sons and grandsons. The age of onset of smoking of a father has also been suggested to be correlated with the son's BMI [15]. These studies while limited, have identified that responses to the environment appear to reflect sex-specific transfer between generations. One must also consider that unlike females, males experience continual gamete (sperm) renewal, which can act to "wash out", or re-set the effect of a given exposure (e.g., diet, exercise, stress level). This capability affords males the unique possibly to marginalize the transfer of a potentially harmful effect to their children [reviewed in 121]. In this model, alterations in sperm methylation or sncRNAs resultant from a lifestyle choice like diet may be at least partially restored by a simple change in caloric intake.

The relationship between the father's diet and smoking status during his prepubescent, slow growth period, on the health of his children [15, 118] suggests there are critical times of exposure. Epidemiological studies in human suggest that a critical time of exposure for males occurs prior to the onset of puberty, between 9 and 12 years of age [118]. Mouse models and

human studies [reviewed in 133] have highlighted the impact of *in utero* environment as a time of relative phenotypic plasticity [reviewed in 134], and suggest this as a critical time of exposure for male-line epigenetic inheritance [134, 135].

The focus of most studies surrounds a single system input, but this does not reflect the intricate network of inputs that constitute allostatic load. For example, the common functional measure of obesity, BMI, will be influenced by multiple factors, including diet and exercise. While animal models provide a means to better control for these cofounding factors [132], this is rarely possible in human studies. A limited number of studies have begun to examine the relationship between age and diet with respect to this paradigm. On the one hand, mouse model studies have identified that undernourishment of males *in utero* can lead to inheritance of altered DNA methylation via sperm [134], yet this effect was not identified if the period of undernourishment followed birth [135]. On the other hand, human epidemiological studies [15, 118] noted that the age at which a male experiences malnutrition impacts the health of multiple generations. This may be in part due to age of exposure impacting the ability for the alteration to be “washed out”.

The fertilizing spermatozoon is essentially a dynamic single cell system that contains half of the information for the next generation in which a single cell is selected in some manner from millions. The field continues to seek to understand how specific paternal components, and their respective mechanisms influence reproductive health. The potential importance of sperm RNAs in the maintenance and transference of biological information between generations underscores the necessity of achieving an understanding. Perhaps this provides a mechanism to assess specific fitness of a new trait prior to adaptive selection, and in part, provide one of the components of genetic resilience [11, 136]. For now, we are hindered by the lack of a comprehensive understanding of how these epigenetic effectors are modulated.

Four fundamental questions remain with respect to paternal RNAs. 1) Do sperm RNAs directly interact with other epigenetic effectors, perhaps directly with the genome (chromatin) or other RNAs or structures? 2) What is the role of RNAs in transmitting specific phenotypes? 3) What is the corresponding mechanism by which this is achieved? 4) How may we modify the future-past in a controlled manner? Addressing each of these questions in this unique human single cell system will require the development of new molecular and computational tools. The answers will likely revolutionize our understanding of reproduction and health, as the totality of the male contribution reflective of his past experiences that impacts the future birth and life course of his child, becomes appreciated.

## **Acknowledgments**

The authors would like to thank Dr. Àlex Clop and Dr. Meritxell Jodar for their critical review and comments. We apologize to those authors whom we were unable to cite given the constraints of the manuscript.

## **References**

1. Ostermeier GC, Miller D, Huntriss JD, Diamond MP, Krawetz SA. Reproductive biology: delivering spermatozoan RNA to the oocyte. *Nature* 2004; 429:154.
2. Krawetz SA. Paternal contribution: new insights and future challenges. *Nat Rev Genet* 2005; 6:633-642.

3. Garrido N, Martinez-Conejero JA, Jauregui J, Horcajadas JA, Simon C, Remohi J, Meseguer M. Microarray analysis in sperm from fertile and infertile men without basic sperm analysis abnormalities reveals a significantly different transcriptome. *Fertil Steril* 2009; 91:1307-1310.
4. Platts AE, Dix DJ, Chemes HE, Thompson KE, Goodrich R, Rockett JC, Rawe VY, Quintana S, Diamond MP, Strader LF, Krawetz SA. Success and failure in human spermatogenesis as revealed by teratozoospermic RNAs. *Hum Mol Genet* 2007; 16:763-773.
5. Chen Q, Yan M, Cao Z, Li X, Zhang Y, Shi J, Feng GH, Peng H, Zhang X, Zhang Y, Qian J, Duan E, et al. Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder. *Science* 2016; 351:397-400.
6. Gapp K, Jawaid A, Sarkies P, Bohacek J, Pelczar P, Prados J, Farinelli L, Miska E, Mansuy IM. Implication of sperm RNAs in transgenerational inheritance of the effects of early trauma in mice. *Nat Neurosci* 2014; 17:667-669.
7. Rodgers AB, Morgan CP, Leu NA, Bale TL. Transgenerational epigenetic programming via sperm microRNA recapitulates effects of paternal stress. *Proc Natl Acad Sci U S A* 2015; 112:13699-13704.
8. Sharma U, Conine CC, Shea JM, Boskovic A, Derr AG, Bing XY, Belleannee C, Kucukural A, Serra RW, Sun F, Song L, Carone BR, et al. Biogenesis and function of tRNA fragments during sperm maturation and fertilization in mammals. *Science* 2016; 351:391-396.

9. Skogen JC, Overland S. The fetal origins of adult disease: a narrative review of the epidemiological literature. *JRSM Short Rep* 2012; 3:59.
10. McEwen BS, Wingfield JC. What is in a name? Integrating homeostasis, allostasis and stress. *Horm Behav* 2010; 57:105-111.
11. Lundberg U. Stress hormones in health and illness: the roles of work and gender. *Psychoneuroendocrinology* 2005; 30:1017-1021.
12. McDade TW. Life history, maintenance, and the early origins of immune function. *Am J Hum Biol* 2005; 17:81-94.
13. Grandjean V, Fourre S, De Abreu DA, Derieppe MA, Remy JJ, Rassoulzadegan M. RNA-mediated paternal heredity of diet-induced obesity and metabolic disorders. *Sci Rep* 2015; 5:18193.
14. de Castro Barbosa T, Ingerslev LR, Alm PS, Versteyhe S, Massart J, Rasmussen M, Donkin I, Sjogren R, Mudry JM, Vetterli L, Gupta S, Krook A, et al. High-fat diet reprograms the epigenome of rat spermatozoa and transgenerationally affects metabolism of the offspring. *Mol Metab* 2016; 5:184-197.
15. Pembrey ME, Bygren LO, Kaati G, Edvinsson S, Northstone K, Sjostrom M, Golding J, Team AS. Sex-specific, male-line transgenerational responses in humans. *Eur J Hum Genet* 2006; 14:159-166.
16. Kierszenbaum AL, Tres LL. Structural and transcriptional features of the mouse spermatid genome. *J Cell Biol* 1975; 65:258-270.
17. Betlach CJ, Erickson RP. A unique RNA species from maturing mouse spermatozoa. *Nature* 1973; 242:114-115.

18. Rejon E, Bajon C, Blaize A, Robert D. RNA in the nucleus of a motile plant spermatozoid: characterization by enzyme-gold cytochemistry and in situ hybridization. *Mol Reprod Dev* 1988; 1:49-56.
19. Pessot CA, Brito M, Figueroa J, Concha, II, Yanez A, Burzio LO. Presence of RNA in the sperm nucleus. *Biochem Biophys Res Commun* 1989; 158:272-278.
20. Kumar G, Patel D, Naz RK. c-MYC mRNA is present in human sperm cells. *Cell Mol Biol Res* 1993; 39:111-117.
21. Miller D, Tang PZ, Skinner C, Lilford R. Differential RNA fingerprinting as a tool in the analysis of spermatozoal gene expression. *Hum Reprod* 1994; 9:864-869.
22. Rohwedder A, Liedigk O, Schaller J, Glander HJ, Werchau H. Detection of mRNA transcripts of beta 1 integrins in ejaculated human spermatozoa by nested reverse transcription-polymerase chain reaction. *Mol Hum Reprod* 1996; 2:499-505.
23. Wykes SM, Visscher DW, Krawetz SA. Haploid transcripts persist in mature human spermatozoa. *Mol Hum Reprod* 1997; 3:15-19.
24. Wykes SM, Miller D, Krawetz SA. Mammalian spermatozoal mRNAs: tools for the functional analysis of male gametes. *J Submicrosc Cytol Pathol* 2000; 32:77-81.
25. Xu HL, Swoboda I, Bhalla PL, Singh MB. Male gametic cell-specific gene expression in flowering plants. *Proceedings of the National Academy of Sciences of the United States of America* 1999; 96:2554-2558.
26. Miller D, Briggs D, Snowden H, Hamlington J, Rollinson S, Lilford R, Krawetz SA. A complex population of RNAs exists in human ejaculate spermatozoa: implications for understanding molecular aspects of spermiogenesis. *Gene* 1999; 237:385-392.

27. Grunewald S, Paasch U, Glander HJ, Anderegg U. Mature human spermatozoa do not transcribe novel RNA. *Andrologia* 2005; 37:69-71.
28. Gur Y, Breitbart H. Mammalian sperm translate nuclear-encoded proteins by mitochondrial-type ribosomes. *Genes Dev* 2006; 20:411-416.
29. Gur Y, Breitbart H. Protein synthesis in sperm: dialog between mitochondria and cytoplasm. *Mol Cell Endocrinol* 2008; 282:45-55.
30. Ostermeier GC, Dix DJ, Miller D, Khatri P, Krawetz SA. Spermatozoal RNA profiles of normal fertile men. *Lancet* 2002; 360:772-777.
31. Engel ML, Chaboud A, Dumas C, McCormick S. Sperm cells of *Zea mays* have a complex complement of mRNAs. *Plant J* 2003; 34:697-707.
32. Martins RP, Krawetz SA. Towards understanding the epigenetics of transcription by chromatin structure and the nuclear matrix. *Gene Ther Mol Biol* 2005; 9:229-246.
33. Ostermeier GC, Goodrich RJ, Moldenhauer JS, Diamond MP, Krawetz SA. A suite of novel human spermatozoal RNAs. *J Androl* 2005; 26:70-74.
34. Velculescu VE, Zhang L, Vogelstein B, Kinzler KW. Serial analysis of gene expression. *Science* 1995; 270:484-487.
35. Zhao Y, Li Q, Yao C, Wang Z, Zhou Y, Wang Y, Liu L, Wang Y, Wang L, Qiao Z. Characterization and quantification of mRNA transcripts in ejaculated spermatozoa of fertile men by serial analysis of gene expression. *Hum Reprod* 2006; 21:1583-1590.

36. Wang H, Zhou Z, Xu M, Li J, Xiao J, Xu ZY, Sha J. A spermatogenesis-related gene expression profile in human spermatozoa and its potential clinical applications. *J Mol Med (Berl)* 2004; 82:317-324.
37. Amanai M, Brahmajosyula M, Perry AC. A restricted role for sperm-borne microRNAs in mammalian fertilization. *Biol Reprod* 2006; 75:877-884.
38. Girard A, Sachidanandam R, Hannon GJ, Carmell MA. A germline-specific class of small RNAs binds mammalian Piwi proteins. *Nature* 2006; 442:199-202.
39. Rassoulzadegan M, Grandjean V, Gounon P, Vincent S, Gillot I, Cuzin F. RNA-mediated non-mendelian inheritance of an epigenetic change in the mouse. *Nature* 2006; 441:469-474.
40. Rando OJ. Daddy issues: paternal effects on phenotype. *Cell* 2012; 151:702-708.
41. Larriba E, del Mazo J. Role of Non-Coding RNAs in the Transgenerational Epigenetic Transmission of the Effects of Reprotoxicants. *Int J Mol Sci* 2016; 17:452.
42. Gapp K, Bohacek J. Epigenetic germline inheritance in mammals: looking to the past to understand the future. *Genes Brain Behav* 2017.
43. Anway MD, Cupp AS, Uzumcu M, Skinner MK. Epigenetic transgenerational actions of endocrine disruptors and male fertility. *Science* 2005; 308:1466-1469.
44. Krawetz SA, Kruger A, Lalancette C, Tagett R, Anton E, Draghici S, Diamond MP. A survey of small RNAs in human sperm. *Hum Reprod* 2011; 26:3401-3412.

45. Peng H, Shi J, Zhang Y, Zhang H, Liao S, Li W, Lei L, Han C, Ning L, Cao Y, Zhou Q, Chen Q, et al. A novel class of tRNA-derived small RNAs extremely enriched in mature mouse sperm. *Cell Res* 2012; 22:1609-1612.
46. Sendler E, Johnson GD, Mao S, Goodrich RJ, Diamond MP, Hauser R, Krawetz SA. Stability, delivery and functions of human sperm RNAs at fertilization. *Nucleic Acids Res* 2013; 41:4104-4117.
47. Margolin G, Khil PP, Kim J, Bellani MA, Camerini-Otero RD. Integrated transcriptome analysis of mouse spermatogenesis. *BMC Genomics* 2014; 15:39.
48. Selvaraju S, Parthipan S, Somashekar L, Kolte AP, Krishnan Binsila B, Arangasamy A, Ravindra JP. Occurrence and functional significance of the transcriptome in bovine (*Bos taurus*) spermatozoa. *Sci Rep* 2017; 7:42392.
49. Card CJ, Anderson EJ, Zamberlan S, Krieger KE, Kaproth M, Sartini BL. Cryopreserved bovine spermatozoal transcript profile as revealed by high-throughput ribonucleic acid sequencing. *Biol Reprod* 2013; 88:49.
50. Das PJ, McCarthy F, Vishnoi M, Paria N, Gresham C, Li G, Kachroo P, Sudderth AK, Teague S, Love CC, Varner DD, Chowdhary BP, et al. Stallion sperm transcriptome comprises functionally coherent coding and regulatory RNAs as revealed by microarray analysis and RNA-seq. *PLoS One* 2013; 8:e56535.
51. Johnson GD, Sendler E, Lalancette C, Hauser R, Diamond MP, Krawetz SA. Cleavage of rRNA ensures translational cessation in sperm at fertilization. *Mol Hum Reprod* 2011; 17:721-726.

52. Capra E, Turri F, Lazzari B, Cremonesi P, Gliootti TM, Fojadelli I, Stella A, Pizzi F. Small RNA sequencing of cryopreserved semen from single bull revealed altered miRNAs and piRNAs expression between High- and Low-motile sperm populations. *BMC Genomics* 2017; 18:14.
53. Jodar M, Sendler E, Moskvtsev SI, Librach CL, Goodrich R, Swanson S, Hauser R, Diamond MP, Krawetz SA. Absence of sperm RNA elements correlates with idiopathic male infertility. *Sci Transl Med* 2015; 7:295re296.
54. Goodrich RJ, Anton E, Krawetz SA. Isolating mRNA and small noncoding RNAs from human sperm. *Methods Mol Biol* 2013; 927:385-396.
55. Jodar M, Selvaraju S, Sendler E, Diamond MP, Krawetz SA, Reproductive Medicine N. The presence, role and clinical use of spermatozoal RNAs. *Hum Reprod Update* 2013; 19:604-624.
56. Johnson GD, Mackie P, Jodar M, Moskvtsev S, Krawetz SA. Chromatin and extracellular vesicle associated sperm RNAs. *Nucleic Acids Res* 2015; 43:6847-6859.
57. Smith S, Hosid S, Scott L. Use of postseparation sperm parameters to determine the method of choice for sperm preparation for assisted reproductive technology. *Fertil Steril* 1995; 63:591-597.
58. Mao S, Goodrich RJ, Hauser R, Schrader SM, Chen Z, Krawetz SA. Evaluation of the effectiveness of semen storage and sperm purification methods for spermatozoa transcript profiling. *Syst Biol Reprod Med* 2013; 59:287-295.

59. Allamaneni SS, Agarwal A, Rama S, Ranganathan P, Sharma RK. Comparative study on density gradients and swim-up preparation techniques utilizing neat and cryopreserved spermatozoa. *Asian J Androl* 2005; 7:86-92.
60. Lalancette C, Platts AE, Johnson GD, Emery BR, Carrell DT, Krawetz SA. Identification of human sperm transcripts as candidate markers of male fertility. *J Mol Med (Berl)* 2009; 87:735-748.
61. Georgiadis AP, Kishore A, Zorrilla M, Jaffe TM, Sanfilippo JS, Volk E, Rajkovic A, Yatsenko AN. High quality RNA in semen and sperm: isolation, analysis and potential application in clinical testing. *J Urol* 2015; 193:352-359.
62. Goodrich R, Johnson G, Krawetz SA. The preparation of human spermatozoal RNA for clinical analysis. *Arch Androl* 2007; 53:161-167.
63. Jodar M, Sendler E, Moskvtsev SI, Librach CL, Goodrich R, Swanson S, Hauser R, Diamond MP, Krawetz SA. Response to Comment on "Absence of sperm RNA elements correlates with idiopathic male infertility". *Sci Transl Med* 2016; 8:353tr351.
64. Mao S, Sendler E, Goodrich RJ, Hauser R, Krawetz SA. A comparison of sperm RNA-seq methods. *Syst Biol Reprod Med* 2014; 60:308-315.
65. Jodar M, Anton E, Krawetz SA. Sperm RNA and Its Use as a Clinical marker. In: De Jonge CJ, Barratt CL (eds.), *The Sperm Cell: Production, Maturation, Fertilization, Regeneration*, 2 ed. Cambridge: Cambridge University Press; 2017: 59-72.
66. Jodar M, Sendler E, Krawetz SA. The protein and transcript profiles of human semen. *Cell Tissue Res* 2016; 363:85-96.

67. Johnson GD, Lalancette C, Linnemann AK, Leduc F, Boissonneault G, Krawetz SA. The sperm nucleus: chromatin, RNA, and the nuclear matrix. *Reproduction* 2011; 141:21-36.
68. Chen Q, Yan W, Duan E. Epigenetic inheritance of acquired traits through sperm RNAs and sperm RNA modifications. *Nat Rev Genet* 2016; 17:733-743.
69. Engreitz JM, Ollikainen N, Guttman M. Long non-coding RNAs: spatial amplifiers that control nuclear structure and gene expression. *Nat Rev Mol Cell Biol* 2016; 17:756-770.
70. Ward WS. Function of sperm chromatin structural elements in fertilization and development. *Mol Hum Reprod* 2010; 16:30-36.
71. Zhang X, Gao F, Fu J, Zhang P, Wang Y, Zeng X. Systematic identification and characterization of long non-coding RNAs in mouse mature sperm. *PLoS One* 2017; 12:e0173402.
72. Gao F, Zhang P, Zhang H, Zhang Y, Zhang Y, Hao Q, Zhang X. Dysregulation of long noncoding RNAs in mouse testes and spermatozoa after exposure to cadmium. *Biochem Biophys Res Commun* 2017; 484:8-14.
73. Schmitz SU, Grote P, Herrmann BG. Mechanisms of long noncoding RNA function in development and disease. *Cell Mol Life Sci* 2016; 73:2491-2509.
74. Zhang C, Gao L, Xu EY. LncRNA, a new component of expanding RNA-protein regulatory network important for animal sperm development. *Semin Cell Dev Biol* 2016; 59:110-117.
75. Cortes-Lopez M, Miura P. Emerging Functions of Circular RNAs. *Yale J Biol Med* 2016; 89:527-537.

76. Dong Y, He D, Peng Z, Peng W, Shi W, Wang J, Li B, Zhang C, Duan C. Circular RNAs in cancer: an emerging key player. *J Hematol Oncol* 2017; 10:2.
77. Dong WW, Li HM, Qing XR, Huang DH, Li HG. Identification and characterization of human testis derived circular RNAs and their existence in seminal plasma. *Sci Rep* 2016; 6:39080.
78. Lau NC. Small RNAs in the animal gonad: guarding genomes and guiding development. *Int J Biochem Cell Biol* 2010; 42:1334-1347.
79. Yuan S, Schuster A, Tang C, Yu T, Ortogero N, Bao J, Zheng H, Yan W. Sperm-borne miRNAs and endo-siRNAs are important for fertilization and preimplantation embryonic development. *Development* 2016; 143:635-647.
80. Carmell MA, Girard A, van de Kant HJ, Bourc'his D, Bestor TH, de Rooij DG, Hannon GJ. MIWI2 is essential for spermatogenesis and repression of transposons in the mouse male germline. *Dev Cell* 2007; 12:503-514.
81. Moazed D. Small RNAs in transcriptional gene silencing and genome defence. *Nature* 2009; 457:413-420.
82. Liu WM, Pang RT, Chiu PC, Wong BP, Lao K, Lee KF, Yeung WS. Sperm-borne microRNA-34c is required for the first cleavage division in mouse. *Proc Natl Acad Sci U S A* 2012; 109:490-494.
83. Yuan S, Tang C, Zhang Y, Wu J, Bao J, Zheng H, Xu C, Yan W. mir-34b/c and mir-449a/b/c are required for spermatogenesis, but not for the first cleavage division in mice. *Biol Open* 2015; 4:212-223.

84. Lee RC, Hammell CM, Ambros V. Interacting endogenous and exogenous RNAi pathways in *Caenorhabditis elegans*. *RNA* 2006; 12:589-597.
85. Nodine MD. Mobile small RNAs: Sperm-companion communication. *Nat Plants* 2016; 2:16041.
86. Modzelewski AJ, Hilz S, Crate EA, Schweidenback CT, Fogarty EA, Grenier JK, Freire R, Cohen PE, Grimson A. Dgcr8 and Dicer are essential for sex chromosome integrity during meiosis in males. *J Cell Sci* 2015; 128:2314-2327.
87. Zimmermann C, Romero Y, Warnefors M, Bilican A, Borel C, Smith LB, Kotaja N, Kaessmann H, Nef S. Germ cell-specific targeting of DICER or DGCR8 reveals a novel role for endo-siRNAs in the progression of mammalian spermatogenesis and male fertility. *PLoS One* 2014; 9:e107023.
88. Yang F, Wang PJ. Multiple LINEs of retrotransposon silencing mechanisms in the mammalian germline. *Semin Cell Dev Biol* 2016; 59:118-125.
89. O'Donnell KA, Boeke JD. Mighty Piwis defend the germline against genome intruders. *Cell* 2007; 129:37-44.
90. Pantano L, Jodar M, Bak M, Ballesca JL, Tommerup N, Oliva R, Vavouri T. The small RNA content of human sperm reveals pseudogene-derived piRNAs complementary to protein-coding genes. *RNA* 2015; 21:1085-1095.
91. Giebler M, Greither T, Muller L, Mosinger C, Behre HM. Altered PIWI-LIKE 1 and PIWI-LIKE 2 mRNA expression in ejaculated spermatozoa of men with impaired sperm characteristics. *Asian J Androl* 2017.

92. Tang W, Tu S, Lee HC, Weng Z, Mello CC. The RNase PARN-1 Trims piRNA 3' Ends to Promote Transcriptome Surveillance in *C. elegans*. *Cell* 2016; 164:974-984.
93. Ding DQ, Liu JL, Dong KZ, Midic U, Hess RA, Xie HR, Demireva EY, Chen C. PNLDC1 is essential for piRNA 3' end trimming and transposon silencing during spermatogenesis in mice. *Nature Communications* 2017; 8.
94. Aravin AA, Sachidanandam R, Girard A, Fejes-Toth K, Hannon GJ. Developmentally regulated piRNA clusters implicate MILI in transposon control. *Science* 2007; 316:744-747.
95. He X, Chen X, Zhang X, Duan X, Pan T, Hu Q, Zhang Y, Zhong F, Liu J, Zhang H, Luo J, Wu K, et al. An Lnc RNA (GAS5)/SnoRNA-derived piRNA induces activation of TRAIL gene by site-specifically recruiting MLL/COMPASS-like complexes. *Nucleic Acids Res* 2015; 43:3712-3725.
96. Law JA, Jacobsen SE. Establishing, maintaining and modifying DNA methylation patterns in plants and animals. *Nat Rev Genet* 2010; 11:204-220.
97. Kirchner S, Ignatova Z. Emerging roles of tRNA in adaptive translation, signalling dynamics and disease. *Nat Rev Genet* 2015; 16:98-112.
98. Jachowicz JW, Bing X, Pontabry J, Boskovic A, Rando OJ, Torres-Padilla ME. LINE-1 activation after fertilization regulates global chromatin accessibility in the early mouse embryo. *Nat Genet* 2017; 49:1502-1510.
99. Miller D. Confrontation, Consolidation, and Recognition: The Oocyte's Perspective on the Incoming Sperm. *Cold Spring Harb Perspect Med* 2015; 5:a023408.


100. Slotkin RK, Martienssen R. Transposable elements and the epigenetic regulation of the genome. *Nat Rev Genet* 2007; 8:272-285.
101. Spadafora C. Sperm-mediated 'reverse' gene transfer: a role of reverse transcriptase in the generation of new genetic information. *Hum Reprod* 2008; 23:735-740.
102. Niakan KK, Han J, Pedersen RA, Simon C, Pera RA. Human pre-implantation embryo development. *Development* 2012; 139:829-841.
103. Vassena R, Boue S, Gonzalez-Roca E, Aran B, Auer H, Veiga A, Izpisua Belmonte JC. Waves of early transcriptional activation and pluripotency program initiation during human preimplantation development. *Development* 2011; 138:3699-3709.
104. Rando OJ. Intergenerational Transfer of Epigenetic Information in Sperm. *Cold Spring Harb Perspect Med* 2016; 6.
105. Ntostis P, Carter D, Iles D, Huntriss JD, Tzetis M, Miller D. Potential sperm contributions to the murine zygote predicted by in silico analysis. *Reproduction* 2017.
106. Saunders CM, Larman MG, Parrington J, Cox LJ, Royse J, Blayney LM, Swann K, Lai FA. PLC zeta: a sperm-specific trigger of Ca(2+) oscillations in eggs and embryo development. *Development* 2002; 129:3533-3544.
107. Practice Committee of the American Society for Reproductive M. Evaluation and treatment of recurrent pregnancy loss: a committee opinion. *Fertil Steril* 2012; 98:1103-1111.
108. Rogenhofer N, Ott J, Pilatz A, Wolf J, Thaler CJ, Windischbauer L, Schagdarsurengin U, Steger K, von Schonfeldt V. Unexplained recurrent miscarriages are associated with an aberrant sperm protamine mRNA content. *Hum Reprod* 2017; 32:1574-1582.

109. Brahem S, Mehdi M, Landolsi H, Mougou S, Elghezal H, Saad A. Semen parameters and sperm DNA fragmentation as causes of recurrent pregnancy loss. *Urology* 2011; 78:792-796.
110. Coughlan C, Clarke H, Cutting R, Saxton J, Waite S, Ledger W, Li T, Pacey AA. Sperm DNA fragmentation, recurrent implantation failure and recurrent miscarriage. *Asian J Androl* 2015; 17:681-685.
111. Eisenberg ML, Sapra KJ, Kim SD, Chen Z, Buck Louis GM. Semen quality and pregnancy loss in a contemporary cohort of couples recruited before conception: data from the Longitudinal Investigation of Fertility and the Environment (LIFE) Study. *Fertil Steril* 2017; 108:613-619.
112. Gil-Villa AM, Cardona-Maya W, Agarwal A, Sharma R, Cadavid A. Assessment of sperm factors possibly involved in early recurrent pregnancy loss. *Fertil Steril* 2010; 94:1465-1472.
113. Nanassy L, Carrell DT. Paternal effects on early embryogenesis. *J Exp Clin Assist Reprod* 2008; 5:2.
114. Aoki VW, Moskovtsev SI, Willis J, Liu L, Mullen JB, Carrell DT. DNA integrity is compromised in protamine-deficient human sperm. *J Androl* 2005; 26:741-748.
115. Szyf M. Nongenetic inheritance and transgenerational epigenetics. *Trends Mol Med* 2015; 21:134-144.
116. Spadafora C. Soma to germline inheritance of extrachromosomal genetic information via a LINE-1 reverse transcriptase-based mechanism. *Bioessays* 2016; 38:726-733.

117. Cossetti C, Lugini L, Astrologo L, Saggio I, Fais S, Spadafora C. Soma-to-germline transmission of RNA in mice xenografted with human tumour cells: possible transport by exosomes. *PLoS One* 2014; 9:e101629.
118. Kaati G, Bygren LO, Edvinsson S. Cardiovascular and diabetes mortality determined by nutrition during parents' and grandparents' slow growth period. *Eur J Hum Genet* 2002; 10:682-688.
119. Donkin I, Versteyhe S, Ingerslev LR, Qian K, Mechta M, Nordkap L, Mortensen B, Appel EV, Jorgensen N, Kristiansen VB, Hansen T, Workman CT, et al. Obesity and Bariatric Surgery Drive Epigenetic Variation of Spermatozoa in Humans. *Cell Metab* 2016; 23:369-378.
120. Kono T, Obata Y, Wu Q, Niwa K, Ono Y, Yamamoto Y, Park ES, Seo JS, Ogawa H. Birth of parthenogenetic mice that can develop to adulthood. *Nature* 2004; 428:860-864.
121. Palmer NO, Bakos HW, Fullston T, Lane M. Impact of obesity on male fertility, sperm function and molecular composition. *Spermatogenesis* 2012; 2:253-263.
122. Herati AS, Zhelyazkova BH, Butler PR, Lamb DJ. Age-related alterations in the genetics and genomics of the male germ line. *Fertil Steril* 2017; 107:319-323.
123. Stone BA, Alex A, Werlin LB, Marrs RP. Age thresholds for changes in semen parameters in men. *Fertil Steril* 2013; 100:952-958.
124. Chavarro JE, Toth TL, Wright DL, Meeker JD, Hauser R. Body mass index in relation to semen quality, sperm DNA integrity, and serum reproductive hormone levels among men attending an infertility clinic. *Fertil Steril* 2010; 93:2222-2231.

125. Campbell JM, Lane M, Owens JA, Bakos HW. Paternal obesity negatively affects male fertility and assisted reproduction outcomes: a systematic review and meta-analysis. *Reprod Biomed Online* 2015; 31:593-604.
126. Alio AP, Salihi HM, McIntosh C, August EM, Weldeselasse H, Sanchez E, Mbah AK. The effect of paternal age on fetal birth outcomes. *Am J Mens Health* 2012; 6:427-435.
127. Bissonnette N, Levesque-Sergerie JP, Thibault C, Boissonneault G. Spermatozoal transcriptome profiling for bull sperm motility: a potential tool to evaluate semen quality. *Reproduction* 2009; 138:65-80.
128. Chen C, Wu H, Shen D, Wang S, Zhang L, Wang X, Gao B, Wu T, Li B, Li K, Song C. Comparative profiling of small RNAs of pig seminal plasma and ejaculated and epididymal sperm. *Reproduction* 2017; 153:785-796.
129. Singh RP, Shafeeq CM, Sharma SK, Singh R, Mohan J, Sastry KV, Saxena VK, Azeez PA. Chicken sperm transcriptome profiling by microarray analysis. *Genome* 2016; 59:185-196.
130. Govindaraju A, Uzun A, Robertson L, Atli MO, Kaya A, Topper E, Crate EA, Padbury J, Perkins A, Memili E. Dynamics of microRNAs in bull spermatozoa. *Reprod Biol Endocrinol* 2012; 10:82.
131. Chen X, Wang Y, Zhu H, Hao H, Zhao X, Qin T, Wang D. Comparative transcript profiling of gene expression of fresh and frozen-thawed bull sperm. *Theriogenology* 2015; 83:504-511.
132. Uhl EW, Warner NJ. Mouse Models as Predictors of Human Responses: Evolutionary Medicine. *Curr Pathobiol Rep* 2015; 3:219-223.


133. Sutton EF, Gilmore LA, Dunger DB, Heijmans BT, Hivert MF, Ling C, Martinez JA, Ozanne SE, Simmons RA, Szyf M, Waterland RA, Redman LM, et al. Developmental programming: State-of-the-science and future directions-Summary from a Pennington Biomedical symposium. *Obesity (Silver Spring)* 2016; 24:1018-1026.
134. Radford EJ, Ito M, Shi H, Corish JA, Yamazawa K, Isganaitis E, Seisenberger S, Hore TA, Reik W, Erkek S, Peters A, Patti ME, et al. In utero effects. In utero undernourishment perturbs the adult sperm methylome and intergenerational metabolism. *Science* 2014; 345:1255903.
135. Shea JM, Serra RW, Carone BR, Shulha HP, Kucukural A, Ziller MJ, Vallaster MP, Gu H, Tapper AR, Gardner PD, Meissner A, Garber M, et al. Genetic and Epigenetic Variation, but Not Diet, Shape the Sperm Methylome. *Dev Cell* 2015; 35:750-758.
136. Gillespie CF, Phifer J, Bradley B, Ressler KJ. Risk and resilience: genetic and environmental influences on development of the stress response. *Depress Anxiety* 2009; 26:984-992.



**Figure 1**

Three decades of sperm RNAs summarized with key studies. Sperm RNAs were first detected in ferns [18], and subsequently in mammals [19]. In the 1990s several independent studies [20-25] validated the presence of mRNAs in sperm with different approaches including RT-PCR, RT-qPCR, and ISH. In the 2000s, with novel technologies, microarray allowed the first human sperm transcriptome profile [30], a technique posteriorly used for studying male infertility [36]. The first species to be sequenced (by sperm RNA cloning and cDNA sequencing) was in maize [31]. In 2004, sperm RNAs were shown to be delivered to the zygote [1]. sncRNAs were firstly detected in human sperm by microarray [33] and corroborated, a few years later, with sncRNA-seq [44]. With the boost of NGS techniques, mRNA-seq studies were carried in sperm RNAs in human [46], and agricultural species as bovine [48, 49] and stallion [50]. NGS has also been used to identify biomarkers related to male infertility in humans [53] or sperm quality (motility) in bulls [52]. Relevant transgenerational epigenetics studies start with the work from


Rassoulzadegan and colleagues [39], were miRNAs were suggested to modify offspring phenotype. Several studies in mice have followed, were parents have been set in extreme environmental conditions, including stress [6,7] and diet [5, 8, 13] and sncRNAs profiles have been studied. ISH (*In Situ* Hybridization); sncRNAs (small non-coding RNAs); miRNAs (micro RNAs); tRNAs (transfer RNAs).



**Figure 2**

Defining sperm RNA Elements (REs) with the discovery analysis approach. Dark blue lines represent mapped sequencing reads. Dashed grey line correspond to an annotated intergenic region. The approach also provides a reliable quantitation method of isoforms, where read abundance would vary as a function of transcript isoform. Since mature sperm are transcriptionally and translationally silent, the term “abundance” is preferred rather than “expression”. **A)** Four REs are detected, corresponding to the 4 annotated exons of Gene 1. Similar read abundance is observed in RE 1, 2, and 3, but lower abundance in RE 4, suggesting

possible 3' degradation of the transcript. **B)** 5' – end degradation. **C)** Only 1 RE is detected, corresponding to the annotated exon 3. **D)** Sperm RE analysis detects all the annotated exons as well as an intronic RE that may correspond to an unannotated exon or other gene biotype. **E)** Two REs are detected in intronic regions and may correspond to intronic retained elements. **F)** RE detected upstream of the 5' UTR, but could also exist 3'. **G)** Two REs are detected in an intergenic region, may correspond to unannotated exons from nearby genes, exons from a new unannotated gene or other gene biotypes.



**Figure 3**

Distribution of sperm RNAs and their roles. **A-B)** The outer nuclear layers indicated in light and medium blue contain ~2/3 of all sperm RNAs (including coding and non-coding), the majority of which are long RNAs (> 200 nt). These RNAs include exogenous RNAs packaged within vesicles and RNAs extruded from the inner nucleus either bound to associated proteins or within vesicles during cytoplasmic extrusion. The remaining ~1/3 of total sperm RNAs reside

within the dark blue inner nuclear fraction. **C)** RNAs found within the inner nuclear fraction are likely directly associated with the DNA or to chromatin bound proteins to influence chromatin structure and may provide an epigenetic signature. Examples include the chromatin associated RNAs, such as lncRNAs, piRNAs, miRNAs, and siRNAs. **D)** Mitochondrial RNAs have been observed in the midpiece of the spermatozoa. lncRNAs (long non-coding RNAs); miRNAs (microRNAs); piRNAs (Piwi-interacting RNAs); pri-miRNAs (primary miRNAs); siRNAs (small interfering RNAs); TE (transposable elements); tRNAs (transfer RNAs); TSS (transcript start site); rRNAs (ribosomal RNAs).