

1
2
3

INVARIANT CURVES AND INTEGRABILITY OF PLANAR C^r DIFFERENTIAL SYSTEMS

ANTONI FERRAGUT AND JAUME LLIBRE

ABSTRACT. We improve the known expressions of the C^r differential systems in the plane having a given C^{r+1} invariant curve, or a given C^{r+1} first integral. Their application to polynomial differential systems having either an invariant algebraic curve, or a first integral also improves the known results on such systems.

4
5

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

A C^r real planar differential system is a system of the form

$$\dot{x} = P(x, y), \quad \dot{y} = Q(x, y), \tag{1}$$

6 where $(x, y) \in D \subseteq \mathbb{R}^2$, D is the domain of definition of the system and $P, Q \in C^r$, where r
 7 is a positive integer, or $r = \infty$, or $r = \omega$ (meaning that the system is analytic). The vector
 8 field associated to system (1) is

$$X(x, y) = P(x, y) \frac{\partial}{\partial x} + Q(x, y) \frac{\partial}{\partial y}.$$

9 In what follows we shall talk indistinctly of the differential system (1) or of its vector field
 10 X .

11 Let $\mathbb{R}[x, y]$ be the ring of polynomials in the variables x, y with real coefficients. If
 12 $P, Q \in \mathbb{R}[x, y]$ then we say that the differential system (1) is *polynomial*. In such a case, we
 13 define the *degree* of this polynomial system as $\max\{\deg P, \deg Q\}$, where $\deg P, \deg Q$ are
 14 the degree of P and Q , respectively.

15 Let U be an open subset of \mathbb{R}^2 . A *first integral* of X in U is a locally non-constant
 16 function $H : U \rightarrow \mathbb{R}$ which is constant on all the solutions of X contained in U , i.e.
 17 $X(H) = 0$ in the points of U . In this case we also say that X is *integrable* on U .

18 Let $g \in C^r$. The curve $g = 0$ is *invariant* under the flow of system (1) if

$$X(g)|_{g=0} = \left(P \frac{\partial g}{\partial x} + Q \frac{\partial g}{\partial y} \right) \Big|_{g=0} = 0.$$

19 The *Nambu bracket* (or the *Lie bracket*, or the *Jacobian*) of two functions $f, g \in C^1$ is

$$\{f, g\} = \frac{\partial f}{\partial x} \frac{\partial g}{\partial y} - \frac{\partial f}{\partial y} \frac{\partial g}{\partial x}.$$

20 We start our results by finding a way (inspired in Corollary 1.3.3 of [9]) for writing all
 21 systems (1) having an invariant curve $g = 0$.

22 **Theorem 1.** *Let $g = 0$ be a C^{r+1} invariant curve of system (1). Then for any C^{r+1} function*
 23 *f such that $\{g, f\} \neq 0$ we have*

$$\dot{x} = P = \frac{X(g)f_y - X(f)g_y}{\{g, f\}}, \quad \dot{y} = Q = \frac{-X(g)f_x + X(f)g_x}{\{g, f\}}. \tag{2}$$

24 The following result is well-known, for a proof see for instance [3].

2010 *Mathematics Subject Classification.* 34A05, 34A34, 34A55, 34C05, 37C10.

Key words and phrases. planar differential system, C^r function, invariant curve, exponential factor, co-factor, Darboux theory of integrability.

1 **Theorem 2.** Assume that a polynomial differential system has an invariant algebraic curve
 2 $g(x, y) = 0$ such that there are no points at which g and its first derivatives all vanish. If
 3 $(g_x, g_y) = 1$, then the system has the following normal form:

$$\dot{x} = Ag - Dg_y, \quad \dot{y} = Bg + Dg_x, \quad (3)$$

4 where A, B, D are suitable polynomials.

5 The hypotheses in the previous theorem are necessary, as the polynomial differential
 6 system given in the next example shows. However applying Theorem 1 we can obtain the
 7 polynomial differential system of the example if the functions A, B, D are not necessarily
 8 polynomials.

9 **Example 1.** Consider the polynomial differential system of degree five appearing in [2]

$$\dot{x} = 2x + y - 3x^4, \quad \dot{y} = 4y + 2x^3 - 9x^3y + 3x^5.$$

10 The algebraic curve $g(x, y) = (y - x^2)(y - x^3) = 0$ is invariant under the flow of this
 11 differential system. Observe that g and its first derivatives vanish at the origin, and also at
 12 $(1, 1)$.

13 Note that we cannot obtain \dot{y} from the expression $Bg + Dg_x$ of Theorem 2 because the
 14 lowest degree of g is 2 and $x|g_x$. Hence the hypotheses of Theorem 2 do not hold for this
 15 example and it is easy to see that there do not exist polynomials A, B, D such that (3)
 16 holds because in $Bg + Dg_x$ the term $4y$ of \dot{y} cannot appear.

17 Theorem 1 has no hypotheses besides $\{g, f\} \neq 0$. If we apply it with $f = y$, then we
 18 have

$$\begin{aligned} \frac{X(g)f_y - X(f)g_y}{\{g, f\}} &= \frac{(2 - 3x)(4 + 5x + 6x^2)}{x(5x^3 - 2y - 3xy)}g - \frac{4y + 2x^3 - 9x^3y + 3x^5}{x(5x^3 - 2y - 3xy)}g_y \\ &= 2x + y - 3x^4 = \dot{x}, \end{aligned}$$

$$\begin{aligned} \frac{-X(g)f_x + X(f)g_x}{\{g, f\}} &= \frac{4y + 2x^3 - 9x^3y + 3x^5}{x(5x^3 - 2y - 3xy)}g_x \\ &= 4y + 2x^3 - 9x^3y + 3x^5 = \dot{y}. \end{aligned}$$

19 Hence we obtain $\dot{x} = Ag - Dg_y, \dot{y} = Bg + Dg_x$ with the rational functions

$$A = \frac{(2 - 3x)(4 + 5x + 6x^2)}{x(5x^3 - 2y - 3xy)}, \quad B = 0, \quad D = \frac{4y + 2x^3 - 9x^3y + 3x^5}{x(5x^3 - 2y - 3xy)}.$$

20

□

21 The arguments used in the previous example to construct the vector field can be general-
 22 ized. The following theorem includes the condition $g_x \neq 0$ on a curve $g = 0$. This condition
 23 is not restrictive. If $g_x \equiv 0$ then $g_y \neq 0$ (otherwise g is a constant), and in this case the
 24 theorem applies swapping x and y .

25 **Theorem 3.** Assume that a polynomial differential system has an invariant algebraic curve
 26 $g = g(x, y) = 0$. If $g_x \neq 0$, then the system has the following normal form:

$$\dot{x} = Ag - Bg_y, \quad \dot{y} = Bg_x, \quad (4)$$

27 where A, B are suitable rational functions. Conversely, if a polynomial differential system
 28 can be written as in (4), then the curve $g = 0$ is invariant under the flow of this differential
 29 system.

30 Note that Theorem 3 provides all the polynomial differential systems having a given
 31 invariant algebraic curve without any assumptions on the curve, while Theorem 2 needs
 32 some assumptions.

33 When system (1) is polynomial and $g = 0$ is an invariant algebraic curve, there exists a
 34 polynomial k , called the *cofactor*, that satisfies the equation $X(g) = kg$. The polynomial

1 cofactors play a main role in the Darboux theory of integrability (see [1, 5, 6, 10]). Of course
 2 we can write this equation as $k = X(g)/g$, and this is a polynomial, so $g|X(g)$. If g is an
 3 invariant curve of system (1) we can define also its *cofactor* as the function $k = X(g)/g$.
 4 We know that in the paper [7] the authors improve the Darboux theory of integrability
 5 using invariant curves which are not necessarily algebraic but having polynomial cofactors.
 6 Note that in our case the cofactors are in general non-polynomial.

7 It is widely known that if a linear combination of cofactors of invariant algebraic curves
 8 of a polynomial differential system is zero, then this system has a Darboux first integral (see
 9 Theorem 8.7 of [6]). We state here our third theorem. It generalizes the classical Darboux
 10 Theorem for polynomial differential systems to C^r differential systems with C^{r+1} invariant
 11 curves.

12 **Theorem 4.** *Consider the C^r differential system (1) and let $g_i = 0$ be a C^{r+1} invariant*
 13 *curve of (1) with cofactor k_i for $i = 1, \dots, M$. Then $H = \prod_{i=1}^M g_i^{\nu_i}$ is a first integral of (1)*
 14 *if and only if $\sum_{i=1}^M \nu_i k_i = 0$, for some convenient $\nu_i \in \mathbb{R}$.*

15 Theorem 4 will be proved in section 4.

16 A function $V(x, y)$ is an *inverse integrating factor* of the differential system (1) if the
 17 differential system

$$\dot{x} = \frac{P}{V} = -H_y, \quad \dot{y} = \frac{Q}{V} = H_x,$$

18 is Hamiltonian, with $H(x, y)$ the Hamiltonian function. For more information about the
 19 inverse integrating factor see [8]. Concerning the differential system (1) and the inverse
 20 integrating factors, we have the following theorem, inspired in Corollary 1.4.4 of [9].

21 **Theorem 5.** *If the differential system (1) has a first integral H , then we can write it as*

$$\dot{x} = \frac{X(f)}{\{H, f\}} \{H, x\}, \quad \dot{y} = \frac{X(f)}{\{H, f\}} \{H, y\},$$

22 where f is an arbitrary function. Moreover,

$$V(x, y) = \frac{X(f)}{\{H, f\}}$$

23 is an inverse integrating factor of the system.

24 We finally deal with differential systems having a first integral of the form $ge^{h/g}$. We
 25 have the following result.

26 **Proposition 6.** *Suppose that the differential system (1) has the first integral*

$$H(x, y) = ge^{h/g},$$

27 where $g, h \in C^{r+1}$ are such that $\{g, h\} \neq 0$. Then the function $V = -X(g)g/\{g, h\}$ is an
 28 inverse integrating factor of the system.

29 2. PRELIMINARY RESULTS

30 We state here some results of [9] that we shall use later on. The first one (see Corollary
 31 1.3.3 of [9]) provides the planar differential systems having a given invariant curve.

32 **Theorem 7.** *Let $g(x, y) = 0$ be a function defined in an open set $D \subseteq \mathbb{R}^2$. Then any*
 33 *differential system defined in D for which $g = 0$ is invariant can be written as*

$$\begin{aligned} \dot{x} &= \phi \frac{\{x, f\}}{\{g, f\}} + \lambda \frac{\{g, x\}}{\{g, f\}}, \\ \dot{y} &= \phi \frac{\{y, f\}}{\{g, f\}} + \lambda \frac{\{g, y\}}{\{g, f\}}, \end{aligned}$$

34 where f, ϕ and λ are arbitrary functions such that $\phi|_{g=0} = 0$ and $\{g, f\} \neq 0$ in D .

1 Next theorem (see Corollary 1.4.4 of [9]) provides the planar differential systems having
2 a given first integral.

3 **Theorem 8.** *Let $H(x, y)$ be a function defined in an open set $D \subseteq \mathbb{R}^2$. Then the most
4 general differential systems defined in D which admit the first integral H are*

$$\dot{x} = \lambda \frac{\{H, x\}}{\{H, f\}}, \quad \dot{y} = \lambda \frac{\{H, y\}}{\{H, f\}},$$

5 where f is an arbitrary function such that $\{H, f\} \neq 0$ in D .

6

3. PROOFS

7 *Proof of Theorem 1.* According to Theorem 7, system (1) can be written in D as

$$\begin{aligned} \dot{x} &= P(x, y) = \phi \frac{\{x, f\}}{\{g, f\}} + \lambda \frac{\{g, x\}}{\{g, f\}}, \\ \dot{y} &= Q(x, y) = \phi \frac{\{y, f\}}{\{g, f\}} + \lambda \frac{\{g, y\}}{\{g, f\}}, \end{aligned} \tag{5}$$

8 where ϕ is an arbitrary function such that $\phi|_{g=0} = 0$, $\{g, f\} \neq 0$ in D , and λ, f are arbitrary
9 functions.

10 We note that we can actually compute an expression for ϕ, λ in terms of g, f just solving
11 the linear system (5) that defines P and Q in the unknowns ϕ and λ , which has a unique
12 solution because its determinant is one. This linear system is

$$\begin{aligned} \{x, f\}\phi + \{g, x\}\lambda &= P\{g, f\}, \\ \{y, f\}\phi + \{g, y\}\lambda &= Q\{g, f\}, \end{aligned}$$

13 or equivalently

$$\begin{aligned} f_y\phi - g_y\lambda &= P\{g, f\}, \\ -f_x\phi + g_x\lambda &= Q\{g, f\}, \end{aligned}$$

14 and by using the Cramer method, we have

$$\phi = \frac{\begin{vmatrix} P\{g, f\} & -g_y \\ Q\{g, f\} & g_x \end{vmatrix}}{\begin{vmatrix} f_y & -g_y \\ -f_x & g_x \end{vmatrix}} = \frac{\{g, f\}(Pg_x + Qg_y)}{\{g, f\}} = X(g),$$

15 and

$$\lambda = \frac{\begin{vmatrix} f_y & P\{g, f\} \\ -f_x & Q\{g, f\} \end{vmatrix}}{\begin{vmatrix} f_y & -g_y \\ -f_x & g_x \end{vmatrix}} = \frac{\{g, f\}(Qf_y + Pf_x)}{\{g, f\}} = X(f).$$

16 Hence the only arbitrary function in (5) is f , and it must satisfy that $\{g, f\} \neq 0$ in D . \square

17 *Proof of Theorem 3.* We assume in this proof that $g_x \neq 0$. If $g_x \equiv 0$ then we must have
18 $g_y \neq 0$, and we can swap x and y .

19 Theorem 1 assures that all the differential systems having $g = 0$ as invariant write
20 as system (2). The function f which appears in Theorem 1 is arbitrary, because the
21 simplification of the quotients in (2) provide directly P and Q . In particular, we can fix
22 $f(x, y) = y$, for which $X(f) = Q$ and $\{g, f\} = g_x \neq 0$. Moreover, system (2) writes as

$$\dot{x} = P = \frac{X(g) - Qg_y}{g_x}, \quad \dot{y} = Q = \frac{Qg_x}{g_x}. \tag{6}$$

23 Since we are assuming that $g = 0$ is invariant, there must exist a polynomial $k(x, y)$ such
24 that $X(g) = kg$. Substituting in (6) we have

$$\dot{x} = P = \frac{kg - Qg_y}{g_x}, \quad \dot{y} = Q. \tag{7}$$

1 In order to obtain a polynomial differential system we have to impose that $(kg - Qg_y)/g_x$
 2 is a polynomial. That is, we need to fix the coefficients of k and Q in such a way that P
 3 is a polynomial. If we succeed, then the differential system (7) is polynomial and moreover
 4 $g = 0$ is invariant for system (7).

5 Clearly there exist two polynomials A and B such that

$$kg - Qg_y = Ag_x + B.$$

6 Then

$$P = \frac{kg - Qg_y}{g_x} = A + \frac{B}{g_x},$$

7 with $\deg B < \deg g - 1$. We need to obtain conditions on the coefficients of k and Q such
 8 that $B = 0$. The equation $B = 0$ can be written as a linear system of equations, one
 9 equation for each monomial of B . The unknowns of this linear system are the coefficients
 10 of k and Q . Of course the degrees of k and Q must be large enough to assure that the
 11 system is compatible.

12 We note that we can always find a solution, i.e. the differential system $\dot{x} = g - g_y, \dot{y} = g_x$
 13 has the curve $g = 0$ invariant; its cofactor is g_x . So the system coming from $B = 0$ is
 14 compatible if we have enough unknowns. More precisely, if there are enough coefficients of
 15 k and Q ; or equivalently if the degrees of k and Q are large enough.

16 Since $\deg B \leq \deg g - 2$, the linear system has at most $\binom{\deg g}{2}$ equations. The number
 17 of unknowns is $\binom{\deg k + 2}{2} + \binom{\deg Q + 2}{2}$. Hence we must have

$$\binom{\deg k + 2}{2} + \binom{\deg Q + 2}{2} \geq \binom{\deg g}{2}.$$

18 Once the linear system is solved, some coefficients of k and Q are fixed, and some other
 19 may remain free. Therefore $B = 0$ and P is a polynomial. Those free coefficients provide
 20 all the polynomial differential systems having the curve g invariant.

21 We have obtained the most general polynomial differential system $\dot{x} = P, \dot{y} = Q$ having
 22 the curve $g = 0$ invariant. Moreover, k is the cofactor of $g = 0$. The degree of the system is
 23 either $\deg Q$ in the case $\deg k < \deg Q$, or $\deg k + 1$ otherwise. So the theorem follows. \square

24

4. INVARIANT OBJECTS

25 **4.1. Darboux first integrals.** Next we prove Theorem 4.

26 *Proof of Theorem 4.* We compute $X(H)$:

$$X(H) = X \left(\prod_{i=1}^M g_i^{\nu_i} \right) = \sum_{i=1}^M \nu_i X(g_i) g_i^{\nu_i - 1} \prod_{j \neq i} g_j^{\nu_j} = \sum_{i=1}^M \nu_i k_i g_i^{\nu_i} \prod_{j \neq i} g_j^{\nu_j} = \left(\sum_{i=1}^M \nu_i k_i \right) H.$$

27 Then the theorem follows. \square

28 In the Darboux theory of integrability for complex polynomial differential systems (and
 29 consequently also for real polynomial differential systems) there exists a minimum number
 30 of invariant algebraic curves that assures a first integral. This is due to the linear combi-
 31 nation of cofactors: these cofactors are polynomials of degree lower than the degree of the
 32 polynomial system, say m ; that is they belong to the vector space of all complex polynomi-
 33 als in the variables x, y of degree at most $m - 1$. A base of this ring has $\binom{m + 1}{2}$ elements.

34 If we have more than $\binom{m + 1}{2}$ cofactors then there must exist a linear combination of
 35 them being zero.

1 For the \mathcal{C}^r differential systems in the real plane this kind of facts do not occur in general,
 2 so a minimum number of curves that assures a first integral cannot be established unless we
 3 restrict the class of differential systems.

4 **4.2. The inverse integrating factor.** We first prove Theorem 5.

5 *Proof of Theorem 5.* If the differential system (1) has a first integral H , then from Theorem
 6 8, we know that we can write it as

$$\dot{x} = \lambda \frac{\{H, x\}}{\{H, f\}}, \quad \dot{y} = \lambda \frac{\{H, y\}}{\{H, f\}}, \quad (8)$$

7 where f is an arbitrary function such that $\{H, f\} \neq 0$. We prove that $X(f) = \lambda$, indeed

$$X(f) = \lambda \frac{\{H, x\}}{\{H, f\}} f_x + \lambda \frac{\{H, y\}}{\{H, f\}} f_y = \lambda \frac{-H_y f_x + H_x f_y}{\{H, f\}} = \lambda.$$

8 From (8), the function $V = X(f)/\{H, f\}$ is an inverse integrating factor, because we
 9 have

$$\dot{x} = P = -VH_y, \quad \dot{y} = Q = VH_x.$$

10 Notice that $V = 0$ is an invariant curve of system (1). Hence the theorem follows. \square

11 When the differential system (1) has an inverse integrating factor, the associated differ-
 12 ential system $\dot{x} = P/V$, $\dot{y} = Q/V$ is Hamiltonian, so the area of any region of the domain
 13 of definition of this Hamiltonian system is the same after it is moved forward or backward
 14 by the flow of the system. In the set $\{V = 0\}$ this system is not Hamiltonian. The orbits
 15 that in their neighborhood do not allow this preservation of the area are contained into
 16 $\{V = 0\}$.

17 In the next example we study a polynomial differential system having a polynomial
 18 inverse integrating factor V . The set $\{V = 0\}$ in this example is formed by a focus and a
 19 limit cycle. The area is not preserved close to these two orbits, hence the corresponding
 20 Hamiltonian system cannot be defined in them.

21 **Example 2.** Consider the polynomial differential system

$$\dot{x} = -y + x(x^2 + y^2 - 1), \quad \dot{y} = x + y(x^2 + y^2 - 1),$$

22 which appears in Example 2 of page 126 of [11]. Using polar coordinates it is easy to
 23 see that this system has a focus at the origin and a limit cycle at the circle $x^2 + y^2 = 1$.
 24 Moreover it has the polynomial inverse integrating factor $V(x, y) = (x^2 + y^2)(x^2 + y^2 - 1)$.
 25 The system has the Darboux first integral

$$H(x, y) = \frac{x^2 + y^2 - 1}{x^2 + y^2} e^{2 \arctan \frac{y}{x}}.$$

26 We can write it in polar coordinates as

$$H(r, \theta) = \frac{r^2 - 1}{r^2} e^{2(\theta + k\pi)}, \quad k \in \mathbb{Z}.$$

27 \square

28 **4.3. No exponential factors.** When system (1) is polynomial and we have an invariant
 29 algebraic curve $g = 0$ and an additional invariant algebraic curve close to $g = 0$, say
 30 $g_\varepsilon = g + \varepsilon f + \mathcal{O}(\varepsilon^2)$, we can define an exponential factor, which is a function $F = e^{f/g}$
 31 such that $X(F)/F$ is a polynomial. So the notion of exponential factor is associated to the
 32 notion of multiplicity of an invariant algebraic curve, see [4] for more details.

33 When dealing with \mathcal{C}^{r+1} invariant curves of a \mathcal{C}^r differential system the notion of expo-
 34 nential factor has no sense, because close to a \mathcal{C}^{r+1} invariant curve there are infinitely many
 35 other \mathcal{C}^{r+1} invariant curves.

1

5. A SPECIAL DARBOUX FUNCTION

2 We consider in this section differential systems (1) having a first integral of the form
 3 $H(x, y) = ge^{h/g}$, where $h, g \in C^{r+1}$. We prove Proposition 6.

4 *Proof of Proposition 6.* We have

$$\frac{\partial \log H}{\partial x} = \frac{gg_x + h_xg - hg_x}{g^2} = \frac{Q}{V}, \quad \frac{\partial \log H}{\partial y} = \frac{gg_y + h_yg - hg_y}{g^2} = -\frac{P}{V},$$

5 where V is the inverse integrating factor associated to the first integral $\log H$. Note that
 6 $V = -P/(\log H)_y = Q/(\log H)_x$. Since $g = 0$ is invariant under the flow of system (1),
 7 from the expressions of P and Q given in (2) we have

$$V = -\frac{(X(g)f_y - X(f)g_y)g^2}{\{g, f\}(g_yg + h_yg - hg_y)} = \frac{(-X(g)f_x + X(f)g_x)g^2}{\{g, f\}(g_xg + h_xg - hg_x)},$$

8 where f is an arbitrary function such that $\{g, f\} \neq 0$. This last equality can be written as

$$gX(f)\{h, g\} + (h - g)X(g)\{g, f\} - gX(g)\{h, f\} = 0. \quad (9)$$

9 Since f is arbitrary and $\{g, h\} \neq 0$, we can set $f = h$. Then

$$(X(g)(h - g) - gX(h))\{g, h\} = 0,$$

10 or equivalently

$$X(h) = \frac{X(g)}{g}(h - g). \quad (10)$$

11 When the differential system (1) is polynomial, equation (10) is equivalent to say that
 12 $-X(g)/g$ is the cofactor of the exponential factor $e^{h/g}$.

13 Back to the expression of V we have

$$V = \frac{Q}{(\log H)_x} = \frac{-X(g)h_x + X(h)g_x}{\frac{\{g, h\}}{g_xg + h_xg - hg_x}} = \frac{X(g)g(-gh_x + (h - g)g_x)}{\{g, h\}(g_xg + h_xg - hg_x)} = -\frac{X(g)g}{\{g, h\}}.$$

14 Then the proposition follows. \square

15 Note that if the differential system of Proposition 6 is polynomial, then $e^{h/g}$ is an expo-
 16 nential factor with cofactor $-X(g)/g$.

17 **Remark 1.** If $\{g, h\} = 0$ in the proof of Proposition 6, then, from (9), we can obtain

$$(h - g)\{g, f\} - g\{h, f\} = 0,$$

for all f such that $\{g, f\} \neq 0$. In particular, for $f = \log H$, we have

$$\begin{aligned} (h - g)\{g, \log H\} - g\{h, \log H\} &= (h - g) \left(-g_x \frac{P}{V} - g_y \frac{Q}{V} \right) - g \left(-h_x \frac{P}{V} - h_y \frac{Q}{V} \right) \\ &= -(h - g) \frac{X(g)}{V} + g \frac{X(h)}{V} = -\frac{X(g)(h - g) - gX(h)}{V} = 0. \end{aligned}$$

18 So we also obtain (10) in the case $\{g, h\} = 0$.

19

ACKNOWLEDGEMENTS

20 The authors are partially supported by the MINECO grant MTM2013-40998-P. A. Ferragut is
 21 also partially supported by the Universitat Jaume I grant P1-1B2015-16. J. Llibre is also partially
 22 supported by the Generalitat de Catalunya grant 2014SGR-568 and the grants FP7-PEOPLE-2012-
 23 IRSES 318999 and 316338.

REFERENCES

1

- 2 [1] C. CHRISTOPHER AND C. LI, *Limit cycles of differential equations*, Advanced Courses in Mathematics.
3 CRM Barcelona. Birkhäuser Verlag, Basel, 2007.
- 4 [2] C. CHRISTOPHER, J. LLIBRE, CH. PANTAZI AND S. WALCHER, *Inverse problems for multiple invariant*
5 *curves*, Proc. R. Soc. Edinburgh **137A** (2007), 1197–1226.
- 6 [3] C. CHRISTOPHER, J. LLIBRE, CH. PANTAZI AND X. ZHANG, *Darboux integrability and invariant alge-*
7 *braic curves for planar polynomial systems*, J. Phys. A: Math. Gen. **35** (2002), 2457–2476.
- 8 [4] C. CHRISTOPHER, J. LLIBRE AND J.V. PEREIRA, *Multiplicity of invariant algebraic curves in polynomial*
9 *vector fields*, Pacific J. Math. **229** (2007), 63–117.
- 10 [5] G. DARBOUX, *Mémoire sur les équations différentielles algébriques du premier ordre et du premier degré*
11 *(Mélanges)*, Bull. Sci. Math. **2** (1878), 60–96; 123–144; 151–200.
- 12 [6] F. DUMORTIER, J. LLIBRE AND J.C. ARTÉS, *Qualitative Theory of Planar Differential Systems*, Uni-
13 versitext, Springer, 2006.
- 14 [7] I.A. GARCÍA AND J. GINÉ, *Generalized cofactors and nonlinear superposition principles*, Appl. Math.
15 Lett. **16** (2003), 1137–1141.
- 16 [8] I.A. GARCÍA AND M. GRAU, *A survey on the inverse integrating factor*, Qual. Theory Dyn. Syst. **9**
17 (2010), 115–166.
- 18 [9] J. LLIBRE AND R. RAMÍREZ, *Inverse problems in ordinary differential equations and applications*,
19 Progress in Mathematics **313**, Birkhäuser Basel, 2016.
- 20 [10] D. SCHLOMIUK, *Algebraic and geometric aspects of the theory of polynomial vector fields. Bifurcations*
21 *and periodic orbits of vector fields*, (Montreal, PQ, 1992), 429–467, NATO Adv. Sci. Inst. Ser. C Math.
22 Phys. Sci., 408, Kluwer Acad. Publ., Dordrecht, 1993.
- 23 [11] Y. YE, *Theory of limit cycles*, American Mathematical Society, Translations of Mathematical Mono-
24 graphs **66**, 1986.

25 A. FERRAGUT: INSTITUT UNIVERSITARI DE MATEMÀTIQUES I APLICACIONS DE CASTELLÓ (IMAC) AND
26 DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT JAUME I, EDIFICI TI (ESTEC), AV. DE VICENT SOS
27 BAYNAT, S/N, CAMPUS DEL RIU SEC, 12071 CASTELLÓ DE LA PLANA, SPAIN
28 *E-mail address:* ferragut@uji.es

29 J. LLIBRE: DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BEL-
30 LATERRA, BARCELONA, CATALONIA, SPAIN
31 *E-mail address:* jllibre@mat.uab.cat