
This is the **accepted version** of the journal article:

Pérez Varela, María [et al.]. «Functional characterization of AbaQ, a novel efflux pump mediating quinolone resistance in *acinetobacter baumannii*». *Antimicrobial Agents and Chemotherapy*, Vol. 62, Num. 9 (September 2018), art. e00906-18 DOI 10.1128/AAC.00906-18, PMID 29941648, PMID PMC6125561

This version is available at <https://ddd.uab.cat/record/334538>

under the terms of the  license.

A Short-Form Paper for Antimicrobial Agents and Chemotherapy

Title: Functional characterization of AbaQ, a novel efflux pump mediating quinolone resistance in *Acinetobacter baumannii*.

Running title: The *A. baumannii* AbaQ efflux pump

Authors: María Pérez-Varela¹, Jordi Corral¹, Jesús Aranda^{1,2*} and Jordi Barbé¹

¹Departament de Genètica i Microbiologia, Facultat de Biociències, Universitat Autònoma de Barcelona, Bellaterra 08193, Cerdanyola del Vallès (Barcelona), Spain

²Serra Hünter Fellow. Generalitat de Catalunya, Spain

*Corresponding author: jesus.aranda@uab.cat, Phone: (34) 935811665, Fax: (34) 935812387

ABSTRACT

Acinetobacter baumannii has emerged as an important multidrug-resistant nosocomial pathogen. In previous work, we identified a putative MFS transporter, AU097_RS17040, involved in the pathogenicity of *A. baumannii* (Pérez-Varela *et al.*, Infect Immun. 2017). In this study, we analyzed the susceptibility to diverse antimicrobial agents of *A. baumannii* cells defective in this transporter, referred to herein as AbaQ. Our results showed that AbaQ is mainly involved in the extrusion of quinolone-type drugs in *A. baumannii*.

Acinetobacter baumannii is a multidrug-resistant (MDR) pathogen that causes hospital-acquired infections (1). In previous work, we identified a new, putative major facilitator superfamily (MFS) transporter in *A. baumannii* strain ATCC 17978, AU097_RS17040. This transporter, referred to herein as AbaQ (*A. baumannii* quinolone resistance transporter), is involved in surface-associated motility as well as the virulence of *A. baumannii* (2).

Sequence analysis of AbaQ, annotated as a MFS transporter in *A. baumannii* strain ATCC 17978 (accession number WP_000345069), indicated an ORF of 1305 nucleotides. According to the deduced amino acid sequence, the protein consists of 434 residues and has a molecular mass of 47.8 kDa and a theoretical isoelectric point (pI) of 9.27. Based on predictions of its secondary structure and transmembrane topology, AbaQ is composed of 12 α -helical transmembrane segments, with both the N- and C-termini located in the cytoplasm (Fig. 1A). Support for this structure came from an independent analysis that revealed the 3D structure of the protein (Fig. 1B). These data indicated that AbaQ is a drug H⁺ antiporter-1 (DHA1), which differs from DHA2-type MFS drug transporters by the presence of 12 rather than 14 transmembrane segments (3). The predicted product of the *abaQ* gene exhibited low amino acid identity and similarity (< 24% and < 38%, respectively) with other MFS transporters involved in drug efflux in *A. baumannii* (Table 1).

To determine whether a lack of AbaQ alters antimicrobial susceptibilities, the response of two *A. baumannii abaQ* mutants obtained in previous work (2) to antimicrobials of different classes was tested. Compared to the wild-type (WT) parental strain, *A. baumannii* ATCC 17978, the *abaQ* mutant had a 2- to 4-fold higher susceptibility to trimethoprim and novobiocin (Table 2). The highest susceptibilities (approximately 8- to 32-fold) occurred in response to the quinolone-type antibiotics ciprofloxacin,

levofloxacin, and nalidixic acid (Table 2). By contrast, the mutant and its WT parent did not differ in their susceptibilities to β -lactams (meropenem and ampicillin), aminoglycosides (amikacin and gentamycin), macrolides (erythromycin), and polymyxines (colistin) or to other antimicrobials (chloramphenicol, tetracycline, minocycline, and rifampin) (Table 2). To further corroborate the relevance of this MFS in other *A. baumannii* strains, we analyzed the *abaQ* knockout mutant derived from *A. baumannii* strain MAR002, a biofilm-hyperproducing strain recently isolated from a wound sample collected from a patient at the Hospital del Mar in Barcelona (2, 4, 5). Assays of the mutant indicated that inactivation of the AbaQ homologue in *A. baumannii* MAR002 (99% amino acid identity) also caused the highest reduction (approximately 8- to 32-fold) in the minimum inhibitory concentration (MIC) of quinolone-type antibiotics (data not shown).

To complement the mutants, the AbaQ gene, including its own promoter, was amplified from the genome of *A. baumannii* strain ATCC 17978 using the AbaQFXbaI and AbaQRXbaI oligonucleotides (5'-ACTGTCTAGAGGAATATCACAGCTTGCAGCG and 5'-ACTGTCTAGATTACAAAGGCTTTTGAATATTC, respectively), cloning them into the *Xba*I restriction site of the pET-RA vector (6). Complementation of the *abaQ*-mutant derived from *A. baumannii* ATCC 17978 completely restored antimicrobial susceptibility to the same levels determined in the WT parental strain (Table 2). The recovery of the WT phenotype was also observed in the complemented *abaQ* MAR002 mutant derivative (data not shown). These results provided clear evidence of the critical role of AbaQ in the efflux of quinolone-type antibiotics in *A. baumannii*.

To assess whether AbaQ confers resistance to quinolones through an active efflux mechanism, the MIC of ciprofloxacin in the presence of the efflux pump inhibitor

carbonyl cyanide 3-chlorophenylhydrazone CCCP (Sigma) was determined in *A. baumannii* strains carrying (WT and $\text{AbaQ}^- + \text{pET-RA+AbaQ}$) or lacking (AbaQ^- and $\text{AbaQ}^- + \text{pET-RA}\emptyset$) the *abaQ* gene. The addition of CCCP (20 mg/L) decreased the ciprofloxacin MIC 16-fold in *AbaQ*-carrying cells (from 0.25 to 0.015 mg/L) but only 4-fold in cells lacking the *abaQ* gene (from 0.0078 to 0.0019 mg/L). The addition of CCCP alone did not inhibit bacterial cell growth in any of the *A. baumannii* strains, indicating that the above results were not due to the toxicity of CCCP itself. Accordingly, when ampicillin, which is not a substrate of the *AbaQ* transporter (Table 2), was used instead of ciprofloxacin, there was no difference in antimicrobial susceptibility in the presence or absence of CCCP in any of the cultured *A. baumannii* strains (data not shown). These results unequivocally demonstrated that the *abaQ* gene product confers decreased susceptibility to quinolones, by encoding an active efflux transporter.

The majority of the MFS transporters involved in drug-efflux described thus far in *A. baumannii* mediate chloramphenicol efflux, with a smaller number mediating the efflux of other antimicrobials, such as erythromycin, tetracycline, minocycline, fosfomycin, and colistin (Table 1). Among the latter group of transporters is *AmvA*, which participates in the efflux of erythromycin and different classes of disinfectants, detergents, and dyes but also confers modest resistance to quinolone-type antimicrobials (7). In *amvA* mutant strains, susceptibility to both ciprofloxacin and norfloxacin is 2-fold higher than in the WT strain whereas susceptibility to nalidixic acid is unchanged (7). Other transporters belonging to different families and involved in the extrusion of a wide range of antimicrobials, including quinolones, have been described in *A. baumannii*: *AdeABC* (8), *AdeDE* (9), *AdeFGH* (10), *AdeIJK* (11), *AdeM* (12), and *AdeT* (13), all of them belonging to the resistance/nodulation/division (RND)

superfamily; AbeM, AbeM2, and AbeM4 transporters, belonging to the MATE (multiple antimicrobial toxin extrusion) family (12, 14); and AbeS, belonging to the small multidrug resistance (SMR) family (15). No role in quinolone efflux has been detected in the only ATP-binding cassette (ABC) transporter described so far in *A. baumannii* (A1S_1503) (16) or in any of the transporters that make up the most recently discovered family of efflux pumps: the proteobacterial antimicrobial compound efflux (PACE) transporters (17).

To our knowledge, AbaQ, which is widely present in *A. baumannii* clinical isolates and involved in both surface-associated motility and virulence (2), is the first MFS efflux-pump shown to play an important role in the extrusion of quinolone-type antimicrobials in this MDR nosocomial pathogen.

ACKNOWLEDGEMENTS

We thank Joan Ruiz (UAB) and Susana Escibano (UAB) for their excellent technical assistance, as well as our UAB students Paula García and Daniel Quiñones for their helpful support. We also thank the Germán Bou lab for the *A. baumannii* MAR002 strain. This study was supported by grant BIO2016-77011-R from the *Ministerio de Economía y Competitividad*. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Table 1. Major facilitator superfamily (MFS) transporters described in *A. baumannii*

MFS	Accession number	Main antimicrobial exported	Identity ^a	Similarity ^a
AmvA (7)	ACQ82816	Erythromycin	17.8%	31.0%
TetA (18)	AAO38186	Tetracycline	17.6%	28.5%
TetB (19)	AFV67369	Minocycline	19%	33.8%
CraA (20)	ABO13543	Chloramphenicol	19%	36.9%
FloR (21)	AQT19056	Chloramphenicol	16.9%	28.2%
CmlA (22)	AMD83513	Chloramphenicol	17.5%	28.3%
AbaF (23)	ABO11759	Fosfomycin	23.7%	37.6%
EmrB (24)	ABO12199	Colistin	17.2%	30.7%
AedC (25)	ABO11341	Tetracycline/ Chloramphenicol	18.5%	32.7%

^aWith respect to AbaQ of *A. baumannii* strain ATCC 17978, using the Basic Local Alignment Search Tool (BLAST; <http://www.ncbi.nlm.nih.gov>) and Clustal Omega (<http://www.ebi.ac.uk/Tools/msa/clustalo/>).

Table 2. Minimum inhibitory concentrations (MICs, mg/L) of various antimicrobials in wild-type *A. baumannii* ATCC 17978 (WT) and the *abaQ* mutant derivative (AbaQ⁻). The MICs for the *abaQ* mutant carrying the empty pET-RA plasmid (pET-RAØ) and the mutant carrying the pET-RA+AbaQ plasmid (complemented mutant) are also shown.

	WT	AbaQ ⁻	Fold change ^a	AbaQ ⁻ + pET-RAØ	AbaQ ⁻ + pET-RA +AbaQ	Fold change ^b
Ciprofloxacin	0.25	0.0078	32	0.0078	0.25	32
Levofloxacin	0.125	0.0078	16	0.0078	0.125	16
Nalidixic acid	8	1	8	1	8	8
Trimethoprim	16	4	4	4	16	4
Novobiocin	8	4	2	4	8	2
Meropenem	2	2	1	2	2	1
Ampicillin	16	16	1	16	16	1
Amikacin	1	1	1	1	1	1
Gentamicin	0.5	0.5	1	0.5	0.5	1
Erythromycin	4	4	1	4	4	1
Colistin	0.5	0.5	1	0.5	0.5	1
Chloramphenicol	16	16	1	16	16	1
Tetracycline	4	4	1	4	4	1
Minocycline	0.062	0.062	1	0.062	0.062	1
Rifampin	4	4	1	NA ^c	NA ^c	NA ^c

^aRatio of the MICs of the WT vs. AbaQ⁻.

^bRatio of the MICs for AbaQ⁻+pET-RA+AbaQ vs. AbaQ⁻+ pET-RAØ.

^cNA, not applicable (the pET-RA plasmid carries rifampin resistance).

FIGURE LEGEND

Fig. 1. (A) Prediction of the structure of AbaQ using Protter (10). The putative protein is shown parallel to the cytoplasmic membrane. **(B)** 3D representation of AbaQ viewed along the plane of the membrane from the periplasmic side using RaptorX Structure Prediction (26) and visualized using PyMOL software (27). The 12 transmembrane α -helices are numbered (1–12); both the N- and C-termini are located in the cytoplasm.

REFERENCES

1. Lynch J, Zhanel G, Clark N. 2017. Infections due to *Acinetobacter baumannii* in the ICU: Treatment options. *Semin Respir Crit Care Med* 38:311–325.
2. Pérez-Varela M, Corral J, Vallejo JA, Rumbo-Feal S, Bou G, Aranda J, Barbé J. 2017. Mutants in the β -subunit of the RNA polymerase impairing the surface-associated motility and virulence of *Acinetobacter baumannii*. *Infect Immun* 85(8).
3. Paulsen IT, Brown MH, Skurray RA. 1996. Proton-dependent multidrug efflux systems. *Microbiol Rev* 60:575–608.
4. Álvarez-Fraga L, López M, Merino M, Rumbo-Feal S, Tomás M, Bou G, Poza M. 2015. Draft genome sequence of the biofilm-hyperproducing *Acinetobacter baumannii* clinical strain MAR002. *Genome Announc* 3:e00824-15.
5. Álvarez-Fraga L, Pérez A, Rumbo-Feal S, Merino M, Vallejo JA, Ohneck EJ, Edelmann RE, Beceiro A, Vázquez-Ucha JC, Valle J, Actis LA, Bou G, Poza M. 2016. Analysis of the role of the LH92_11085 gene of a biofilm hyper-producing *Acinetobacter baumannii* strain on biofilm formation and attachment to eukaryotic cells. *Virulence* 7:443–455.

6. Aranda J, Poza M, Pardo BG, Rumbo S, Rumbo C, Parreira JR, Rodriguez-Velo P, Bou G, Rodríguez-Velo P, Bou G. 2010. A rapid and simple method for constructing stable mutants of *Acinetobacter baumannii*. BMC Microbiol 10:279.
7. Rajamohan G, Srinivasan VB, Gebreyes WA. 2010. Molecular and functional characterization of a novel efflux pump, AmvA, mediating antimicrobial and disinfectant resistance in *Acinetobacter baumannii*. J Antimicrob Chemother 65:1919–1925.
8. Magnet S, Courvalin P, Lambert T. 2001. Resistance-Nodulation-Cell Division-type efflux pump involved in aminoglycoside resistance in *Acinetobacter baumannii* strain BM4454. Antimicrob Agents Chemother 45:3375–3380.
9. Chau S-L, Chu Y-W, Houang ETS. 2004. Novel resistance-nodulation-cell division efflux system AdeDE in *Acinetobacter* genomic DNA Group 3. Antimicrob Agents Chemother 48:4054–4055.
10. Coyne S, Rosenfeld N, Lambert T, Courvalin P, Périchon B. 2010. Overexpression of resistance-nodulation-cell division pump AdeFGH confers multidrug resistance in *Acinetobacter baumannii*. Antimicrob Agents Chemother 54:4389–4393.
11. Damier-Piolle L, Magnet S, Bremont S, Lambert T, Courvalin P. 2008. AdeIJK, a resistance-nodulation-cell division pump effluxing multiple antibiotics in *Acinetobacter baumannii*. Antimicrob Agents Chemother 52:557–562.
12. Eijkelkamp BA, Hassan KA, Paulsen IT, Brown MH. 2011. Development of a high-throughput cloning strategy for characterization of *Acinetobacter baumannii* drug transporter proteins. J Mol Microbiol Biotechnol 20:211–219.

13. Srinivasan VB, Rajamohan G, Pancholi P, Marcon M, Gebreyes WA. 2011. Molecular cloning and functional characterization of two novel membrane fusion proteins in conferring antimicrobial resistance in *Acinetobacter baumannii*. J Antimicrob Chemother 66:499–504.
14. Su X-Z, Chen J, Mizushima T, Kuroda T, Tsuchiya T. 2005. AbeM, an H⁺-coupled *Acinetobacter baumannii* multidrug efflux pump belonging to the MATE family of transporters. Antimicrob Agents Chemother 49:4362–4364.
15. Srinivasan VB, Rajamohan G, Gebreyes WA. 2009. Role of AbeS, a novel efflux pump of the SMR family of transporters, in resistance to antimicrobial agents in *Acinetobacter baumannii*. Antimicrob Agents Chemother 53:5312–5316.
16. Li L, Hassan KA, Brown MH, Paulsen IT. 2016. Rapid multiplexed phenotypic screening identifies drug resistance functions for three novel efflux pumps in *Acinetobacter baumannii*. J Antimicrob Chemother 71:1223–1232.
17. Hassan KA, Liu Q, Henderson PJF, Paulsen IT. 2015. Homologs of the *Acinetobacter baumannii* AceI transporter represent a new family of bacterial multidrug efflux systems. MBio 6(1).
18. Ribera A, Roca I, Ruiz J, Gibert I, Vila J. 2003. Partial characterization of a transposon containing the *tet(A)* determinant in a clinical isolate of *Acinetobacter baumannii*. J Antimicrob Chemother 52:477–480.
19. Martí S, Fernández-Cuenca F, Pascual A, Ribera A, Rodríguez-Baño J, Bou G, Miguel Cisneros J, Pachón J, Martínez-Martínez L, Vila J, Grupo de estudio de infección hospitalaria (GEIH). 2006. Prevalence of the *tetA* and *tetB* genes as mechanisms of resistance to tetracycline and minocycline in *Acinetobacter baumannii* clinical isolates. Enferm Infecc Microbiol Clin 24:77–80.


- 210 20. Roca I, Marti S, Espinal P, Martínez P, Gibert I, Vila J. 2009. CraA, a major
211 facilitator superfamily efflux pump associated with chloramphenicol resistance in
212 *Acinetobacter baumannii*. Antimicrob Agents Chemother 53:4013–4014.
- 213 21. Hamidian M, Nigro SJ, Hartstein RM, Hall RM. 2017. RCH51, a multiply
214 antibiotic-resistant *Acinetobacter baumannii* ST103IP isolate, carries resistance
215 genes in three plasmids, including a novel potentially conjugative plasmid
216 carrying *oxa235* in transposon Tn6252. J Antimicrob Chemother 72:1907–1910.
- 217 22. Fournier P-E, Vallenet D, Barbe V, Audic S, Ogata H, Poirel L, Richet H, Robert
218 C, Mangenot S, Abergel C, Nordmann P, Weissenbach J, Raoult D, Claverie J-
219 M. 2006. Comparative genomics of multidrug resistance in *Acinetobacter*
220 *baumannii*. PLoS Genet 2:e7.
- 221 23. Sharma A, Sharma R, Bhattacharyya T, Bhando T, Pathania R. 2017. Fosfomycin
222 resistance in *Acinetobacter baumannii* is mediated by efflux through a major
223 facilitator superfamily (MFS) transporter—AbaF. J Antimicrob Chemother
224 72:68–74.
- 225 24. Lin M-F, Lin Y-Y, Lan C-Y. 2017. Contribution of EmrAB efflux pumps to
226 colistin resistance in *Acinetobacter baumannii*. J Microbiol 55:130–136.
- 227 25. Hassan KA, Brzoska AJ, Wilson NL, Eijkelkamp BA, Brown MH, Paulsen IT.
228 2011. Roles of DHA2 family transporters in drug resistance and iron homeostasis
229 in *Acinetobacter* spp. J Mol Microbiol Biotechnol 20:116–124.
- 230 26. Wang S, Li W, Liu S, Xu J. 2016. RaptorX-Property: a web server for protein
231 structure property prediction. Nucleic Acids Res 44:W430-435.
- 232 27. Schrödinger, L. L. C. (2010). The PyMOL Molecular Graphics System, Version

233 1.3r1. Available at: <http://www.pymol.org>.

234

Fig. 1

(A)



(B)

