



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Changes in salivary analytes in canine parvovirus: A high-resolution quantitative proteomic study



Lorena Franco-Martínez^a, Asta Tvarijonavičiute^a, Anita Horvatić^b, Nicolas Guillemain^b, José Joaquín Cerón^a, Damián Escribano^{a,c}, David Eckersall^{b,d}, Meriç Kocatürk^e, Zeki Yilmaz^e, Elsa Lamy^f, Silvia Martínez-Subiela^{a,*}, Vladimir Mrljak^b

^a Interdisciplinary Laboratory of Clinical Analysis Interlab-UMU, Regional Campus of International Excellence Mare Nostrum, University of Murcia, Espinardo, Murcia 30100, Spain

^b ERA Chair FP7, Internal Diseases Clinic, Faculty of Veterinary Medicine, University of Zagreb, Heinzelova 55, 10 000 Zagreb, Croatia

^c Department of Animal and Food Science, School of Veterinary Medicine, Universitat Autònoma de Barcelona, 08193, Bellaterra, Spain

^d Institute of Biodiversity, Animal Health and Comparative Medicine, College of Medicine, Veterinary Medicine and Life Sciences, University of Glasgow, Glasgow G61 1QH, United Kingdom

^e Department of Internal Medicine, Faculty of Veterinary Medicine, Uludağ University, 16059, Bursa, Turkey

^f ICAAM - Institute of Mediterranean Agricultural and Environmental Sciences, University of Evora, Portugal

ARTICLE INFO

Keywords:

Bioinformatics
Biomarker
Canine parvovirus
Dog
Label-based proteomics
Saliva

ABSTRACT

The present study evaluated the changes in salivary proteome in parvoviral enteritis (PVE) in dogs through a high-throughput quantitative proteomic analysis. Saliva samples from healthy dogs and dogs with severe parvovirus that survived or perished due to the disease were analysed and compared by Tandem Mass Tags (TMT) analysis. Proteomic analysis quantified 1516 peptides, and 287 (corresponding to 190 proteins) showed significantly different abundances between studied groups. Ten proteins were observed to change significantly between dogs that survived or perished due to PVE.

Bioinformatics' analysis revealed that saliva reflects the involvement of different pathways in PVE such as catalytic activity and binding, and indicates antimicrobial humoral response as a pathway with a major role in the development of the disease. These results indicate that saliva proteins reflect physiopathological changes that occur in PVE and could be a potential source of biomarkers for this disease.

1. Introduction

Parvoviruses are small icosahedral viruses that infect many animal species including humans worldwide. Canine parvovirus type-2 is the causal agent of canine parvoviral enteritis (PVE), a severe disease that causes nearly 100% morbidity and up to 10–90% mortality in unvaccinated adults and puppies, respectively [1]. The clinical presentation of PVE ranges from mild to severe, based on clinical signs and physical examination [2,3]. The most common clinical signs in puppies include acute severe vomiting and bloody diarrhoea, fever, dehydration and lethargy [4]. Complications such as cardiac damage, which may result in death or permanent myocardial damage [5], and systemic inflammatory response syndrome (SIRS) prior to sepsis have been described [6].

In dogs, saliva has proven to be a useful biofluid for the diagnostic of several conditions such as leishmaniosis [7] and helicobacter

infection [8], as well as for evaluation of stress [9]. The use of non-invasive specimens such as saliva provides several advantages compared to invasive methods, being safer for the personal and the patient, easier to collect, pain-free and causes reduced sampling stress [10]. Saliva can have potential advantages versus others non-invasive samples such as faeces, which usually need a laborious pre-treatment prior to its analysis. In particular, in the case of PVE, it has been described a poor sensitivity of faeces when used in rapid antigen ELISA test [11]. For this, it is envisioned that saliva can play an increasingly important role in the early diagnosis and monitoring of diseases [12].

However, to the author's best knowledge, there are no reported studies about changes in saliva composition in PVE. The knowledge of the changes that occur in saliva in dogs with parvovirus could help to detect possible biomarkers in this fluid for PVE disease.

Proteomics sciences focus on the study of the proteome content of cells, organs and organisms [13]. Proteomic techniques have become

* Corresponding author.

E-mail address: silviam@um.es (S. Martínez-Subiela).

<https://doi.org/10.1016/j.cimid.2018.09.011>

Received 5 July 2018; Received in revised form 17 September 2018; Accepted 18 September 2018

0147-9571/© 2018 Elsevier Ltd. All rights reserved.

widely used with the aim of discovering disease-related biomarkers in diverse biological fluids that could be used in routine practice for early diagnosis, prognosis or treatment monitoring [14,15]. Over the last several years, novel technologies such as the use of isobaric tags have greatly increase the sensitivity of the proteomic analysis [16–18]. Tandem Mass Tags (TMT) allow for relative simultaneous quantification of differentially labelled peptides [19] since each sample is marked and distinguished by differences on their reporter ion masses [20,21]. However, there is still a scarcity of reports using TMT technology in saliva samples, and no reports have been found describing the use of TMT to detected possible biomarkers of diagnosis and prognosis in PVE.

The objective of the present study is to evaluate possible changes in saliva proteins associated to canine parvovirus infection. This would help to clarify if saliva components can reflect the physiopathological changes associated to the disease and to evaluate the possibility of identifying potential biomarkers of PVE in saliva samples. For this, saliva samples were collected from healthy dogs (control group) and from dogs with PVE that survived or died because the infection. Saliva samples were analysed by quantitative high-throughput TMT-based proteomic approach enabling the identification of significantly varying proteins among different groups. Subsequently, the differentially expressed proteins were used as a starting point for creating protein interaction networks.

2. Materials and methods

2.1. Animals

A total of 11 client-owned dogs that were presented to the Veterinary Teaching Hospital, Small Animal Clinics, Bursa Uludag University, Bursa, Turkey, of different breeds were involved in this study (Table 1). Parvovirus infection was diagnosed by compatible clinical (acute bloody diarrhoea, vomiting, anorexia, dehydration etc.) and haematological signs (leukopenia, neutropenia and lymphopenia) in combination with the positive test results of the commercially available faecal diagnostic test (Antigen Rapid PVE kit, Animal Genetics, Inc., Suwon, Korea). Selected clinical (body temperature and heart and respiratory rates) and haematological (leukogram, eritrogram and thrombogram) findings were used to assess the general health status of the animal. Dogs were excluded if they were co-infected with other viral (canine distemper or coronavirus), parasite (coccidiosis, giardiasis, ascaridiosis etc.), or vector borne diseases by use of faecal screening test, fecal microscopic examinations, or speed test for ehrlichiosis, anaplasmosis, lyme, and dirofilariasis (Anigen Rapid CaniV-4, Bionote, Korea), respectively.

The dogs were divided into three groups according to their clinical condition and development of the disease. A group with 5 healthy dogs was considered as control group. Virus is shed for a few days before the onset of clinical signs. Therefore, even if all dogs in control group were

considered healthy based on the clinical, haematological and serum biochemistry findings (comprehensive profile, VetScan, Abaxis), faecal screening test for CPV-Ag were applied. Dogs with negative test result were enrolled to study. All dogs in test group were treated as described previously [22]. Briefly, fluid replacement therapy (dextrose 5% with a balanced crystalloid solution - lactated ringer, and hydroxyethyl starch), anti-emetics (metoclopramid 0.2-0.4 mg/kg, BID, subcutaneously, and ranitidin 2–4 mg/kg, BID, intramuscularly), parenteral antibiotics (ampicillin 30 mg/kg, BID + amikasin 5–10 mg/kg, BID + metronidazol 25 mg/kg, BID), immune stimulator (imunex), probiotics and vitamins (E and C) were used. Dogs with clinical diagnosis of severe parvovirus were divided into two groups according to the progress of the disease: 3 dogs that survived (survival group) (age 3.67 ± 0.94 months, 1 male and 2 female) and 3 that died because of the disease (non-surviving group) (age 6.67 ± 1.25 months, 2 males and 1 female). Therefore, dogs were classified according to the severity of their clinical and haematological signs (Table 1). There was a similarity of presented clinical signs and its duration (2–4 days) in dogs within test group. Despite the treatment, three dogs died within 5 days (one on day 3, one on day 4 and other on day 5).

All the procedures were approved by the Local Ethical Committee of the University of Uludag.

2.2. Saliva sampling

At least 0.5 ml of unstimulated saliva specimens were collected from each patient by placing small cotton swabs around the mouth. When the cotton swabs were thoroughly moist, they were placed in collection devices (Salivette saliva collection tube / V-Bottom, Sarstedt, Aktiengesellschaft & Co, Nümbrecht, Germany), centrifuged (3000 x g for 10 min, 4 °C), and the supernatant was stored at -80 °C until analysis [23]. Due to dehydration and subsequent dry mouth, for some animals the sampling protocol was performed twice in order to obtain at least 0.5 ml of unstimulated saliva.

2.3. Proteomic study of saliva samples and LC-MS/MS analysis

From each sample, 35 µg of acetone-precipitated proteins were subjected to reduction, alkylation, digestion and labelled using 6-plex Tandem Mass Tag reagents according to manufacturer instructions (Thermo Scientific) with some modification, as described previously [24]. In short, 35 µg of sample and internal standards were reduced with 200 mM DTT (Sigma-Aldrich), alkylated with 375 mM iodoacetamide (Sigma- Aldrich) and precipitated with ice-cold acetone (VWR, Pennsylvania, USA) overnight. Samples were then centrifuged and acetone was decanted. Pellets were resuspended with 50 µl of 100 mM TEAB buffer and digested with trypsin (Promega) overnight at 37 °C (2.5 µg of trypsin per 100 µg of protein). TMT label reagents were equilibrated to room temperature, resuspended in anhydrous

Table 1
Main exploratory and haematological findings in healthy and dogs with PVE.

ID	Breed	Gender / age (months)	T °C	P /min	R /min	WBC / N / L X 10 ³ /µL	Hct %	Total protein gr/dL	Group
1	Mixed	M / 2	39.2	120	22	11.5 / 8.7 / 2.1	27.6	ND	Healthy
2	Mixed	M / 2	39.3	110	24	10.7 / 6.8 / 3.2	36.1	ND	Healthy
3	Mixed	F / 2	38.9	112	18	12.9 / 7.8 / 4.2	32.8	5.3	Healthy
4	Mixed	M / 5	38.7	124	16	13.5 / 10.2 / 2.0	40.6	451	Healthy
5	Mixed	M / 10	38.5	114	16	15.0 / 12.8 / 1.5	48.8	6.9	Healthy
6	Rottweiler	F / 5	39.8	128	44	2.1 / 0.6 / 1.4	31.8	4	Survival
7	Cocker	M / 3	38.3	200	44	2.5 / 1.0 / 1.3	39.1	4.7	Survival
8	Mixed	F / 3	37.5	148	20	1.6 / 0.6 / 0.8	29.7	4.1	Survival
9	Anatolian sheepdog	M / 5	39.8	120	44	0.3 / 0.1 / 0.1	34.2	3.7	Not-survival
10	Rottweiler	M / 7	40.7	108	28	1.2 / 0.3 / 0.8	43.2	5.6	Not-survival
11	Cocker	F / 8	37.0	108	32	0.3 / 0.0 / 0.2	37.5	3.3	Not-survival

M: male, F: female, ND: not determined, P: pulsation, R: respiration, Hct: Hematocrite, WBC: white blood cell count, N: Neutrophil count, L: Lymphocyte count.

acetonitrile LC–MS grade (Thermo Scientific) and added to each sample. Labelling reaction was incubated for 1 h at room temperature and then quenched by adding 5% hydroxylamine (Thermo Scientific) for 15 min. Samples were then combined at equal amounts and 6 µg of each mixed sample set was placed in a well of a microplate, vacuum-dried and stored at -20 °C before further LC–MS/MS analysis. The LC–MS/MS analysis was performed on Dionex Ultimate 3000 RSLC nano flow system (Dionex, Camberley, UK) and Orbitrap Elite mass spectrometer (Thermo Fisher Scientific) as described elsewhere [25].

2.4. Statistical analysis

In order to compare the abundances of peptides detected in proteomic analysis between 2 groups in 3 comparisons (Control versus Survival, Control versus Non-surviving and Survival versus Non-surviving), data were normalized by logarithmic transformation and Student's *t*-test was used (one-tailed, unpaired) were performed. In all cases, values of $P < 0.05$ were considered to be significant. Fold changes (FC) have been calculated as follow $FC = \log_2(\text{Group 1} / \text{Group 2})$. Statistics were performed with RStudio (v1.0.143) (R Studio Team. RStudio [26] “RStudio Team. RStudio: Integrated Development Environment for R [Internet]. Boston, MA: RStudio, Inc.; 2015. Available from: <http://www.rstudio.com/>,” 2015).

2.5. GO pathways

The data obtained in the proteomic study were used for the Gene Ontology (GO) analysis. Canine genes encoding proteins differentially expressed were converted to their human orthologs using the Ensembl orthologs database and its BioMart tool for data mining (www.ensembl.org). Obtained genes were used to determine the GO terms over-represented in two conditions (Survival and Non-surviving), by the utilization of the Cytoscape (v3.6.1) plug-in ClueGO (v2.5.0) [27,28] on the *Homo sapiens* GO-biological process (22/01/2018). GO terms over-represented for each group were submitted to analysis by REVIGO (allowed similarity = 0.7, SimRel) to remove redundant GO terms and groups related GO terms based on their functional description [29]. Finally, pathways interactomes were designed in Cytoscape using the radial layout.

3. Results

3.1. High resolution quantitative proteomic analysis

High-resolution quantitative proteomic analysis allowed the identification of 1516 canine peptides from the eleven non-depleted canine saliva samples. For the comparison Control versus Survival, 148 peptides were differentially expressed, corresponding to 90 unique proteins. For the comparison Control versus Non-surviving, 125 peptides were differentially expressed, corresponding to 90 unique proteins. For the comparison Survival versus Non-surviving, 14 peptides were differentially expressed, corresponding to 10 unique proteins. Therefore a total of 190 proteins were identified that differed between groups.

Of the 90 differentially expressed proteins between survival and control groups, the proteins most down-regulated in the survival group were cathelicidin antimicrobial peptide (CAMP), rho-GDP dissociation inhibitor beta (ARHGDIB), apolipoprotein A-1 (APO-A1), neutrophil elastase (ELANE), matrix metalloproteinase-9 (MMP9), EF-hand domain containing protein D2 (EFHD2), CD177 antigen (CD177), plastin-2 (LCP1), retinol binding protein 4 (RBP4), and maltase-glucoamylase intestinal (MGAM). Epididymal – specific lipocalin-9 (LCN9), BPI fold-containing family B member 2 (BPIFB2), lymphocyte antigen 6D (LY6D), desmoglein (-1 and -3, DSG1 and DSG3, respectively), alpha-2-macroglobulin-like protein 1 (A2ML1), democollin-2 (DSC2), leucine-rich-alpha-2-glycoprotein (LRG1), polymeric immunoglobulin receptor (PIGR), and L-amino acid oxidase (IL4I1) were the proteins most

regulated in the survival group. Of these, CAMP and ARHGDIB were the proteins most down-regulated (-2.29 and -2.26-fold lower expression, respectively), while LCN9 and BPIFB2 were the most up-regulated (2.28 and 1.62-fold higher) in the survival group compared to the control group.

When non-surviving and control groups were compared, 90 proteins showed differences in abundance. The proteins most down-regulated were ARHGDIB, CAMP, APOA1, hemoglobin subunit beta (HBB), RBP4, CD177, MMP9, vitamin D binding protein (GC) and ELANE. On the other hand, immunoglobulin heavy chain (IGH), LCN9, clusterin (CLU), trefoil factor 1 (TFF1), LY6D, olfactomedin 4 (OLFM4), LRG1, glyoxalase I (GLO1), BPIFB2, and DSG3 were the proteins most up-regulated.

Finally, when the two groups of dogs with parvovirus were compared, 10 proteins showed different abundances. Ezrin, allergen Dog 1, lactoperoxidase, L-lactate dehydrogenase A, cystatin-M and macrophage-capping protein were higher in survival group, whereas alpha-actinin-1, peptidyl-prolyl cis-trans isomerase A, ribosylidihydronicotinamide dehydrogenase and lactoylglutathione lyase were up-regulated in the dogs that died due to the disease.

3.2. Bioinformatics

The GO analysis performed using Cytoscape plug-in ClueGO identified 34 GO terms between survival and controls, which were filtered for redundancies and then grouped into 12 main groups using REVIGO. The functional groups included antimicrobial humoral response, entry of bacterium into host cell, tissue homeostasis, heterotypic cell-cell adhesion and endoplasmic reticulum mannose trimming (Fig. 1).

When comparing non-surviving and control groups, 42 GO terms were identified, which were grouped into 11 main functional groups including negative regulation of cellular response to growth factor stimulus, antimicrobial humoral response, transition metal ion homeostasis, endoplasmic reticulum mannose trimming and glutathione derivative biosynthesis (Fig. 2).

Comparison of survival versus non-surviving groups did not identify enough proteins significantly expressed to generate a reliable GO analysis using only experimental results.

Finally, GO terms most represented in this study after being filtered by REVIGO and depicted by Cytoscape are shown in Figs. 3–6. When the network from control and survival group was analysed, a negative regulation of wound healing (related with the down-regulation of fibrinogen FGA, FGB and FGG genes) and negative regulation of cellular response to growth factor stimulus (related to the up-regulation of the ubiquitin complex composed by RPS27 A, UBC, UBB and UBA52 genes) nodes demonstrated central roles in the development of the disease since they are involved in many pathways.

The network obtained from comparison of non-surviving and control groups suggested that the over-expressed UBC gene (which expresses ubiquitin C protein) appears to have a central role by affecting or being affected by numerous pathways identified as related with non-surviving group in parvovirus. Interestingly, this ubiquitin C protein is related only with GO terms over-expressed in non-surviving group. This indicates that, in the present study, all cellular pathways which use ubiquitin form a pertinent group, which is positively correlated with the non-survival status. Similar to the previously mentioned network, fibrinogen expression is down-regulated in saliva of dogs with parvovirus.

When merging fold changes data and over-represented pathways in both non-surviving and survival groups, the obtained network highlights that antimicrobial humoral response was shown as central term. This node is the only one which is over-represented in both parvovirus groups (survival and non-surviving) but with different expression when compared to controls: it was lower-expressed in the non-surviving group, but was not differentially expressed in survival group as compared to controls. Antibacterial humoral response, negative regulation of response to external stimulus, and killing of cells of other organisms




Fig. 1. Over-represented GO terms in dogs that survived canine parvovirus, grouped by REVIGO, based on their description. Leading GO terms (N = 12) for each group was defined as the highest significance inside the group.

are the three down-regulated nodes which are directly linked with antimicrobial humoral response. The network also showed 14 proteins surrounding this central node, 9 of which were down-regulated (ELANE, kininogen 1 [KNG-1], FGA, FGB, S100 calcium binding protein

A8 [S100A8], S100 calcium binding protein A12 [S100A12], CAMP, lysozyme [LYZ] and HRG), and 5 up-regulated (beta-2 microglobulin [B2M], deleted in malignant brain tumours 1 [DMBT1], BPI fold-containing family B member 1 [BPIFB1], BPIFB2, and CLU) when




Fig. 2. Over-represented GO terms in dogs that died due to canine parvovirus, grouped by REVIGO, based on their description. Leading GO terms (N = 12) for each group was defined as the highest significance inside the group.




Fig. 3. Over-represented GO terms generated by the comparison of dogs that survived and dogs that died due to parvovirus, grouped by REVIGO, based on their description. Leading GO term (N = 9) for each group was defined as the highest significance inside the group.

compared to controls.

4. Discussion

This paper was focused in the identification of the differentially expressed proteins in saliva and in describing the biological roles which may be affected due to PVE. The present study identified for the first time 190 differentially expressed proteins in saliva from healthy dogs and dogs with severe parvovirus using TMT technology, 10 of which were differentially expressed between animals that survived and perish due to the disease. These differences in proteins indicate that saliva can reflect several physiological pathways in canine parvovirus.

Overall changes in protein expression in saliva from dogs with parvovirus obtained in this study by high-resolution quantitative proteomic analysis suggested alterations in coagulation and inflammation systems, which are closely related pathways since the activation of one mechanism may lead to the activation of the other [30]. Coagulation products such as thrombin can promote inflammatory responses, while inflammation is known to suppresses anticoagulation mechanisms [31]. The immune system provides protection against infection; however, its activation needs to be tightly regulated in order to prevent excessive inflammation and subsequent damage to the host [32].

From all the proteins that change in saliva of dogs with parvovirus, two that are most down-regulated and the two proteins that increased the most in the canine patients are worthy of specific mention. The two most down-regulated proteins in the dogs with parvovirus that survived, cathelicidin antimicrobial peptide (CAMP) and Rho-GDP dissociation inhibitor beta (ARHGDI), have protective role against infections and are involved in the inflammatory response. Cathelicidins are a group of short cationic antimicrobial peptides with an important role in the protection against infections [33–35]. The main source of CAMPs are the specific granules of neutrophils, which are released upon neutrophil activation [36,37], although they can be produced by other cells such as epithelial cells at mucosal surfaces [38]. CAMPs are known

to regulate the activation of a wide variety of toll-like receptors, which plays important roles in the modulation of the inflammatory response during infections [39,40]. In addition, antimicrobial activity has been described in CAMPs, and the loss of cathelicidin expression has been reported to increase the susceptibility to *Escherichia coli* infection in mice [33,34]. Rho-GDP dissociation inhibitor beta (ARHGDI) is primarily expressed in cells of a hematopoietic lineage and belongs to the RhoGDI family, which are known to regulate Rho-GTPases [41]. ARHGDI showed an ability to decrease human immunodeficiency virus type 1 (HIV-1) replication in cell lines and can be used as a biomarker of HIV-1 infection [42]. ARHGDI expression has been also proposed as suppressor of metastasis and predictor of prognosis in human patients with bladder cancer, in which ARHGDI positive tumours are related with longer disease-free and disease-specific survival rates [43].

On the other hand, the two most up-regulated proteins in dogs with parvovirus that survived, were lipocalin-9 (LCN9) and BPI fold-containing family B member 2 (BPIFB2). Lipocalins are a family of proteins that usually carry lipids or other hydrophobic or amphiphilic compounds accommodated in a cavity within their conformation, including steroids, hormones and metabolites such as vitamins and cofactors [44]. Furthermore, lipocalins have been reported to be modulators of cell growth and metabolism, and regulators of the immune response [45,46]. Although little is known about LCN9, a related protein, neutrophil gelatinase-associated lipocalin, NGAL, has been associated with bacterial infection in humans [47–49]. BPIFB2 belongs to a family of proteins considered as a key component of the innate immune response against Gram-negative bacteria [50], but the physiological functions of the BPIFB family still remain largely unknown [51]. Despite this, a previous study on BPIFB2 has reported the involvement of BPIFB proteins regulating enterovirus infection, with BPIFB6 assisting the enterovirus replication by controlling secretory pathway trafficking and Golgi complex morphology [51].

In addition to these 4 proteins, 86 other proteins were significantly differentially expressed in saliva of parvovirus infected dogs compared




Fig. 4. *In silico* inferred interactome network of identified GO terms over-represented in canine parvovirus (survival versus control groups). Differentially expressed proteins interacting with at least 1 term were added. Radial layout was applied and the GO group leader terms are in dark blue text. Nodes colours represent the group of GO terms (determined by ReviGO) and border are represented in green and red for over-expressed and lower-expressed proteins, respectively, when compared to control group.




Fig. 5. *In silico* inferred interactome network of identified GO terms over-represented in canine parvovirus (dead versus control groups). Differentially expressed proteins interacting with at least 1 term were added. Radial layout was applied and the GO group leader terms are in dark blue text. Nodes colours represent the group of GO terms (determined by ReviGO) and border are represented in green and red for over-expressed and lower-expressed proteins, respectively, when compared to control group.

to healthy dogs. By analysing the over-represented cellular pathways defined by those proteins by gene ontology analyses, it could be observed that they are involved in other physiopathological pathways such as catalytic activity, binding, transporter activity, antioxidant activity, structural molecule activity, receptor activity and translation regulator activity. Since more than 85% of the genes differentially expressed belong to the proteins related with catalytic activity and binding, the discussion will focus on these and related proteins.

In regard to the proteins related to catalytic activity, some were down-regulated, including, apolipoprotein A-1 (APO-A1), neutrophil esterase (ELANE), complement C3 (C3) and histidine-rich glycoprotein (HRG). APO-A1 is the major protein component of high density lipoproteins in human and other mammals [52] and the decreased levels APO-A1 observed in the present study are in concordance with other responses described in different canine diseases such as leishmaniosis [53] or canine idiopathic dilated cardiomyopathy [25], suggesting that APO-A1 could be a feasible biomarker in dogs. Lower expression of neutrophil esterase was observed in both groups of dogs with PVE when compared to control group. Neutrophil esterase is a serine proteinase with important antimicrobial effects that are released by neutrophils and enhances inflammatory responses. Genetic mutations in its gene (ELANE) has been identified in forms of hereditary neutropenia in humans and dogs [54]. There is a case reported of a mutation in ELANE gene as the cause of chronic parvovirus in a woman [55]. The activation of complement C3 supports local inflammatory responses against pathogens and initiates the humoral immune response, playing a central role in the activation of the complement system [56]. Increased C3

expression has also been reported in dogs with canine babesiosis [57]. In our study, different hypotheses could explain the down-regulation of complement C3 in saliva of dog with PVE such as a reduction due to cleavage of the protein in order to activate complement cascade, or as a consequence of an immunodeficiency [58]. Histidine-rich glycoprotein is a relatively abundant plasma protein with a wide range of targets and functions, such as regulating clotting and fibrinolysis, and angiogenesis [59,60]. HRG roles in anti-inflammatory effects and pathogen control have been proposed since, after experimental *Streptococcus pyogenes* infection, HRG-deficient mice showed higher bacterial burden and pro-inflammatory cytokines than wild-type mice [61].

Thirty-four of the proteins differentially expressed in saliva between controls and dogs with parvovirus have binding activity, including many of the proteins described above such as APO-A1, A2M, C3 and LCN9. Other important proteins differentially expressed in saliva between healthy and dogs with parvovirus with binding activity are desmoglein-1 and -3 (DSG1 and DSG3, with 1.39 and 1.43- fold higher in dogs with parvovirus, respectively), and clusterins (CLU, 0.71-fold higher). Desmogleins are a family of calcium-binding transmembrane glycoproteins which play an important role in the desmosomes transmembrane component [62]. In humans, salivary levels of DSG1 and DSG3 are correlated with serum anti DSG titres, proposing salivary DSG as suitable markers of diagnosis of *Pemphigus vulgaris* [63]. Clusterins are secretory glycoproteins that are involved in multiple physiopathological processes, including lipid metabolism, tissue remodelling, reproduction, transport and cell apoptosis [64]. These proteins have been classically considered as markers of cell death [65]; however, protective




Fig. 6. *In silico* inferred interactome network of identified GO terms over-represented in canine parvovirus (survival and dead versus control groups). Differentially expressed proteins interacting with at least 1 term were added. Radial layout was applied. Nodes fill colour figure protein and GO terms regulation (red for negatively and green for positively regulated proteins) and node border represent the presence of the term in dead only (black), survival only (white) or in both groups (grey).

roles such as anti-apoptotic or prevention of complement attacking complex functions of clusterins have been suggested [66,67].

The bioinformatics analysis of the *in silico* inferred proteins network allowed the generation of more information from the experimental results, including data on intracellular pathways. This revealed the impact on processes such as an inhibition of humoral and cellular responses, and a positive effect in the entry of bacterium into host cell. These results are in concordance with the pathogenicity of PVE in which, after the ingestion, the virus replicates in the lymphoid tissue in the dog's throat and then spreads to the bloodstream infecting rapidly dividing cells, resulting in depletion of lymphocytes in lymph nodes and necrosis of the intestinal crypts [68]. These mucosal intestinal lesions may give rise to secondary bacterial invasions from the intestines, leading to systemic inflammatory response syndrome (SIRS), sepsis and endotoxemia [69–72].

Bioinformatics allowed several pathways and proteins to be highlighted which could be useful in the diagnosis of PVE with saliva samples. For example, the network created by comparison of survival and control groups revealed that fibrinogen and ubiquitin complexes could play important roles in the development of the disease. The down-regulation of fibrinogen observed in the present study is in contrast with other reports that showed increased levels of fibrinogen in dogs with parvovirus [73], leishmaniosis [74] or babesiosis [75]; although dogs with uncomplicated babesiosis showed higher levels of fibrinogen than those with SIRS or multiple organ dysfunction syndrome (MODS) [75]. This could be explained by the fact that, although fibrinogen is considered as a positive acute phase protein in dogs, in disseminated intravascular coagulation (DIC) and hyperfibrinolysis the

concentration of fibrinogen falls due to fibrinogen cleavage to form soluble fibrin monomers [76,77]. Progressive decrease of fibrinogen levels was also reported in dogs during acute haemorrhage [78], as occurs in parvovirus. On the other hand, ubiquitins have been shown to be involved in numerous intracellular processes such as cell cycle control, apoptosis, DNA repair, regulation of transcription, stress responses, and targeting cellular proteins for degradation by the 26S proteasome [79]. As described in previous reports, viruses often make use of the ubiquitin conjugation to target for degradation of cellular proteins that might otherwise affect the replication of the virus [80].

One of the limitations of the present study may be the sample size, which could have limited the potential to identify proteins that are reproducibly modulated in concentration in dogs with PVE due to the biological inter-individual variations. Therefore this could be considered as a pilot study, and further validation studies with a larger number of dogs should be made to confirm these findings. The changes in protein expression and the subsequent bioinformatics analysis described in the present study were based on the analysis of saliva and, therefore, further studies are desirable to discern if they reflect systemic alterations. Nevertheless, despite these limitations, our results indicate that PVE can produce changes in saliva proteome that can reflect an inflammatory response and changes in coagulation system and immune response – among other biological processes.

This paper was focused on the identification of the differentially expressed proteins in saliva and in the description of the biological roles of the proteins affected due to the disease. Further large-scale clinical studies with larger populations would be desirable to elucidate if the proteins identified in this report can be of help in the earlier detection,

predicting the outcome, or giving information to the clinician to adjust the treatment. If these studies are successful, these proteins can be used for developing rapid diagnostic tests such as lateral flow kits in saliva for clinical practice.

5. Conclusions

TMT-based proteomic analysis allowed the identification of 190 proteins with different abundances between healthy dogs and dogs with PVE, most of them not previously reported in canine parvovirus. These proteins are involved in various physiopathological processes such as coagulation, inflammation and defence mechanisms, and could be considered as potentially suitable biomarkers of the disease. This study demonstrated that saliva proteome is a suitable biofluid for both study and diagnostic of canine parvovirus.

Disclosure statement

The authors report no other conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Acknowledgements

LFM was granted with predoctoral contract 'FPU' of University of Murcia, Spain. AT has a post-doctoral fellowship "Juan de la Cierva Incorporación" and DE has a post-doctoral fellowship "Juan de la Cierva Formación" supported by the "Ministerio de Economía y Competitividad", Spain. This work was supported by a grant from the Program for Research Groups of Excellence of the Seneca Foundation, Murcia, Spain (grant 19894/GERM/15) and European Commission ERA chair FP7 grant (VetMedZg #621394).

References

- [1] S. Nandi, M. Kumar, Canine parvovirus: current perspective, *Indian J. Virol.* 21 (2010) 31–44, <https://doi.org/10.1007/s13337-010-0007-y>.
- [2] V. Matijatko, V. Mrljak, I. Kiš, N. Kučer, J. Foršek, T. Živičnjak, Ž. Romić, Z. Šimec, J.J. Ceron, Evidence of an acute phase response in dogs naturally infected with *Babesia canis*, *Vet. Parasitol.* 144 (2007) 242–250, <https://doi.org/10.1016/j.vetpar.2006.10.004>.
- [3] D. Panda, R.C. Patra, S. Nandi, D. Swarup, Oxidative stress indices in gastroenteritis in dogs with canine parvoviral infection, *Res. Vet. Sci.* 86 (2009) 36–42, <https://doi.org/10.1016/j.rvsc.2008.05.008>.
- [4] M. Kocaturk, A. Tvarijonavičiute, S. Martínez-Subiela, F. Tecles, O. Eralp, Z. Yilmaz, J.J. Ceron, Inflammatory and oxidative biomarkers of disease severity in dogs with parvoviral enteritis, *J. Small Anim. Pract.* 56 (2015) 119–124, <https://doi.org/10.1111/jsap.12250>.
- [5] J. Ford, L. McEndaffer, R. Renshaw, A. Molesan, K. Kelly, Parvovirus infection is associated with myocarditis and myocardial fibrosis in young dogs, *Vet. Pathol.* 54 (2017) 964–971, <https://doi.org/10.1177/0300985817725387>.
- [6] A.J. Mohr, A.L. Leisewitz, L.S. Jacobson, J.M. Steiner, C.G. Ruaux, Da Williams, Effect of early enteral nutrition on intestinal permeability, intestinal protein loss, and outcome in dogs with severe parvoviral enteritis, *J. Vet. Intern. Med.* 17 (2003) 791–798, <https://doi.org/10.1111/j.1939-1676.2003.tb02516.x>.
- [7] A. Cantos-Barreda, D. Escribano, L.J. Bernal, J.J. Ceron, S. Martínez-Subiela, Quantification of anti-Leishmania antibodies in saliva of dogs, *Vet. Parasitol.* 242 (2017) 54–58, <https://doi.org/10.1016/j.vetpar.2017.05.017>.
- [8] M. Jankowski, J. Spuzak, K. Kubiak, K. Glińska-Suchocka, M. Biernat, An evaluation of the usefulness of invasive and non-invasive methods used to diagnose *Helicobacter* spp. Infections in dogs, *Pol. J. Vet. Sci.* 20 (2017) 491–499, <https://doi.org/10.1515/pjvs-2017-0059>.
- [9] F. Di Nardo, L. Anfossi, L. Ozella, A. Sacconi, C. Giovannoli, G. Spano, C. Baggiani, Validation of a qualitative immunochromatographic test for the noninvasive assessment of stress in dogs, *J. Chromatogr. B* 1028 (2016) 192–198, <https://doi.org/10.1016/j.jchromb.2016.06.019>.
- [10] A. Lindsay, J.T. Costello, Realising the potential of urine and saliva as diagnostic tools in sport and exercise medicine, *Sport. Med.* 47 (2016) 1–21, <https://doi.org/10.1007/s40279-016-0558-1>.
- [11] C. Desario, N. Decaro, M. Campolo, A. Cavalli, F. Cirone, G. Elia, V. Martella, E. Lorusso, M. Camero, C. Buonavoglia, Canine parvovirus infection: which diagnostic test for virus? *J. Virol. Methods* 126 (2005) 179–185, <https://doi.org/10.1016/j.jviromet.2005.02.006>.
- [12] L.A. Tabak, A revolution in biomedical assessment: the development of salivary diagnostics, *J. Dent. Educ.* 65 (12) (2001) 1335–1339.
- [13] N. Guillemin, A. Horvatić, J. Kuleš, A. Galan, V. Mrljak, M. Bhide, Omics approaches to probe markers of disease resistance in animal sciences, *Mol. Biosyst.* 12 (2016) 2036–2046, <https://doi.org/10.1039/C6MB00220J>.
- [14] H. Li, C. Li, H. Wu, T. Zhang, J. Wang, S. Wang, J. Chang, Identification of Apo-A1 as a biomarker for early diagnosis of bladder transitional cell carcinoma, *Proteome Sci.* 9 (21) (2011), <https://doi.org/10.1186/1477-5956-9-21>.
- [15] C. Ruiz-Romero, F.J. Blanco, Proteomics role in the search for improved diagnosis, prognosis and treatment of osteoarthritis, *Osteoarthr. Cartil.* 18 (2010) 500–509, <https://doi.org/10.1016/j.joca.2009.11.012>.
- [16] D. Baeumlisberger, T.N. Arrey, B. Rietschel, M. Rohmer, D.G. Papisotiriou, B. Mueller, T. Beckhaus, M. Karas, Labeling elastase digests with TMT: informational gain by identification of poorly detectable peptides with MALDI-TOF/TOF mass spectrometry, *Proteomics* 10 (2010) 3905–3909, <https://doi.org/10.1002/pmic.201000288>.
- [17] L. Dayon, N. Turck, T. García-Berrococo, N. Walter, P.R. Burkhard, A. Vilalta, J. Sahuquillo, J. Montaner, J.-C. Sanchez, Brain extracellular fluid protein changes in acute stroke patients, *J. Proteome Res.* 10 (2011) 1043–1051, <https://doi.org/10.1021/pr101123t>.
- [18] P. Giron, L. Dayon, N. Turck, C. Hoogland, J.-C. Sanchez, Quantitative analysis of human cerebrospinal fluid proteins using a combination of cysteine tagging and amine-reactive isobaric labeling, *J. Proteome Res.* 10 (2011) 249–258, <https://doi.org/10.1021/pr100535f>.
- [19] P.L. Ross, Y.N. Huang, J.N. Marchese, B. Williamson, K. Parker, S. Hattan, N. Khainovski, S. Pillai, S. Dey, S. Daniels, S. Purkayastha, P. Juhasz, S. Martin, M. Bartlett-Jones, F. He, A. Jacobson, D.J. Pappin, Multiplexed protein quantitation in *Saccharomyces cerevisiae* using amine-reactive isobaric tagging reagents, *Mol. Cell Proteomics* 3 (2004) 1154–1169, <https://doi.org/10.1074/mcp.M400129-MCP200>.
- [20] C. Baldini, L. Giusti, F. Ciregia, Y. Da Valle, C. Giacomelli, E. Donadio, F. Ferro, S. Galimberti, V. Donati, L. Bazzichi, S. Bombardieri, A. Lucacchini, Correspondence between salivary proteomic pattern and clinical course in primary Sjögren syndrome and non-Hodgkin's lymphoma: a case report, *J. Transl. Med.* 9 (188) (2011), <https://doi.org/10.1186/1479-5876-9-188>.
- [21] P. Beckett, The basics of 2D DIGE, *Methods Mol. Biol.* 854 (2012) 9–19, https://doi.org/10.1007/978-1-61779-573-2_2.
- [22] Z. Yilmaz, S. Senturk, Characterisation of lipid profiles in dogs with parvoviral enteritis, *J. Small Anim. Pract.* 48 (2007) 643–650, <https://doi.org/10.1111/j.1748-5827.2007.00391.x>.
- [23] M.D. Contreras-Aguilar, F. Tecles, S. Martínez-Subiela, D. Escribano, L.J. Bernal, J.J. Ceron, Detection and measurement of alpha-amylase in canine saliva and changes after an experimentally induced sympathetic activation, *BMC Vet. Res.* 13 (2017) 266, <https://doi.org/10.1186/s12917-017-1191-4>.
- [24] S. Martínez-Subiela, J.J. Ceron, Z. Yilmaz, S. Martínez-Subiela, A. Horvatić, D. Escribano, L. Pardo-Marin, M. Kocaturk, V. Mrljak, R. Burchmore, J.J. Ceron, Z. Yilmaz, Identification of novel biomarkers for treatment monitoring in canine leishmaniasis by high-resolution quantitative proteomic analysis, *Vet. Immunol. Immunopathol.* 191 (2017) 60–67, <https://doi.org/10.1016/j.vimid.2016.10.002>.
- [25] P. Bilić, N. Guillemin, A. Kovačević, B. Beer Ljubić, I. Jović, A. Galan, P.D. Eckersall, R. Burchmore, V. Mrljak, Serum proteome profiling in canine idiopathic dilated cardiomyopathy using TMT-based quantitative proteomics approach, *J. Proteomics* 179 (2018) 110–121, <https://doi.org/10.1016/j.jprot.2018.03.007>.
- [26] RStudio Team. RStudio, Integrated Development Environment for R, 2015. [Internet]. Available from: RStudio, Inc., Boston, MA, 2015 <http://www.rstudio.com/>.
- [27] G. Bindea, B. Mlecnik, H. Hackl, P. Charoentong, M. Tosolini, A. Kirilovsky, W.H. Fridman, F. Pagès, Z. Trajanoski, J. Galon, ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks, *Bioinformatics* 25 (2009) 1091–1093, <https://doi.org/10.1093/bioinformatics/btp101>.
- [28] P. Shannon, A. Markiel, O. Ozier, N.S. Baliga, J.T. Wang, D. Ramage, N. Amin, B. Schwikowski, T. Ideker, Cytoscape: a software Environment for integrated models of biomolecular interaction networks, *Genome Res.* 13 (2003) 2498–2504, <https://doi.org/10.1101/gr.1239303>.
- [29] F. Supek, M. Bošnjak, N. Škunca, T. Šmuc, Revigo summarizes and visualizes long lists of gene ontology terms, *PLoS One* 6 (2011), <https://doi.org/10.1371/journal.pone.0021800>.
- [30] V.W.M. Van Hinsbergh, Endothelium - Role in regulation of coagulation and inflammation, *Semin. Immunopathol.* 34 (1) (2012) 93–106, <https://doi.org/10.1007/s00281-011-0285-5>.
- [31] C.T. Esmon, K. Fukudome, T. Mather, W. Bode, L.M. Regan, D.J. Stearns-Kurosawa, S. Kurosawa, Inflammation, sepsis, and coagulation, *Haematologica* 84 (1999) 254–259, <https://doi.org/10.1189/392>.
- [32] M. Coorens, V.A.F. Schneider, A.M. de Groot, A. van Dijk, M. Meijerink, J.M. Wells, M.R. Scheenstra, E.J.A. Veldhuizen, H.P. Haagsman, Cathelicidins inhibit *Escherichia coli*-induced TLR2 and TLR4 activation in a viability-dependent manner, *J. Immunol.* 199 (2017) 1418–1428, <https://doi.org/10.4049/jimmunol.1602164>.
- [33] M. Chromek, I. Arvidsson, D. Karpman, The antimicrobial peptide cathelicidin protects mice from *Escherichia coli* O157:H7-Mediated disease, *PLoS One* 7 (2012), <https://doi.org/10.1371/journal.pone.0046476>.
- [34] M. Chromek, Z. Slamová, P. Bergman, L. Kovács, L. Podracká, I. Ehrén, T. Hökfelt, G.H. Gudmundsson, R.L. Gallo, B. Agerberth, A. Brauner, The antimicrobial peptide cathelicidin protects the urinary tract against invasive bacterial infection, *Nat. Med.* 12 (2006) 636–641, <https://doi.org/10.1038/nm1407>.
- [35] V. Nizet, T. Ohtake, X. Lauth, J. Trowbridge, J. Rudisill, R.A. Dorschner, V. Pestonjamas, J. Piraino, K. Huttner, R.L. Gallo, Innate antimicrobial peptide protects the skin from invasive bacterial infection, *Nature* 414 (2001) 454–457, <https://doi.org/10.1038/35106587>.
- [36] N.J. Jann, M. Schmalzer, S.A. Kristian, K.A. Radek, R.L. Gallo, V. Nizet, A. Peschel, R. Landmann, Neutrophil antimicrobial defense against *Staphylococcus aureus* is mediated by phagolysosomal but not extracellular trap-associated cathelicidin, *J. Leukoc. Biol.* 86 (2009) 1159–1169, <https://doi.org/10.1189/jlb.0209053>.
- [37] A. van Dijk, M.H.G. Tersteeg-Zijderderveld, J.L.M. Tjeerdema-van Bokhoven, A.J.M. Jansman, E.J.A. Veldhuizen, H.P. Haagsman, Chicken heterophils are

- recruited to the site of Salmonella infection and release antibacterial mature Cathelicidin-2 upon stimulation with LPS. *Mol. Immunol.* 46 (2009) 1517–1526, <https://doi.org/10.1016/j.molimm.2008.12.015>.
- [38] J. Schaubert, C. Svanholm, S. Term??n, K. Iffland, T. Menzel, W. Scheppach, R. Melcher, B. Agerberth, H. L??hrs, G.H. Gudmundsson, Expression of the cathelicidin LL-37 is modulated by short chain fatty acids in colonocytes: relevance of signalling pathways, *Gut* 52 (2003) 735–741, <https://doi.org/10.1136/gut.52.5.735>.
- [39] M.G. Scott, D.J. Davidson, M.R. Gold, D. Bowdish, R.E.W. Hancock, The human antimicrobial peptide LL-37 is a multifunctional modulator of innate immune responses, *J. Immunol.* 169 (2002) 3883–3891, <https://doi.org/10.4049/jimmunol.169.7.3883>.
- [40] A. Van Dijk, M. Van Eldik, E.J.A. Veldhuizen, H.L.M. Tjeerdma-Van Bokhoven, M.R. De Zoete, F.J. Bikker, H.P. Haagsman, Immunomodulatory and anti-inflammatory activities of chicken cathelicidin-2 derived peptides, *PLoS One* 11 (2016), <https://doi.org/10.1371/journal.pone.0147919>.
- [41] A. Togawa, J. Miyoshi, H. Ishizaki, M. Tanaka, A. Takakura, H. Nishioka, H. Yoshida, T. Doi, A. Mizoguchi, N. Matsuura, Y. Niho, Y. Nishimune, S.I. Nishikawa, Y. Takai, Progressive impairment of kidneys and reproductive organs in mice lacking Rho GDI α , *Oncogene* 18 (1999) 5373–5380, <https://doi.org/10.1038/sj.onc.1202921>.
- [42] T. Watanabe, E. Urano, K. Miyauchi, R. Ichikawa, M. Hamatake, N. Misawa, K. Sato, H. Ebina, Y. Koyanagi, J. Komano, The hematopoietic cell-specific rho GTPase inhibitor ARHGDI B/D4GDI limits HIV type 1 replication, *AIDS Res. Hum. Retroviruses* 28 (2012) 913–922, <https://doi.org/10.1089/aid.2011.0180>.
- [43] D. Theodorescu, L.M. Sapinoso, M.R. Conaway, G. Oxford, G.M. Hampton, H.F. Frierson, Reduced expression of metastasis suppressor RhoGDI2 is associated with decreased survival for patients with bladder cancer, *Clin. Cancer Res.* 10 (2004) 3800–3806, <https://doi.org/10.1158/1078-0432.CCR-03-0653>.
- [44] E. Jensen-Jarolim, L.F. Pacios, R. Bianchini, G. Hofstetter, F. Roth-Walter, Structural similarities of human and mammalian lipocalins, and their function in innate immunity and allergy, *Allergy* 71 (2016) 286–294, <https://doi.org/10.1111/all.12797>.
- [45] B. Åkerstrom, D.R. Flower, J.P. Salier, Lipocalins: unity in diversity, *Biochim. Biophys. Acta - Protein Struct. Mol. Enzymol* 1482 (1–2) (2000) 1–8, [https://doi.org/10.1016/S0167-4838\(00\)00137-0](https://doi.org/10.1016/S0167-4838(00)00137-0).
- [46] L. Lögdberg, L. Wester, Immunocalins: a lipocalin subfamily that modulates immune and inflammatory responses, *Biochim. Biophys. Acta - Protein Struct. Mol. Enzymol.* 1482 (1–2) (2000) 284–297, [https://doi.org/10.1016/S0167-4838\(00\)00164-3](https://doi.org/10.1016/S0167-4838(00)00164-3).
- [47] G. Fjaertoft, T. Foucard, S. Xu, P. Venge, Human neutrophil lipocalin (HNL) as a diagnostic tool in children with acute infections: a study of the kinetics, *Acta Paediatr.* 94 (2005) 661–666, <https://doi.org/10.1080/08035250510031610>.
- [48] J. Mårtensson, M. Bell, A. Oldner, S. Xu, P. Venge, C.R. Martling, Neutrophil gelatinase-associated lipocalin in adult septic patients with and without acute kidney injury, *Intensive Care Med.* 36 (2010) 1333–1340, <https://doi.org/10.1007/s00134-010-1887-4>.
- [49] E. Parravicini, S.L. Nemerofsky, K.A. Michelson, T.K. Huynh, M.E. Sise, D.A. Bateman, J.M. Lorenz, J.M. Barasch, Urinary neutrophil gelatinase-associated lipocalin is a promising biomarker for late onset culture-positive sepsis in very low birth weight infants, *Pediatr. Res.* 67 (2010) 636–640, <https://doi.org/10.12103/PDR.0b013e3181da75c1>.
- [50] L. Bingle, F.A. Barnes, H. Lunn, M. Musa, S. Webster, C.W.I. Douglas, S.S. Cross, A.S. High, C.D. Bingle, Characterisation and expression of SPLUNC2, the human orthologue of rodent parotid secretory protein, *Histochem. Cell Biol.* 132 (2009) 339–349, <https://doi.org/10.1007/s00418-009-0610-4>.
- [51] S. Morosky, N.J. Lennemann, C.B. Coyne, BPIFB6 Regulates Secretory Pathway Trafficking and Enterovirus Replication, *J. Virol.* 90 (2016) 5098–5107, <https://doi.org/10.1128/JVI.00170-16>.
- [52] Y. Huang, J.A. DiDonato, B.S. Levison, D. Schmitt, L. Li, Y. Wu, J. Buffa, T. Kim, G.S. Gerstenecker, X. Gu, C.S. Kadiyala, Z. Wang, M.K. Culley, J.E. Hazen, A.J. Didonato, X. Fu, S.Z. Berisha, D. Peng, T.T. Nguyen, S. Liang, C.-C. Chuang, L. Cho, E.F. Plow, P.L. Fox, V. Gogonea, W.H.W. Tang, J.S. Parks, E.A. Fisher, J.D. Smith, S.L. Hazen, An abundant dysfunctional apolipoprotein A1 in human atheroma, *Nat. Med.* 20 (2014) 193–203, <https://doi.org/10.1038/nm.3459>.
- [53] D. Escribano, A. Tvarijonavičiute, M. Kocaturk, J.J. Cerón, L. Pardo-Marín, A. Torrecillas, Z. Yilmaz, S. Martínez-Subiela, Serum apolipoprotein-A1 as a possible biomarker for monitoring treatment of canine leishmaniasis, *Comp. Immunol. Microbiol. Infect. Dis.* 49 (2016) 82–87, <https://doi.org/10.1016/j.cimid.2016.10.002>.
- [54] M. Horwitz, K.F. Benson, Z. Duan, F.-Q. Li, R.E. Person, Hereditary neutropenia: dogs explain human neutrophil elastase mutations, *Trends Mol. Med.* 10 (2004) 163–170, <https://doi.org/10.1016/j.molmed.2004.02.002>.
- [55] U. Thanarajasingam, M.A. Jensen, J.M. Dorschner, T. Wampler Muskardin, Y. Ghodke-Puranik, M. Purmalek, E. Eliopoulos, M.I. Zervou, G.N. Goulielmos, M. Howard, M.J. Kaplan, T.B. Niewold, Brief report: a novel ELANE mutation associated with inflammatory arthritis, defective NETosis, and recurrent parvovirus infection, *Arthritis Rheumatol.* (Hoboken, N.J.) 69 (2017) 2396–2401, <https://doi.org/10.1002/art.40314>.
- [56] C.S. Chee, K.M. Chang, M.F. Loke, V.P. Angela Loo, V. Subrayan, Association of potential salivary biomarkers with diabetic retinopathy and its severity in type-2 diabetes mellitus: a proteomic analysis by mass spectrometry, *PeerJ* 4 (2016) e2022, <https://doi.org/10.7717/peerj.2022>.
- [57] J. Kuleš, V. Mrljak, R. Barić Rafaj, J. Selanec, R. Burchmore, P.D. Eckersall, Identification of serum biomarkers in dogs naturally infected with *Babesia canis canis* using a proteomic approach, *BMC Vet. Res.* 10 (2014) 111, <https://doi.org/10.1186/1746-6148-10-111>.
- [58] K.M. O'Neil, H.D. Ochs, S.R. Heller, L.C. Cork, J.M. Morris, J.A. Winkelstein, Role of C3 in humoral immunity. Defective antibody production in C3-deficient dogs, *J. Immunol.* 140 (1988) 1939–1945.
- [59] C. Lee, E. Bongcam-Rudloff, E. Sollner, W. Jahnen-Dechent, L. Claesson-Welsh, Type 3 cystatins; fetuins, kininogen and histidine-rich glycoprotein, *Front Biosci (Landmark Ed.)* 1 (2009) 2911–2922, <https://doi.org/10.1111/febs.12167>.
- [60] I.K.H. Poon, K.K. Patel, D.S. Davis, C.R. Parish, M.D. Hulett, Histidine-rich glycoprotein: the Swiss Army knife of mammalian plasma, *Blood* 117 (7) (2011) 2093–2101, <https://doi.org/10.1182/blood-2010-09-303842>.
- [61] O. Shannon, V. Rydengård, A. Schmidtchen, M. Mörgelin, P. Alm, O.E. Sørensen, L. Björk, Histidine-rich glycoprotein promotes bacterial entrapment in clots and decreases mortality in a mouse model of sepsis, *Blood* 116 (2010) 2365–2372, <https://doi.org/10.1182/blood-2010-02-271858>.
- [62] Ellebrecht, C.T., Payne, A.S., n.d. Plakophilins, desmogleins, and pemphigus: the tail wagging the dog Desmosomes and disease. <https://doi.org/10.1038/jid.2013.491>.
- [63] Z. Hallaji, H. Mortazavi, V. Lajevardi, B. Tamizifar, A. Amirzargar, M. Daneshpazhooh, C. Chams-Davatchi, Serum and salivary desmoglein 1 and 3 enzyme-linked immunosorbent assay in pemphigus vulgaris: correlation with phenotype and severity, *J. Eur. Acad. Dermatol. Venereol.* 24 (2010) 275–280, <https://doi.org/10.1111/j.1468-3083.2009.03408.x>.
- [64] M.E. Rosenberg, J. Silkensen, Clusterin: Physiologic and pathophysiologic considerations, *Int. J. Biochem. Cell Biol.* 27 (7) (1995) 633–645, [https://doi.org/10.1016/1357-2725\(95\)00027-M](https://doi.org/10.1016/1357-2725(95)00027-M).
- [65] Wong, P., Borst, D.E., Farber, D., Danciger, J.S., Tenniswood, M., Chader, G.J., van Veen, T., n.d. Increased TRPM-2/clusterin mRNA levels during the time of retinal degeneration in mouse models of retinitis pigmentosa. *Biochem. Cell Biol.* 72, 439–446.
- [66] E.V. Martynova, A.N. Maksudova, V.G. Shakirova, S.R. Abdulkhakov, I.M. Khaertynova, V.A. Anokhin, V.V. Ivanova, I.M. Abiola, E.E. Garanina, L.G. Tazetdinova, A.H. Valiullina, S.F. Khaiboullina, Urinary clusterin is upregulated in Nephropathia Epidemica, *Dis. Markers* 2018 (2018) 1–7, <https://doi.org/10.1155/2018/8658507>.
- [67] F. Zhang, M. Kumano, E. Beraldi, L. Fazli, C. Du, S. Moore, P. Sorensen, A. Zoubeidi, M.E. Gleave, Clusterin facilitates stress-induced lipidation of LC3 and autophagosome biogenesis to enhance cancer cell survival, *Nat. Commun.* 5 (2014), <https://doi.org/10.1038/ncomms6775>.
- [68] M. Kocaturk, S. Martinez, O. Eralp, A. Tvarijonavičiute, J. Ceron, Z. Yilmaz, Prognostic value of serum acute-phase proteins in dogs with parvoviral enteritis, *J. Small Anim. Pract.* 51 (2010) 478–483, <https://doi.org/10.1111/j.1748-5827.2010.00965.x>.
- [69] D.K. Macintire, S. Smith-Carr, *Canine parvovirus. Part II. Clinical signs, diagnosis, and treatment*, *Compend. Contin. Vet. Educ. Pract.* 19 (1997) 291–302.
- [70] N.L. Mantione, C.M. Otto, Characterization of the use of antiemetic agents in dogs with parvoviral enteritis treated at a veterinary teaching hospital: 77 cases (1997–2000), *J. Am. Vet. Med. Assoc.* 227 (2005) 1787–1793, <https://doi.org/10.2460/javma.2005.227.1787>.
- [71] J. Turk, W. Fales, M. Miller, L. Pace, J. Fischer, G. Johnson, J. Creeger, S. Turnquist, L. Pittman, A. Rottinghaus, Enteric *Clostridium perfringens* infection associated with parvoviral enteritis in dogs: 74 cases (1987–1990), *J. Am. Vet. Med. Assoc.* 200 (1992) 991–994.
- [72] J. Turk, M. Miller, T. Brown, W. Fales, J. Fischer, H. Gosser, S. Nelson, D. Shaw, R. Solorzano, Coliform septicemia and pulmonary disease associated with canine parvoviral enteritis: 88 cases (1987–1988), *J. Am. Vet. Med. Assoc.* 196 (1990) 771–773.
- [73] C.M. Otto, T.M. Rieser, M.B. Brooks, M.W. Russell, Evidence of hypercoagulability in dogs with parvoviral enteritis, *J. Am. Vet. Med. Assoc.* 217 (2000) 1500–1504, <https://doi.org/10.2460/javma.2000.217.1500>.
- [74] D. Britti, M. Gaspari, G. Massimini, F. Casalnuovo, V.M. Morittu, G. Cuda, Proteomic analysis in canine leishmaniasis, *Veterinary Research Communications*, (2010), pp. 91–96, <https://doi.org/10.1007/s11259-010-9410-5>.
- [75] J. Kuleš, C. de Torre-Minguela, R. Barić Rafaj, J. Gotić, P. Nižić, J.J. Ceron, V. Mrljak, Plasma biomarkers of SIRS and MODS associated with canine babesiosis, *Res. Vet. Sci.* 105 (2016) 222–228, <https://doi.org/10.1016/j.rvsc.2016.02.011>.
- [76] E. Carretón, R. Morchón, F. Simón, M.C. Juste, J.C. Méndez, J.A. Montoya-Alonso, Cardiopulmonary and inflammatory biomarkers in the assessment of the severity of canine dirofilariosis, *Vet. Parasitol.* 206 (2014) 43–47, <https://doi.org/10.1016/j.vetpar.2014.08.019>.
- [77] J.J. Cerón, P.D. Eckersall, S. Martínez-Subiela, Acute phase proteins in dogs and cats: current knowledge and future perspectives, *Vet. Clin. Pathol.* 34 (2) (2005) 85–99, <https://doi.org/10.1111/j.1939-165X.2005.tb00019.x>.
- [78] A.M. Lynch, A.M. DeLaforcade, D. Meola, A. Shih, C. Bandt, N.H. Guerrero, C. Riccò, Assessment of hemostatic changes in a model of acute hemorrhage in dogs, *J. Vet. Emerg. Crit. Care San Antonio (San Antonio)* 26 (2016) 333–343, <https://doi.org/10.1111/vec.12457>.
- [79] S.C. Johnston, C.N. Larsen, W.J. Cook, K.D. Wilkinson, C.P. Hill, Crystal structure of a deubiquitinating enzyme (human UCH-L3) at 1.8 angstrom resolution, *EMBO J.* 16 (1997) 3787–3796.
- [80] F. Randow, P.J. Lehner, Viral avoidance and exploitation of the ubiquitin system, *Nat. Cell Biol.* 11 (5) (2009) 527–534, <https://doi.org/10.1038/ncb0509-527>.