

# Delegating Sex: Differential Gene Expression in Stolonizing Syllids Uncovers the Hormonal Control of Reproduction

Patricia Álvarez-Campos<sup>1,2,3,8,\*</sup>, Nathan J. Kenny<sup>3</sup>, Aida Verdes<sup>1,4,5</sup>, Rosa Fernández<sup>6</sup>, Marta Novo<sup>7</sup>, Gonzalo Giribet<sup>2,†</sup>, and Ana Riesgo<sup>4,†</sup>

<sup>1</sup>Facultad de Ciencias, Departamento de Biología (Zoología), Universidad Autónoma de Madrid, Spain

<sup>2</sup>Museum of Comparative Zoology, Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts

<sup>3</sup>Department of Life Sciences, The Natural History Museum of London, London, United Kingdom

<sup>4</sup>Department of Biology, The Graduate Center, City University of New York

<sup>5</sup>Sackler Institute for Comparative Genomics, American Museum of Natural History, New York, New York

<sup>6</sup>Bioinformatics & Genomics Unit, Center for Genomic Regulation, Barcelona, Spain

<sup>7</sup>Facultad de Biología, Departamento de Biodiversidad, Ecología y Evolución, Universidad Complutense de Madrid, Spain

<sup>8</sup>Present address: Department of Biological & Medical Sciences, Oxford Brookes University, Headington Campus, Gypsy Lane, Oxford, United Kingdom

<sup>†</sup>These authors contributed equally as Senior authors.

\*Corresponding author: E-mail: patriciaalvarezcampos@gmail.com.

Accepted: December 9, 2018

Data deposition: This project has been deposited at Sequence Read Archive (SRA, NCBI) under the accession SRP133371.

## Abstract

Stolonization in syllid annelids is a unique mode of reproduction among animals. During the breeding season, a structure resembling the adult but containing only gametes, called stolon, is formed generally at the posterior end of the animal. When stolons mature, they detach from the adult and gametes are released into the water column. The process is synchronized within each species, and it has been reported to be under environmental and endogenous control, probably via endocrine regulation. To further understand reproduction in syllids and to elucidate the molecular toolkit underlying stolonization, we generated Illumina RNA-seq data from different tissues of reproductive and nonreproductive individuals of *Syllis magdalena* and characterized gene expression during the stolonization process. Several genes involved in gametogenesis (*ovochymase*, *vitellogenin*, *testis-specific serine/threonine-kinase*), immune response (*complement receptor 2*), neuronal development (*tyrosine-protein kinase Src42A*), cell proliferation (*alpha-1D adrenergic receptor*), and steroid metabolism (*hydroxysteroid dehydrogenase 2*) were found differentially expressed in the different tissues and conditions analyzed. In addition, our findings suggest that several neurohormones, such as methyl farnesoate, dopamine, and serotonin, might trigger stolon formation, the correct maturation of gametes and the detachment of stolons when gametogenesis ends. The process seems to be under circadian control, as indicated by the expression patterns of *r-opsins*. Overall, our results shed light into the genes that orchestrate the onset of gamete formation and improve our understanding of how some hormones, previously reported to be involved in reproduction and metamorphosis processes in other invertebrates, seem to also regulate reproduction via stolonization.

**Key words:** transcriptomics, stolonizing syllids, reproduction, hormonal control.

## Introduction

Annelids in the family Syllidae have a remarkable reproductive strategy, which has attracted the attention of many biologists (e.g., Nygren 1999 and references herein). Syllids exhibit epitoky, which largely implies morphological changes

associated with reproduction (Malaquin 1893), and can be further divided into a variety of reproductive modes. In all epitokous modes, there are two states: the sexually immature worm, called an “atoke,” and the sexually mature worm, or “epitoke. after.” Among the epitokous types of reproduction,

© The Author(s) 2018. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

one of the most common is epigamy, which is not exclusive to syllids, where the entire atoke transforms into the epitoke, developing swimming chaetae, enlarging its eyes and undergoing changes in musculature (Wissocq 1970; Daly 1975; Garwood 1991). One of the most extreme types of epitokous reproduction is squizogamy or stolonization, where only a part of the individual transforms into an epitokal sexual stage, either by generating new segments or by differentiating pre-existing ones (Franke 1999). When the breeding season approaches, the syllid atoke (or stock) starts to develop a peculiar structure at the end of its body, that resembles the adult and is known as the stolon (Agassiz 1863) (fig. 1). The stolons possess several features similar to the stock, such as eyes and antennae, but are filled with gametes (figs. 1 and 2A–E), as their brief existence is exclusively devoted to mating, followed by death (Franke 1999). The stock produces and transfers the gametes to the stolon, which is released from the stock when mature (with developed eyes and antennae) (figs. 1 and 2E), and swims to the surface to spawn (Potts 1911; Mesnil and Caullery 1919). The pelagic stolon releases gametes into the water column, via the nephridiopores in the case of sperm, and through rupture of the body wall for the eggs (Okada 1937; Durchon 1951, 1952, 1959; Wissocq 1966, 1970; Schroeder and Hermans 1975; Franke 1980). Finally, before or after stolon detachment (depending on the species), the stock regenerates the lost final segments (e.g., Marion and Bobretsky 1875; Michel 1898; Okada 1929) (figs. 1 and 2F).

The molecular toolkit involved in annelid reproduction is still far from being understood, although studies in several annelid species have shed some light into the matter. For instance, genes involved in pheromone production that are essential for mate recognition and spawning, such as *Temptin* and *Attractin*, and those involved in gametogenesis or fertilization, such as *Fertilin* or *Acrosin*, have been identified in *Spirobranchus (Pomatoceros) lamarckii*, *Hormogaster samnitica* and *H. elisae* (e.g., Kang et al. 2002; Rivera et al. 2005; Takahashi et al. 2009; Novo et al. 2013). It is also well-known that the germline specification in the marine annelids *Alitta virens*, *Platynereis dumerilii*, *Capitella teleta*, or *Hermodice carunculata* involves the expression of several genes including *vasa*, *nanos*, and *pivi* during embryogenesis, and that *vitellogenin (Vtg)* is required for yolk formation in the oocyte (Hafer et al. 1992; Rebscher et al. 2007; Dill and Seaver 2008; Thamm and Seaver 2008; Giani et al. 2011; Mehr et al. 2015; Schenk et al. 2016). Interestingly, a recent study has reported the potential involvement of the sesquiterpenoid methyl farnesoate (MF), the brain neurohormone that directly regulates *Vtg* in yolk production of *P. dumerilii* females, therefore influencing the correct development of oocytes (Schenk et al. 2016). Particularly, a decrease in MF levels in the brain of *P. dumerilii* during reproduction allowed oocyte maturation but suppressed normal somatic functions and caudal regenerative capacities (Schenk et al. 2016). In crustaceans, MF has been showed to play essential roles in development and

reproduction (Xie et al. 2016), similar to the role of juvenile hormone (JH) in insects (Riddiford 1994; Wyatt and Davey 1996). Other hormones have also been proposed to play essential roles in annelid reproduction, such as the prostomium (i.e., first preoral segment of the animal) hormone 8, 11, 14-eicosatrienoic acid, which seems to be responsible for sperm maturation and spawning in *Arenicola marina* males (Bentley 1985; Bentley et al. 1990; Pacey and Bentley 1992).

Similarly, it has been proposed that the stolonization process in syllids is under hormonal control, following endogenous circadian and circalunar rhythms influenced by exogenous factors, including annual photoperiod, temperature, or moon cycles (Franke 1986a, 1999). It has been hypothesized that during the summer time, with long days and high temperatures, a stolonization-promoting hormone produced in the prostomium is secreted to control a second stolonization-suppressing hormone produced in the proventricle (i.e., specialized structure of the digestive tract), allowing the initiation of stolonization (Franke 1999). In contrast, during winter, when days are short and temperatures low at high latitudes, the proventricle is not controlled by the prostomium, and the proventricular stolonization-suppressing hormone then inhibits stolonization (e.g., Abeloos 1950; Durchon 1952, 1959; Durchon and Wissocq 1964; Franke 1980, 1981, 1983a, 1983b, 1985, 1999; Heacox 1980; Heacox and Schroeder 1982; Franke and Pfannenstiel 1984; Verger-Bocquet 1984). Hormonal factors have also been suggested to drive the sexual differentiation of the stolon (Franke 1980; Heacox and Schroeder 1982), in particular the female stolon, given that it seems that male stolon differentiation occurs autonomously, whereas female stolon differentiation may depend on hormone release by male stolons (Franke 1999). However, no candidate hormone has been proposed to control reproduction and regeneration processes in syllids, although it seems clear that there might be several involved, not only in the brain, but also in the proventricle (e.g., Schroeder and Hermans 1975; Franke 1999; Weidhase et al. 2016).

In summary, although molecular mechanisms underlying reproduction are relatively well studied in a few annelids (e.g., Kang et al. 2002; Thamm and Seaver 2008; Giani et al. 2011; Novo et al. 2013; Schenk et al. 2016), the molecular toolkit involved in the stolonization process of syllids has not been examined yet. Thus, our aim in the present study is to provide a first glimpse into the gene expression patterns occurring during the stolonization process in the syllid species *Syllis magdalena*. To achieve this goal, we have pursued four main objectives: 1) to characterize molecularly and morphologically the stolonization process in the target species; 2) to provide a detailed description of the genes potentially involved in the triggering of stolonization and the formation/releasing of stolons and gametes, through differential gene expression analyses of reproductive and nonreproductive individuals in different tissues; 3) to understand the evolution of selected




FIG. 1.—Syllinae schizogamous reproductive cycle (stolonization) using light microscope pictures of *Syllis magdalena*.

candidate genes with major roles in the reproductive processes of the phylum Annelida; and 4) to investigate if the molecular signal that determines when to divert resources from somatic functions to reproduction is the same across annelids (i.e., synthesis of MF).


## Results and Discussion

### General Morphology and Ultrastructure of the Stolons in *S. magdalena*

The stolons of *S. magdalena* were dicerous, with two pairs of red eyes and one pair of antennae formed at the beginning of the stolonization process (figs. 2A–E and 3A, 3B), similar to the process observed in *Syllis amica* (see Wissocq 1970) but different to the late formation of head structures in *Syllis gracilis* (see Pettibone 1963) or *Syllis hyalina* (see Malaquin 1893). Natatory capillary chaetae were not developed during the stages in which the stolon was attached to the stock. Before stolon detachment, the stock completely regenerated the final part of the body that was transformed during the stolon formation (fig. 2F). Female stolons were purple, completely full of oocytes arranged around the through-gut (figs. 2A, 2C, 2E, 3A and B). Male stolons were white, completely full of spermatogonia, and also arranged around the gut (fig. 2B and D).

The epithelium of the female and male stolons was columnar, comprised by large epithelial cells ( $>10\ \mu\text{m}$  in maximum length) with basal non-nucleolated nuclei, and large globular glandular cells with electrondense material (fig. 3C). In both stolons, below the epithelia, there was a thick layer of muscle fibers, then the germinative epithelium, and finally the digestive epithelium (fig. 3C–F). The muscle fibers of both female and male stolons presented the regular morphology of muscle fibers of the adults, with a double striation and 25–35 myofibrils and clusters of mitochondria near the tips (fig. 3C and E). We did not observe the “stolonial” muscle fibers described in *S. amica* with the mitochondria toward the middle of the fiber (Wissocq 1967) while attached to the stock. It is possible that the reorganization of the muscle fibers takes place later in the stolonization process, but it is improbable, given that it occurs during head formation in the stolon of *S. amica* (see Wissocq 1967), a process that we observed in *S. magdalena*.

In the female germinative epithelium, large yolky oocytes ( $50\ \mu\text{m}$  approximately) were surrounded by non-nucleolated nurse cells (fig. 3D). Oocytes were connected by microvillar processes (fig. 3D). The male germinative epithelium only contained two large sacs of spermatogonia in the specimens collected (fig. 3E and F). Spermatogonia (ca.  $1\ \mu\text{m}$  in diameter) were densely packed and possessed a non-nucleolated nucleus with




**FIG. 2.**—Light microscopy pictures of *Syllis magdalena* stolonizing female (A) and male (B). Confocal micrographs of *S. magdalena* stolonizing female (C), male (D), female stolon (E), and male stolons (F). Arrows in (A)–(D) pointing to the eyes of stolons (e). Arrows in (E) pointing to antennae (a). Arrow in (F) pointing to the regeneration of the final segments in the stock (rfs).

chromatin condensation processes (fig. 3E and F). The digestive epithelium was comprised of large ( $>10\ \mu\text{m}$  in maximum length) convoluted multiflagellated cells (fig. 3F). We did not observe digestive material in the lumen of the stolon gut (fig. 3F). There were no differences in the developmental stage of gametes between the anterior and posterior parts of stolons (see also differential expression results).

#### General Characterization of the *De Novo* Transcriptomes

Out of the 32 libraries generated, we assembled the REFSOM transcriptome (reference transcriptome for somatic parts of reproductive and non-reproductive individuals) using only somatic tissues of nonreproductive (NON-REPRO) and reproductive (REPRO) specimens (23 RNA-seq libraries in total). For the REFTOTREPRO assembly (reference transcriptome for the all



**FIG. 3.**—Light and electron microscopy pictures of the anterior part of the female and male stolons of *Syllis magdalena*. (A, B) Location of antennae (a) and the two pairs of eyes (e) in the female stolon. (C) Transmission electron micrographs of the epithelium of the female stolon showing the muscle fibers (mf), granular cells (gc), and epithelial cells (ec). (D) Developing oocytes showing nucleolate (nu) nucleus (n), ooplasm filled with yolk platelets, and microvilli (m) contacting close oocytes. Note the muscle fibers (mf), nurse cells (nc), and the digestive epithelium (dc) surrounding the germinal epithelium. (E–F) Germinal epithelium (ge) in the male stolon. The stolon epithelium is comprised by a layer of epithelial cells (ec) with interspersed granular cells (gc), and a layer of muscle fibers (mf); spermatogonia develop in the germinal epithelium (gc) below. The digestive cells (dc) lay below the germinal epithelium.

the parts in reproductive individuals), we used 18 RNA-seq libraries of both somatic and reproductive tissues of reproductive (REPRO) specimens (further details in Material and Methods). Assembly statistics for both reference transcriptomes are summarized in [supplementary file S1, Supplementary Material](#) online alongside read mapping results for each tissue and specimen, but overall they represent well assembled transcriptomes with N50 values over 750 in both cases ([supplementary file S1, Supplementary Material](#) online). The coverage of our assemblies is similar or slightly higher than those in other studies on marine invertebrates (e.g., Meyer et al. 2009; Riesgo et al. 2012; Pérez-Portela et al. 2016).

A summary of the assessment of both transcriptomes assemblies and their annotation completeness (Simão et al. 2015) is shown in [supplementary file S3, Supplementary Material](#) online. Out of the 978 metazoan single copy orthologs, our REFSOM assembly is 97% complete (950 complete BUSCOs, 267 single-copy BUSCOs, and 683 complete duplicated BUSCOs), while 2.7% of BUSCOs are fragmented (26 BUSCOs) and only 0.2% are missing (2 BUSCOs). On the other hand, our REFTOTREPRO assembly is 94% complete (918 complete BUSCOs, 316 single-copy BUSCOs and 316 complete duplicated BUSCOs), while 5.6% of BUSCOs are fragmented (55 BUSCOs) and 0.5% are missing (5 BUSCOs). In comparison to other annelid transcriptomes, which found around 80% of complete BUSCOs in *Pygospio elegans* (Heikkinen et al. 2017) and *Urechis unicinctus* (Park et al. 2018), and approximately 60% in *Sabellaria alveolata* and *Phragmatopoma caudata* (Buffet et al. 2018), the completeness of our transcriptomes was exceptionally high ([supplementary file S3, Supplementary Material](#) online).

An overview of the assigned GO terms for each transcriptome [including three different categories: cellular component (CC), biological process (BP), and molecular function (MF)] and GO enrichment analyses using Fisher's tests are shown in [supplementary file S2A, Supplementary Material](#) online. The GO enrichment results for the comparisons of both transcriptomes showed 36 GO terms overrepresented in REFSOM related to cellular organization and regulation, metabolism and binding, among others ([supplementary fig. S2B, Supplementary Material](#) online). In contrast, only eight categories appeared enriched in REFTOTREPRO, mainly related to signaling activity ([supplementary fig. S2C, Supplementary Material](#) online). Interestingly, one of these enriched categories is the activity of G-protein coupled receptors, which bind light-sensitive compounds, pheromones, hormones, neurotransmitters and other ligands involved in secretory processes or cell development, among other functions (e.g., Li et al. 1999; Iversen et al. 2002; Hauser et al. 2006; Asahara et al. 2013). The results of several of these G-protein coupled receptor

expression levels on the different tissues and conditions analyzed are discussed below.


### Differential Gene Expression Analyses

#### *Pairwise Comparisons of Somatic Tissues (Anterior Part, Proventricle, Final Segments) between REPRO and NON-REPRO Individuals (REFSOM Transcriptome)*

We detected 792 differentially expressed genes in the comparison between REPRO and NON-REPRO somatic tissues, 494 of them being upregulated in REPRO (178 in females and 316 in males) and 298 in NON-REPRO (fig. 4; [supplementary files S4, S5A, and S6, Supplementary Material](#) online). Of these 792 genes, only 292 (~37%) had a BLAST hit and, therefore only the putative annotations for those genes ([supplementary file S6, Supplementary Material](#) online) are discussed below. Among the pairwise comparisons of REPRO and NON-REPRO tissues, the final segment tissues are the ones that showed more differentially expressed genes (fig. 4C), with 223 differentially expressed in the comparison of female final segments and NON-REPRO final segments (152 upregulated in female) and 460 differentially expressed genes in the comparison of male final segments and NON-REPRO final segments (304 of those upregulated in male). The pairwise comparisons of anterior part and proventricle between reproductive and nonreproductive individuals showed low numbers of differentially expressed genes (fig. 4A and B). Among them, the highest number of differentially expressed genes was found in the proventricle, with 7 differentially expressed genes upregulated in both females and males when compared with nonreproductive, and 20 and 36 differentially expressed genes upregulated in the proventricle of nonreproductive individuals (fig. 4B).

In the anterior part and the proventricle of females, the genes upregulated ([supplementary file S6, Supplementary Material](#) online) were related mostly to immune processes (*complement receptor 2*) or food processing (*trefoil factor 2*, *cubilin*, *serine protease 27* and *chitinase*). Similarly, in the male anterior part and proventricle ([supplementary file S6, Supplementary Material](#) online), most genes were involved in nutrient transport (*sugar transporter STL1* and *glycogen phosphorylase*), as well as development of the nervous system (*tyrosine-protein kinase Src42A*).

Several genes related to gametogenesis were found differentially expressed in the final segments of female and male REPRO individuals compared with NONREPRO ([supplementary file S6, Supplementary Material](#) online), including *vitellogenin* (*Vtg*) and *ovochymase* (*OVCH*) in females, and *testis-specific serine/threonine-kinase* (*TSSK*) in males, which indicates an important role of the final segments during the gametogenesis process in both stolonizing females and males. *Vitellogenin* has been already reported to be involved in annelid gametogenesis, specifically as a yolk precursor (e.g., Hafer et al. 1992), but *OVCH*, an ovary-specific gene involved



**FIG. 4.**—Heatmaps of differentially expressed genes (annotated and not annotated genes) from pairwise comparisons of somatic tissues between reproductive (both female and male) and nonreproductive individuals. Anterior part tissue comparisons (A), proventricule comparisons (B), and final segments comparisons (C). Different colors indicate relative expression levels based on raw read counts (see color key and histogram on each). Similarity in expression patterns between genes and individuals is represented by clustering. A, anterior part; P, proventricule; F, final segments.

in egg development of several animals (e.g., Lindsay and Hedrick 1995; Gao and Zhang 2009; Mino and Sawada 2016), is here reported for the first time in annelids. The same occurs for *TSSK*, whose expression, confined almost exclusively to testes, has largely been studied in several mammals (Hao et al. 2004), but never in annelids. Remarkably, two hormone receptors for *relaxin* and *follistatin* were found differentially expressed in the final segments of reproductive females (supplementary file S6, Supplementary Material online). The insulin-related peptide *relaxin* is important for the growth and remodeling of reproductive tissues during mammal pregnancy (e.g., Gunnarsen et al. 1995; Hsu et al. 2002) and is active in the ovary and during embryogenesis of zebrafish (e.g., Donizetti et al. 2008, 2010; Wilson et al. 2009). *Relaxin* activity has also been reported in invertebrates, including in the tunicate *Ciona intestinalis* (e.g., Ivell and Anand-Ivell 2005; Olinski et al. 2006), and in the starfish *Asterina pectinifera* (Mita 2013; Mita et al. 2014), where it takes part in oocyte release from the ovary, but this is the first time that it is described in annelids. Likewise, *follistatin*, reported as a follicle-stimulating hormone, with several additional regulatory functions both in reproductive and nonreproductive tissues (Phillips and de Kretser 1998), has been already found in the transcriptome of other annelids such as *C. teleta* and *S. lamarckii* (Kenny et al. 2015), but without a particular association with any biological process. In our case, it seems that both *relaxin* and *follistatin* are important during oocyte development in *S. magdalena*, as they are expressed in tissues where oogenesis is taking place before oocytes are transferred into the stolon (see also Results and Discussion).

#### Pairwise Comparisons of Somatic (Anterior Part, Proventricle, Final Segments) between REPRO Females and Males (REFTOTREPRO Transcriptome)

We detected 234 genes differentially expressed in the comparison between female and male somatic tissues, 85 of them being upregulated in female (0 in anterior part, 27 in proventricle, 58 in final segments) and 149 in males (only in final segments) (see details in fig. 5A and B; supplementary file S7, Supplementary Material online). Of these 234 genes, only 84 (~35%) of transcripts were annotated (supplementary file S7, Supplementary Material online). No differential expression was found in the comparisons of the female and male anterior parts, and in the proventricle comparisons, we only found differentially expressed genes in the females (fig. 5A; supplementary file S7, Supplementary Material online; see Results and Discussion). Similar to the previous comparisons (see above), the somatic tissue sample that showed more differentially expressed genes was the final segments, with 149 genes upregulated in males and 58 in females (fig. 5B; supplementary file S7, Supplementary Material online).

As in the previous comparisons (see section above), several gametogenesis-related genes, such as *vitellogenin*,


*ovochymase* (*OVOCH*) in females, and *TSSK* in males, were differentially expressed in F (fig. 5B; supplementary file S7, Supplementary Material online). In addition, we also found *NOTCH* differentially expressed in F of REPRO males (fig. 5B; supplementary file S7, Supplementary Material online). This gene has been reported to have a role in segment formation and adult regeneration in annelids (e.g., Thamm and Seaver 2008), and therefore may also be involved in segment formation of stolons and pygidium regeneration of *S. magdalena* (fig. 2F). However, the *NOTCH* pathway has been also reported to be essential for the correct development of gametes in *Drosophila melanogaster* and mammals (Xu et al. 1992; Hayashi et al. 2001; Murta et al. 2014), and therefore it could also be playing such role during spermatogenesis in *S. magdalena*.

Two different transcripts of *ovochymase* were differentially expressed in final segments (*OVOCH1*) and proventricle (*OVOCH2*) female tissues (fig. 5A; supplementary file S7, Supplementary Material online). *Ovochymases* are involved in the oogenesis in other invertebrates, where they help avoid self-fertilization and are localized in the vitelline coat of oocytes (Mino and Sawada 2016). In the ascidian *Halocynthia roretzi*, *ovochymase* has a signal peptide, three trypsin-like serine protease domains and six CUB domains (Mino and Sawada 2016). We found 3 *ovochymases* (two DE, *OVOCH1* and *OVOCH2*, and one non-DE, *OVOCH3*) in *S. magdalena*, none of them containing a signal peptide and all containing significantly fewer trypsin-like serine protease and CUB domains (supplementary file S8, Supplementary Material online). The trypsin-like serine protease domain is not exclusive to *ovochymases*, because it also occurs in chymotrypsins (supplementary file S8, Supplementary Material online), which are digestive enzymes. Given the digestive function of the proventricle in syllids, *OVOCH1* and *OVOCH2* may be performing different functions in *S. magdalena* F and P tissues, respectively. Our molecular phylogeny of *ovochymases* and chymotrypsins in animals confirmed that *OVOCH1* and *OVOCH3* are homologs of other animal *ovochymases*, whereas *OVOCH2* (the one differentially expressed in the proventricle) is, in fact, homolog of mollusk chymotrypsin (supplementary file S8, Supplementary Material online). *OVOCH1* in *S. magdalena* could be assisting in the maturation of the oocyte, creating an envelope that could further prevent self-fertilization during gamete release in the water column.

#### Pairwise Comparisons of Stolons between REPRO Females and Males (REFTOTREPRO Transcriptome)

We detected 1,150 differentially expressed genes in the comparison between reproductive tissues of female and male individuals, 872 upregulated in female stolons and 278 in male stolons (fig. 5C; supplementary file S7, Supplementary Material online). This comparison showed the largest differences, with ~75% of genes upregulated in females (872) and






**Fig. 5.**—Heatmaps based on differentially expressed genes (annotated and not annotated genes) from pairwise comparisons of somatic tissues between females and males (A, B) and reproductive tissues (stolons) (C, D). Proventricle comparisons (A), final segments comparisons (B), female and male stolons comparisons (C), and anterior and posterior parts of stolons (female and male together) (D). Different colors indicate relative expression levels based on raw read counts (see color key and histogram on each). Similarity in expression patterns between genes and individuals is represented by clustering. A, anterior part; P, proventricle; F, final segments; AS, anterior half of stolon; FS, posterior part of stolon.

~25% in males (278) (fig. 5C; [supplementary file S7, Supplementary Material](#) online). In addition, we also compared the anterior and posterior halves of stolons, finding only seven genes upregulated in the anterior half (fig. 5D; [supplementary file S7, Supplementary Material](#) online), most of them related to eye (*rhabdomeric opsin*, *retinal-binding protein*) or brain (*TRPC channel protein*) functioning.

Among the most upregulated Biological Process categories in female stolons, we found Nicotinamide metabolism (fig. 6). Cells need to accommodate the bioenergetic demands during oogenesis, nicotinate and nicotinamide are essential for organisms as the precursors for generation of the coenzymes NAD<sup>+</sup> and NADP<sup>+</sup>, which are fundamental in redox reactions and carry electrons from one reaction to another, being



**Fig. 6.**—Gene ontology treemaps for annotated differentially expressed genes in female stolons versus male stolons. The GO terms downregulated in female stolons are upregulated accordingly in male stolons.

the pillars of many metabolic pathways. The gene *nicotinamide mononucleotide adenylyltransferase 1-like*, which catalyzes the formation of NAD<sup>+</sup>, was upregulated in the female stolon when compared with the male stolon (supplementary file S7, Supplementary Material online). Other metabolic pathways upregulated in the female stolons include both fructose and carbohydrate metabolism, illustrating the high energetic requirements of oogenesis (fig. 6). In male stolons, the major upregulated process related to the high energetic demands of spermatogenesis is Purine metabolism, a pathway required for nucleotide biosynthesis (fig. 6). Interestingly, the MAPK

cascade (included in the category “Styrene catabolism”), which is central to cell proliferation, is upregulated in female stolons (fig. 6). Similarly, the gene *alpha-1D adrenergic receptor-like*, which also regulates cell proliferation is upregulated in female stolons.

As in the case of final segments (see section above), *Vtg* and *OVOCH* in females, and *TSSK* and *NOTCH* in males, were also differentially expressed in stolons of females and males (fig. 5C; supplementary file S7, Supplementary Material online). These results indicate an important role of the stolons in the maturation of gametes, in contrast to what has been

traditionally suggested, where the stolons are thought to be only a place to keep and later spread the gametes. However, no genes related to gamete maturation were found differentially expressed in the comparison between the anterior and posterior halves of stolons, which suggest that there is no sequential anteroposterior maturation of gametes within the stolons (fig. 5D; [supplementary file S7, Supplementary Material](#) online), in agreement with our results from the morphological and ultrastructural study.


*Relaxin* was also found differentially expressed in female stolons, reinforcing the hypothesis of its implication in annelid oogenesis and its potential role in the release of oocytes into the water column, as it has been suggested for *relaxin* in *A. pectinifera* (Mita et al. 2014). Other genes involved in gametogenesis of annelids (e.g., Rebscher et al. 2007; Dill and Seaver 2008; Novo et al. 2013) were also found differentially expressed in female stolons ([supplementary file S7, Supplementary Material](#) online), including the member of the DEAD-box helicase protein family, *vasa*. We found two paralogs of the gene *vasa* (the DE *vasa1* and the non-DE *vasa2*) among our transcripts, in contrast to what is found in other annelids that only present one (see [supplementary file S9, Supplementary Material](#) online). While *vasa2* grouped with all *vasa* orthologs obtained in annelids, *vasa1* branched out from the annelids and appeared basal to other *vasa* orthologs from metazoans ([supplementary file S9, Supplementary Material](#) online), being more similar to *ATP-dependent RNA helicase vasa-like* proteins in arthropods than to *vasa* proteins of annelids when blasted. These results may suggest that different paralogs may be performing different functions in *S. magdalena* ([supplementary file S9, Supplementary Material](#) online). While *vasa2* could be playing a role in the female germline determination localized in the oocytes of *S. magdalena*, *vasa1* could be participating in the maintenance of totipotency of the stem cells (Juliano and Wessel 2010), although *ATP-dependent RNA helicase vasa-like* proteins are also known to be involved in oogenesis. Interestingly, we also found the category Steroid biosynthesis upregulated in female stolons (fig. 6). In addition, our study shows the upregulation of the gene *hydroxysteroid dehydrogenase 2 isoform X2*, that could potentially mediate steroid hormone metabolism (Seckl and Walker 2001), and suggests hormonal control over the final stages of stolonization in *S. magdalena*.

In male stolons, most of the upregulated genes were involved in the construction of the flagellar apparatus (Inaba 2011), including *dyneins*, *cilia-* and the *flagella-associated proteins*, *ropporin*, *radial spoke 3*, and *kinesins*. This is unsurprising, given the presence of sperm in these tissues, but is an excellent positive control.

### Hormonal Control of Stolonization

Because MF was discovered to be produced by mandibular organs of numerous crustaceans, this form of the insect JH (JH

III), has been commonly considered as the crustacean equivalent of insect JH (Laufer and Biggers 2001; Miyakawa et al. 2013). Comparably to JH in insects, MF regulates many aspects of crustacean physiology, including reproduction (Xie et al. 2016). In this context, MF is more actively synthesized by females during vitellogenesis, and higher levels of MF are associated with large reproductive systems and aggressive mating behavior in males of the spider crab *Libinia emarginata* (Laufer et al. 1992). In the annelid *C. teleta*, exogenous extracts of MF were found to affect larval metamorphosis and settlement (Laufer and Biggers 2001), and MF has been recently demonstrated to be directly involved in *P. dumerilii* regeneration and female sexual maturation (Schenk et al. 2016). This latter study not only showed that the decrease of MF levels in the brain induces reproduction and suppresses regenerative capacities in *P. dumerilii*, but it also reported an ortholog of the MF receptor of arthropods (*bHLH-PAS-domain-containing transcription factor methoprene-tolerant receptor, MTr*) in the leucocytes (coelomic cells that synthesize yolk via production of *Vtg* protein), demonstrating that this hormone is not restricted to arthropods, as it was assumed (Schenk et al. 2016). Because detection of MF is not possible in RNAseq data, in order to assess whether *S. magdalena* could use a similar molecular signal to determine when to divert resources from somatic functions to reproduction, we investigated if *S. magdalena* also possessed an ortholog of *MTr*, identified as the arthropod and lophotrochozoan sesquiterpenoid receptor (e.g., Konopova and Jindra 2007; Miyakawa et al. 2013; Jindra et al. 2015; Schenk et al. 2016). In our de novo transcriptomes, we identified two transcripts encoding bHLH-PAS-domain-containing transcription factor that showed strong similarity to *P. dumerilii MTr*. In fact, our molecular phylogeny of *MTr* revealed that the *S. magdalena* ortholog is closely related to *MTr* orthologs of *P. dumerilii* and *C. teleta* (fig. 7A). In agreement with Schenk et al. (2016), our results also confirmed that annelid *MTr* is clearly an ortholog of insects and crustaceans *MTrs* (fig. 7A). These findings allow us to suggest that MF may be one of the hormones responsible for syllid stolonization. If the MF is involved in syllid reproduction, we would expect to find differences in the levels of expression of MF receptors (*MTr*) among the stolonizing and nonstolonizing syllid samples (higher in the latter), similar to what has been reported during oocyte maturation and male reproductive behavior in crustaceans and other annelids (e.g., Laufer et al. 1992; Schenk et al. 2016). Surprisingly, higher expression levels (albeit not statistically significant) of *MTr* were found only in anterior and posterior tissues of female, therefore REPRO individuals (fig. 7B), but not in the NON-REPRO specimens as it was postulated by Schenk et al. (2016). In addition, we also found high expression levels (albeit not statistically significant) of the *Farnesoid nuclear X receptor (FXr)* (Forman et al. 1995) in the anterior tissue of females and in the anterior and the proventricle of males (fig. 7B; [supplementary file S10, Supplementary](#)




**Fig. 7.**—(A) Phylogenetic reconstruction of the protein alignment for methoprene-tolerant receptor (MTr) found in our samples. (B) Heatmap showing the relative levels of expression in the different tissues and conditions analyzed of the transcripts that putatively may be involved in the synthesis of the neurohormone methyl farnesoate (MF): MTr, Farnesol oxidase/dehydrogenase (SDR11), Farnesal dehydrogenases (ALDH3), the differentially expressed transcript Farnesyl pyrophosphate synthase (FPPS) and putative methyl transferase (Mtase). Different colors indicate relative expression levels based on raw read counts (see color key and histogram on each). (C) Phylogenetic reconstruction of the differentially expressed *MTases* in the female stolon. (D) Synthesis pathway of MF and JH in arthropods. A, anterior part; P, proventricle; F, final segments; AS, anterior half of stolon; FS, posterior part of stolon.

Material online). Thus, in contrast to what was found in *P. dumerilii* but similar to what has been reported for arthropods, an increase in MF (or a similar putative sesquiterpenoid) may be necessary to initiate the reproductive process in stolonizing syllids (fig. 7B) (Laufer et al. 1992; Gäde et al. 1997; Wyatt 1997; Hansen et al. 2014). The fact that the differences between conditions are not statistically significant can be explained because the NON-REPRO specimens were collected only one week before the beginning of the stolonization process, and therefore they might have already entered the initial stages of reproduction without visible morphological changes. On the other hand, as in the case of *A. marina* (e.g., Pacey and Bentley 1992), it is also possible that a non-identified hormone, sesquiterpenoid or otherwise, is orchestrating the important metamorphic changes that occur during syllid stolon development, similarly to what MF and JHs do in arthropods (e.g., Hui et al. 2010; Maruzzo et al. 2012; Wen et al. 2015). However, the presence of sesquiterpenoids is further suggested by other DE gene results, as discussed further below.


Interestingly, other neurotransmitter receptors were found to be upregulated in the posterior end of NON-REPRO specimens: *dopamine receptor (DAR)*, belonging to the large family of G-protein coupled receptors, was downregulated in the final segments of females, and *serotonin transporter (SERT or 5-HTT)*, which terminates the action of serotonin, was downregulated in the final segments of males (supplementary file S7, Supplementary Material online; fig. 8A). Our molecular phylogeny corroborates that these proteins are orthologs of the *C. teleta DAR type 2 (DAR2; fig. 8B)* and *C. teleta* and *Helobdella robusta SERT* genes (fig. 8B). Dopamine (DA) and Serotonin (SER) are biogenic amines that act as neurotransmitters and hormones, regulating an array of important physiological functions both in vertebrates and invertebrates (e.g., Winberg et al. 1997; Neckameyer, 1998a; Gingrich et al. 2000; Wicker-Thomas and Hamann 2008; Dufour et al. 2010; Giang et al. 2011). In *D. melanogaster* DA and SER control a wide range of behavioral processes such as circadian rhythms, sleep, mating behavior, learning or aggression (e.g., Nichols 2007; Giang et al. 2011), and also stimulate fertility and female receptivity (Neckameyer 1998b; Marican et al. 2004). In *C. elegans*, male mating behavior and egg deposition are also induced by DA and SER (Sulston et al. 1975; Weinshenker et al. 1995; Dempsey et al. 2005). In addition, both hormones have been reported to be involved in larval metamorphosis in cnidarians, molluscs, and echinoderms (Couper and Leise 1996; McCauley 1997; Matsuura et al. 2009). In annelids, dopaminergic and serotonergic systems have been found in several species (Grothe et al. 1987; Dietzel and Gottmann 1988; Schlawny et al. 1991; Spörhase-Eichmann et al. 1998; Krajniak and Klohr 1999; Zaccardi et al. 2004; Lawrence and Soame 2009; Helm et al. 2014; Rinskaya-Korsakova et al. 2016; Bauknecht and Jékely 2017; Verasztó et al. 2017). However, the participation of

DA and SER in annelid reproduction has only been demonstrated in a handful of studies. Although it was thought that DA played an important role in sexual differentiation in *Ophryotrocha puerilis* (Grothe and Pfannenstiel 1986; Grothe et al. 1987; Pfannenstiel and Spiehl 1987), it was later demonstrated that the catecholaminergic system of this species was involved in mechano- and/or chemoreception (Schlawny et al. 1991). In contrast, both SER and DA in nereids seem to have a positive effect on oocyte development, the first by directly inducing their maturation and the second by switching off the action of the JH (Lawrence and Soame 2009). Similarly, in the decapod *Penaeus merguensis* SER induces ovarian maturation through MF production (Makkapan et al. 2011). In this sense, increased levels of both hormones, as indicated by the upregulation of their receptors and/or transporters (*DAR* and *SERT*) just before the beginning of stolonization (NON-REPRO individuals), could be the stimulus required to initiate oocyte and sperm development during syllid stolonization, with a decrease in the levels afterwards during the course of gametogenesis. In addition to this suggested putative direct role in gametogenesis per se, DA could also be the putative hormone in the brain and/or proventricle inducing the production of MF (or other sesquiterpenoid) to regulate stolonization in *S. magdalena*, as found for DA and the JH of nereids and decapods (Lawrence and Soame 2009; Makkapan et al. 2011). Our results thus indicate a possible role of several hormonal factors in the sexual differentiation of stolons, in agreement with previous studies (Franke 1980; Heacox and Schroeder 1982).


In addition, if DA and SER were the neurohormones regulating stolonization in syllids, our results do not support the traditional view in which male stolons differentiate autonomously and female stolons differentiate upon hormone release by the male stolon (Franke 1999). We found upregulation of the receptors of these two neurohormones in both female and male individuals at the beginning of stolonization. DA and SER have been reported to be under the influence of photoperiodic and circadian rhythms, which are essential for synchronizing several processes in animals (Andretic and Hirsh 2000; Doyle et al. 2002; Lawrence and Soame 2009). Therefore, we suggest that both female and male stolon differentiation are triggered by environmental cues regulating the production of DA and SER. As in other annelids, the main external signals that may be controlling the synchronicity of the reproductive period in syllids are light and seawater temperature (e.g., Franke 1986b). In the Adriatic Sea, the breeding season of *Syllis prolifera* is restricted from late March to early October, when the temperature ranges from 14 to 19 °C, and there are around 12–13 h of light per day (Franke 1986b). Similar results were observed in *S. magdalena*, which seems to breed during the southern hemisphere summer (see sampling methods) with a mean seawater temperature around 15 °C and around 13 h of light per day.



**Fig. 8.**—Phylogenetic reconstruction (A) and heatmap of relative levels of expression in all the tissues and conditions (B) of the genes *dopamine receptor* (*DAR*) and *serotonin transporter* (*SERT*). Different colors indicate relative expression levels based on raw read counts (see color key and histogram on each). A, anterior part; P, proventricle; F, final segments; AS, anterior half of stolon; FS, posterior part of stolon.



**Fig. 9.**—Phylogenetic reconstruction of the protein alignment for the different *opsin* genes (*rhabdomeric* and *ciliary*) found in our samples (A) and levels of expression of all of them in the different tissues and conditions analyzed (B). *Rhabdomeric opsin 5* appeared differentially expressed in the anterior part of stolons. A, anterior part; P, proventricle; F, final segments; AS, anterior half of stolon; FS, posterior part of stolon. Different colors indicate relative expression levels based on raw read counts (see color key and histogram on each).



**FIG. 10.**—Proposed multihormonal model for stolonization control. During the breeding season, DA and SER levels increase in response to external stimuli triggering gamete production in the final segments (up-regulation of DA and SER) (A). Once stolonization has begun, a variety of other hormones and proteins are produced for the correct development and maturation of gametes (up-regulation of Vtg, OVOCH Relaxin, Follistatin, and TSSK) (B). Finally, when gametes are completely mature and also as a response to external stimuli (up regulation of r-opsins), MF or a similar hormone (up-regulation of FPP and Mtransf) is produced to allow stolon release (C). Dashed lines represent hypothesized involvement of molecules, whereas solid lines represent molecule expression results observed in our study.



In addition to steroid hormone control, we found some differentially expressed genes in the female stolons, potentially involved in the production of pheromones (specifically the sesquiterpenoid MF; see section above): *Farnesyl pyrophosphate synthase (FPPS)* and several *methyl transferases (MTases)* (fig. 7B and C; [supplementary file S7, Supplementary Material](#) online), which could synthesize sesquiterpenoids similar to MF and JHIII in arthropods (e.g., Tobe and Bendena 1999; Hui et al. 2010). Specifically, *FPPS* is required at the beginning of the process to catalyze the reaction, generating Farnesyl Diphosphate, the raw material for sesquiterpenoid production, which is then transformed into Farnesol (through *Farnesol phosphatase, FP*), then Farnesal (via the *Farnesol oxidase/dehydrogenase, SDR11*), later into FA (through *Farnesal dehydrogenases, ALDHE3*), and, in the canonical pathway, finally into MF in crustaceans (through *Farnesoic acid methyl transferase, FAMEt*), or into JH in insects (through an *epoxidase, FAMEt* and *Juvenile hormone acid O-methyltransferase, JHAMT*) (e.g., Hui et al. 2010) (fig. 7D).

Following Schenk et al. (2016) and given our results (including those for *methoprene-tolerant receptor*, and *Farnesoid X receptor*, above), a similar pathway seems to occur in annelids, with the synthesis of some form of sesquiterpenoid regulating reproduction, as occurs in arthropods (Xie et al. 2016). In fact, our phylogenetic results confirmed that the differentially expressed transcripts annotated as *FPPS* and of a variety *MTases* (fig. 7C; [supplementary file S10, Supplementary Material](#) online) are orthologs, and thus the beginning and end of the synthesis cascade, and the likely bottleneck, are differentially expressed. In addition, orthologs of *FPP*, *SDR11*, and *ALDHE3* of spiralian were clearly found in our samples ([supplementary file S10, Supplementary Material](#) online), although these are not differentially expressed themselves. These differentially expressed *MTases* are of a variety of annotations, with some possessing homologs across the Bilateria. None possess clear homology to known arthropod *FAMEt* or *JHAMT* sequences. However, all could potentially be performing a similar role in vivo, and one apparent Spiralia novelty is present, which we posit as an excellent candidate for future functional investigation.

However, despite this persuasive circumstantial evidence, we still cannot confirm that the final product of this biosynthetic pathway in *S. magdalena* is MF or another sesquiterpenoid, until functional analyses are performed to test this hypothesis. Besides the putative involvement of sesquiterpenoids in the beginning of syllid stolonization, which is reinforced by the high expression of *SDR11* and *ALDH3* in somatic tissues of both male and female individuals (fig. 7B), it seems that in our case it may also affect later stages, because *FPPS* and *MTases* are differentially expressed in female stolons ([supplementary file S7, Supplementary Material](#) online). Thus, the increase of MF levels could also be regulating the *vitellogenin* levels necessary for yolk formation, as it commonly occurs with JH in arthropods (Laufer et al. 1992; Gäde et al. 1997;

Wyatt 1997; Hansen et al. 2014). In fact, the overexpression of this hormone in stolons could be the triggering signal for the stolon release from the stock. We did not find any enzyme necessary to synthesize hormones or neuropeptides differentially expressed in the male stolons, which might indicate that the synchronicity in the release of female and male stolons might be directly controlled by the female via the production of MF, as it has been also reported during spawning in *A. marina* (Hardege and Bentley 1997).

In addition, as discussed above, MF production has been shown to be influenced by external stimuli (e.g., Shin et al. 2012; Girish et al. 2015; Toyota et al. 2015), which could trigger the stolonization process simultaneously in syllid species according to the traditional hypothesis (e.g., Franke 1999). One of these external stimulus is ambient light variation, which is detected via photosensitive pigments such as opsin proteins and represents a common mechanism mediating the synchronization of gamete release or spawning in a variety of marine invertebrates (Kaniewska et al. 2015; Siebert and Juliano 2017). We have identified several *opsin* homologs in *S. magdalena*, including a *rhabdomeric opsin* previously characterized in other annelids (e.g., Arendt et al. 2004; Randel et al. 2013; Gühmann et al. 2015), that was found differentially expressed in the anterior part of stolons ([supplementary file S7, Supplementary Material](#) online), but not in the anterior part of the stock. Our molecular phylogeny including all *opsins* found in *S. magdalena* (fig. 9A) revealed that the differentially expressed *rhabdomeric opsin (r-opsin 5)* and two other nondifferentially expressed *opsins (r-opsin 3 and 4)* are homologs of the *P. dumerilii opsin* found in larval eyes (Arendt et al. 2002). Differences on expression levels among tissues and conditions were observed in the different *opsins* found in our samples (fig. 9B), which suggest several roles of *opsins* at different stages of syllids development, as it has been already established in other marine annelids (e.g., Arendt et al. 2004). Specifically, the upregulation of *r-opsin 5* in the anterior part of the stolons, where the stolon eyes are located (figs. 2A, 2B and 3A, 3B) suggests that this *opsin* copy in particular might be responsible for detecting the light changes that would trigger MF production, and the subsequent synchronous stolon release and spawning in *S. magdalena*. A similar mechanism has been recently demonstrated in the hydrozoan jellyfish *Clytia hemisphaerica*, in which spawning is mediated by oocyte maturation-inducing neuropeptide hormones, whose release is triggered as a response to blue-cyan light detected by a gonad photosensory *opsin* (Artigas et al 2018).

## Conclusions

Using Illumina RNA-seq data, we provide the first transcriptomic characterization of the reproductive process in a species of the family Syllidae. Here, we performed a series of pairwise comparisons of gene expression patterns in different tissues

and conditions that allowed us to identify the molecular mechanisms underlying the stolonization process of *S. magdalena*. We found an array of differentially expressed genes involved in immune response, neuronal development, gametogenesis, cell proliferation, and steroid metabolism playing different roles in the reproductive process of *S. magdalena*. Among the most striking results of our study was the continuous gamete maturation occurring in both the final segments and the stolons and the hormonal regulation of the reproduction. Thus, following previous hypotheses proposed for other annelids, including syllids (e.g., Franke and Pfannenstiel 1984; Pacey and Bentley 1992; Franke 1999; Lawrence and Soame 2009; Schenk et al. 2016), we suggest a multihormonal model for the control of syllid stolonization, influenced by environmental signals affecting the anterior part (prostomium) and proventricle of the animal, as it was traditionally hypothesized (e.g., Franke 1999), but also influencing the posterior end of the animals (and thus, the gonads) (fig. 10). When the breeding season approaches, both DA and SER levels increase triggered by photoperiod and circadian rhythms (Andretic and Hirsh 2000; Lawrence and Soame 2009) and they directly influence the gonads of prereproductive individuals (upregulation of DA/SER in final segments of NON-REPRO), initiating gamete production (fig. 10A and B). The increase of DA and SER could also positively regulate the production of the putative brain and/or proventricle hormones (such as MF or similar), as in several other invertebrates (Couper and Leise 1996; McCauley 1997; Matsuura et al. 2009) regulating the gamete production (and the metamorphosis to produce stolons), as observed in crustaceans and insects (e.g., Shin et al. 2012; Girish et al. 2015; Toyota et al. 2015). At this point, a variety of other hormones and proteins, such as *Vtg*, *OVCH*, *relaxin*, *follicle-stimulating hormone*, and *TSSK*, play their role in the correct development of gametes (fig. 10B) until maturation is completed. During gamete and stolon maturation, high levels of MF may be required for yolk formation (upregulation in female stolon of *Vtg*, *FPPS*, and *MTases*), and the presence of MF could additionally trigger stolon release from the stock as a response to external stimuli (as indicated by the upregulation of photosensitive *r-opsins*) (fig. 10C). We also suggest that the synchronicity of the stolon and gamete release may not only be mediated by exogenous factors such as light and water temperature, but also by chemical cues provided by the female stolons, as demonstrated in other annelids (Hardege and Bentley 1997).

Overall, our results illuminate the process of stolonization in syllids, improving our understanding of how some putative hormones and gametogenesis-related genes regulate the reproduction in stolonizing syllids. However, the transcriptomic approach adopted here does not allow us to locate the specific expression of these genes, and further functional studies are needed to provide a more complete overview of the expression patterns and the proper functioning of specific pathways during reproduction in *S. magdalena*. In addition, RNAi

or CRISPR/Cas9 experiments to inhibit the expression of G-protein coupled receptors and other hormones and neuropeptides would provide promising routes to understand their role during stolonization in syllids, allowing us to elucidate once and for all how these annelids delegate sex to their stolons.

## Materials and Methods

### Sample Collection and Preservation

Eight individuals of *S. magdalena* were collected in intertidal algal turfs of *Ulva rigida* and *Perumytilus purpuratus* beds, in Las Cruces, Central Chile (33°30'06"S, 71°37'55"W) in January 2014. Four specimens were collected during full moon, two of which were developing female stolons and the other two male stolons (REPRO specimens); the other four specimens were sampled before the full moon and were not engaged in reproduction (NON-REPRO specimens). All samples were immediately fixed in RNA<sub>later</sub> and stored at -80 °C until RNA extraction. Two additional male and female stolons were preserved complete in 2.5% glutaraldehyde in 0.4 M PBS for electron and confocal microscopy.

### Confocal and Transmission Electron Microscopy

Whole specimens preserved in 2.5% glutaraldehyde were mounted in slides to obtain images of autofluorescent tissues during stolonization with a Nikon Eclipse upright with A1-Si confocal microscope at the Image Analysis Center (IAC) of the Natural History Museum of London. No stain was applied, but images were obtained in DAPI 488, 555, and 647 channels, under gentle laser excitation. For transmission electron microscopy (TEM), specimens fixed in 2.5% glutaraldehyde were later postfixed in 1% osmium tetroxide and rinsed twice in PBS before dehydration with an increasing series of acetone (from 50% to 100%). Samples were further embedded in epoxy resin, serially sectioned with an ULTRACUT ultramicrotome at 64 nm, poststained with uranyl acetate and lead citrate, and observed with a JEOL JEM1010 microscope at the Serveis Científic-Tècnics (SCT) at the Universitat de Barcelona and at the Servicio Interdepartamental de Investigación (SIDI) of the Universidad Autónoma de Madrid.

### RNA Extraction

Our biological replicates (same biologic samples taken from different specimens,  $n = 8$ , 4 REPRO—two males and two females—and four NON-REPRO) were as follows: three somatic parts were chosen for RNA extraction from all specimens: anterior part (A = prostomium + first two segments), proventricle (P = all segments containing the proventricle), and final part (F = pygidium + two final segments). In addition, we sequenced the stolons (S) from specimens engaged in stolonization (REPRO): both the anterior (AS) and posterior

half parts (FS). Each tissue sample was transferred to a microcentrifuge tube containing 500  $\mu$ l of TRIzol (Invitrogen), and ground with a RNase-free plastic pestle to break down the tissue, and isolate RNA and DNA. Then, another 500  $\mu$ l of TRIzol and 10  $\mu$ l of glycogen were added. After 10 min incubating the mixture at room temperature (RT), 100  $\mu$ l of the RNA-isolating reagent bromochloropropane was mixed in by vortexing. After 10 min incubation at RT, samples were centrifuged at 16,000 relative centrifugal force (rcf) units for 15 min at 4 °C to separate the solution into three layers. The upper aqueous layer, which contained total RNA, was recovered and mixed with 500 ml of isopropanol, and incubated at –20 °C overnight. Afterwards, the sample was centrifuged at 16,000 rcf for 15 min at 4 °C, and the supernatant was removed. Total RNA precipitation was performed by washing the remaining pellet twice by adding 1 ml of 75% ethanol and centrifuging it at 16,000 rcf at 4 °C for 5 min. The dried pellet was eluted in 100  $\mu$ l of RNA Storage solution (Invitrogen). mRNA purification was performed with a Dynabeads mRNA Purification Kit (Invitrogen), following manufacturer's instructions. After incubation of total RNA at 65 °C for 5 min, the samples were incubated for 30 min with 200 ml of magnetic beads in a rocker and washed twice with washing buffer.

Thirteen microliters of 10 mM Tris–HCl were added to the eluate and the mixture was incubated at 80 °C for 2 min. The supernatant was immediately transferred to a 0.5 ml microcentrifuge tube and stored at –80 °C. Quality of mRNA was measured with a pico RNA assay in the Agilent 2100 BioAnalyzer (Agilent Technologies). Quantity was measured with an RNA assay in a Qubit fluorometer (Life Technologies). Further details about RNA prep protocols can be found in Fernández et al. (2014).

### cDNA Library Construction and Next-Generation Sequencing

cDNA libraries were constructed from extracted mRNA in the Apollo 324 automated system using the PrepX mRNA 8 Protocol Kit (IntegenX) set to 200 base pairs (bp) and stranded mRNA, under the Library Prep Illumina setting. A polymerase chain reaction (PCR) was run to amplify cDNA libraries, using the KAPA Library Amplification Kit. PCR was run as follows: Denaturation (45 s at 98 °C), cycling (15 s at 98 °C, 30 s at 60 °C, and 15 s at 72 °C, for 16 cycles), and final extension (1 min at 72 °C). During the PCR process, the samples were marked with a different index to allow pooling for sequencing. cDNA library quality and size were measured through a dsDNA high sensitivity (HS) assay in an Agilent 2100 BioAnalyzer (Agilent Technologies). A quantitative real-time PCR (qPCR) was run to measure cDNA library concentration using the KAPA Library Quantification Kit. qPCR settings were as follows: Initial denaturation (5 min at 95 °C for 1 cycle), then denaturation (30 s at 95 °C) and annealing/extension/

data acquisition (45 s at 60 °C) combined for 35 cycles. The libraries were then run on the Illumina HiSeq 2500 sequencing platform, with output of paired-end reads of 150 bp by the FAS Center for Systems Biology at Harvard University.

### Sequence Processing and De Novo Assembly

Demultiplexed Illumina HiSeq 2500 sequencing data sets of the 30 tissue samples, in FASTQ format, were retrieved; the quality of the raw reads was assessed and visualized using FASTQC v. 0.11.5 ([www.bioinformatics.babraham.ac.uk](http://www.bioinformatics.babraham.ac.uk)). Adapter sequences and bases with low-quality phred scores (<30) were trimmed off, and a length filter was applied retaining sequences of >25 bases using TRIMGALORE v. 0.4.2 ([www.bioinformatics.babraham.ac.uk](http://www.bioinformatics.babraham.ac.uk)).

Two de novo transcriptome assemblies for *S. magdalena* were constructed with the software Trinity to streamline further differential gene expression analyses (Grabherr et al. 2011; Haas et al. 2013): A reference transcriptome (REFSOM assembly) containing reads from only the somatic parts (anterior part, proventricle, final segments) of each individual of both REPRO and NON-REPRO specimens (23 libraries), and a reference transcriptome including the 5 different parts (anterior part, proventricle, final segments, anterior half part of stolon, and posterior half of stolon) of each individual (13 libraries) for only the reproductive specimens (REFTOTREPRO assembly). We did not obtain enough RNA from two of the female tissue samples, proventricle of specimen 0 and anterior part of stolon of specimen 1, to build a library, and therefore conditions “proventricle” and “anterior half of stolon” were represented by a single library in females. Given the large number of raw reads obtained in our study (>500 million reads), we assembled two different reference transcriptomes, because assembling a single reference transcriptome with the available computational resources would have proved computationally impossible. Raw reads have been deposited in the Sequence Read Archive (BioProject ID PRJNA434571; SRA accession: SRP133371).

For further quantitative assessment of the assembly and annotation completeness we applied the software tool BUSCO (Benchmarking Universal Single-Copy Orthologs; Simão et al. 2015), with default settings using the metazoan database (metazoan\_odb9, dated February 13, 2016). This method is based on evolutionarily informed expectations of gene content and is broadly used as a benchmark for testing completeness of genomes and transcriptomes.

### Transcriptome Characterization: Blast and Annotation

Annotation of transcriptome contigs or transcripts (containing all isoforms) for both de novo assemblies were done separately using BlastX against a selection of nonredundant (nr) database from NCBI containing only proteins from Metazoa, with an expected value (*E*-value) cutoff of  $1e^{-5}$  (Altschul

et al. 1997). BLAST results of the two de novo assemblies were used to retrieve Gene Ontology (GO) terms with BLAST2GO 4.0.2 (Conesa et al. 2005) under the three different categories: CC, BP, and MF. In addition, GO enrichment analyses using Fisher's test were done in BLAST2GO, to assess which GO terms were significantly overrepresented in pairwise comparisons between both REFSOM and REFTOTREPRO transcriptomes. The *P*-value for the reciprocal comparisons was adjusted to a 0.05 false discovery rate (FDR) (Benjamini and Hochberg 1995). The Galaxy web-based platform (<http://usegalaxy.org>) was used to align the RSEM results of each sample with BlastX results for the de novo assemblies for display.

### Estimation of Expression Levels

In order to obtain expression levels, as read counts, of genes (with all isoforms collapsed) for each tissue type of *S. magdalena* specimens in both reproductive and nonreproductive conditions, trimmed paired reads after trimming were mapped against the reference transcriptome, using BOWTIE2 v. 2.2.1 (Langmead and Salzberg 2012), as implemented in Trinity (Grabherr et al. 2011). The software RSEM v. 1.2.11 (Li and Dewey 2011) was used to generate a table containing read counts.

### Differential Gene Expression Analyses

Differential gene expression analyses were computed in pairwise comparisons of different tissues and conditions using the R package DESeq2, which allows analyses to be performed with low numbers of replicates (Anders and Huber 2010). Before analyzing differential gene expression, read counts were normalized by estimating a scaling factor for each transcript in DESeq2 (Dillies et al. 2013). The significance value for multiple comparisons was FDR adjusted to 0.01 (Benjamini and Hochberg 1995). Visualization of the significant outcomes of genes differentially expressed (upregulated and downregulated) between the tissues and conditions was obtained with a heatmap performed with the "GPLOTS" package of R (<http://www.r-project.org>). Using the GO annotation results for the "reference" transcriptome, we obtained the GO terms associated with the differentially expressed isoforms in both pairwise comparisons, which were then implemented together with their *P*-value (adjusted) associated in REVIGO web server (Supek et al. 2011), and graphically represented with the "TREETMAP" function in R. Size of the rectangles was adjusted to reflect the *P*-value using the `abs_log_pvalue` option in REVIGO.

### Phylogenetic Analyses

The evolutionary history of specific genes that could potentially be involved in the stolonization process was also assessed through phylogenetic inference. The translated amino acid sequences of these genes were aligned with ortholog of the

same genes in other metazoans obtained from GenBank using MUSCLE ver. 3.6 (Edgar 2004). The G-protein coupled receptors *DAR2* and *SERT* were analyzed together. Both *vasa* and *PL10* are *DEAD-box* helicases and were analyzed together. Other genes were examined in their individual gene families. We selected the best-fit model of amino acid substitution (LG +  $\Gamma$  + G, WAG, as indicated in Figure legends) with ProtTest ver. 2.4 (Abascal et al. 2005) under the Akaike Information Criterion (Posada and Buckley 2004) and later fed into the software for phylogenetic reconstruction. Maximum likelihood analyses of all the genes were conducted in RAxML ver. 7.2.7 (Stamatakis 2006) with 500 independent searches and 1000 bootstrap replicates (Stamatakis et al. 2008).

### Supplementary Material

*Supplementary data* are available at *Genome Biology and Evolution* online.

### Acknowledgments

The authors are indebted to many members of the Giribet Lab at Harvard University for their help during sample processing, specially to Dr Sarah Lemer and Dr David Combosch (currently at University of Guam). Special thanks go to Dr Greg Rouse (Scripps Institution of Oceanography, UCSan Diego) and Dr Carlos Sentís (UAM) who provided comments and advice at the beginning of the research. The first author is also very grateful to Dr Raquel Pérez-Palacios, Dr Jorge Barbazán (Institut Curie, Paris), and members of Dr Michel Vervoort lab (Institut Jacques Monod) for their support and useful comments to improve the last version of the manuscript. We also thank also to Milagros Guerra (CBM, CSIC) for his help with TEM observations at UAM. This research received funding from the European Union's (European Atomic Energy Community's) Seventh Framework Program (FP7/2007–2013; FP7/2007–2011) under grant agreement 227799 to P.A.-C. Sequencing and analyses were conducted with internal MCZ funds to G.G. and with the support of the Center for Systems Biology and the Research Computing group, both from the Faculty of Arts and Sciences (Harvard University), and with internal funds from the Department of Life Sciences (Natural History Museum of London) to AR. One anonymous reviewer and Associate Editor Mandë Holford provided constructive criticism which helped to improve this study.

### Literature Cited

- Abascal F, Zardoya R, Posada D. 2005. ProtTest: selection of best-fit models of protein evolution. *Bioinformatics* 21(9):2104–2105.
- Abeloos M. 1950. Régénération et stolonisation épigame chez l'Annélide *Syllis prolifera* Krohn. *C R Acad Sci (Comptes rendus de l'Académie des Sciences)*. 230:1899–1900.

- Agassiz A. 1863. On alternate generation in annelids, and the embryology of *Autolytus cornutus*. Boston J Nat Hist. 7:384–409.
- Altschul SF, et al. 1997. Gapped Blast and PSIBlast: a new generation of protein database search programs. Nucleic Acids Res. 25(17):3389–3402.
- Anders S, Huber W. 2010. Differential expression analysis for sequence count data. Genome Biol. 11(10):R106.
- Andretic R, Hirsh J. 2000. Circadian modulation of dopamine receptor responsiveness in *Drosophila melanogaster*. Proc Natl Acad Sci U S A. 97(4):1873–1878.
- Arendt D, Tessmar K, de Campos-Baptista MI, Dorresteyn A, Wittbrodt J. 2002. Development of pigment-cup eyes in the polychaete *Platynereis dumerilii* and evolutionary conservation of larval eyes in Bilateria. Development 129(5):1143–1154.
- Arendt D, Tessmar RK, Snyman H, Dorresteyn AW, Wittbrodt J. 2004. Ciliary photoreceptors with a vertebrate-type opsin in an invertebrate brain. Science 306(5697):869–871.
- Artigas GQ, et al. 2018. A gonad-expressed opsin mediates light-induced spawning in the jellyfish *Clytia*. eLife 7:e29555.
- Asahara S, et al. 2013. *Ras-related C3 botulinum toxin substrate 1 (RAC1)* regulates glucose-stimulated insulin secretion via modulation of F-actin. Diabetologia 56(5):1088–1097.
- Bauknecht P, Jékely G. 2017. Ancient coexistence of norepinephrine, tyramine, and octopamine signaling in bilaterians. BMC Biol. 15(1):6.
- Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. J R Stat Soc Ser B Stat Methodol. 57:289–300.
- Bentley MG. 1985. Sperm maturation response in *Arenicola marina* L.: an in vitro assay for sperm maturation factor and its partial purification. Int J Invertebrate Reprod Dev. 8(3):139–148.
- Bentley MG, Clark S, Pacey AA. 1990. The role of arachidonic acid and eicosatrienoic acids in the activation of spermatozoa in *Arenicola marina* L. (Annelida: polychaeta). Biol Bull. 178(1):1–9.
- Buffet JP, Corre E, Duvernois-Berthet E, Fournier J, Lopez PJ. 2018. Adhesive gland transcriptomics uncovers a diversity of genes involved in glue formation in marine tube-building polychaetes. Acta Biomater. 72:316–328.
- Conesa A, et al. 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. Bioinformatics 21(18):3674–3676.
- Couper JM, Leise EM. 1996. Serotonin injections induce metamorphosis in larvae of the gastropod mollusc *Ilyanassa obsoleta*. Biol Bull. 191(2):178–186.
- Daly JM. 1975. Reversible epitoky in the life history of the polychaete *Odontosyllis polycera* (Schmarda, 1861). J Mar Biol Assoc UK. 55(02):327–344.
- Dempsey CM, Mackenzie SM, Gargus A, Blanco G, Sze JY. 2005. Serotonin (5HT), fluoxetine, imipramine and dopamine target distinct 5HT receptor signaling to modulate *Caenorhabditis elegans* egg-laying behavior. Genetics 169(3):1425–1436.
- Dietzel ID, Gottmann K. 1988. Development of dopamine-containing neurons and dopamine uptake in embryos of *Hirudo medicinalis*. Dev Biol. 128(2):277–283.
- Dill KK, Seaver EC. 2008. *Vasa* and *nanos* are coexpressed in somatic and germ line tissue from early embryonic cleavage stages through adulthood in the polychaete *Capitella* sp. I. Dev Genes Evol. 218(9):453–463.
- Dillies MA, et al. 2013. A comprehensive evaluation of normalization methods for Illumina high-throughput RNA sequencing data analysis. Brief Bioinform. 14(6):671–683.
- Donizetti A, et al. 2008. Two neuron clusters in the stem of postembryonic zebrafish brain specifically express *relaxin-3* gene: first evidence of nucleus incertus in fish. Dev Dyn. 237(12):3864–3869.
- Donizetti A, et al. 2010. Characterization and developmental expression pattern of the *relaxin receptor rxfp1* gene in zebrafish. Dev Growth Differ. 52(9):799–806.
- Doyle SE, Grace MS, McIvor WI, Menaker M. 2002. Circadian rhythms of dopamine in mouse retina: the role of melatonin. Vis Neurosci. 19(5):593–601.
- Dufour S, Sebert ME, Weltzien FA, Rousseau K, Pasqualini C. 2010. Neuroendocrine control by dopamine of teleost reproduction. J Fish Biol. 76(1):129–160.
- Durchon M. 1951. Stolonisation et hermaphroditisme succesif chez *Syllis amica*. Arch Zool Exp Gen. 88:96–100.
- Durchon M. 1952. Recherches expérimentales sur deux aspects de la reproduction chez les Annélides Polychètes: l'épitoquie et la stolonisation. Ann Sci Nat Zool Biol Anim. 14:117–206.
- Durchon M. 1959. Contribution à l'étude de la stolonisation chez les Syllidiens (Annélides, Polychètes): i. Syllinae. Bull Biol Fr Bel. 93:155–219.
- Durchon M, Wissocq J-C. 1964. Contribution à l'étude de la stolonisation chez les Syllidiens (Annélides Polychètes): II. Autolytinae. Ann Sci Nat Zool Biol Anim. 6:159–208.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res. 32(5):1792–1797.
- Fernández R, et al. 2014. Evaluating topological conflict in centipede phylogeny using transcriptomic data sets. Mol Biol Evol. 31(6):1500–1513.
- Forman BM, et al. 1995. Identification of a nuclear receptor that is activated by farnesol metabolites. Cell 81(5):687–693.
- Franke H-D. 1980. Zur Determination der zeitlichen Verteilung von Fortpflanzungsprozessen in Laborkulturen des Polychaeten *Typosyllis prolifera*. Helgol Meeresunters. 34(1):61–84.
- Franke H-D. 1981. Der Einfluß von Temperatur und Tageslänge auf das endokrine System der Fortpflanzungskontrolle bei dem Polychaeten *Typosyllis prolifera*. Verh Dtsch Zool Ges. 190:1–190.
- Franke H-D. 1983a. Endocrine mechanisms mediating light temperature effects on male reproductive activity in *Typosyllis prolifera* (Polychaeta, Syllidae). Roux Arch Dev Biol. 192(2):95–102.
- Franke H-D. 1983b. Endocrine control of reproductive periodicity in male *Typosyllis prolifera* (Polychaeta, Syllidae). Int J Invertebr Reprod. 6(4):229–238.
- Franke H-D. 1985. On a clocklike mechanism timing lunar rhythmic reproduction in *Typosyllis prolifera* (Polychaeta). J Comp Physiol A. 156(4):553–561.
- Franke H-D. 1986a. Resetting a circalunar reproduction rhythm with artificial moonlight signals: phase-response curve and 'moon-off' effect. J Comp Physiol A. 159(4):569–576.
- Franke HD. 1986b. The role of light and endogenous factors in the timing of the reproductive cycle of *Typosyllis prolifera* and some other polychaetes. Am Zool. 26(2):433–445.
- Franke HD. 1999. Reproduction of the Syllidae (Annelida: polychaeta). Hydrobiologia 402:39–55.
- Franke H-D, Pfannenstiel H-D. 1984. Some aspects of endocrine control of polychaete reproduction. Fortschr Zool. 29: 53–72.
- Gäde G, Hoffmann KH, Spring JH. 1997. Hormonal regulation in insects: facts, gaps, and future directions. Physiol Rev. 77(4):963–1032.
- Gao K, Zhang S. 2009. Ovocymase in amphioxus *Branchiostoma belcheri* is an ovary-specific trypsin-like serine protease with an antibacterial activity. Dev Comp Immunol. 33(12):1219–1228.
- Garwood PR. 1991. Reproduction and the classification of the family Syllidae (Polychaeta). Ophelia 5(Suppl):81–87.
- Giang T, Rauchfuss S, Ogueta M, Scholz H. 2011. The serotonin transporter expression in *Drosophila melanogaster*. J Neurogenet. 25(1–2):17–26.
- Giani VC, Yamaguchi E, Boyle MJ, Seaver EC. 2011. Somatic and germline expression of *pivi* during development and regeneration in the marine polychaete annelid *Capitella teleta*. Evodevo 2:10.

- Gingrich B, Liu Y, Cascio C, Wang Z, Insel TR. 2000. Dopamine D2 receptors in the nucleus accumbens are important for social attachment in female prairie voles (*Microtus ochrogaster*). *Behav Neurosci*. 114(1):173.
- Girish BP, Swetha CH, Reddy PS. 2015. Induction of ecdysteroidogenesis, methyl farnesoate synthesis and expression of ecdysteroid receptor and retinoid X receptor in the hepatopancreas and ovary of the giant mud crab, *Scylla serrata* by melatonin. *Gen Comp Endocrinol*. 217:37–42.
- Grabherr MG, et al. 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol*. 29(7):644–652.
- Grothe C, Pfannenstiel HD. 1986. Cytophysiological study of neurosecretory and pheromonal influences on sexual development in *Ophryotrocha puerilis* (Polychaeta, Dorvilleidae). *Int J Invertebr Reprod Dev*. 10(2):227–239.
- Grothe C, Seidl K, Pfannenstiel HD. 1987. Cytochemical and biochemical characterization of neurosecretory material in the brain of an annelid, *Ophryotrocha puerilis* (Polychaeta). *Gen Comp Endocrinol*. 68(1):1–5.
- Gunnerson JM, Crawford RJ, Tregear GW. 1995. Expression of the *relaxin* gene in rat tissues. *Mol Cell Endocrinol*. 110(1-2):55–64.
- Gühmann M, et al. 2015. Spectral tuning of phototaxis by a go-opsin in the rhabdomeric eyes of *Platynereis*. *Curr Biol*. 25(17):2265–2271.
- Haas BJ, et al. 2013. De novo transcript sequence reconstruction from RNA-Seq: reference generation and analysis with Trinity. *Nat Protoc*. 8(8):1494–1512.
- Hafer J, Fischer A, Ferenz HJ. 1992. Identification of the yolk receptor protein in oocytes of *Nereis virens* (Annelida, Polychaeta) and comparison with the locust vitellogenin receptor. *J Comp Physiol B*. 162(2):148–152.
- Hansen IA, Attardo GM, Rodriguez SD, Drake LL. 2014. Four-way regulation of mosquito yolk protein precursor genes by juvenile hormone-, ecdysone-, nutrient-, and insulin-like peptide signaling pathways. *Front Physiol*. 5:103.
- Hao Z, et al. 2004. Expression analysis of the human testis-specific serine/threonine kinase (TSSK) homologues. A TSSK member is present in the equatorial segment of human sperm. *Mol. Hum Reprod*. 10(6):433–444.
- Hardege JD, Bentley MG. 1997. Spawning synchrony in *Arenicola marina*: evidence for sex pheromonal control. *Proc R Soc Lond B Biol Sci*. 264(1384):1041–1047.
- Hauser F, Williamson M, Cazzamali G, Grimmelikhuijzen CJ. 2006. Identifying neuropeptide and protein hormone receptors in *Drosophila melanogaster* by exploiting genomic data. *Brief Funct Genomic Proteomic*. 4(4):321–330.
- Hayashi T, et al. 2001. Requirement of *Notch 1* and its ligand *jagged 2* expressions for spermatogenesis in rat and human testes. *J Androl*. 22(6):999–1011.
- Heacox AE. 1980. Reproduction and development of *Typosyllis pulchra* (Berkeley and Berkeley) (Polychaeta: syllidae). *Pac Sci*. 34:245–259.
- Heacox AE, Schroeder PC. 1982. The effects of prostomial and proventriculus removal on sex determination and gametogenesis in *Typosyllis pulchra* (Polychaeta: syllidae). *Roux Arch Dev Biol*. 191(2):84–90.
- Heikkinen LK, Kesäniemi JE, Knott KE. 2017. *De novo* transcriptome assembly and developmental mode specific gene expression of *Pygospio elegans*. *Evol Dev*. 19(4-5):205–217.
- Helm C, Stevenson PA, Rouse GW, Bleidorn C. 2014. Immunohistochemical investigations of *Myzostoma cirriferum* and *Mesomyzostoma cf. katoi* (Myzostomida, Annelida) with implications for the evolution of the myzostomid body plan. *Zoomorphology* 133(3):257–271.
- Hsu SY, et al. 2002. Activation of orphan receptors by the hormone relaxin. *Science* 295(5555):671–674.
- Hui JH, Hayward A, Bendena WG, Takahashi T, Tobe SS. 2010. Evolution and functional divergence of enzymes involved in sesquiterpenoid hormone biosynthesis in crustaceans and insects. *Peptides* 31(3):451–455.
- Inaba K. 2011. Sperm flagella: comparative and phylogenetic perspectives of protein components. *Mol Hum Reprod*. 17(8):p524–538.
- Ivell R, Anand-Ivell R. 2005. Understanding *relaxin* in the female reproductive system. *Curr Med Chem Immunol Endocr Metab Agents* 5(5):383–389.
- Iversen A, Cazzamali G, Williamson M, Hauser F, Grimmelikhuijzen CJ. 2002. Molecular cloning and functional expression of a *Drosophila* receptor for the neuropeptides *capa-1* and *and-2*. *Biochem Biophys Res Commun*. 299(4):628–633.
- Jindra M, Uhlířova M, Charles JP, Smykal V, Hill RJ. 2015. Genetic evidence for function of the bHLH-PAS protein Gce/Met as a juvenile hormone receptor. *PLoS Genet*. 11(7):e1005394.
- Juliano C, Wessel G. 2010. Versatile germline genes. *Science* 329(5992):640–641.
- Kang D, Pilon M, Weisblat DA. 2002. Maternal and zygotic expression of a *nanos*-class gene in the leech *Helobdella robusta*: primordial germ cells arise from segmental mesoderm. *Dev Biol*. 245(1):28–41.
- Kaniewska P, et al. 2015. Signaling cascades and the importance of moonlight in coral broadcast mass spawning. *eLife* 4:e09991.
- Kenny NJ, et al. 2015. The Lophotrochozoan TGF- $\beta$  signalling cassette-diversification and conservation in a key signalling pathway. *Int J Dev Biol*. 58(6-7-8):533–549.
- Konopova B, Jindra M. 2007. Juvenile hormone resistance gene Methoprene-tolerant controls entry into metamorphosis in the beetle *Tribolium castaneum*. *Proc Natl Acad Sci USA*. 104(25):10488–10493.
- Krajniak KG, Klohr RW. 1999. The effects of FMRFamide, serotonin, and acetylcholine on the isolated crop-gizzard of the earthworm, *Lumbricus terrestris*. *Comp Biochem Physiol A Mol Integr Physiol*. 123(4):409–415.
- Langmead B, Salzberg S. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods*. 9(4):357–359.
- Laufer H, Sagi A, Ahl J, Homola E. 1992. Methyl farnesoate appears to be a crustacean reproductive hormone. *Invertebr Reprod Dev*. 22(1-3):17–19.
- Laufer H, Biggers WJ. 2001. Unifying concepts learned from methyl farnesoate for invertebrate reproduction and post-embryonic development. *Am Zool*. 41:442–457.
- Lawrence AJ, Soame JM. 2009. The endocrine control of reproduction in Nereidae: a new multi-hormonal model with implications for their functional role in a changing environment. *Philos Trans R Soc Lond B Biol Sci*. 364(1534):3363–3376.
- Li B, Dewey C. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinform*. 12:323.
- Li C, Kim K, Nelson LS. 1999. FMRFamide-related neuropeptide gene family in *Caenorhabditis elegans*. *Brain Res*. 848(1-2):26–34.
- Lindsay LL, Hedrick JL. 1995. Isolation and characterization of ovochymase, a chymotrypsinlike protease released during *Xenopus laevis* egg activation. *Dev Biol*. 167(2):513–516.
- Makkapan W, Maikao L, Miyazaki T, Chotigeat W. 2011. Molecular mechanism of serotonin via methyl farnesoate in ovarian development of white shrimp: *Fenneropenaeus merguensis* de Man. *Aquaculture* 321(1-2):101–107.
- Malaquin A. 1893. Recherches sur les Syllidiens, morphologie, anatomie, reproduction, développement. *Mem Soc Sci Agric Arts Lille*. 18:1–477.
- Marican C, Duportets L, Birman S, Jallon JM. 2004. Female-specific regulation of cuticular hydrocarbon biosynthesis by dopamine in *Drosophila melanogaster*. *Insect Biochem Mol Biol*. 34(8):823–830.

- Marion AF, Bobretsky NV. 1875. Etude des Annelides du golfe de Marseille. *Ann Sci Nat Zool Paleontol.* 2:2–46.
- Maruzzo D, Aldred N, Clare AS, Høeg JT. 2012. Metamorphosis in the cirripede crustacean *Balanus amphitrite*. *PLoS One* 7(5):e37408.
- Matsuura H, Yazaki I, Okino T. 2009. Induction of larval metamorphosis in the sea cucumber *Apostichopus japonicus* by neurotransmitters. *Fish Sci.* 75(3):777–783.
- McCaughey DW. 1997. Serotonin plays an early role in the metamorphosis of the hydrozoan *Phialidium gregarium*. *Dev Biol.* 190(2):229–240.
- Mehr S, et al. 2015. Transcriptome sequencing and annotation of the polychaete *Hermodice carunculata* (Annelida, Amphinomidae). *BMC Genomics* 16(1):445.
- Mesnil F, Caullery M. 1919. Sur un processus normal de fragmentation, suivie de régénération, chez un Annelide polychète *Syllis gracilis* Gr.” *Compt Rend Acad Sci* 169:926–929.
- Meyer E, et al. 2009. Sequencing and *de novo* analysis of a coral larval transcriptome using 454 GSFLX. *BMC Genomics* 10(1):1.
- Michel A. 1898. Recherches sur la régénération chez les Annelides. *Bull Sci Fr Bel.* 31:245–420.
- Mino M, Sawada H. 2016. Follicle cell trypsin-like protease Ovochymase: Its cDNA cloning, localization, and involvement in the late stage of oogenesis in the ascidian *Halocynthia roretzi*. *Mol Reprod Dev.* 83(4):347–358.
- Mita M. 2013. *Relaxin*-like gonad-stimulating substance in an echinoderm, the starfish: a novel relaxin system in reproduction of invertebrates. *Gen Comp Endocrinol.* 181:241–245.
- Mita M, Takeshige Y, Nakamura M. 2014. Effect of relaxin-like gonad-stimulating substance on gamete shedding and 1-methyladenine production in starfish ovaries. In: Sawada H., Inoue N., Iwano M. (eds) *Sexual Reproduction in Animals and Plants*. Tokyo, Japan: Springer. p. 115–122.
- Miyakawa H, et al. 2013. A mutation in the receptor Methoprene-tolerant alters juvenile hormone response in insects and crustaceans. *Nat Commun.* 4:1856.
- Murta D, et al. 2014. In vivo *Notch* signaling blockade induces abnormal spermatogenesis in the mouse. *PLoS One* 9(11):e113365.
- Neckameyer WS. 1998a. Dopamine and mushroom bodies in *Drosophila*: experience-dependent and-independent aspects of sexual behavior. *Learn Mem.* 5(1):157–165.
- Neckameyer WS. 1998b. Dopamine modulates female sexual receptivity in *Drosophila melanogaster*. *J Neurogenet.* 12(2):101–114.
- Nichols CD. 2007. 5-HT<sub>2</sub> receptors in *Drosophila* are expressed in the brain and modulate aspects of circadian behaviors. *Dev Neurobiol.* 67(6):752–763.
- Novo M, Riesgo A, Fernández-Guerra A, Giribet G. 2013. Pheromone evolution, reproductive genes, and comparative transcriptomics in Mediterranean earthworms (Annelida, Oligochaeta, Hormogastridae). *Mol Biol Evol.* 30 (7):1614–1629.
- Nygren A. 1999. Phylogeny and reproduction in Syllidae (Polychaeta). *Zool J Linn Soc.* 126(3):365–386.
- Okada YK. 1929. Regeneration and fragmentation in the syllidean polychaetes. (Studies on the Syllidae II). *Wilhelm Roux Arch Entwickl.* 115(3):542–600.
- Okada YK. 1937. La stolonisation et les caractères sexuels du stolon chez les Syllidiens polychètes (Études sur les Syllidiens III). *Jpn J Zool.* 7:441–490.
- Oliniski RP, Lundin L-G, Hallbook F. 2006. Conserved synteny between the *Ciona* genome and human paralogs identifies large duplication events in the molecular evolution of the *insulin-relaxin* gene family. *Mol Biol Evol.* 23(1):10–22.
- Pacey AA, Bentley MG. 1992. The fatty acid 8, 11, 14-eicosatrienoic acid induces spawning in the male lugworm *Arenicola marina*. *J Exp Biol.* 173(1):165–179.
- Pfannenstiel HD, Spiehl D. 1987. Dopamine induces sex reversal in females of *Ophryotrocha puerilis* (Polychaeta). *Cell Differ.* 20:84.
- Park C, et al. 2018. The developmental transcriptome atlas of the spoon worm *Urechis unicinctus* (Echiurida: Annelida). *Gigascience* 7(3):gij007.
- Pérez-Portela R, Turon X, Riesgo A. 2016. Characterization of the transcriptome and gene expression of four different tissues in the ecologically relevant sea urchin *Arbacia lixula* using RNA-seq. *Mol Ecol Resour.* 16(3):794–808.
- Pettibone MH. 1963. Marine polychaete worms of the New England region. 1. Aphroditidae through Trochochaetidae. *Bull US Natl Mus.* 227:1–346.
- Phillips DJ, de Kretser DM. 1998. Follistatin: a multifunctional regulatory protein. *Front Neuroendocrinol.* 19(4):287–322.
- Potts FA. 1911. Methods of reproduction in the syllids. *Ergebnisse Fortschritte Zool.* 3(1):1–72.
- Posada D, Buckley TR. 2004. Model selection and model averaging in phylogenetics: advantages of Akaike information criterion and Bayesian approaches over likelihood ratio tests. *Syst Biol.* 53(5):793–808.
- Randel N, Bezares-Calderón LA, Gühmann M, Shahidi R, Jékely G. 2013. Expression dynamics and protein localization of rhabdomeric opsins in *Platynereis* larvae. *Integr Comp Biol.* 53(1):7–16.
- Rebscher N, Zelada-González F, Banisch TU, Raible F, Arendt D. 2007. *Vasa* unveils a common origin of germ cells and of somatic stem cells from the posterior growth zone in the polychaete *Platynereis dumerilii*. *Dev Biol.* 306(2):599–611.
- Riddiford LM. 1994. Cellular and molecular actions of juvenile hormone I. General considerations and premetamorphic actions. *Adv Insect Physiol.* 24:213–274 [Database].
- Riesgo A, et al. 2012. Comparative transcriptomics of newly sequenced invertebrates and efficiency estimation of genomic sampling in non-model taxa. *Front Zool.* 9(1):33.
- Rimskaya-Korsakova NN, Kristof A, Malakhov VV, Wanninger A. 2016. Neural architecture of *Galathowenia oculata* Zach, 1923 (Oweniidae, Annelida). *Front Zool.* 13(1):5.
- Rivera AS, Gonsalves FC, Song MH, Norris BJ, Weisblat DA. 2005. Characterization of *Notch*-class gene expression in segmentation stem cells and segment founder cells in *Helobdella robusta* (Lophotrochozoa; Annelida; Clitellata; Hirudiniida; Glossiphoniidae). *Evol Dev.* 7(6):588–599.
- Schenk S, Krauditsch C, Frühauf P, Gerner C, Raible F. 2016. Discovery of methylfarnesoate as the annelid brain hormone reveals an ancient role of sesquiterpenoids in reproduction. *eLife* 5:e17126.
- Schlawny A, Hamann T, Müller MA, Pfannenstiel HD. 1991. The catecholaminergic system of an annelid (*Ophryotrocha puerilis*, Polychaeta). *Cell Tissue Res.* 265(1):175–184.
- Schroeder PC, Hermans CO. 1975. *Annelida: polychaeta*. In: Giese AC, Pearse JS, editors. *Reproduction of marine invertebrates*, Vol. 3. New York: Academic Press. p. 1–213.
- Seckl JR, Walker BR. 2001. Minireview: 11 $\beta$ -hydroxysteroid dehydrogenase type I—a tissue-specific amplifier of glucocorticoid action. *Endocrinology* 142(4):1371–1376.
- Siebert S, Juliano CE. 2017. Sex, polyps, and medusae: determination and maintenance of sex in cnidarians. *Mol Reprod Dev.* 84(2):105–119.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31(19):3210–3212.
- Shin SW, Zou Z, Saha TT, Raikhel AS. 2012. bHLH-PAS heterodimer of methoprene-tolerant and Cycle mediates circadian expression of juvenile hormone-induced mosquito genes. *Proc Natl Acad Sci U S A.* 109(41):16576–16581.

- Spörhase-Eichmann U, Winkler M, Schürmann FW. 1998. Dopaminergic sensory cells in the epidermis of the earthworm. *Naturwissenschaften* 85(11):547–550.
- Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22(21):2688–2690.
- Stamatakis A, Hoover P, Rougemont J. 2008. A rapid bootstrap algorithm for the RAxML web servers. *Syst Biol*. 57(5):758–771.
- Sulston J, Dew M, Brenner S. 1975. Dopaminergic neurons in the nematode *Caenorhabditis elegans*. *J Comp Neurol*. 163(2):215–226.
- Supek F, Bošnjak M, Škunca N, Šmuc T. 2011. REVIGO summarizes and visualizes long lists of gene ontology terms. *PloS one* 6(7):p.e21800.
- Takahashi T, et al. 2009. An EST screen from the annelid *Pomatoscerus lamarckii* reveals patterns of gene loss and gain in animals. *BMC Evol Biol*. 9(1):240.
- Thamm K, Seaver EC. 2008. *Notch* signaling during larval and juvenile development in the polychaete annelid *Capitella* sp. I. *Dev Biol*. 320(1):304–318.
- Tobe SS, Bendena WG. 1999. The regulation of juvenile hormone production in arthropods: functional and evolutionary perspectives. *Ann NY Acad Sci* 897(1 NEUROPEPTIDES):300–310.
- Toyota K, et al. 2015. Methyl farnesoate synthesis is necessary for the environmental sex determination in the water flea *Daphnia pulex*. *J Insect Physiol*. 80:22–30.
- Verasztó C, et al. 2017. Ciliomotor circuitry underlying whole-body coordination of ciliary activity in the *Platynereis* larva. *eLife* 6:1–25.
- Verger-Bocquet M. 1984. Mise en évidence de l'influence de la photopériode sur la stolonisation, par l'intermédiaire du prostomium chez *Syllis amica* Quatrefages (Annelide, Polychète). *Arch Biol*. 95:301–306.
- Weidhase M, Beckers P, Bleidorn C, Aguado MT. 2016. On the role of the proventricule region in reproduction and regeneration in *Typosyllis antoni* (Annelida: syllidae). *BMC Evol Biol*. 16(1):196.
- Weinshenker D, Gian G, Thomas JH. 1995. Genetic and pharmacological analysis of neurotransmitters controlling egg laying in *C. elegans*. *J Neurosci*. 15(10):6975–6985.
- Wen D, et al. 2015. Methyl farnesoate plays a dual role in regulating *Drosophila* metamorphosis. *PLoS Genet*. 11(3):e1005038.
- Wicker-Thomas C, Hamann M. 2008. Interaction of dopamine, female pheromones, locomotion and sex behavior in *Drosophila melanogaster*. *J Insect Physiol*. 54(10–11):1423–1431.
- Wilson BC, Burnett D, Rappaport R, Parry LJ, Fletcher EK. 2009. *Relaxin-3* and *RXFP3* expression, and steroidogenic actions in the ovary of teleost fish. *Comp Biochem Physiol A Mol Integr Physiol*. 153(1):69–74.
- Winberg S, Nilsson A, Hylland P, Söderstöm V, Nilsson GE. 1997. Serotonin as a regulator of hypothalamic-pituitary-interrenal activity in teleost fish. *Neurosci Lett*. 230(2):113–116.
- Wissocq JC. 1966. Rôle du proventricule dans le déterminisme de la stolonisation de *Syllis amica* Quatrefages (Annelide Polychète). *C R Acad Sci*. 262:2605–2608.
- Wissocq JC. 1970. Évolution de la musculature longitudinale dorsale et ventrale au cours de la stolonisation de *Syllis amica* Quatrefages (Annelide polychète): Muscles du ver asexué et muscles du stolon. I. *J Microsc*. 9:355–358.
- Wyatt GR. 1997. Juvenile hormone in insect reproduction - a paradox? *Eur J Entomol*. 94:323–333.
- Wyatt GR, Davey KG. 1996. Cellular and molecular actions of juvenile hormone. II. Roles of juvenile hormone in adult insects. *Adv Insect Physiol*. 26:1–155.
- Xie X, et al. 2016. The potential role of juvenile hormone acid methyltransferase in methyl farnesoate (MF) biosynthesis in the swimming crab *Portunus trituberculatus*. *Anim Reprod Sci*. 168:40–49.
- Xu T, Caron LA, Fehon RG, Artavanis-Tsakonas S. 1992. The involvement of the Notch locus in *Drosophila* oogenesis. *Development* 115(4):913–922.
- Zaccardi ML, Traina G, Cataldo E, Brunelli M. 2004. Sensitization and dishabituation of swim induction in the leech *Hirudo medicinalis*: role of serotonin and cyclic AMP. *Behav Brain Res*. 153(2):317–326.

Associate Editor: Mandë Holford