


---

This is the **accepted version** of the article:

Mármol-Sánchez, Emilio; Quintanilla Aguado, Raquel; Jordana i Vidal, Jordi; [et al.]. «An association analysis for 14 candidate genes mapping to meat quality quantitative trait loci in a Duroc pig population reveals that the ATP1A2 genotype is highly associated with muscle electric conductivity». *Animal Genetics*, Vol. 51, Issue 1 (February 2020), p. 95-100. DOI 10.1111/age.12864

---

This version is available at <https://ddd.uab.cat/record/216804>

under the terms of the  **CC BY** license

1 **RUNNING HEAD:** Candidate gene analysis for pig meat quality

2

3 **An association analysis for 14 candidate genes mapping to meat quality QTL in a**  
4 **Duroc pig population reveals that the *ATPIA2* genotype is highly associated with**  
5 **muscle electric conductivity**

6

7 Emilio Mármol-Sánchez<sup>1</sup>, Raquel Quintanilla<sup>2</sup>, Jordi Jordana<sup>3</sup>, Marcel Amills<sup>1,3</sup>

8

9 <sup>1</sup>Department of Animal Genetics, Centre for Research in Agricultural Genomics (CRAG),  
10 CSIC-IRTA-UAB-UB, Campus de la Universitat Autònoma de Barcelona, Bellaterra,  
11 Spain.

12 <sup>2</sup>Animal Breeding and Genetics Programme, Institute for Research and Technology in Food  
13 and Agriculture (IRTA), Torre Marimon, Caldes de Montbui, Spain.

14 <sup>3</sup>Departament de Ciència Animal i dels Aliments, Universitat Autònoma de Barcelona,  
15 Bellaterra, Spain.

16

17 Address for correspondence: Marcel Amills, Department of Animal Genetics, Centre for  
18 Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB, Campus de la  
19 Universitat Autònoma de Barcelona, Bellaterra 08193, Spain. E-mail:  
20 marcel.amills@uab.cat. Tel. +34 93 563 6600

21

22

23

24

## 25 **Summary**

26

27         In previous genome-wide association studies carried out in a Duroc commercial line  
28 (Lipgen population), we detected on pig chromosomes 3, 4 and 14 several QTL for *gluteus*  
29 *medius* muscle redness (GM a\*), electric conductivity in the *longissimus dorsi* muscle (LD  
30 CE) and vaccenic acid content in the LD muscle (LD C18:1 n-7), respectively. We have  
31 genotyped, in the Lipgen population, 19 single nucleotide polymorphisms (SNP) mapping  
32 to 14 genes located within these three QTL. Subsequently, association analyses have been  
33 performed. After correction for multiple testing, two SNPs in the *TGFBRAP1*  
34 (rs321173745) and *SELENOI* (rs330820437) genes were associated with GM a\*, while  
35 *ACADSB* (rs81449951) and *GPR26* (rs343087568) genotypes displayed significant  
36 associations with LD vaccenic content. Moreover, the polymorphism of the *ATPIA2*  
37 (rs344748241), *ATP8B2* (rs81382410) and *CREB3L4* (rs321278469 and rs330133789)  
38 genes showed significant associations with LD CE. We made a second round of association  
39 analyses including the SNPs mentioned above as well as other SNPs located in the  
40 chromosomes to which they map to. After performing a correction for multiple testing, the  
41 only association that remained significant at the chromosome-wide level was that between  
42 the *ATPIA2* genotype and LD CE. From a functional point of view, this association is  
43 meaningful because this locus encodes a subunit of the Na<sup>+</sup>/K<sup>+</sup>-ATPase responsible of  
44 maintaining an electrochemical gradient across the plasma membrane.

45

46 **Keywords:** Pig, single nucleotide polymorphism, meat quality, Na<sup>+</sup>/K<sup>+</sup>-ATPase.

47

48

49 Meat quality traits are of paramount importance for the pig industry because they  
50 determine, to a great extent, consumer acceptance and financial profit. Once pigs are  
51 slaughtered, there is a decline of the pH of the skeletal muscle due to the production of  
52 lactic acid through anaerobic glycolysis (Rosenvold & Andersen 2003). The rate of muscle  
53 acidification has a strong effect on meat color and water-holding capacity. In this way, a  
54 low ultimate pH (5.4-5.3) is associated with pale, soft and exudative (PSE) meat as well as  
55 with an increased electrical conductivity (CE) and elevated drip and cooking losses (Lee *et*  
56 *al.* 2000; Rosenvold & Andersen 2003). In contrast, a high ultimate pH (6.3 or higher)  
57 results in dark, firm and dry (DFD) meat with a high water-holding capacity and a lower  
58 CE (Lee *et al.* 2000; Kim *et al.* 2016). Adverse effects on meat quality are influenced by  
59 both genetic and environmental factors. Recessive and dominant genotypes in the porcine  
60 ryanodine receptor 1 (*RYR1*) and the protein kinase AMP-activated non-catalytic subunit  $\gamma_3$   
61 (*PRKAG3*) genes, respectively, are strong predisposing factors to the occurrence of PSE  
62 meats (Fujii *et al.* 1991; Milan *et al.* 2000). On the other hand, there are multiple factors  
63 related with pig management and transportation (pre-slaughter stress), stunning method at  
64 slaughter, carcass chilling and pelvic suspension of carcasses that influence pork quality  
65 (Rosenvold & Andersen 2003). Another important parameter that determines meat quality  
66 is intramuscular fat (IMF) composition. In this regard, it is well known that fatty acid  
67 composition can have important consequences on the oxidative stability of meat during  
68 processing and retail display as well as on fat firmness (Wood *et al.* 2008).

69 In previous genome-wide association studies, we identified several genomic regions  
70 containing quantitative trait loci (QTL) for meat Minolta a\* value (redness), CE (González-  
71 Prendes *et al.* 2017) and IMF composition (González-Prendes *et al.* 2019) traits measured  
72 in the *longissimus dorsi* (LD) and *gluteus medius* (GM) muscle samples of 350 Duroc

73 barrows (Lipgen population). Details about the rearing of the pigs can be found in Gallardo  
74 *et al.* (2009), while a thorough description of QTL mapping methods is reported in  
75 González-Prendes *et al.* (2017). The measurement of CE was done 24 hours after slaughter  
76 by using a Pork Quality Meter (Intek GmbH), while Minolta a\* value was determined with  
77 a Minolta Chroma-Meter CR-200 (Konica Minolta) equipment at the same time point.  
78 Muscle fatty acid composition was measured as previously described by Quintanilla *et al.*  
79 (2011). In the current work, we have selected 14 candidate genes located within QTL  
80 regions for GM a\* on SSC3, LD CE on SSC4, and LD vaccenic content on SSC14 (**Table**  
81 **1**). These genes were: phosphorylase kinase catalytic subunit  $\gamma$  1 (*PHKG1*), transforming  
82 growth factor  $\beta$  receptor associated protein 1 (*TGFBRAP1*), selenoprotein I (*SELENOI*),  
83 hydroxyacyl-CoA dehydrogenase trifunctional multienzyme (*HADHA*), coatomer protein  
84 complex subunit  $\alpha$  (*COPA*), proliferation and apoptosis adaptor protein 15 (*PEA15*),  
85 calsequestrin 1 (*CASQ1*), ATPase Na<sup>+</sup>/K<sup>+</sup> transporting  $\alpha_2$  subunit (*ATPIA2*), ATPase  
86 phospholipid transporting 8B2 (*ATP8B2*), cAMP responsive element binding protein 3 like  
87 4 (*CREB3L4*), CREB regulated transcription coactivator 2 (*CRTC2*), acyl-CoA  
88 dehydrogenase short/branched chain (*ACADSB*), G protein-coupled receptor 26 (*GPR26*)  
89 and C-terminal binding protein 2 (*CTBP2*).

90 Genes were selected based on bibliographic information about their biological  
91 functions which suggested that they could be involved in the determinism of any of the  
92 three traits under study (GM a\*, LD CE and LD C18:1 n-7). Based on available RNA-Seq  
93 (Cardoso *et al.* 2017) and whole-genome data (our unpublished results), we called 19 SNPs  
94 mapping to these 14 genes by using the GATK Best Practices workflow for SNP calling  
95 (<https://software.broadinstitute.org/gatk/best-practices/workflow?id=11145>) in accordance  
96 with protocols reported by Mármol-Sánchez *et al.* (2019). Nineteen SNPs were finally

97 selected because the SnpEff software predicted that they might have functional effects  
98 (Cingolani *et al.* 2012), as reported in **Supplementary Table 1**. The 19 selected SNPs  
99 (**Table 1**) were genotyped at the Servei Veterinari de Genètica Molecular of the Universitat  
100 Autònoma de Barcelona (<http://sct.uab.cat/svgm/en>) by using a QuantStudio 12K Flex  
101 Real-Time PCR System (ThermoFisher Scientific). Association analyses between SNPs  
102 and phenotypes were performed with the Genome-wide Efficient Mixed-Model Association  
103 (GEMMA) software (Zhou & Stephens 2012). The following statistical model was used:

104

$$105 \mathbf{y} = \mathbf{W}\boldsymbol{\alpha} + \mathbf{x}\boldsymbol{\delta} + \mathbf{u} + \boldsymbol{\varepsilon}$$

106

107 where  $\mathbf{y}$  is the vector of phenotypic observations for every individual;  $\boldsymbol{\alpha}$  corresponds  
108 to a vector including the intercept plus the fixed effects, *i.e.* batch effect with 4 categories  
109 (all traits), and farm origin effect with 3 categories (all traits). The  $\boldsymbol{\alpha}$  vector also contains  
110 the regression coefficients of the following covariates: (1) Carcass weight at slaughterhouse  
111 for meat quality traits, and (2) IMF content in the LD muscle for LD fatty acid  
112 composition;  $\mathbf{W}$  is the incidence matrix relating phenotypes with the corresponding effects;  
113  $\mathbf{x}$  is the vector of the genotypes corresponding to the set of selected polymorphisms;  $\boldsymbol{\delta}$  is the  
114 allele substitution effect for each polymorphism;  $\mathbf{u}$  is a vector of random individual effects  
115 with a n-dimensional multivariate normal distribution  $MVN_n(0, \lambda \tau^{-1} \mathbf{K})$ , where  $\tau^{-1}$  is the  
116 variance of the residual errors,  $\lambda$  is the ratio between the two variance components and  $\mathbf{K}$  is  
117 a known relatedness matrix derived from the SNPs; and  $\boldsymbol{\varepsilon}$  is the vector of residual errors.  
118 Results were corrected for multiple testing by using the false discovery rate (FDR) method  
119 reported by Benjamini & Hochberg (1995). The correction for multiple testing took into  
120 account the number of candidate SNPs (2<sup>nd</sup> column of **Table 1**) mapping to each one of the

121 SSC3 GM a\* (5 SNPs), SSC4 LD CE (11 SNPs) and SSC14 LD (C18:1) n-7 (3 SNPs)  
122 QTL.

123 Performance of association analyses with the methodology described above  
124 revealed the existence of several associations that remained significant even after correction  
125 for multiple testing. We have found, for instance, an association between GM Minolta a\*  
126 value and missense mutations in the *TGFBRAP1* and *SELENOI* genes, which map to two  
127 different GM a\* QTL on SSC3 (**Table 1**). The inactivation of the *TGFBRAP1* gene results  
128 in the suppression of aerobic glycolysis and increased levels of mitochondrial respiration  
129 and fatty acid oxidation (Yoshida *et al.* 2013), while *SELENOI* encodes a selenoprotein  
130 fundamental for the synthesis of phosphatidylethanolamine, a molecule with important  
131 effects on the oxidation of lipid membranes, oxidative phosphorylation and mitochondrial  
132 morphology (Tasseva *et al.* 2013; Poyton *et al.* 2016). We have also detected significant  
133 associations between LD CE and SNPs in the *ATPIA2*, *ATP8B2* and *CREB3L4* genes,  
134 which map to SSC4 LD CE QTL covering two regions spanning from 85.6 to 91 Mb and  
135 from 95.2 to 97.8 Mb. These findings are very suggestive because the *ATPIA2* gene, the  
136 one showing the most significant association, is preferentially expressed in the skeletal and  
137 heart muscle and brain and it encodes the  $\alpha_2$  subunit of the ion pump  $\text{Na}^+/\text{K}^+$  ATPase  
138 (Clausen *et al.* 2017). Noteworthy,  $\text{Na}^+/\text{K}^+$ -ATPases provide the energy necessary for the  
139 maintenance of  $\text{Na}^+$  and  $\text{K}^+$  electrochemical gradients across the plasma membrane by  
140 hydrolyzing ATP (Clausen *et al.* 2017; Sampedro *et al.* 2018). These gradients are essential  
141 for the preservation of the resting membrane potential as well as for the generation of  
142 electrical impulses in the skeletal muscle and nervous system (Clausen *et al.* 2017;  
143 Sampedro *et al.* 2018). The *ATP8B2* protein is also an ATPase with flippase activity

144 towards phosphatidyl choline, a key component of phospholipid membranes with important  
145 effects on the functioning of the sarcoendoplasmic reticulum  $\text{Ca}^{2+}$ ATPase pumps (Shin &  
146 Takatsu 2018; Fajardo *et al.* 2018), while CREB3L4 is a transmembrane bZip transcription  
147 factor involved in the modulation of endoplasmic reticulum stress (Kim *et al.* 2014). Our  
148 association analysis has also revealed the existence of significant associations between the  
149 phenotypic variation of LD vaccenic (C18:1 n-7) content and SSC14 SNPs located in the  
150 *ACADSB* gene, which catalyzes the oxidation of branched-chain fatty acids (Porta *et al.*  
151 2019) and the *GPR26* gene, whose inactivation leads to hyperphagia, glucose intolerance,  
152 hyperinsulinemia, dyslipidemia and obesity in mice (Chen *et al.* 2012).

153 We have made a second round of association analyses in which the SNPs that  
154 previously showed evidence of statistical significance were compared against the whole  
155 sets of the Porcine SNP60 BeadChip SNPs co-localizing to the same chromosome  
156 (chromosome-wide analysis) *i.e.* 3,123 SNPs on SSC3, 3,899 SNPs on SSC4 and 4,203  
157 SNPs on SSC14. These 11,225 SNPs were obtained from previously published porcine  
158 SNP60 BeadChip data reported by González-Prendes *et al.* (2017). In this case, the  
159 correction for multiple testing took into account the number of SNPs mentioned above for  
160 each one of the three chromosomes under analysis, *i.e.* 3,128, 3,910 and 4,206 independent  
161 tests were taken into consideration when performing association analyses for pig  
162 chromosomes SSC3, SSC4 and SSC14. Interestingly, the rs344748241 SNP in the *ATPIA2*  
163 gene was the only one that surpassed the chromosome-wide threshold of significance (q-  
164 value < 0.05) (**Table 1, Figure 1**). Noteworthy, this SNP was not significant when we  
165 made an association analysis at the genome-wide level (data not shown). Additionally, we  
166 used the *LD* function of *gaston* R package (v1.5.5; Perdry & Dandine-Roulland 2019) to  
167 evaluate the presence of linkage disequilibrium among the SNP markers that showed



168 significant associations with LD CE after correction for multiple testing at the  
169 chromosome-wide level (**Supplementary Figure 1**). The amount of linkage disequilibrium  
170 was expressed as  $r^2$  in accordance with the definition of Hill & Robertson (1968). As  
171 shown in **Supplementary Figure 1**, we observed a high degree of linkage disequilibrium  
172 between the rs344748241 (*ATPIA2* gene) and the rs80782100 (*IGSF8* gene) markers.  
173 Noteworthy, the rs80782100 SNP, which maps to an intronic position within the  
174 immunoglobulin superfamily member 8 gene, displays the highest association with the LD  
175 CE phenotype, as described in González-Prendes *et al.* (2017).

176 As previously discussed, we consider that the *ATPIA2* gene is a strong positional  
177 and functional candidate to explain the CE QTL found on SSC4 because  $\text{Na}^+$ ,  $\text{K}^+$  ATPases  
178 are fundamental to induce an electrochemical gradient across the plasma membrane of cells  
179 (Suhail 2010), and their kinetics are modulated by the extracellular pH (Salonikidis *et al.*  
180 2000), a parameter which also displays strong effects on muscle electrical conductivity. In  
181 pigs, the *ATPIA2* gene has been sequenced (Henriksen *et al.* 2013) and its polymorphisms  
182 have been associated with fat cut percentage (Fontanesi *et al.* 2012). A next step would be  
183 to re-sequence the whole gene in Lipgen pigs with alternative genotypes (QQ vs qq) for the  
184 LD CE QTL on SSC4, to build a complete catalogue of SNPs with potential effects on  
185 protein activity and expression and to investigate their association with CE in the Lipgen  
186 population. Subsequently, functional tests should be applied to ascertain whether any of the  
187 mutations in the pig *ATPIA2* gene with highly significant  $q$ -values also have causal effects  
188 on muscle conductivity.

189

## 190 **Acknowledgments**

191

192           The authors are indebted to Selecció Batallé S.A. for providing the animal material  
193 used in this study. We gratefully acknowledge J. Reixach and J. Soler (IRTA) for their  
194 collaboration in the experimental protocols. This research was partially funded with grants  
195 AGL2013-48742-C2-1-R and AGL2013-48742-C2-2-R awarded by the Spanish Ministry  
196 of Economy and Competitiveness and grant 2014 SGR 1528 from the Agency for  
197 Management of University and Research Grants of the Generalitat de Catalunya. Emilio  
198 Mármol-Sánchez was funded with a FPU Ph.D. grant from the Spanish Ministry of  
199 Education (FPU15/01733). We also acknowledge the support of the Spanish Ministry of  
200 Economy and Competitiveness for the Center of Excellence Severo Ochoa 2016-2019  
201 (SEV-2015-0533) grant awarded to the Center for Research in Agricultural Genomics.  
202 Thanks also to the CERCA Programme of the Generalitat de Catalunya.

203

#### 204 **Data availability**

205

206 These 11,225 SNPs included in this study were obtained from published Porcine SNP60  
207 BeadChip data reported by González-Prendes *et al.* (2017), which can be accessed at the  
208 Figshare public repository ([https://figshare.com/s/ 2e636697009360986794](https://figshare.com/s/2e636697009360986794)).

209

#### 210 **Conflict of interest**

211

212 The authors declare that they have no conflict of interest.

213

214

215

216 **References**

217

218 Benjamini Y. & Hochberg Y. (1995) Controlling the false discovery rate: a practical and  
219 powerful approach to multiple testing. *Journal of the Royal Statistical Society*  
220 *Series B (Methodological)* **57**, 289-300.

221 Cardoso T.F., Cánovas A., Canela-Xandri O., González-Prendes R., Amills M. &  
222 Quintanilla R. (2017) RNA-seq based detection of differentially expressed genes in  
223 the skeletal muscle of Duroc pigs with distinct lipid profiles. *Scientific Reports* **7**,  
224 40005.

225 Chen D., Liu X., Zhang W. & Shi Y. (2012) Targeted inactivation of GPR26 leads to  
226 hyperphagia and adiposity by activating AMPK in the hypothalamus. *PLoS One* **7**,  
227 e40764.

228 Cingolani P., Platts A., Wang Ie L., Coon M., Nguyen T., Wang L., Land S.J., Lu X. &  
229 Ruden D.M. (2012) A program for annotating and predicting the effects of single  
230 nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila*  
231 *melanogaster* strain w1118; iso-2; iso-3. *Fly* **6**, 80-92.

232 Clausen M.V., Hilbers F. & Poulsen H. (2017) The structure and function of the Na,K-  
233 ATPase isoforms in health and disease. *Frontiers in Physiology* **8**, 371.

234 Fajardo V.A., Mikhaeil J.S., Leveille C.F., Tupling A.R. & LeBlanc P.J. (2018) Elevated  
235 whole muscle phosphatidylcholine: phosphatidylethanolamine ratio coincides with  
236 reduced SERCA activity in murine overloaded plantaris muscles. *Lipids in Health*  
237 *and Disease* **17**, 47.

238 Fontanesi L., Galimberti G., Calò D.G., Fronza R., Martelli P.L., Scotti E., Colombo M.,  
239 Schiavo G., Casadio R., Buttazzoni L. & Russo V. (2012) Identification and

240 association analysis of several hundred single nucleotide polymorphisms within  
241 candidate genes for back fat thickness in Italian Large White pigs using a selective  
242 genotyping approach. *Journal of Animal Science* **90**, 2450-64.

243 Fujii J., Otsu K., Zorzato F., de Leon S., Khana V.K., Weiler J.E., O' Brien P.J. &  
244 McLennan D.H. (1991) Identification of a mutation in porcine ryanodine receptor  
245 associated with malignant hyperthermia. *Science* **253**, 448-451.

246 Gallardo D., Quintanilla R., Varona L., Díaz I., Ramírez O., Pena R.N., & Amills M.  
247 (2009) Polymorphism of the pig *acetyl-coenzyme A carboxylase  $\alpha$*  gene is  
248 associated with fatty acid composition in a Duroc commercial line. *Animal Genetics*  
249 **40**, 410-419.

250 González-Prendes R., Quintanilla R., Cánovas A., Manunza A., Figueiredo Cardoso T.,  
251 Jordana J., Noguera J.L., Pena R.N. & Amills M. (2017) Joint QTL mapping and  
252 gene expression analysis identify positional candidate genes influencing pork  
253 quality traits. *Scientific Reports* **7**, 39830.

254 González-Prendes R., Quintanilla R., Mármol-Sánchez E., Pena R.N., Ballester M.,  
255 Cardoso T.F., Manunza A., Casellas J., Cánovas A., Diaz I., Noguera J.L., Castelló  
256 A., Mercade A. & Amills M. (2019) Comparing the mRNA expression profile and  
257 the genetic determinism of intramuscular fat traits in the porcine *gluteus medius* and  
258 *longissimus dorsi* muscles. *BMC Genomics* **20**, 170.

259 Henriksen C., Kjaer-Sorensen K., Einholm A.P., Madsen L.B., Momeni J., Bendixen C.,  
260 Oxvig C., Vilsen B. & Larsen K. (2013) Molecular cloning and characterization of  
261 porcine Na<sup>+</sup>/K<sup>+</sup>-ATPase isoforms  $\alpha_1$ ,  $\alpha_2$ ,  $\alpha_3$  and the ATP1A3 promoter. *PLoS One* **8**,  
262 e79127.

263

264 Hill W.G. & Robertson A. (1968) Linkage disequilibrium in finite populations. *Theoretical*  
265 *and Applied Genetics* **38**, 226-31.

266 Kim T.H., Jo S.H., Choi H., Park J.M., Kim M.Y., Nojima H., Kim J.W. & Ahn Y.H.  
267 (2014) Identification of *Creb3l4* as an essential negative regulator of adipogenesis.  
268 *Cell Death & Disease* **5**, e1527.

269 Kim T.W., Kim C.W., Yang M.R., No G.R., Kim S.W. & Kim I.S. (2016) Pork quality  
270 traits according to postmortem pH and temperature in Berkshire. *Korean Journal*  
271 *for Food Science of Animal Resources* **36**, 29-36.

272 Lee S., Norman J.M., Gunasekaran S., van Laack R.L., Kim B.C. & Kauffman R.G. (2000)  
273 Use of electrical conductivity to predict water-holding capacity in post-rigor pork.  
274 *Meat Science* **55**, 385-9.

275 Mármol-Sánchez E., Quintanilla R., Cardoso T.F., Jordana J. & Amills M. (2019)  
276 Polymorphisms of the cryptochrome 2 and mitoguardin 2 genes are associated with  
277 the variation of lipid-related traits in Duroc pigs. *Scientific Reports*. In press. doi:  
278 10.1038/s41598-019-45108-z.

279 Milan D., Jeon J.T., Looft C., Amarger V., Robic A., Thelander M., Rogel-Gaillard C.,  
280 Paul S., Iannuccelli N., Rask L., Ronne H., Lundström K., Reinsch N., Gellin J.,  
281 Kalm E., Le Roy P., Chardon P. & Andersson L. (2000) A mutation in *PRKAG3*  
282 associated with excess glycogen content in pig skeletal muscle. *Science* **288**, 1248-  
283 1251.

284 Perdry H. & Dandine-Roullard C. (2019) gaston: Genetic Data Handling (QC, GRM, LD,  
285 PCA) & Linear Mixed Models. R package version 1.5.5. [https://CRAN.R-](https://CRAN.R-project.org/package=gaston)  
286 [project.org/package=gaston](https://CRAN.R-project.org/package=gaston)

287 Porta F., Chiesa N., Martinelli D. & Spada M. (2019) Clinical, biochemical and molecular  
288 spectrum of short/branched-chain acyl-CoA dehydrogenase deficiency: two new  
289 cases and review of literature. *Journal of Pediatric Endocrinology and Metabolism*  
290 **32**, 101-108.

291 Poyton M.F., Sendeki A.M., Cong X. & Cremer P.S. (2016) Cu<sup>2+</sup> binds to  
292 phosphatidylethanolamine and increases oxidation in lipid membranes. *Journal of*  
293 *the American Chemical Society* **138**, 1584-90.

294 Quintanilla R., Pena R.N., Gallardo D., Cánovas A., Ramírez O., Díaz I., Noguera J.L. &  
295 Amills M. (2011) Porcine intramuscular fat content and composition are regulated  
296 by quantitative trait loci with muscle-specific effects. *Journal of Animal Science* **89**,  
297 2963-71.

298 Rosenvold K. & Andersen H.J. (2003) Factors of significance for pork quality-a review.  
299 *Meat Science* **64**, 219-37.

300 Salonikidis P.S., Kirichenko S.N., Tatjanenko L.V., Schwarz W. & Vasilets L.A. (2000)  
301 Extracellular pH modulates kinetics of the Na<sup>+</sup>,K<sup>+</sup>-ATPase. *Biochimica et*  
302 *Biophysica Acta (BBA) - Biomembranes* **1509**, 496-504.

303 Sampedro Castañeda M., Zanolte E., Scalco R.S., Scaramuzzi V., Marques Caldas V.,  
304 Conti Reed U., da Silva A.M.S., O'Callaghan B., Phadke R., Bugiardini E., Sud R.,  
305 McCall S., Hanna M.G., Poulsen H., Männikkö R. & Matthews E. (2018) A novel  
306 ATP1A2 mutation in a patient with hypokalaemic periodic paralysis and CNS  
307 symptoms. *Brain* **141**, 3308-3318.

308 Shin H.W. & Takatsu H. (2018) Substrates of P4-ATPases: beyond aminophospholipids  
309 (phosphatidylserine and phosphatidylethanolamine). *The FASEB Journal* **3**, 3087-  
310 3096.

311 Suhail M. (2010) Na, K-ATPase: Ubiquitous multifunctional transmembrane protein and its  
312 relevance to various pathophysiological conditions. *Journal of Clinical Medicine*  
313 *Research* **2**, 1-17.

314 Tasseva G., Bai H.D., Davidescu M., Haromy A., Michelakis E. & Vance J.E. (2013)  
315 Phosphatidylethanolamine deficiency in mammalian mitochondria impairs oxidative  
316 phosphorylation and alters mitochondrial morphology. *Journal of Biological*  
317 *Chemistry* **288**, 4158-4173.

318 Wood J.D., Enser M., Fisher A.V., Nute G.R., Sheard P.R., Richardson R.I., Hughes S.I. &  
319 Whittington F.M. (2008) Fat deposition, fatty acid composition and meat quality: A  
320 review. *Meat Science* **78**, 343-58.

321 Yoshida S., Tsutsumi S., Muhlebach G., Sourbier C., Lee M.J., Lee S., Vartholomaïou E.,  
322 Tatokoro M., Beebe K., Miyajima N., Mohny R.P., Chen Y., Hasumi H., Xu W.,  
323 Fukushima H., Nakamura K., Koga F., Kihara K., Trepel J., Picard D. & Neckers L.  
324 (2013) Molecular chaperone TRAP1 regulates a metabolic switch between  
325 mitochondrial respiration and aerobic glycolysis. *Proceedings of the National*  
326 *Academy of Sciences of the United States of America* **110**, E1604-12.

327 Zhou X. & Stephens M. (2012) Genome-wide efficient mixed-model analysis for  
328 association studies. *Nature Genetics* **44**, 821-824.

329 **Table 1.** An association analysis between 19 SNPs mapping to 14 candidate genes and meat quality traits recorded in a Duroc pig  
 330 population (significant associations are shown in bold)<sup>1</sup>.

Gene	SNP	Type	Trait	P-value	q-value	P-value*	q-value*	$\delta \pm SE$	A <sub>1</sub>	MAF
<i>PHKG1</i>	rs697732005 (3:16.830 Mb)	Splice region variant (G/A)	<b>GM a*</b>	0.88661	0.88661	0.68325	0.96577	-0.02 (0.142)	A	0.3443
<i>TGFBRAP1</i>	<b>rs321173745</b> <b>(3:49.516 Mb)</b>	<b>Missense variant</b> <b>(A/G)</b>		<b>0.00361</b>	<b>0.00902</b>	<b>0.03108</b>	0.67220	0.549 (0.186)	G	0.1875
<i>SELENOI</i>	<b>rs330820437</b> <b>(3:112.635 Mb)</b>	<b>Missense variant</b> <b>(A/G)</b>		<b>0.00039</b>	<b>0.00196</b>	<b>0.01307</b>	0.51778	0.643 (0.181)	G	0.1757
<i>HADHA</i>	rs81215086 (3:112.794 Mb)	Missense variant (G/A)		0.53993	0.67491	0.62966	0.96577	-0.102 (0.169)	A	0.2899
	rs344578723 (3:112.796 Mb)	Missense variant (G/A)		0.53466	0.67491	0.67980	0.96577	-0.104 (0.169)	A	0.2866
<i>COPA</i>	rs340853721 (4:90.163 Mb)	Splice region variant (T/C)	<b>LD CE</b>	0.90735	0.95684	0.79005	0.99942	0.014 (0.091)	T	0.4351
	rs333099339 (4:90.183 Mb)	Splice region variant (T/C)		0.87813	0.95684	0.88586	0.99942	0.017 (0.090)	T	0.4381
	rs80949931 (4:90.186 Mb)	Missense variant (A/G)		0.95684	0.95684	0.68990	0.99942	-0.002 (0.091)	A	0.4335
<i>PEA15</i>	rs329681990 (4:90.266 Mb)	Splice region variant (G/A)		0.85666	0.95684	0.58021	0.99942	-0.014 (0.091)	G	0.433
<i>CASQ1</i>	rs334946278 (4:90.280 Mb)	Splice region variant (G/A)		0.95267	0.95684	0.92240	0.99942	0.005 (0.104)	A	0.1304
<i>ATPIA2</i>	<b>rs344748241</b> <b>(4:90.356 Mb)</b>	<b>Splice region</b> <b>variant (G/A)</b>		<b>6.515E-06</b>	<b>7.167E-05</b>	<b>0.00006</b>	<b>0.02518</b>	-0.325 (0.066)	G	0.497
<i>ATP8B2</i>	<b>rs81382410</b>	<b>Splice region</b>		<b>0.00285</b>	<b>0.01565</b>	<b>0.00256</b>	0.21113	-0.233	T	0.3345



	(4:95.435 Mb)	variant (T/C)						(0.077)		
<b>CREB3L4</b>	rs329686514 (4:95.717 Mb)	Missense variant (C/T)		0.08043	0.17695	0.22592	0.97957	-0.155 (0.088)	T	0.3063
	<b>rs321278469</b> (4:95.717 Mb)	<b>Missense variant</b> (C/A)		<b>0.00639</b>	<b>0.01757</b>	<b>0.00554</b>	0.30475	-0.228 (0.083)	C	0.3084
	<b>rs330133789</b> (4:95.721 Mb)	<b>Missense variant</b> (G/A)		<b>0.00493</b>	<b>0.01757</b>	<b>0.01769</b>	0.57188	0.254 (0.075)	A	0.3373
<i>CRTC2</i>	rs330198768 (4:95.740 Mb)	Intron variant (C/T)		0.32931	0.60373	0.56310	0.99942	-0.083 (0.085)	T	0.3687
<i>ACADSB</i>	rs81449951 (14:132.588 Mb)	Missense variant (C/A)	<b>LD (C18:1) n-7</b>	<b>0.04036</b>	0.08073	<b>0.0424837</b>	0.8322423	0.093 (0.045)	A	0.2109
<b>GPR26</b>	<b>rs343087568</b> (14:133.182 Mb)	<b>Splice region</b> <b>variant (A/G)</b>		<b>0.00333</b>	<b>0.01334</b>	0.1269422	0.9956111	-0.096 (0.032)	G	0.4632
<i>CTBP2</i>	rs339956077 (14:134.334 Mb)	Splice region variant (G/A)		0.88166	0.88166	0.1269422	0.9956111	0.007 (0.046)	A	0.2094

331 <sup>1</sup> The *P*-value and the *q*-value terms define the statistical significance of the association analysis before and after correcting for  
332 multiple testing with a false discovery rate approach, respectively. The correction for multiple testing took into account the number of  
333 candidate SNPs (2<sup>nd</sup> column of Table 1) mapping to each one of the SSC3 GM a\* (5 SNPs), SSC4 CE (11 SNPs) and SSC14 LD  
334 (C18:1) n-7 (3 SNPs) QTL. The *P*-value\* and the *q*-value\* terms define the statistical significance of the chromosome-wide  
335 association analysis before and after correcting for multiple testing with a false discovery rate approach, respectively. In this case, the  
336 correction for multiple testing took into account the number of markers in the Porcine SNP60 BeadChip mapping to pig chromosomes  
337 SSC3 (3,123 SNPs), SSC4 (3,899 SNPs) and SSC14 (4,203 SNPs). Other terms that need to be defined are:  $\delta$ , estimated allele  
338 substitution effect and its standard error (SE); A<sub>1</sub>, minor allele; MAF, minor allele frequency; GM a\*, Minolta a\* value (redness) in the

339 *gluteus medius* muscle; LD CE, electric conductivity in the *longissimus dorsi* muscle; and LD (C18:1) n-7, vaccenic acid content in the  
340 *longissimus dorsi* muscle.

341 **LEGENDS TO FIGURES**

342 **Figure 1:** Manhattan plot depicting associations between electrical conductivity in the  
343 *longissimus dorsi* muscle and the genotypes of markers in the *ATPIA2* (rs344748241),  
344 *ATP8B2* (rs81382410) and *CREB3L4* (rs321278469 and rs330133789) loci plus 3,899  
345 additional SNPs mapping to pig chromosome 4 (SSC4). The positions of these three genes  
346 are SSC4: 90.292-90.371 Mb (*ATPIA2*), SSC4: 95.426-95.446 Mb (*ATP8B2*) and SSC4:  
347 95.714-95.723 Mb (*CREB3L4*). The green line represents the nominal *P-value* of  
348 significance, while the blue line indicates the *P-value* of significance after correcting for  
349 multiple testing with a false discovery rate approach (*q-value*). The rs344748241 SNP in  
350 the *ATPIA2* gene is located 23 kb away from the peak of the LD CE QTL, *i.e.*  
351 ALGA0026686 (rs80782100; 4:90.378 Mb) SNP, as reported by González-Prendes *et al.*  
352 (2017).

353

354

355

356


357

358

359

360

### SSC4



361

362

363

364

365

366

367

368 **SUPPLEMENTARY DATA**

369 **Supplementary Table 1:** Additional information about selected SNP and their potential  
370 impact and deleteriousness (SIFT).

371 **Supplementary Figure 1:** Graph depicting the magnitude of linkage disequilibrium among  
372 SNPs that showed significant associations with *longissimus dorsi* electric conductivity after  
373 correction for multiple testing at the chromosome-wide level. Here, the amount of linkage  
374 disequilibrium is expressed as  $r^2$  as defined by Will & Robertson (1968) and such  
375 parameter was calculated with the *LD* function of *gaston* R package.

376

377

378

379

380

381

382

383

384

385

386

SNP_ID	Allele	Consequence	Impact	Symbol	SIFT
rs697732005	A	splice_region_variant,intron_variant	LOW	<i>PHKG1</i>	-
rs321173745	G	missense_variant	MODERATE	<i>TGFBRAP1</i>	tolerated(0.22)
rs330820437	G	missense_variant	MODERATE	<i>SELENOI</i>	tolerated(1)
rs81215086	A	missense_variant	MODERATE	<i>HADHA</i>	deleterious(0.04)
rs344578723	A	missense_variant	MODERATE	<i>HADHA</i>	tolerated(0.81)
rs340853721	C	splice_region_variant,intron_variant	LOW	<i>COPA</i>	-
rs333099339	C	splice_region_variant,intron_variant	LOW	<i>COPA</i>	-
rs80949931	G	missense_variant	MODERATE	<i>COPA</i>	tolerated(0.15)
rs329681990	A	splice_region_variant,intron_variant	LOW	<i>PEA15</i>	-
rs334946278	A	splice_region_variant,synonymous_variant	LOW	<i>CASQ1</i>	-
rs344748241	A	splice_region_variant,synonymous_variant	LOW	<i>ATP1A2</i>	-
rs81382410	G	splice_region_variant,intron_variant	LOW	<i>ATP8B2</i>	-
rs329686514	T	missense_variant	MODERATE	<i>CREB3L4</i>	tolerated_low_confidence(0.05)
rs321278469	A	missense_variant	MODERATE	<i>CREB3L4</i>	tolerated(0.24)
rs330133789	A	missense_variant	MODERATE	<i>CREB3L4</i>	tolerated(0.32)
rs330198768	T	intron_variant	MODIFIER	<i>CRTC2</i>	-
rs81449951	A	missense_variant	MODERATE	<i>ACADSB</i>	tolerated(0.32)
rs343087568	G	splice_region_variant,intron_variant	LOW	<i>GPR26</i>	-
rs339956077	T	splice_region_variant,intron_variant	LOW	<i>CTBP2</i>	-

387

388

