
This is the **accepted version** of the article:

Hu, Minjie; Peñuelas, Josep; Sardans i Galobart, Jordi; [et al.]. «Denitrification rates in tidal marsh soils : the roles of soil texture, salinity and nitrogen enrichment». European journal of soil science, 2020. DOI 10.1111/ejss.12956

This version is available at <https://ddd.uab.cat/record/224214>

under the terms of the  **CC BY** COPYRIGHT license

1 **Denitrification rates in tidal marsh soils: The roles of soil texture, salinity, and**
2 **nitrogen enrichment**

3 **M.J. HU^{a,b}, J. PEÑUELAS^{c,d}, J. SARDANS^{c,d}, J.F. HUANG^a, K. XU^a & C. TONG^a**

4 *^aKey Laboratory of Humid Sub-tropical Eco-geographical Process of Ministry of Education, Fujian*
5 *Normal University, Fuzhou 350007, China; ^bCollege of the Environment and Ecology, Xiamen*
6 *University, Xiamen 361102, China; ^cCSIC, Global Ecology CREAM-CSIC-UAB, Bellaterra, 08193*
7 *Barcelona, Catalonia, Spain; and ^dCREAF, Cerdanyola del Vallès, 08193 Barcelona, Catalonia,*
8 *Spain*

9 **Correspondence:** C. TONG. E-mail: tongch@fjnu.edu.cn

10 **Running title:** *Denitrification rates in tidal marsh soils*

11 **Summary**

12 The denitrification rates of freshwater and oligohaline tidal marsh soils with different textures (loam and sandy
13 soils) in a subtropical estuary, and their responses to nitrogen (N) loading, were investigated. In both marshes,
14 the denitrification rates varied significantly with the season only in loam soil. The denitrification rates were
15 highest in oligohaline marsh loam soil and lowest in freshwater marsh sand soil. NH_4NO_3 addition significantly
16 increased the denitrification rates of all the marsh soils. Our findings suggest that soil texture, soil organic
17 matter (SOM) content and low-level increases in salinity all had large effects on denitrification, indicating that
18 the dynamics of denitrification rates in estuarine marshes with low-level salinity were controlled by the
19 interaction of salinity and soil texture but mainly depended on SOM content. We propose that denitrification
20 in tidal marshes plays an important role in regulating current and future N loading into estuary and inshore
21 coastal waters, especially for tidal freshwater marshes, which introduces great uncertainty to the N dynamics
22 of estuaries under global changes.

23 *Keywords: soil texture, salinity, nitrogen enrichment, SOM, denitrification, estuarine tidal marsh*

24 **Highlights:**

- 25 ➤ Denitrification rates were higher in loam than sandy soil, independent of salinity.
- 26 ➤ Denitrification rates were higher in oligohaline marsh loam than freshwater marsh loam.
- 27 ➤ Exogenous N input enhanced denitrification in estuarine marsh soil.
- 28 ➤ Loam with high-N enhanced N removal, especially in freshwater environments.

29 **Introduction**

30 Denitrification is the dominant natural pathway for N-cycling transformation in wetlands. However,
31 the factors that control this pathway are multifaceted and interactive (Day *et al.*, 2018; Neubauer *et*
32 *al.*, 2019; Zhang *et al.*, 2019). Estuarine marshes play a vital role in denitrification because they
33 consist of a combination of anaerobic environments and include carbonaceous substrates in the soil
34 (Marks *et al.*, 2016). Previous studies confirmed that denitrification in estuaries is variable and
35 potentially regulated by various factors, such as the soil texture, salinity and N loading (Lee *et al.*,
36 2017; Wang *et al.*, 2017; Hinshaw *et al.*, 2019). Owing to the influence of upstream runoff and tidal
37 saltwater intrusion, the salinity and soil texture change significantly in estuary systems, and, at the
38 same time, large amounts of exogenous N derived from fertilization are introduced into estuaries from
39 runoff, tides and deposition (Hu *et al.*, 2019; Neubauer *et al.*, 2019). However, few studies have
40 explicitly linked potential denitrification with the above factors simultaneously, and this knowledge
41 gap currently limits the understanding of the geochemical processes that drive N cycling and the
42 associated environmental responses.

43 The Min River estuary tidal marsh undergoes a clear shift from a freshwater to an oligohaline
44 environment (Tong *et al.*, 2017) and has a high N input from agricultural and industrial activities (Hu
45 *et al.*, 2019). Such an estuarine system provides an ideal environment for studying the responses of
46 soil denitrification to the roles of soil texture, salinity and N enrichment. Herein, we conducted a
47 seasonal incubation experiment to assess the interactive effects of soil texture, salinity and N loading
48 on the denitrification of marsh soils. We hypothesized (1) that the denitrification rate of freshwater
49 marshes is higher than that of oligohaline marshes, and (2) in both freshwater and oligohaline marshes,
50 the denitrification rate is higher in loamy than in sandy soils, especially in the soils with additional N.

51 **Materials and methods**

52 Soils were collected from a subtropical estuarine marsh in the Min River estuary in southeastern
53 China ([Figure S1](#)). We selected two tidal marshes: the freshwater Longxiangdao Marsh (26°1.8'52.8"
54 N, 119°18'17.8" E) and the oligohaline Shanyutan Marsh (26°1.8'13" N, 119°37'46" E), with average
55 salinities of 0.08±0.02 and 2.70±0.12 ppt, respectively. In the Longxiangdao marsh, we collected soil
56 samples with different textures from stands of the same plant species (*Cyperus malaccensis*). In the
57 Shanyutan Marsh, the loam soil samples were also collected in *C. malaccensis* stands, but the sandy
58 soil samples were only collected in *Spartina alterniflora* stands near the sea. The biomass of *C.*
59 *malaccensis* was not significantly different between the two marshes (Wang *et al.*, 2017; Luo *et al.*,
60 2019). In the Shanyutan marsh, the total above- and belowground biomasses were 4230 g m⁻² for the
61 *C. malaccensis* stand and 4620 g m⁻² for the *S. alterniflora* stand, which were nearly identical. Two
62 soils (0–15 cm; in triplicate) of contrasting textures were selected at each marsh, namely sandy
63 (coarse-textured) and loam (fine-textured) soils. The soils were sieved to 2 mm, homogenized, and
64 then divided into two subsamples for measuring the denitrification rate and physicochemical
65 properties. Tidal water was also collected simultaneously for incubation and analysis. The main
66 properties of the soils and tidal water, along with their values, are listed in [Table 1](#).

67 The denitrification potential was measured using the modified chloramphenicol-amended
68 acetylene (C₂H₂) inhibition technique (Magalhães *et al.*, 2005; Marton & Craft, 2012; Ballantine *et*
69 *al.*, 2014). Although the acetylene inhibition technique may result in an underestimated ambient
70 denitrification rate due to an inhibitory effect on nitrification (Rudolph *et al.*, 2006; McCrackin &
71 Elser, 2010; Palta *et al.*, 2016), this technique has nevertheless been employed by many previous
72 studies (Ullah & Zinati, 2006; McCrackin & Elser, 2010; Marton & Craft, 2012; Lishawa *et al.*, 2014;

73 Tomasek *et al.*, 2017) to compare denitrification rates among different sites and treatments. Briefly,
74 30 g of fresh soil was transferred to 140 mL flasks; chloramphenicol (0.3 mg) and glucose (1.2 mg)
75 were added as substrate and enzyme inhibitor, respectively. Then, 30 mL of the treatment solutions
76 were injected into each flask, in which N was added as an NH_4NO_3 solution ($0.5 \text{ mg}\cdot\text{g}^{-1}$ dry soil);
77 tidal water was added instead of the NH_4NO_3 solution in the control samples (six replicates for each
78 treatment). Each treatment sample was sealed and made anoxic by filling with pure N_2 for 5 min.
79 Then, the samples were divided into two subgroups (triplicate): with and without acetylene (10% vol:
80 vol). All samples were then incubated at four temperatures (21, 11, 17 and 28 °C representing the
81 seasonal conditions of the autumn, winter, spring and summer samples, respectively) in a dark
82 shaking incubator for 6 h. Gas samples were taken after 0.5, 1.5, 3.5 and 6 h of incubation, and the
83 N_2O concentrations were analyzed using a gas chromatograph (Shimadzu Corporation, Kyoto, Japan)
84 equipped with an electron-capture detector. The denitrification rates were calculated as the difference
85 between the N_2O produced with and without acetylene (Magalhães *et al.*, 2005) and were expressed
86 in $\mu\text{g N}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ based on their dry weight. The physicochemical properties of soils and tidal waters
87 were measured as described in our previous study (Tong *et al.*, 2017; Hu *et al.*, 2019).

88 When necessary, data (i.e., denitrification rates and environment variables) were log-
89 transformed to meet the analysis of variance (ANOVA) assumption of normality and
90 homoscedasticity. The differences in denitrification rates and environment variables between the
91 different seasons and soils were tested using a one-way ANOVA. In cases where significant fixed
92 effects were detected, pairwise comparisons among groups were conducted via Tukey's post hoc test.
93 A two-way ANOVA was used to identify the effects of soil texture and season on the denitrification
94 rates using R version 3.5.1 (R Development Core Team, 2008). Overall distributions and variations
95 in soil denitrification rates and environmental parameters among the study sites were summarized

96 using a principal components analysis (PCA) in R platform.

97 **Results and discussion**

98 The pH of the soils varied significantly but irregularly with the season (Table S1). Soil ammonium-
99 N (NH_4^+ -N) concentrations varied substantially with the season but were less variable in response to
100 salinity and texture. Soil nitrate-N (NO_3^- -N) concentrations had similar seasonal patterns among the
101 sites and were larger in the loam than the sandy soil. First, these larger nitrate concentrations should
102 drive increased nitrification in the fine-texture soils because they continue to have larger nitrate
103 concentrations despite having greater nitrate losses due to increased denitrification rates. Second, the
104 larger total C and N contents in the loam soils together with the higher soil C/N ratios (Table 1) clearly
105 indicates less mineralization of organic N, allowing more substrates for nitrification (Janssen, 1996).
106 The concentrations of dissolved organic carbon (DOC) and organic matter (OM) in the oligohaline
107 marsh with loam soil were the largest and had clear seasonal patterns (Table S1). The high DOC and
108 OM concentrations in the loam soil were assumed to be caused by the physical protection and large
109 surface area of the bigger fine fraction in the loam compared to that of the sand, which protected OM
110 against decomposition and improved adhesion (Perryman *et al.*, 2011).

111 The denitrification rates of both freshwater and oligohaline marsh soils varied significantly with
112 the season ($F=12.59$, $P<0.01$, and $F=17.23$, $P<0.01$, respectively; Figure 1; Table S2), which is
113 consistent with the results of a previous study in which optimum temperatures triggered high
114 denitrification rates by stimulating substrate availability and denitrification potential (Wang *et al.*,
115 2017). The denitrification rates were higher during the winter than during the spring and autumn,
116 probably owing to the increased NO_3^- -N availability in the winter (Table S1), which can supply more
117 N substrates to denitrifiers. The denitrification rates were significantly higher in the loam soil than in

118 the sandy soil (Figure 1). These findings may be mainly attributed to a variety of circumstances. First,
119 the lower aeration in the fine-textured loam soil, due to the high level of soil moisture and the low
120 bulk density (Table 1), favours denitrification processes by reducing redox potentials during
121 anaerobic incubation. Second, the larger concentrations of NO₃⁻-N, DOC and OM in the loam soil
122 than in the sand (Table S1) accelerate denitrifier growth and enzyme synthesis by supplying organic
123 substrates and inorganic electron acceptors (Gu *et al.*, 2013; Palta *et al.*, 2016), thereby enhancing
124 soil denitrification capacity. Soil OM provides not only electrons for NO₃⁻-N denitrification through
125 mineralization, but also decreases soil redox potential, thus resulting in high substrate availability and
126 suitable environment for the growth and activity of denitrifiers (McLain & Martens, 2006; Xu & Cai,
127 2007). Moreover, NO₃⁻-N leaching, which is necessary for microbial growth, is less in loam soil,
128 which contributes to denitrification (Lee *et al.*, 2014).

129 The denitrification rate of the loam soil was significantly higher in the oligohaline marsh than
130 the freshwater marsh ($F=52.01$, $P<0.001$; Figure 1). This finding is inconsistent with earlier studies
131 where denitrification decreased strongly as salinity increased (Osborne *et al.*, 2015; Marks *et al.*,
132 2016). This is most likely because the small variation of salinity in this estuary (0-3‰) cannot
133 substantially affect the activity of denitrifiers, which in turn also implies that the denitrifiers were
134 acclimated to the soils with low-level increases in salinity. Moreover, the concentrations of NO₃⁻-N,
135 DOC and OM in the loam soil were greater in the oligohaline than the freshwater marsh (Table S1),
136 which provides sufficient substrate for denitrifiers. A review of the literature also indicated that
137 denitrification rate patterns change with fluctuations in salinity (salinity range 0‰–30‰; Table S3),
138 suggesting that the effect of salinity is site-dependent. The overall PCA indicated that the groups of
139 samples of each soil type are clearly separated in the 2-dimension layout generated by the two main
140 axes (Figure 2). Soil denitrification rates consistently correlate positively with soil N, DOC, OM and

141 pH and negatively with bulk density. Overall, our data indicate that the variable salinity and texture,
142 and the interaction between them, have a significant effect on denitrification ($F=21.92$, 271.54 , and
143 5.27 ; $P<0.001$, <0.001 , and <0.05 , respectively).

144 The addition of NH_4NO_3 substantially changed the denitrification rates, but the impact varied
145 with the soil texture and salinity (Figure 3). Specifically, NH_4NO_3 addition significantly increased the
146 denitrification rate relative to the control in the sandy soil ($F=9.58$, $P<0.05$) and loam soil ($F=7.33$,
147 $P<0.05$) of the freshwater marsh, and in the loam soil ($F=1.99$, $P<0.01$) of the oligohaline marsh by
148 212, 102, and 125%, respectively. Nitrogen is often the limiting nutrient in this estuary system (Wang
149 *et al.*, 2014); therefore, NH_4NO_3 addition provided abundant NO_3^- -N as the substrate for direct
150 denitrification (Figure S2b) and also contributed to nitrification by increasing NH_4^+ -N availability
151 (Figure S2a), which can lead to high rates of coupled nitrification-denitrification (Gu *et al.*, 2013).

152 Our findings initially indicated that soil texture might be a critical factor controlling N cycling
153 in wetland systems, regardless of salinity. They suggested that fine-textured soils with low porosity
154 and large contents of NO_3^- -N and OM would fuel the denitrification process and would be
155 accompanied by the removal of N. Further, different soil textures in the estuary system are formed at
156 different sedimentation times due to sea-land interactions (Wallace *et al.*, 2005). The longer the
157 sedimentation time, the finer the soil texture, and the associated increases in the content of substrates
158 such as OM eventually leads to increases in denitrification capacity. Our data also clearly
159 demonstrated that an increased N loading could potentially promote the denitrification rates of soils
160 with contrasting textures and salinities. These results indicate that denitrification could be an
161 important pathway to regulate current and future N loading into estuarine and inshore coastal waters,
162 which reduces the negative effects of exogenous N on water eutrophication and acidification but, on

163 the other hand, may increase the risk of global warming. Further, we conclude that tidal marshes with
164 fine-textured loam soil had relatively higher N removal potential, especially for high N-enrichment
165 environments. This partly supports our hypothesis that much uncertainty is introduced in the N
166 dynamics of estuaries under longer-term, climate change-mediated sea level rise and N deposition.
167 Thus, the detailed mechanisms and processes that control estuarine denitrification and, subsequently,
168 N cycling need further consideration.

169 **Acknowledgments**

170 This research was jointly funded by the National Science Foundation of China (41877335, 41801062),
171 the Key Science Foundation of Fujian Province (2019J02008), China Postdoctoral Science
172 Foundation (2018M630731, 2019T120556), European Research Council Synergy grant (ERC-SyG-
173 2013-610028 IMBALANCE-P), Spanish Government grant (CGL2016-79835-P), and Catalan
174 Government grant (SGR 2017-1005).

175 **Data Availability Statement**

176 The data that support the findings of this study are available from the corresponding author upon
177 reasonable request.

178 **Conflict of interest**

179 The authors declare that they have no conflicts of interest in this work.

180 **References**

- 181 Ballantine, K.A., Groffman, P.M., Lehmann, J. & Schneider, R.L. 2014. Stimulating nitrate removal processes
182 of restored wetlands. *Environmental Science and Technology*, **48**, 7365-7373.
- 183 Day, J.W., DeLaune, R.D., White, J.R., Lane, R.R., Hunter, R.G. & Shaffer, G.P. 2018. Can denitrification
184 explain coastal wetland loss: a review of case studies in the Mississippi Delta and New England. *Estuarine,
185 Coastal and Shelf Science*, **213**, 294-304.
- 186 Gu, J., Nicoullaud, B., Rochette, P., Gossel, A., Hénault, C., Cellier, P. & Richard, G. 2013. A regional
187 experiment suggests that soil texture is a major control of N₂O emissions from tile-drained winter wheat
188 fields during the fertilization period. *Soil Biology and Biochemistry*, **60**, 134-141.
- 189 Hinshaw, S.E., Zhang, T., Harrison, J.A. & Dahlgren, R.A. 2019. Excess N₂ and denitrification in hyporheic
190 porewaters and groundwaters of the San Joaquin River, California. *Water Research*, 115161.
- 191 Hu, M., Peñuelas, J., Sardans, J., Huang, J., Li, D. & Tong, C. 2019. Effects of nitrogen loading on emission
192 of carbon gases from estuarine tidal marshes with varying salinity. *Science of the Total Environment*, **667**,
193 648-657.
- 194 Janssen, B.H. 1996. Nitrogen mineralization in relation to C:N ratio and decomposability of organic materials.
195 *Plant and Soil*, **181**, 39-45.
- 196 Lee, M.J., Hwang, S.I. & Ro, H.M. 2014. Interpreting the effect of soil texture on transport and removal of
197 nitrate-N in saline coastal tidal flats under steady-state flow condition. *Continental Shelf Research*, **84**,
198 35-42.
- 199 Lee, P.O., Cherry, J.A. & Edmonds, J.W. 2017. Organic nitrogen runoff in coastal marshes: Effects on
200 ecosystem denitrification. *Estuaries and Coasts*, **40**, 437-446.
- 201 Lishawa, S.C., Jankowski, K., Geddes, P., Larkin, D.J., Monks, A.M. & Tuchman, N.C. 2014. Denitrification

- 202 in a Laurentian Great Lakes coastal wetland invaded by hybrid cattail (*Typha×glauca*). *Aquatic sciences*,
203 **76**, 483-495.
- 204 Luo, M., Zhu, W., JiaFang, H., Liu, Y., Duan, X., Wu, J. & Tong, C. 2019. Anaerobic organic carbon
205 mineralization in tidal wetlands along a low-level salinity gradient of a subtropical estuary: Rates,
206 pathways, and controls. *Geoderma*, **337**, 1245-1257.
- 207 Magalhães, C.M., Joye, S.B., Moreira, R.M., Wiebe, W.J. & Bordalo, A.A. 2005. Effect of salinity and
208 inorganic nitrogen concentrations on nitrification and denitrification rates in intertidal sediments and
209 rocky biofilms of the Douro River estuary, Portugal. *Water Research*, **39**, 1783-1794.
- 210 Marks, B.M., Chambers, L. & White, J.R. 2016. Effect of fluctuating salinity on potential denitrification in
211 coastal wetland soil and sediments. *Soil Science Society of America Journal*, **80**, 516-526.
- 212 Marton, J.M. & Craft, C.B. 2012. Effects of salinity on denitrification and greenhouse gas production from
213 laboratory-incubated tidal forest soils. *Wetlands*, **32**, 347-357.
- 214 McCrackin, M.L. & Elser, J.J. 2010. Atmospheric nitrogen deposition influences denitrification and nitrous
215 oxide production in lakes. *Ecology*, **91**, 528-539.
- 216 McLain, J.E.T. & Martens, D.A. 2006. N₂O production by heterotrophic N transformations in a semiarid soil.
217 *Applied Soil Ecology*, **32**, 253-263.
- 218 Neubauer, S.C., Piehler, M.F., Smyth, A.R. & Franklin, R.B. 2019. Saltwater intrusion modifies microbial
219 community structure and decreases denitrification in tidal freshwater marshes. *Ecosystems*, **22**, 912-928.
- 220 Osborne, R.I., Bernot, M.J. & Findlay, S.E. 2015. Changes in nitrogen cycling processes along a salinity
221 gradient in tidal wetlands of the Hudson River, New York, USA. *Wetlands*, **35**, 323-334.
- 222 Palta, M.M., Ehrenfeld, J.G., Giménez, D., Groffman, P.M. & Subroy, V. 2016. Soil texture and water retention
223 as spatial predictors of denitrification in urban wetlands. *Soil Biology and Biochemistry*, **101**, 237-250.
- 224 Perryman, S.E., Rees, G.N., Walsh, C.J. & Grace, M.R. 2011. Urban stormwater runoff drives denitrifying

- 225 community composition through changes in sediment texture and carbon content. *Microbial Ecology*, **61**,
226 932-940.
- 227 R Development Core Team. 2008. R: A language and environment for statistical computing. R Foundation for
228 Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.
- 229 Rudolph, J., Frenzel, P. & Pfennig, N. 2006. Acetylene inhibition technique underestimates in situ
230 denitrification rates in intact cores of freshwater sediment. *Fems Microbiology Letters*, **85**, 101-106.
- 231 Tomasek, A., Kozarek, J.L., Hondzo, M., Lurndahl, N., Sadowsky, M.J., Wang, P. & Staley, C. 2017.
232 Environmental drivers of denitrification rates and denitrifying gene abundances in channels and riparian
233 areas. *Water Resources Research*, **53**, 6523-6538.
- 234 Tong, C., Cadilloquiroz, H., Zeng, Z., She, C., Yang, P. & Huang, J. 2017. Changes of community structure
235 and abundance of methanogens in soils along a freshwater-brackish water gradient in subtropical estuarine
236 marshes. *Geoderma*, **299**, 101-110.
- 237 Ullah, S. & Zinati, G.M. 2006. Denitrification and nitrous oxide emissions from riparian forests soils exposed
238 to prolonged nitrogen runoff. *Biogeochemistry*, **81**, 253-267.
- 239 Wallace, K.J., Callaway, J.C. & Zedler, J.B. 2005. Evolution of tidal creek networks in a high sedimentation
240 environment: A 5-year experiment at Tijuana Estuary, California. *Estuaries and Coasts*, **28**, 795-811.
- 241 Wang, W.Q., Sardans, J., Zeng, C.S., Zhong, C., Li, Y. & Peñuelas, J. 2014. Responses of soil nutrient
242 concentrations and stoichiometry to different human land uses in a subtropical tidal wetland. *Geoderma*,
243 **232-234**, 459-470.
- 244 Wang, X.M., Hu, M.J., Ren, H.C., Li, J.B., Tong, C. & Musenze, R.S. 2017. Seasonal variations of nitrous
245 oxide fluxes and soil denitrification rates in subtropical freshwater and brackish tidal marshes of the Min
246 River estuary. *Science of the Total Environment*, **616**, 1404-1413.
- 247 Xu, Y.B. & Cai, Z.C. 2007. Denitrification characteristics of subtropical soils in China affected by soil parent

248 material and land use. *European Journal of Soil Science*, **58**, 1293-1303.

249 Zhang, Y., Ji, G., Wang, C., Zhang, X. & Xu, M. 2019. Importance of denitrification driven by the relative

250 abundances of microbial communities in coastal wetlands. *Environmental Pollution*, **244**, 47-54.

251

Figure captions

Figure 1 Changes in potential denitrification rates from different soil textures and salinities based on dry weights. Different lowercase letters on the bars indicate significant differences between seasons ($P < 0.05$); the absence of letters indicates an absence of significant differences; the asterisks (*) indicate significant differences between the soil types ($P < 0.05$). FS and FL represent sand and loam soil in the freshwater marsh, respectively; OS and OL represent sandy and loam soil in the oligohaline marsh, respectively.

Figure 2 Principal component analysis of soil denitrification rates and main environmental parameters in different soil texture and salinity conditions. The means of PC scores of each one of the four soil types are depicted by small circles and the corresponding area around these circles depicts confidence intervals at 95%. BD: bulk density; EC: electrical conductivity; NH_4^+ -N: ammonium nitrogen; NO_3^- -N: nitrate nitrogen; DOC: dissolved organic carbon; SM: soil moisture; OM: organic matter. TC: total carbon; TN: total nitrogen. FS and FL represent sandy and loam soil in the freshwater marsh, respectively; OS and OL represent sandy and loam soil in the oligohaline marsh, respectively.

Figure 3 Responses of the potential denitrification rate to nitrogen addition in soils with different textures and salinities. Different letters on the bars indicate significant differences between soils ($P < 0.05$). Asterisks indicate significant differences between the control and N treatments (* $P < 0.05$ and ** $P < 0.01$). C: control; N: NH_4NO_3 addition. FS and FL represent sand and loam soil in the freshwater marsh, respectively; OS and OL represent sandy and loam soil in the oligohaline marsh, respectively.

Figure 1




Figure 2




Figure 3

