

**FIRST XIPHOSURAN TRACEWAY IN THE MIDDLE MUSCHELKALK
FACIES (MIDDLE TRIASSIC) OF THE CATALAN BASIN (NE IBERIAN
PENINSULA)**

**PRIMER RASTRO DE XIFOSURO EN LAS FACIES MUSCHELKALK MEDIO
(TRIÁSICO MEDIO) DE LA CUENCA CATALANA (NE DE LA PENÍNSULA
IBÉRICA)**

Chabier De Jaime-Soguero^{1*}, Eudald Muijal^{1,2} & Josep Fortuny¹

¹ Institut Català de Paleontologia Miquel Crusafont, ICTA-ICP Building, c/de les
columnes, s/n, E-08193 Cerdanyola del Vallès, Catalonia, Spain;
chabierdejaim@gmail.com; josep.fortuny@icp.cat

² Staatliches Museum für Naturkunde Stuttgart, Rosenstein 1, D-70191 Stuttgart,
Germany; eudald.muijalgrane@smns-bw.de

* Corresponding author

Abstract

In the last decade, the first ichnoassemblages from the middle Muschelkalk facies (upper Anisian–middle Ladinian) of the Catalan Basin (NE Iberian Peninsula) have been discovered. Herein, the first xiphosuran trace fossils are described from the locality of Penya Rubí, a newly discovered ichnosite from the Catalan Basin. The finding opens a window into peri-Tethys ecosystems with coastal influence. The traceway is referred to the ichnogenus *Kouphichnium*: a locomotion trace attributed to xiphosurans. The traceway preserves telson grooves and different imprint morphologies from the various

appendages. The traceway pattern and arrangement of the different traces suggest a crawling locomotion style. The sedimentology suggests a coastal zone with areas influenced by tides (intertidal flat). The morphological variations of the ichnites are correlated to substrate rheology, the locomotion of the tracemaker and environmental conditions.

Keywords

Invertebrate ichnology, *Kouphichnium*, telson, Middle Triassic, coastal palaeoenvironment

Resumen

En la última década, se han descubierto las primeras icnoasociaciones de las facies Muschelkalk medio (Anisiense superior–Ladiniense medio) de la Cuenca Catalana (NE de la Península Ibérica). Aquí se describe el primer rastro fósil de xifosuro de la localidad de Peña Rubí, un icnoyacimiento descubierto recientemente en la Cuenca Catalana. El hallazgo abre una ventana dentro de los ecosistemas del peri-Tethys con influencia costera. El rastro ha sido identificado como *Kouphichnium*: una traza de locomoción asociada a xifosuros fósiles. El rastro preserva surcos del telson e impresiones de los distintos apéndices, que resultan en morfologías diferentes. El patrón del rastro y la disposición de las distintas trazas sugiere un estilo de locomoción de reptación (“crawling”). La sedimentología sugiere una zona costera, con áreas de influencia mareal (intermareal). Las variaciones morfológicas de las icnitas están correlacionadas con la reología del sustrato, la locomoción del productor de las huellas y las condiciones ambientales originales.

Palabras clave

Ichthyología de invertebrados, *Kouphichnium*, telson, Triásico Medio, paleoambiente costero

1. INTRODUCTION

The order Xiphosura Latreille, 1802, generally known as horseshoe crabs, consists of aquatic chelicerate arthropods with a fossil record extending from the Ordovician to the present (Rudkin *et al.*, 2008; Lamsdell, 2016). Although xiphosurans have inhabited in many different environments through most of the Phanerozoic, and the group has survived several mass extinction events, currently only four marine species of the family Limulidae survive (Gaillard, 2011). Because, the morphology of the group remained largely unchanged through much of its evolution, the extant species are considered as “living fossils” (Gaillard, 2011; Kin & Błazejowski, 2014). The xiphosurans show a maximum diversity during the Carboniferous and Triassic periods (Moore *et al.*, 2007). Xiphosuran trace fossils provide additional information about the palaeobiology of these animals and their habitats (Gaillard, 2011; Fernández & Pazos, 2013; Mujal *et al.*, 2018a; Bicknell *et al.*, 2019; Shibata & Varricchio, 2020).

The Triassic fossil record of Xiphosura is particularly interesting as it helps to elucidate how this group survived after the end-Permian mass extinction (Błazejowski *et al.*, 2017; Mujal *et al.*, 2018a). Even more important, in general terms, after the end-Permian mass extinction, the diversity of life shows a general diminishment, with a very poor terrestrial record, particularly in central Pangea (Benton & Newell, 2014). Biodiversity recovery in marine ecosystems was faster than in terrestrial ecosystems due

to a high increase of marine productivity after the mass extinction event (Shen *et al.*, 2015).

Considering on these concepts, the discovery of a new fossil traceway referred to Xiphosura from the north-eastern Iberian Peninsula (Western Tethys) is of special interest, because it was recovered from the middle Muschelkalk facies of the Catalan Basin. Triassic facies are diachronic at European level due to the different influence of sea level (i.e., different transgressions and regressions) among the Western Tethys basins (Virgili *et al.*, 1977; Escudero-Mozo *et al.*, 2015; Franz *et al.*, 2015). In the particular case of the Catalan Basin, a regression episode occurred during the late Anisian–middle Ladinian affected coastal environments, creating numerous coastal river floodplains (Solé de Porta *et al.*, 1987; Calvet & Marzo, 1994; Ortí *et al.*, 2018). This regression is represented in the sedimentary record as red beds and evaporite deposits between the marine Muschelkalk facies (Calvet & Marzo, 1994; Ortí *et al.*, 2018).

In the NE Iberian Peninsula few arthropod remains have been recovered (both direct and indirect evidences). Three taxa from the Muschelkalk facies (Ladinian, upper Middle Triassic) have been described in the Catalan Basin to date (Vía-Boada & Villata, 1966; Romero & Vía-Boada, 1977; Vía, 1987). Also, several xiphosuran traceways have been described from one locality of the terrestrial lower Buntsandstein Pyrenean Basin dated from the Lower Triassic (upper Olenekian) (Mujal *et al.*, 2018a).

The present study analyses the first record of a xiphosuran traceway from the middle Muschelkalk of the Iberian Peninsula. From this unit of the Catalan Basin different ichnoassociations of tetrapod trace fossils have been reported so far (Mujal *et al.*, 2015, 2018b). Thus, the newly discovered traceway adds relevant data on the palaeodiversity of these ecosystems. The main objective of this study is the ichnotaxonomy of the traceway recovered from the ichnosite of Penya Rubí (Vallirana, Barcelona province).

Further, this traceway provides information about the palaeoenvironment and palaeogeography of its tracemaker.

2. GEOLOGICAL SETTING

The outcrop of Peña Rubí is located in the Catalan Basin, currently forming the Catalan Coastal Ranges (Fig. 1). The Catalan Basin is situated at the northeast of the Iberian Peninsula and began to develop during the middle to late Permian due to the extension of the Iberian plate (Galán-Abellán *et al.*, 2013). The Triassic sedimentary succession of the Catalan Basin is composed of the so-called Germanic facies, recognised at European scale, which comprise the following lithostratigraphic units: Buntsandstein, lower (M1), middle (M2) and upper (M3) Muschelkalk, Keuper and Imón Formation (Calvet & Marzo, 1994). The Triassic succession presents a thickness of 500–800 meters (Calvet *et al.*, 1990).

During the Middle Triassic the sedimentary environmental conditions changed drastically in different areas of the Iberian massif, but especially affecting the Catalan Basin. Such changes are reflected in the three different Muschelkalk facies successions within the Catalan Coastal Ranges: They represent marine environments (M1 and M3) separated by the coastal and alluvial environments (M2) (Calvet & Marzo, 1994; Ortí *et al.*, 2018). Following these events, during the Early Triassic, remains of braided and meandering river settings under arid and semiarid climatic conditions are represented by the Buntsandstein facies (Galán-Abellán *et al.*, 2013; Mujal *et al.*, 2017). During the early Anisian (Middle Triassic) there were abundant fluvial systems that gradually change into coastal and marine settings (lower Muschelkalk facies – M1), but later emerging again during the late Anisian–middle Ladinian as a result of a short regression episode, as

represented by the middle Muschelkalk (M2) facies; afterwards, a major transgression adjusted coastal and marine settings (upper Muschelkalk facies – M3) (Calvet & Marzo, 1994; Ortí *et al.*, 2018). This is the reason why the continental environments gain presence in the Catalan Basin (but not in other Iberian basins), and the upper Anisian and lower Ladinian are characterised by the predominance of distal alluvial, coastal and shallow marine successions (Escudero-Mozo *et al.*, 2015; Ortí *et al.*, 2018). Herein, the studied outcrop corresponds to the middle Muschelkalk facies, dated as late Anisian–middle Ladinian based on palynological and biostratigraphic analysis (Solé de Porta *et al.*, 1987; Márquez-Aliaga *et al.*, 2000; Mujal *et al.*, 2015, 2018b).

3. MATERIAL AND METHODS

An *in-situ* slab with a xiphosuran traceway was discovered in the Penya Rubí ichnosite (Vallirana, Catalan Basin). The slab was recovered from a reddish sandstone stratum (50–60 cm thick). The slab dimensions are: 50.8 cm long, 29.45 cm wide and 4.13 cm thick. The ichnological study was carried out in the lab. The slab is stored at the Institut Català de Paleontologia Miquel Crusafont (Sabadell, Catalonia, Spain) and labelled as IPS110272.

A stratigraphic log was performed using a Jacob staff and meter stick. The slab was located in its exact stratigraphic level. After measuring, the data were translated into a stratigraphic sequence using the software Inkscape (v. 0.92.3).

Once in the lab, the traces were measured by caliper and with digital measuring tools from the image manipulating software GIMP (v. 2.10.0). Also, light was shone from different directions to appreciate all the details of the impressions. The traceway was digitised photogrammetrically (see Supplementary Information). This method consists of

creating a three dimensional model using photographs. The photographs were obtained with a digital reflex camera Panasonic Lumix DMC-FZ200 following the procedures of Falkingham (2012) and Mallison & Wings (2014). The three dimensional models was processed with three programs, following the procedures of Mujal *et al.* (2020, see references therein for a review): Agisoft Photoscan (standard version 1.1.4), to generate a dense point cloud, MeshLab (v. 2016.12) to edit the mesh (cleaning, scale and orientation) and ParaView (v. 5.5.0) to generate a coloured depth map and contours. The final three dimensional meshes were used to complement the morphological study of each trace and their deepness. These meshes, refined and improved following some procedures (Marcé-Nogué *et al.*, 2015), can be also used in biomechanical studies to evaluate tracemaker locomotion.

4. OUTCROP OF PENYA RUBÍ (VALLIRANA, CATALAN BASIN)

The Triassic outcrops in the studied area include Buntsandstein, Muschelkalk (M1, M2, M3) and Keuper facies, and the recovered traceway is located in the M2 unit (Figs. 2, 3). The middle Muschelkalk facies are composed of clays and sandstones interbedded with evaporite rocks (Ramón & Calvet, 1987; Ortí *et al.*, 2018). The sandstones are fine-grained, with a clayey matrix and limestone cement. Sandstone grains are, in decreasing order, composed of quartz, feldspar and mica. The mudstones are mainly fine detrital sediments and appear alternated with sandstone strata (Calvet & Marzo, 1994; Ortí *et al.*, 2018).

In the Penya Rubí ichnosite a repichnial traceway of a xiphosuran has been recognised (Figs. 3, 4). These imprints are preserved in a reddish sandstone stratum (50–60 cm), near to the upper limit between sandstones and a stratified mudstone interval

(Figs. 2, 3). The trace-bearing sandstone presents parallel stratification in its lower part and wave ripples in the topmost part, including the surface where the traceway is impressed. In the overlying strata, in a greyish interval of dolostones and mudstones, tetrapod footprints appear. They are currently under study, preliminarily referred to *Rhynchosauroides*, *Procolophonichnium* and *Rotodactylus*. This interval presents a millimetric stratification with alternating thick (greyish) and thin (reddish) laminae building up centimetric sequences. The thick levels are composed of grains coarser than those of the thin levels, and from the bottom to the top, the grain size decreases (fining-upwards sequences).

The overlying interval displays similar stratification than the previous one but it presents reddish colouration; on top of it the first medium-grained sandstone horizons are present. These sandstone beds contain *Chirotherium* footprints, currently under study, and desiccation cracks at their base. In many of the strata ripples are present. In the finely laminated sequences climbing ripples appear, whereas the coarser and thicker sandstones are characterised by unidirectional ripples.

As a whole, the succession can be interpreted as a coastal area, with bidirectional flows and a certain cyclic sedimentation, as indicated by successive layers of fine grained-sandstones with ripples indicating opposite palaeoflows. Most probably, the succession was deposited in an intertidal area with relatively low energy, as already stated by Ortí *et al.* (2018) (see also references therein).

5. SYSTEMATIC PALAEOICHOLOGY

5.1. Ichnogenus *Kouphichnium*

Referred material. The ichnites belong to a relatively regular repetition of symmetrical imprints (IPS110272). This set of imprints is considered to be a traceway preserved in concave epirelief (Fig. 4a-e) (see three-dimensional model in Supplementary Information).

Description. One straight and symmetric traceway, reaching up to 240 mm long and 26.8 mm wide. The traceway is composed of five nearly straight parallel rows of ichnites (Fig. 4a-4b), which show opposite symmetry. The central row consists of six rectilinear grooves (mean length: 20.6 mm). These linear traces are the deepest imprints and present a profound region in the middle of the groove. On both sides, with a symmetric display, numerous smaller prints close to the straight grooves are arranged in two rows in each side. These rows tend to interchange up to four types of imprints (Fig. 4c-4d). The first type shows a bifid shape (mean length: 3.3 mm), composed of a central point from which two curved scratches emerge. The second type is consist of linear scratches, much shorter than the central grooves (mean length: 4.2 mm). The shape of these imprints ranges from straight to curved lines. Despite their linear shape, the extreme anterior of the imprint displays a deep and punctuated impression. The third type of imprints shows pointed ichnites (mean length: 2.0 mm). Finally, at the edges of the traceway arrow-shaped ichnites make up the fourth type (mean length: 5.4 mm). This last imprint type shows the direction of the traceway, starting in the middle of the slab and finishing at one of the ends, i.e., the traceway advances towards the bifid and arrowed end of the traces.

The position of each type of imprint is not completely regular along the rows, but it seems to follow a trend. In both lateral rows, the pointed and bifid imprints tend to be in the inner part of the row, while the lineal and arrow-shaped frequently appear in the outer side. The arrow shaped and linear traces tend to appear in the posterior part of the

telson trace while the bifid and pincer-like imprints are located in the medium part, and the pointed imprints tend to be in the anterior part, near to the telson trace; the inner traces are relatively aligned, outlining a diagonal approaching the central linear groove from the posterior to the anterior. Also, in the upper part of the anterior side is also frequent to find a bifid imprint. The long linear traces are always the central row (Fig. 4a-4e).

Discussion. The morphological characters of the imprints reveal that the traceway corresponds to locomotion traces (repichnia) of an arthropod. The presence of a “tail trace”, corresponding to the central linear grooves, and the disposition and morphology of the imprints in each side allow to assign the traceway to *Kouphichnium*. This ichnogenus is interpreted as a repichnion, likely produced by xiphosurans (Caster, 1938; Goldring & Seilacher, 1971; Seilacher, 2007; Fernández & Pazos, 2013; Rindsberg & Martin, 2015; Mujal *et al.*, 2018a; Shu *et al.*, 2018; Shi *et al.*, 2019). The body of the xiphosurans is composed of three main parts: the prosoma (anterior part), the abdomen (central part) and a relatively long telson (posterior part) (Gaillard, 2011) (Fig. 4e). In the abdomen, xiphosurans present six pairs of appendages that contribute to various functions. The first pair of appendages are the chelicere, which end with pincers, being useful for grasping food. The second to fifth pair of appendages are chelate walking legs, all of which display bifid ends (Shi *et al.*, 2019; Shibata & Varricchio, 2020). The principal function of these appendages is for locomotion. Finally, the sixth pair is known as the pusher legs. They are the strongest appendages and push the animal forward over (commonly soft) substrates (Caster, 1938). This strong heteropody is the cause of the large variability of the resulting traceway (Caster, 1938; Goldring & Seilacher, 1971; Rindsberg & Martin, 2015). The anterior five pairs consist in small pincer-like chelipeds, leaving characteristic V-shaped impressions seen in xiphosuran traceways (Shi *et al.*,

2019). On the other hand, the “pusher” legs produce many variable imprints with digitate and flabellar forms (Gaillard, 2011).

As indicated in the description, the position of each type of imprint tends to be regular. The imprints show a descending line from the posterior part of the central row to the anterior part, being the linear and arrow shaped imprints the furthest ones from the telson at the anterior part, whereas the bifid, pincer-like and pointed imprints are the closest at the posterior part. This trend has been represented in an ideal reconstruction of the traceway (Fig. 4e).

Comparing the anatomy of living limulids and the different types of imprints, we can establish a correlation between trace and tracemaker. The central row of straight grooves would be produced by the telson. According to Caster (1938), nowadays xiphosurans would use the telson to raise the posterior part of the prosoma when the rest of the appendix is buried. This behaviour would describe the telson as a key appendix in the locomotion of the xiphosurans. Similarly, Gaillard (2011) suggested such traceways with discontinuous telson impressions as crawling behaviour. Also, according to Caster (1938) and Goldring and Seilacher (1971), due to the shallow drag of the telson, its impression is only preserved in the superficial laminae of the substrate, hence these imprints usually do not appear as “undertraces” (see also Seilacher, 2007). Otherwise, the two sets of imprints from each side (Fig. 4a, 4b) of the traceway most likely correspond to the walking appendages. The multiple variation of these traces, herein grouped in four types, would represent the locomotion stages of the appendages (Shi *et al.*, 2019).

6. DISCUSSION

6.1. Palaeoecology and palaeoenvironment

The trace fossils were found in a laminar interval composed by reddish sandstones and mudstones (Fig. 3a-b). The presence of sandstone layers with ripples showing opposite palaeoflows, the finely laminated mudstones alternated with sandstones, as well as the presence of climbing ripples, give support to the interpretation of the Penya Rubí outcrop as an intertidal setting, as suggested in other middle Muschelkalk successions (Ortí *et al.*, 2018) and other areas elsewhere (Davis & Dalrymple, 2012). As explained in the previous section, due to the shallow drag of the telson in xiphosurans, these imprints most commonly do not appear in “undertraces”, thus the traceway is most probably preserved in the originally trampled surface.

In addition, the ethology and ecology of extant limulids provides an excellent model for determining the behavioural characteristics and palaeoecology of the ancient xiphosurans (Martin & Rindsberg, 2007; Seilacher, 2007). Limulids can live in a wide range of environments from freshwater to marginal marine settings (Lamsdell, 2016; Shi *et al.*, 2019). But during their life, limulids inhabit different environmental niches at different ontogenetic stages. Mature individuals mostly live in distal areas of the continental shelf and return to the shore and tidal areas to mate and to lay their eggs in the breeding season (Caster, 1938, 1944; Shuster, 1982; Shi *et al.*, 2019). Juveniles are benthic and live in coastal areas for two years before moving into deeper waters as they grow older (Shi *et al.*, 2019). Thus, if it is assumed that the herein studied xiphosuran traceway is the result of similar behaviour, the tracemaker could be a juvenile that lived in the intertidal area or mature specimen that returned to the coast for the breeding season. According to Martin & Rindsberg (2007) on the studies of recent juvenile limulid traceways, likely producing traces assignable to *Nereites*, we favour the latter interpretation.

6.2. Palaeogeographic implications

In recent years, the fossil record from the Muschelkalk facies of the Catalan Basin and other Iberian basins has been enhanced, particularly due to the discovery of tetrapod footprints (Gand *et al.*, 2010; Fortuny *et al.*, 2011; Mujal *et al.*, 2015, 2018b; Berrocal-Casero *et al.*, 2018). The Peña Rubí ichnosite, where the studied traceway was recovered, is not an exception and several tetrapod footprints have been also recovered here. Otherwise, xiphosuran remains have been historically restricted to the upper Muschelkalk (M3) of Alcover (Catalan Basin, Tarragona province) outcrops (Vía-Boada *et al.*, 1977 and references therein). More recently, xiphosuran trace fossils were also discovered in the lower part of the Buntsandstein facies, in non-marine fluvial deposits from the Pyrenean Basin, also stratigraphically and geographically close to tetrapod footprints (Mujal *et al.*, 2017, 2018a).

In the Catalan Basin remains of three limulid taxa have been recovered from the upper Muschelkalk: *Tachypleus gadeai* (Vía-Boada & Villalta, 1966) (first described as *Heterolimulus gadeai*), *Tarracolimulus rieki* Romero & Vía-Boada, 1977 and *Mesolimulus crespelli* Vía, 1987. These species have been described from the quarry of Alcover-Montral (Tarragona, Spain), in Ladinian dolomites. The outcrops of Alcover-Montral (Vía, 1987) make up the first area where Triassic Iberian xiphosuran remains were discovered. Nevertheless, these specimens are slightly younger than the herein reported ichnites.

Outside the Iberian Peninsula, other xiphosuran remains have been recovered in Middle Triassic deposits from peri-Tethys basins. At the German locality of Bernburg, in the Central European Basin, limulid traceways related to shallow marine and intertidal successions have been documented (Diedrich, 2011). A new species of limulid from the

Anisian–lower Ladinian of Tunisia has also been described (Błażejowski *et al.*, 2017). This species, *Limulitella tejraensis*, probably lived in freshwater environments, and would be phylogenetically related with another lower Middle Triassic (Anisian) limulid from France, *Limulitella bronni* (see Błażejowski *et al.*, 2017).

Beyond Europe and North Africa, some Triassic xiphosuran remains have been discovered. Recently in China, ichnites from the Daye Formation of Guiyang (South China) have been described. Shi *et al.* (2019) described two different types of traceways referred as *Kouphichnium* from the Lower Triassic. The palaeoenvironmental conclusions suggest that the tracemakers would inhabit the seafloor at 50-60 m deep, in a complex ichnoassociation with other invertebrates. In North China, Shu *et al.* (2018) described 44 well-preserved traceways attributed to *Kouphichnium* from the upper Permian–Lower Triassic successions. Also, in upper Permian–Lower Triassic terrestrial successions (freshwater settings) from the Central Transantarctic Mountains (Antarctica) several horseshoe crab traceways have been attributed to the ichnospecies *Kouphichnium natalensis* by Hasiotis (2012). Finally, in Madagascar were recovered two specimens of a Lower Triassic limulid from the Ankitokazo Basin, tentatively determined as the genus *Limulitella* by Hauschke *et al.* (2004), and in New South Wales (Australia), the limuloid *Dubbolimulus peetae*, which would have inhabited freshwater settings during the Middle Triassic, was described (Pickett, 1984).

All these fossil remains suggest that, after the Permian mass extinction, xiphosurans recovered and radiated during the Middle Triassic, with species occupying both marine environments and, of special note, continental freshwater environments (cf. Błażejowski *et al.*, 2017; Mujal *et al.*, 2018a). Most of the known Triassic fossil remains of xiphosurans from the present Northern Hemisphere have been described from Europe, with few additional records from North Africa and China.

6.3. Size of the tracemaker

The size of the tracemaker can be inferred from the external width of the traceway, following the formula $I = W \times 1.5$ (Malz, 1964; Shi *et al.*, 2019), where I is the width of the prosoma of the tracemaker and W is the external width of the traceway. In the herein reported specimen, the traceway width is 26.8 mm, so the estimated prosoma width is 40.2 mm. Also, using the formula $L = I \times 2.12$ (see Shi *et al.*, 2019), the length of the tracemaker (L) can be calculated, being 85.2 mm for the tracemaker of the studied specimen. The mean L value obtained in *Kouphichnium* specimens from the Lower Triassic in Daye Formation (China) is 192.7 mm (Shi *et al.*, 2019). Mújal *et al.* (2018a) described *Kouphichnium* specimens from the upper Lower Triassic of the Catalan Pyrenees of up to 70 mm wide. In comparison, the size of the tracemaker from Penya Rubí would be among the smallest found in the Triassic ichnofossil record (Table 1). It suggests that the traceway might correspond to a juvenile specimen or a small xiphosuran taxon. However, according to Martin & Rindsberg (2007), juvenile specimens of xiphosurans might produce traces attributable to *Nereites*, thus the option of a small xiphosuran taxon as the producer of the studied traceway is preferred.

The Iberian species *Mesolimulus crespelli* (width measures: 35 mm for the holotype, and 31 and 53 mm for the other specimens) presents the most similar size to the studied traceway (Table 2). Also, *Tarracolimulus reiki* shows a similar size, although it is slightly smaller. Otherwise, *T. gadeai* shows the most different size, being larger than 140 mm wider than the Penya Rubí specimen. Therefore, *M. crespelli* shows the most similar size to the inferred size of the tracemaker of Penya Rubí *Kouphichnium* traceway, but we cannot ignore that *T. reiki* had an almost identical size.

7. CONCLUSIONS

The arthropod traceway recovered from the Penya Rubí ichnosite of the Catalan Basin has been identified as *Kouphichnium* isp., a locomotion trace fossil (repichnia) attributed to xiphosurans. This is the first invertebrate traceway described from the middle Muschelkalk (Middle Triassic) of the Catalan Basin. The traceway is composed of three parallel rows of imprints, the central one represented by straight grooves (telson impressions) and the lateral rows (two in each side of the telson impression) being composed of straight and pincer-shaped (bifid) imprints (internal row of each side) and pointed and arrow-shaped imprints (external row of each side). The lateral rows evidently correspond to the impressions of the different appendages. According to the general traceway pattern and the arrangement of the different appendages and telson traces, a crawling behaviour was the most probable locomotion style of the tracemaker.

The animal that produced this trace fossil was probably roaming an intertidal area with shallow, quiet water. Considering the size of the imprints, the tracemaker would be a small xiphosuran of 40.2 mm width, which could possibly be an adult of a small-sized taxon in a breeding season. Comparing the size of the traceway, the Triassic species *Mesolimulus crespelli* and *Tarracolimulus rieki* present the most similar size to the theoretical size of the tracemaker of Penya Rubí ichnites, but it is not possible to confidently correlate the traceway with a specific taxon.

8. ACKNOWLEDGMENTS

First of all, thanks to the Spanish Society of Palaeontology (SEP), for granting financial aid to carry out this investigation. We are deeply indebted to Dr. Oriol Oms (Universitat Autònoma de Barcelona), discoverer of the slab, and also to Dr. Arnau Bolet (University of Bristol and ICP), Manel Méndez (ICP) and Marc Riccetto for helping us during the fieldtrips. Also to Dr. Chabier de Jaime for logistics related to the project. J.F. is supported by the Spanish Agencia Estatal de Investigación and the European Regional Development Fund of the European Union (AEI/FEDER EU, project CGL2017-82654-P). J.F. is member of the consolidated research group 2017 SGR 86 GRC of the Generalitat de Catalunya. We acknowledge support from the CERCA programme (ICP) from the Generalitat de Catalunya, and the project “Evolució dels ecosistemes durant la transició Paleozoic–Mesozoic a Catalunya” (ref. CLT009/18/00066), financially supported by the Departament de Cultura (Generalitat de Catalunya). We are grateful to Dr. Andrew K. Rindsberg and Dr. Alberto Pérez-López, who reviewed the manuscript and provided valuable and constructive comments that substantially improved a previous version of this work.

9. BIBLIOGRAPHY

- Benton, M.J. & Newell, A.J. 2014. Impacts of global warming on Permo-Triassic terrestrial ecosystems. *Gondwana Research*, 25, 1308-1337. DOI: 10.1016/j.gr.2012.12.010
- Berrocal-Casero, M., Arribas, M. & Moratalla, J.J. 2018. Didactic and divulgative resources of the Middle Triassic vertebrate Tracksite of Los Arroturos (Province of Guadalajara, Spain). *Geoheritage*, 10(3), 375-384. DOI: 10.1007/s12371-017-0244-1

424 Bicknell, R.D.C., Brougham, T., Charbonnier, S., Sautereau, F., Hitij, T. & Campione,
 425 N. 2019. On the appendicular anatomy of the xiphosurid *Tachypleus syriacus* and
 426 the evolution of fossil horseshoe crab appendages. *The Science of Nature*, 106, 38.
 427 DOI: 10.1007/s00114-019-1629-6

428 Błażejowski, B., Niedźwiedzki G., Boukhalf, K. & Soussi, M. 2017. *Limulitella*
 429 *tejraensis*, a new species of limulid (Chelicerata, Xiphosura) from the Middle
 430 Triassic of southern Tunisia (Saharan Platform). *Journal of Paleontology* 91(5),
 431 960-967. DOI: 10.1017/jpa.2017.29

432 Calvet, F., Tucker, M.E. & Henton, J.M. 1990. Middle Triassic carbonate ramp systems
 433 in the Catalan Basin, northeast Spain: facies, systems tracts, sequences and controls.
 434 *Special Publications International Association of Sedimentology*, 9, 79-108.

435 Calvet, F. & Marzo, M. 1994. El Triásico de las Cordilleras Costero Catalanas:
 436 Estratigrafía, Sedimentología y Análisis Secuencial. Cuaderno de Excursión. III
 437 Coloquio de Estratigrafía y Paleoestratigrafía del Pérmico y Triásico de España.
 438 *Field Guide*, 1-53.

439 Caster, K.E. 1938. A restudy of the Tracks of *Paramphibius*. *Journal of Paleontology*,
 440 12, 3-60.

441 Caster, K.E. 1944. Limuloid trails from the upper Triassic (Chinle) of the Petrified Forest
 442 National Monument, Arizona. *American Journal of Science*, 242, 74-84.

443 Davis, R.A. Jr. & Dalrymple, R.W. (eds.) 2012. *Principles of tidal sedimentology*.
 444 Springer, London.

445 Diedrich, C. G. 2011. Middle Triassic horseshoe crab reproduction areas on intertidal
 446 flats of Europe with evidence of predation by archosaurs. *Biological Journal of the*
 447 *Linnean Society*, 103, 76-105.

448 Escudero-Mozo, M.J., Márquez-Aliaga, A., Goy, A., Martín-Chivelet, A., López-Gómez,
 449 J., Márquez, L., Arche, A., Plasencia, P., Pla, C., Marzo, M. & Sánchez-Fernández,
 450 D. 2015. Middle Triassic carbonate platforms in eastern Iberia: Evolution of their
 451 fauna and palaeogeographic significance in the Western Tethys. *Palaeogeography*,
 452 *Palaeoclimatology*, *Palaeoecology*, 417, 236-260. DOI:
 453 10.1016/j.palaeo.2014.10.041

454 Falkingham, P.L. 2012. Acquisition of high resolution 3D models using free, open-
 455 source, photogrammetric software. *Paleontologia Electronica*, 15, 1-15.
 456 DOI:10.26879/264

457 Fernández, D.E. & Pazos, P.J. 2013. Xiphosurid trackways in a Lower Cretaceous tidal
 458 flat in Patagonia: Palaeoecological implications and the involvement of microbial
 459 mats in tracefossil preservation. *Palaeogeography*, *Palaeoclimatology*,
 460 *Palaeoecology*, 375, 16-29. DOI: 10.1016/j.palaeo.2013.02.008

461 Fortuny, J., Bolet, A., Sellés, A. G., Cartanyà, J. & Galobart, À. 2011. New insights on
 462 the Permian and Triassic vertebrates from the Iberian Peninsula with emphasis on
 463 the Pyrenean and Catalanian basins. *Journal of Iberian Geology*, 37 (1), 65-86.
 464 DOI: 10.5209/rev_JIGE.2011.v37.n1.5

465 Franz, M., Kaiser, S.I., Fischer, J., Heunisch, C., Kustatscher, E., Luppold, F.W., Berner,
 466 U. & Röhlhng, H.-G. 2015. Eustatic and climatic control on the Upper Muschelkalk
 467 Sea (late Anisian/Ladinian) in the Central European Basin. *Global and Planetary*
 468 *Change*, 135, 1-27. DOI: 10.1016/j.gloplacha.2015.09.014

469 Gand, G., De La Horra, R., Galán-Abellán, B., López-Gómez, J., Barrenechea, J.F.,
 470 Arche, A. & Benito, I. 2010. New ichnites from the Middle Triassic of the Iberian
 471 Ranges (Spain) paleoenvironmental and paleogeographical implications. *Historical*
 472 *Biology*, 22, 40-56. DOI: 10.1080/08912961003644096

473 Gaillard, C. 2011. A giant limulid trackway (*Kouphichnium lithographicum*) from the
474 lithographic limestones of Cerin (late Kimmeridgian, France): Ethological and
475 environmental implications. *Swiss Journal of Geosciences*, 104(Suppl. 1), 57-72.
476 DOI: 10.1007/s00015-010-0032-2

477 Galán-Abellán, B., López-Gómez, J., Barrenecha, J.F., Marzo, M., De la Horra, R. &
478 Arche, A. 2013. The beginning of the Buntsandstein cycle (Early-Middle Triassic)
479 in the Catalan Ranges, NE Spain: Sedimentary and palaeogeographic implications.
480 *Sedimentary Geology*, 296, 86-102. DOI: 10.1016/j.sedgeo.2013.08.006

481 Goldring, R. & Seilacher, A. 1971. Limulid undertracks and their sedimentological
482 implications. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*,
483 137(3), 422-442.

484 Hasiotis, S. T. 2012. Horseshoe crabs lived in Permo-Triassic Antarctic freshwater rivers
485 and lakes: Trace fossil evidence from the Buckley and Fremouw Formations
486 Beardmore Glacier Area, Central Transantarctic Mountains (CTAM), Antarctica.
487 *Geological Society of America, Abstracts with Programs*, 44(7), 500.

488 Hauschke, N., Wilde, V. & Brauckmann, C. 2004. Triassic limulids from Madagascar
489 missing links in the distribution of Mesozoic Limulacea. *Neues Jahrbuch für*
490 *Geologie und Paläontologie*, 87-94.

491 Kin, A. & Błazejowski, B. 2014. The horseshoe crab of the genus *Limulus*: Living fossil
492 or stabilomorph? *PLoS ONE*, 9, e108036. DOI: 10.1371/journal.pone.0108036

493 Lamsdell, J.C. 2016. Horseshoe crab phylogeny and independent colonizations of fresh
494 water: Ecological invasion as a driver for morphological innovation.
495 *Palaeontology*, 59(2), 181-194. DOI: 10.1111/pala.12220

496 Mallison, H. & Wings, O. 2014. Photogrammetry in paleontology, a practical guide.
497 *Journal of Paleontological Techniques*, 12, 1-31.

498 Malz, H. 1964. *Kouphichnium walchi*, die Geschichte einer Fährte und ihres Tieres: *Natur*
499 *und Museum*, 94, 81–97.

500 Marcé-Nogué, J., Fortuny, J., Gil, L. & Sánchez, M. 2015. Improving mesh generation in
501 finite element analysis for functional morphology approaches. *Spanish Journal of*
502 *Palaeontology*, 30(1), 117-132.

503 Márquez-Aliaga, A., Valenzuela-Rios, J.I., Calvet, F. & Budurov, K. 2000. Middle
504 Triassic conodonts from northeastern Spain; biostratigraphic implications. *Terra*
505 *Nova*, 12, 77-83.

506 Martin, A.J. & Rindsberg, A.K. 2007. Arthropod tracemakers of *Nereites*? -
507 Neoichnological observations of juvenile limulids and their paleoichnological
508 applications. In *Trace fossils: Concepts, problems, prospects* (Ed. W. Miller), 478-
509 491.

510

511 Moore, R.A., McKenzie, S.C. & Lieberman, B.S. 2007. A Carboniferous synxiphosurine
512 (Xiphosura) from the Bear Gulch Limestone, Montana, USA. *Palaeontology*, 50(4),
513 1013-1019. DOI: 10.1111/j.1475-4983.2007.00685.x

514 Mujal, E., Fortuny, J., Rodríguez-Salgado, P., Diviu, M., Oms, O. & Galobart, À. 2015.
515 First footprints occurrence from the Muschelkalk detrital unit of the Catalan
516 Basin: 3D analyses and palaeoichnological implications. *Spanish Journal of*
517 *Palaeontology*, 30, 97-107.

518 Mujal, E., Fortuny, J., Bolet, A., Oms, O. & López, J.Á. 2017. An archosauromorph
519 dominated ichnoassemblage in fluvial settings from the late Early Triassic of the
520 Catalan Pyrenees. *PLoS ONE*, 12(4), e0174693. DOI:
521 10.1371/journal.pone.0174693

522 Mujal, E., Belaústegui, Z., Fortuny, J., Bolet, A., Oms, O. & López, J.Á. 2018a.
523 Ichnological evidence of a horseshoe crab hot-spot in the Early Triassic
524 Buntsandstein continental deposits from the Catalan Pyrenees (NE Iberian
525 Peninsula). *Journal of Iberian Geology*, 44, 139-153. DOI: 10.1007/s41513-017-
526 0026-2

527 Mujal, E., Iglesias, G., Oms, O., Fortuny, J., Bolet, A. & Méndez, J.M. 2018b.
528 *Rhynchosauroides* footprint variability in a Muschelkalk detrital interval late
529 Anisian–middle Ladinian) from the Catalan Basin (NE Iberian Peninsula). *Ichnos*,
530 25(2-3), 150-161. DOI: 10.1080/10420940.2017.1337571

531 Mujal, E., Marchetti, L., Schoch R.R. & Fortuny, J. 2020. Upper Paleozoic to Lower
532 Mesozoic tetrapod ichnology revisited: Photogrammetry and relative depth pattern
533 inferences on functional prevalence of autopodia. *Frontiers in Earth Science*, 8,
534 248. DOI: 10.3389/feart.2020.00248

535 Ortí, F., Salvany, J.M., Rosell, L., Castelltort, X., Inglès, M. & Playà, E. 2018. Middle
536 Triassic evaporite sedimentation in the Catalan Basin: implications for the
537 paleogeographic evolution in the NE Iberian Platform. *Sedimentary Geology*, 374,
538 158-178. DOI: 10.1016/j.sedgeo.2018.07.005

539 Pickett, J.W. 1984. A new freshwater limuloid from the Middle Triassic of New South
540 Wales. *Palaeontology*, 27(3), 609-621.

541 Ramón, X. & Calvet, F. 1987. Estratigrafía y sedimentología del Muschelkalk inferior
542 del dominio Monserrat-Llobregat (Catalanides). *Estudios Geológicos*, 43, 471-487.

543 Rindsberg, A.K. & Martin, A.J. 2015. Caster's Plasters: Neoichnological Experiments by
544 Kenneth Caster on Limulids in 1937. In *ICHTHOLOGY: Papers from ICHNIA III:*
545 *Geological Association of Canada, Miscellaneous Publications* (Eds. McIlroy, D.),
546 197-210.

547 Romero, A. & Vía-Boada, L. 1977. *Tarracolimulus riei*, nov. gen., nov. sp., nuevo
548 limúlido del Triásico de Montral-Alcover. *Cuadernos de Geología Ibérica*, 4, 239-
549 246.

550 Rudkin, D.M., Young, G.A. & Nowlan, G.S. 2008. The oldest horseshoe crab: A new
551 xiphosurid from Late Ordovician Konservat-Lagerstätten deposits, Manitoba.
552 Canada. *Palaeontology*, 51(1), 1-9. DOI: 10.1111/j.1475-4983.2007.00746.x

553 Seilacher, A. 2007. *Trace fossil analysis*. Berlin, Springer.

554 Shen, J., Schoepfer, S. D., Feng, Q., Zhou, L., Yu, J., Song, H., Wei, H. & Algeo, T.J.
555 2015. Marine productivity changes during the end-Permian crisis and Early Triassic
556 recovery. *Earth-Science Reviews*, 149, 136-162. DOI:
557 10.1016/j.earscirev.2014.11.002

558 Shi, G., Woods, A. D., Yu, M.-Y., Li, X.-W., Wei, H.-Y. & Qiao, D. 2019. Lower Triassic
559 limulid trackways (*Kouphichnium*) from the southwestern margin of the Yangtze
560 carbonate platform: paleoenvironmental and paleoecological implications. *Palaios*,
561 34(4), 229-243. DOI:10.2110/palo.2018.081

562 Shibata, M. & Varricchio, D.J. 2020. Horseshoe crab trace fossils from the Upper
563 Cretaceous Two Medicine Formation of Montana, USA, and a brief review of the
564 xiphosurid ichnological record. *Journal of Paleontology*, 1-19. DOI:
565 10.1017/jpa.2020.16

566 Shu, W., Tong, J., Tian, L., Benton, M.J., Chu, D., Yu, J. & Guo, W. 2018. Limuloid
567 trackways from Permian-Triassic continental successions of North China.
568 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 508, 71-90. DOI:
569 10.1016/j.palaeo.2018.07.022

570 Shuster, C.N. Jr. 1982. A pictorial review of the natural history and ecology of the
571 horseshoe crab *Limulus Polyphemus*, with reference to other Limulidae. In:

Physiology and biology of horseshoe crabs. Studies on normal and environmentally stressed animals (Eds. Bonaventura, J., Bonaventura, C. & Tesh, S.). New York: Alan R. Liss, Inc, 1-55.

Sole de Porta, N., Calvet, F. & Torrento, L. 1987. Análisis palinológico del Triásico de los Catalanides (NE España). *Cuadernos de Geología Ibérica*, 11, 237-254.

Vía, L. 1987. Artrópodos fósiles triásicos de Alcover-Montral. II. Limúlidos. *Cuadernos de Geología Ibérica*, 11, 281-282.

Vía-Boada, L. & Villata, J.F. 1966. *Heterolimulus gadeai* nov. gen., nov. sp., representante de una nueva familia de LIMULACEA, en el Triásico español. *Acta geológica hispánica*, 1, 9-11.

Vía-Boada, L., Villalta, J.F. & Cerdá, M.E. 1977. Paleontología y paleoecología de los yacimientos fosilíferos del Muschelkalk Superior entre Alcover y Mont-Ral (Montañas de Prades, Provincia de Tarragona). *Cuadernos de Geología Ibérica*, 4, 247-256.

Virgili C., Sopena, A., Ramos, A. & Hernando, S. 1977. Problemas de la cronoestratigrafía del Trías en España. *Cuadernos de Geología Ibérica*, 4, 57-87.

10. TABLES

Table 1.

Outcrop	W value (mm) Traceway width	I value (mm) Prosoma width	L value (mm) Tracemaker length
Penya Rubí (this work)	26.8	40.2	85.2
Catalan Pyrenees	70 (maximum)	105	222.6

(Mujal <i>et al.</i> , 2018a)			
Daye Formation (Shi <i>et al.</i> , 2019)	46.9–79.2 mm	89 (average)	192.7 (average)

Table 2.

Taxon of fossil limulid	Width measure data (mm)	Mean width measure (mm)
<i>Tachypleus gadeai</i>	181, 168	174.5
<i>Tarracolimulus rieki</i>	36.9	36.9
<i>Mesolimulus crespelli</i>	35 (holotype), 31, 53	39.7
<i>Kouphichnium</i> isp. (IPS 110272)	40.2	40.2

11. TABLE AND FIGURE CAPTIONS

Table 1. Size comparison. Obtaining the width of the prosoma (I value knowing the width of the traceway, W, using the formula $I = W \times 1.5$) and the length of the tracemakers (L value, using the formula $L = I \times 2.12$) in the xiphosuran traceways from the Lower and Middle Triassic.

Table 2. Width of the Middle Triassic Iberian limulid fossils (all from the upper Muschelkalk of the Catalan Basin) and inferred width of the potential tracemaker for the Peña Rubí traceway. *Tachypleus gadeai* (measures from Vía-Boada & Villata, 1966). *Tarracolimulus rieki* (measures from Romero & Vía-Boada, 1987). *Mesolimulus crespelli* (measures from Vía, 1987).

Figure 1. Geographic location and simplified geological maps of studied areas. a. Location of the Catalan Basin in the Iberian Peninsula. b. Geological map of the Triassic around Peña Rubí (Vallirana) in the Catalan Basin. Maps modified from the Institut Cartogràfic i Geològic de Catalunya (ICGC).

Figure 2. Stratigraphic log of the middle Muschelkalk facies from Peña Rubí (Vallirana, Catalan Basin) where tetrapod footprints and invertebrate traces appear (uppermost part of the outcropping section). Md: Mudstone; Ss: Sandstone; vf:very fine; f:fine; m:medium; c:coarse; vc:very coarse; Cgl: Conglomerate.

Figure 3. a. Overview of the middle Muschelkalk sedimentary succession at Peña Rubí (Vallirana) outcrop. b. Detail of the stratigraphic interval where the sandstone stratum preserving the traceway was found. The arrow points to the exact level.

622

623 **Figure 4.** *Kouphichnium* isp. (IPS110272) from the middle Muschelkalk facies of Penya
624 Rubí, advancing from left to right in the photos. a. Photo of the whole traceway. b.
625 Interpretative outline of the traceway. c. Detail of the traceway showing the linear ichnites
626 produced by appendages II-V. The arrow points to a curvilinear ichnite. d. Detail of the
627 traceway showing the arrow-shaped imprints produced by appendages VI (upper arrow),
628 and pointed, lineal and bifid- imprints produced by appendages II-V (lower arrow). e.
629 Ideal traceway using *Mesolimulus crespelli* (redrawn from Vía, 1987; scaled to fit the
630 traceway width) as potential tracemaker.

631

632 **12. SUPPLEMENTARY INFORMATION**

633 A three-dimensional photogrammetric model of the slab (IPS110272) that contains
634 *Kouphichnium* isp. is available from MorphoBank
635 <http://morphobank.org/permalink/?P3770>.
636







