

RESEARCH

Open Access



International variation in the management of severe COVID-19 patients

Elie Azoulay^{1*} , Jan de Waele², Ricard Ferrer^{3,4}, Thomas Staudinger⁵, Marta Borkowska², Pedro Povoas⁶, Katerina Iliopoulou⁷, Antonio Artigas⁸, Stefan J. Schaller⁹, Manu Shankar-Hari¹⁰, Mariangela Pellegrini^{11,12,13}, Michael Darmon¹, Jozef Kesecioglu¹⁴ and Maurizio Cecconi¹⁵

Abstract

Background: There is little evidence to support the management of severe COVID-19 patients.

Methods: To document this variation in practices, we performed an online survey (April 30–May 25, 2020) on behalf of the European Society of Intensive Care Medicine (ESICM). A case vignette was sent to ESICM members. Questions investigated practices for a previously healthy 39-year-old patient presenting with severe hypoxemia from COVID-19 infection.

Results: A total of 1132 ICU specialists (response rate 20%) from 85 countries (12 regions) responded to the survey. The survey provides information on the heterogeneity in patient's management, more particularly regarding the timing of ICU admission, the first line oxygenation strategy, optimization of management, and ventilatory settings in case of refractory hypoxemia. Practices related to antibacterial, antiviral, and anti-inflammatory therapies are also investigated.

Conclusions: There are important practice variations in the management of severe COVID-19 patients, including differences at regional and individual levels. Large outcome studies based on multinational registries are warranted.

Keywords: Coronavirus, Acute respiratory distress syndrome, Viral infection, Remdesivir

Background

There is little evidence to support the optimal management of severe COVID-19 patients [1, 2]. To document whether there is a variation in practices, we performed an online survey (April 30–May 25, 2020) on behalf of ESICM.

Methods

In this online survey, a case vignette (<https://www.surveymonkey.com/r/F2FFC6S>) was sent to ICU specialists who are members of ESICM. Questions investigated

practices for a previously healthy 39-year-old patient presenting with severe hypoxemia from COVID-19 infection (Table 1). The 85 participating countries were grouped into 12 different regions [3]: continuous variables are described as median (interquartile range [IQR]) and are compared between groups using the non-parametric Wilcoxon rank-sum test. Categorical variables are described as frequency (percentages) and are compared between groups using Fisher's exact test. Statistical analyses were performed with R statistical software, version 3.4.3 (available online at <http://www.r-project.org/>). A *p* value < 0.05 was considered significant.

* Correspondence: elie.azoulay@aphp.fr

¹Médecine Intensive et Réanimation, Department of the St-Louis Hospital, APHP, Hôpital Saint-Louis, Paris University, 1 avenue Claude Vellefaux, 75010 Paris, France

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Results

Response rate was 20% ($N = 1132$ intensive care (ICU) specialists from 85 countries, including 1001 complete answers). Respondents (median 45 years [IQR, 39–53], 34% women) were from Middle Europe (25%), South Europe (23%), the United Kingdom (UK) (12%),

South America (9%), North Europe (8.1%), Eastern Europe (5.3%), Middle-East (5%), North America (4.7%), Asia (3.3%), India (2.7%), Australia-New Zealand (1.3%), or Africa (0.6%); 54% were living in a large city (>1 million inhabitants), and 55% were working in university-affiliated hospitals. The median

Table 1 Distribution of the responses to the case vignette

Numbers (%) or median (interquartile ranges)	Total, 1001 respondents
1. Admission to the ICU of a previously healthy 39-year-old man with severe COVID-19	
Direct admission to the ICU	55%
Admission in an intermediate care unit	34.2%
Delayed admission to the ICU because of lack of bed	1.3%
Patient stays in the emergency department	0.8%
Patient admitted in the wards	8.8%
2. Initial oxygenation strategy	
I increase the oxygen flow to 15 l/min keeping the face mask	24.2%
I change the mask for a Venturi mask	17.5%
I start CPAP or noninvasive ventilation	25.5%
I start high flow nasal oxygen	47.1%
I intubate the patient right away	7.4%
I add prone positioning on spontaneous breathing	37.9%
3. Optimizing oxygenation in a patient with a PF ratio of 84 4 h after intubation	
I will give neuromuscular blockade for 24–48 h	50.9%
I increase and titrate PEEP to optimize recruitment	61.4%
I prone the patient immediately	73.2%
I am considering ECMO immediately	4.7%
Let us wait a little bit	9.9%
4. Regarding the initial antibiotics	
All my patients receive a broad anti-bacterial agent	45.3%
I only give broad anti-bacterial agent to febrile patients	11%
I only give broad anti-bacterial agent if CRP or PCT are high	4.2%
I only give broad anti-bacterials to patients with structural lung diseases	35.8%
I never give broad anti-bacterial agent in severe viral infections	3.7%
5. Regarding initial anti-viral therapy, several options are possible	
The level of evidence is so low that there is nothing I can give	48.9%
I prescribe (hydroxy)chloroquine	42.7%
I prescribe lopinavir/ritonavir	17.0%
I prescribe remdesivir	15%
I prescribe another anti-viral drug	4.6%
6. Are you starting an anti-inflammatory therapy?	
No	52.4%
Yes IL-1 or IL-6 blockade	24.8%
Yes, complement blockade	1.4%
Yes, steroids	31.5%
Yes, another anti-inflammatory drug	2.4%

(IQR) number of ICU beds was increased from 20 (11–36) to 35 (20–60) during the pandemic surge.

As the patient had 88 (peripheral oxygen saturation) SpO_2 on 9 l/min of oxygen, direct ICU admission was reported in 56% (30–90%) of the respondents, with significant variation across regions (Fig. 1, $P < 0.0001$). Most intensivists not directly admitting patients to the ICU would admit them to an intermediate care unit managed by intensivists. However, the issue of bed availability was reported in South Europe (4.5%), South America (2.9%), Scandinavia (1.6%), Middle Europe (1.6%), and the UK (1.1%). Should the patient be not admitted to the ICU, a rapid response team would be involved in 29% of the cases, the ICU specialist would make the outreach her/himself in 24% of the cases, or an ICU nurse would be involved in 7% of the cases. In all other cases, ward or ED physicians would manage the patients. Direct ICU admission was significantly associated with baseline number of ICU beds (22 [12–40] vs. 18 [10–30] beds, $P < 0.0001$) and with the number of COVID-19 patients managed (30 [11–52] vs. 38 [20–70] patients, $P = 0.001$), as well as with management of patients in large cities (56.9% vs. 49.6%, $P = 0.04$).

Respondents were then asked about the first-line oxygenation strategy, which varied significantly across regions (Fig. 2, $P < 0.0001$). First-line high flow nasal cannula (HFNC) was used by 22.9% of the respondents (0% in Australia-New Zealand, 38% in Eastern Europe). Noninvasive ventilation was used by 25.5% of the respondents (5.4% in North America, 43.6% in the UK). Interestingly, 8% of the respondents were using first-line intubation (0% in Australia-New Zealand, 23% in Asia). Women less frequently initiated HFNC (32% vs. 42%, $P = 0.02$). The availability of an intermediate care unit influenced the use of HFNC or non-invasive ventilation (NIV) (32.8% vs. 21.7%, $P = 0.03$). Along this line, a higher number of ICU beds (24 [12–40] vs. 18 [10–30] beds, $P = 0.0009$) was associated with the use of HFNC and NIV. Interestingly, 37.5% were using prone positioning in awake non-ventilated patients. To assess whether HFNC or NIV should be continued, ICU specialists relied on SpO_2 (85.7%), respiratory rate (71.4%), followed by dyspnea (47.1%), and comfort (45.4%). Criteria for intubation included clinical signs of respiratory distress (94%), high oxygen flow to maintain a SpO_2 of 95% (33.5%), or low SpO_2 only (25.6%).

Rate of Direct Admission to the Intensive Care Unit or the Intermediate Care Unit



Rate of Direct Admission to the Intensive Care Unit



Fig. 1 World map displaying practice variations across regions regarding direct admission to the ICU or the intermediate care unit

Following intubation, the patient had a partial pressure of oxygen/fraction of inspired oxygen (P/F) ratio of 84 mmHg. Although prone positioning (71.2%) and neuromuscular blockade (59.7%) were often used to optimize oxygenation, the practice varied significantly across countries. For instance, prone positioning was performed in 70–85% of the cases in Asia, India, Eastern Europe, Middle Europe, South America, South Europe, and the UK, whereas Africa, Australia-New Zealand, Middle East, North America, and Scandinavia were in the 50–70% range (Fig. 3, $P < 0.0001$). Respondent's age was associated with the use of prone positioning (46 [39–54] vs. 44 [37–51] years, $P = 0.007$). Older respondent's age (45 [37–52] vs. 47 [40–55] years, $P = 0.0001$), living in a large city (54.2% vs. 46.8%, $P = 0.03$), and a higher number of COVID-19 patients managed (35 [15–65] vs. 30 [12–55] patients, $P = 0.02$) were associated with the use of neuromuscular blockade.

Antibiotic prescribing was routine for all patients in 44.2% of the respondents and biomarker-guided in 36.5%, without significant variation across regions. Routine antibiotics were more frequently used by respondents working in university-affiliated hospitals (48.3% vs. 40.9%, $P = 0.03$) and those living in large cities (49.3% vs. 40.2%, $P = 0.01$). Biomarker-guided antibiotic therapy was less frequent in large cities (47.3% vs. 57.4%, $P = 0.007$). Regarding antiviral therapy, 48.9% reported not prescribing antivirals, 42.6% were giving hydroxychloroquine, 17% lopinavir-ritonavir, and 15% remdesivir. Figure 4 displays significant variation in antiviral prescriptions across regions ($P < 0.0001$). Physicians not prescribing antivirals were older (47 [40–54] vs. 44 [37–51] years, $P < 0.0001$), and more frequently men (55.4% vs. 39.9%, $P < 0.0001$). Conversely, those prescribing hydroxychloroquine were younger (43 [37–50] vs. 47 [40–54] years, $P < 0.0001$), and more frequently

Use of Non-Invasive Mechanical Ventilation




Use of High Flow Nasal Cannula



Fig. 2 World map displaying practice variations across regions regarding the use of noninvasive oxygenation strategies

Use of Neuromuscular Blockade



Use of Prone Positioning




Fig. 3 World map displaying practice variations across regions regarding the use of neuromuscular blockades and prone positioning

women (41.7% vs. 28.2%, $P < 0.0001$). There was significant variation in the use of interleukin-6 (IL-6)/IL-1 blockade or of corticosteroids across countries ($P < 0.0001$ for both tests). Other collected variables were not associated with the use of anti-inflammatory drugs.

Discussion

This survey highlights important practice variations in the management of severe COVID-19 patients, including differences at regional and individual levels. This illustrates that neither IDSA nor Surviving Sepsis Guidelines did recommend any of these treatments, but instead encouraged inclusion of patients into trials [1, 4, 5]. Since the publication of these guidelines, no more evidence has been made available to ascertain that these specific COVID-19 therapies should be included in the standard of

care. Learning from this heterogeneity will not only raise hypothesis on optimal patient's management, but also serves as a tool to suggest personalized management for each clinical phenotype [6, 7].

This study has several limitations. First, the study suffers from a nonresponse bias of 80%. Second, even though only physicians have responded, we cannot ascertain that all of them had the clinical expertise and the experience of managing COVID-19 patients. Last, questions about specific treatments did not take into account the fact that the level of evidence has changed over time.

Conclusion

As no management guidelines have allowed to guide practices for the COVID-19 pandemic, heterogeneous behaviors are reported. Large

Proportion of Respondents Considering No Antiviral Agent



Proportion of Respondents Considering Remdesivir



Proportion of Respondents Considering Hydroxychloroquine



Fig. 4 World map displaying practice variations across regions regarding the use of antiviral agents

outcome studies based on multinational registries are warranted.

Acknowledgements

We are indebted to Dominique de Boom, Nicolas Vander Elst, and Guy Francois from the ESICM office for their fast and high-quality support to prepare this survey and its diffusion.

Authors' contributions

All authors contributed to the design, the development of the instrument, and interpretation of data. All authors have approved the submitted manuscript.

Funding

None

Availability of data and materials

Fully available upon request.

Ethics approval and consent to participate

All participants agreed online to complete the survey.

Consent for publication

All participants consented.

Competing interests

The authors declare no conflict of interest in relation to this survey.

Author details

¹Médecine Intensive et Réanimation, Department of the St-Louis Hospital, APHP, Hôpital Saint-Louis, Paris University, 1 avenue Claude Vellefaux, 75010 Paris, France. ²Department of Critical Care Medicine, Ghent University Hospital, 9000 Gent, Netherlands. ³Shock, Organ Dysfunction, and Resuscitation Research Group (SODIR), Instituto de Investigación de Vall d'Hebron, Barcelona, Spain. ⁴Departamento de Medicina Intensiva, Hospital Universitario de Vall d'Hebron, Centro de Investigación Biomédica en Red (CIBER) de Enfermedades Respiratorias, Barcelona, Spain. ⁵Department of Medicine I, Intensive Care Unit, Medical University of Vienna, Vienna General Hospital, Vienna, Austria. ⁶NOVA Medical School, CHRC, New University of Lisbon, Unidade de Cuidados Intensivos Polivalente, Hospital de São Francisco Xavier, CHLO, Estrada Do Forte Do Alto Do Duque, 1449-005 Lisbon, Portugal. ⁷Hellenic Army, ICU Nurse Manager General Military Hospital, Athens, Greece. ⁸Critical Care Center, Sabadell Hospital, University Institute Parc Taulí, Autonomous University of Barcelona, Ciberes, Barcelona,

Spain. ⁹Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Department of Anesthesiology and Surgical Intensive Care, Berlin, Germany. ¹⁰School of Immunology and Microbial Science, Kings College London. Guy's and St Thomas' NHS Foundation Trust, ICU Support Offices, St Thomas' Hospital, London, UK. ¹¹Department of Surgical Sciences and Central Intensive Care Unit, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden. ¹²Department of Anesthesia, Operation, and Intensive Care; Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden. ¹³Department of Anesthesiology and Intensive Care Medicine, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden. ¹⁴Department of Intensive Care Medicine, Division of Anesthesiology, Intensive Care and Emergency Medicine, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands. ¹⁵Humanitas Clinical and Research Center, Humanitas University, Milan, Italy.

Received: 17 June 2020 Accepted: 20 July 2020

Published online: 05 August 2020

References

1. Alhazzani W, Möller MH, Arabi YM, et al. Surviving Sepsis Campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). *Intensive Care Med* Published online March 28, 2020:1–34. doi:<https://doi.org/10.1007/s00134-020-06022-5>.
2. Bauchner H, Fontanarosa PB. Randomized clinical trials and COVID-19: managing expectations. *JAMA* Published online May 4, 2020. doi:<https://doi.org/10.1001/jama.2020.8115>.
3. Sprung CL, Ricou B, Hartog CS, et al. Changes in end-of-life practices in European intensive care units from 1999 to 2016. *JAMA*. 2019;322(17):1–12. <https://doi.org/10.1001/jama.2019.14608>.
4. Bhimraj A, Morgan RL, Shumaker AH, et al. Infectious diseases Society of America guidelines on the treatment and management of patients with COVID-19. *Clin Infect Dis Off Publ Infect Dis Soc Am* Published online April 27, 2020. doi:<https://doi.org/10.1093/cid/cia478>.
5. Aziz S, Arabi YM, Alhazzani W, et al. Managing ICU surge during the COVID-19 crisis: rapid guidelines. *Intensive Care Med* Published online June 8, 2020. doi:<https://doi.org/10.1007/s00134-020-06092-5>.
6. Helms J, Severac F, Merdji H, Anglés-Cano E, Meziani F. Prothrombotic phenotype in COVID-19 severe patients. *Intensive Care Med*. Published online May 20, 2020:1–2. doi:<https://doi.org/10.1007/s00134-020-06082-7>.
7. Azoulay E, Zafrani L, Mirouse A, Lengliné E, Darmon M, Chevret S. Clinical phenotypes of critically ill COVID-19 patients. *Intensive Care Med* Published online May 28, 2020. doi:<https://doi.org/10.1007/s00134-020-06120-4>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

