



Contents lists available at ScienceDirect

Science of the Total Environment

journal homepage: www.elsevier.com/locate/scitotenv

Assessing the environmental behavior of alternative fertigation methods in soilless systems: The case of *Phaseolus vulgaris* with struvite and rhizobia inoculation

Verónica Arcas-Pilz^a, Martí Ruffi-Salís^{a,b}, Felipe Parada^a, Xavier Gabarrell^{a,b,*}, Gara Villalba^{a,b}

^a Sostenipra Research Group (2017 SGR 1683), Institut de Ciència i Tecnologia Ambientals (CEX2019-000940-M), Z Building, Universitat Autònoma de Barcelona (UAB), Campus UAB, 08193 Bellaterra, Barcelona, Spain


^b Department of Chemical, Biological and Environmental Engineering, Universitat Autònoma de Barcelona (UAB), Campus UAB, 08193 Bellaterra, Barcelona, Spain



HIGHLIGHTS

- Inorganic fertilizers are a great constrain in urban agriculture due to their emissions.
- Struvite and nitrogen fixing bacteria can be considered to avoid additional impacts.
- While results show lower yields, life cycle impacts are significantly reduced.
- Struvite and nitrogen fixing bacteria rhizobium reduce N and P emitted to water.
- A yield reduction can be assumed depending on complexity of the infrastructure.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 14 October 2020

Received in revised form 23 December 2020

Accepted 24 December 2020

Available online 23 January 2021

Editor: Damia Barcelo

Keywords:

Urban agriculture

Life cycle assessment

Alternative fertilization

Phosphorus recovery

ABSTRACT

Urban agriculture, while being a promising solution to increase food sovereignty in cities, can lead to an unprecedented discharge of nutrient and fertilizer-related emissions into the urban environment. Especially relevant are nitrogen (N) and phosphorus (P), due to their contribution to marine and freshwater eutrophication. Therefore, alternative methods of fertilization need to be put into practice to avoid such impacts to the surrounding environment. Struvite, has been studied as a potential slow releasing fertilizer due to its high P content, while the bacteria rhizobium has been used to fix N directly from the atmosphere. Legumes, like the common bean are N-demanding crops capable of symbiosis with the bacteria rhizobium and have previously shown positive responses to fertilization with struvite. This study aims to analyze the environmental performance of plant production in hydroponic systems combining rhizobium inoculation and struvite (2 g, 5 g, 10 g, 20 g) irrigated with a N and P deficient nutrient solution, using life cycle analysis (LCA). The nutrient content of in- and out-going irrigation was analyzed as well as in plants and beans. The functional unit for the LCA was 1 kg of fresh beans. The results obtained indicate a yield reduction of 60% to 50% in comparison to the control which was irrigated with a full nutrient solution. The impacts from operational stage are less in all impact categories, where most significant reductions up to 69% and 59% are seen in marine-eutrophication and global warming respectively. Although the infrastructure does not change between treatments, its impacts increase due to the lower yields. We determine

* Corresponding author at: Sostenipra Research Group (2017 SGR 1683), Institut de Ciència i Tecnologia Ambientals (MDM-2015-0552), Z Building, Universitat Autònoma de Barcelona (UAB), Campus UAB, 08193 Bellaterra, Barcelona, Spain.

E-mail address: xavier.gabarrell@uab.cat (X. Gabarrell).

that below a 10% of the control yield, the alternative systems have more impact than the use of conventional mineral fertilizers in almost all impact categories, thus pointing to the importance of infrastructure to truly reduce environmental impacts for urban agriculture.

© 2021 Elsevier B.V. All rights reserved.

1. Introduction

Urban Agriculture (UA) has the potential to replace traditional food supply chains to some degree, thereby reducing transportation, packaging and food losses while increasing food sovereignty of cities (Sanyé et al., 2012; Tornaghi, 2017; Sanjuan-Delmás et al., 2018; Siegner et al., 2020). However, the additional need of inorganic chemical fertilizers inevitably results in greater discharge of these chemicals into the environment as well as an increase of the resource depletion potential (Rufi-Salís et al., 2020b). This is especially relevant considering the emission of nitrogen and phosphorus species, substantially contributing to marine and freshwater eutrophication, causing oxygen deprivation in aquatic environments. Specifically, urban water cycles and runoff are a great concern with their high implication in water eutrophication damaging ecosystems close to cities as well as close to intensely fertilized agricultural sites (John H. Ryther and Dunstan, 1971; Lewis et al., 2011). The integration of agriculture within city boundaries could therefore further increase the potential of emissions into the urban water cycles.

It is important to find ways for UA to be highly resource-efficient so that urban areas are able to expand these production practices without incurring significant environmental impacts associated with the additional water, energy, and nutrient requirements. To mitigate these environmental impacts, alternative and more environmentally friendly fertilizers have to be applied to attain competitive yields without causing great impacts to the surrounding environment (Lewis et al., 2011) as well as avoiding further extraction of phosphorous for agricultural purposes (Linderholm et al., 2012).

Recent work has been focused on the recovery of nutrients from wastewater treatment plants (Harder et al., 2019; Lam et al., 2020; Shaddel et al., 2020), showing a great range of possible alternatives for fertilization generated in urban areas as well as their constraints. One of the available options showing great potential for its use in agriculture, is struvite. The struvite crystal is formed by a spontaneous precipitation in wastewater treatment plants and is regarded as a slow releasing fertilizer due to its low dissolution and high content of phosphorous (12.5%), magnesium (9.9%) and nitrogen (5.7%) (Rahman et al., 2014; Talboys et al., 2016; Degryse et al., 2017). It has been reported that the formation of struvite can recover up to 90% of phosphate in wastewater sludge, reaching even higher percentages depending on the precipitation process and source (Kataki et al., 2016).

Studies on the use of struvite in agriculture (Massey et al., 2007; Ackerman et al., 2013; Talboys et al., 2016; Degryse et al., 2017) point out that the use of these recovered nutrients can reduce mineral fertilizer requirements while implying little to no cost for farmers (Karak and Bhattacharyya, 2011). The use of struvite has been shown to successfully substitute the use of mineral phosphorous fertilizers, while reducing nutrient losses to the environment due to its slow dissolution rate (Ahmed et al., 2018). Agricultural production with struvite as the main source of P has been tested on a variety of crops ranging from ryegrass (*Lolium perenne*) and broad beans (*Vicia faba*), which experience an increase of fresh yields of 76% and 54% respectively, to canola (*Brassica napus*) and wheat (*Triticum aestivum* L.), that suffer a reduction of the nutrient uptake and therefore a reduction of the plant yield (Ahmed et al., 2018). Other crops, however, have been seen to experience no significant changes with the use of struvite like the case of maize (Uysal et al., 2014) and corn (Thompson, 2013).

The environmental performance of the struvite extraction as P fertilizer has been previously studied (Ishii and Boyer, 2015) and while its benefits in comparison to virgin phosphorous have been identified in

the reduction of nutrient emissions and offsets commercial fertilizer production, its total environmental performance depends on the chemical inputs used for the struvite precipitation as well as the infrastructure and the recovery accounted in the life cycle inventory (Linderholm et al., 2012; Ishii and Boyer, 2015; Lam et al., 2020).

Whereas these studies have mainly substituted phosphorous based fertilizers, the use of nitrogen in the form of ammonium nitrate, urea and monoammonium is still given to the crops. As previously established, the emissions of nitrogen in the environment are greatly damaging and especially crucial for high N demanding crops like legumes.

The inoculation of legume crops with the bacteria rhizobium has been explored as a way for the plant to fix its nitrogen directly from the air without boosting its environmental footprint (Olivera et al., 2004; Gopalakrishnan et al., 2015; Kontopoulou et al., 2015, 2017; Savvas et al., 2018; Araujo et al., 2020; Sammauria et al., 2020; Sanyal et al., 2020). This bacteria forms an endosymbiotic interaction with the plant, profiting from compounds generated through photosynthesis while fixing atmospheric N₂ that is then given to the plant in form of ammonia (NH₃) (Long, 1989; Fisher and Long, 1992). As a result of these previous studies, it has been seen that in terms of the obtained yields, rhizobium tends to diminish the crop production in comparison to synthetic nitrogen fertilizers (Kontopoulou et al., 2015, 2017; Savvas et al., 2018; Sanyal et al., 2020) while its use on soil for common bean reduces about 19% per ha of the environmental burden when mineral N fertilization is replaced (Araujo et al., 2020).

To summarize the state of the art in nutrient recovery for mineral fertilizer substitution, the above-mentioned studies have shown that struvite can reduce and substitute a significant amount of P fertilizer while recovering great amounts of phosphate nutrients from wastewater treatment plants (WWTP). On the other hand, rhizobium can reduce the need for nitrogen mineral fertilizers, only partially reducing the environmental impact associated to the use of fossil-dependent mineral fertilizers. However, no study has attempted to use both struvite and rhizobium to completely avoid the application of nitrogen and phosphorous fertilizer, thereby reducing environmental impacts even further.

This paper aims to fill this gap by exploring the feasibility and environmental impact of applying struvite combined with rhizobium inoculation as alternative fertilizers of a UA system. To do so, we use the Life Cycle Assessment (LCA) tool to quantify the environmental impacts of a common bean crop in which the seeds were inoculated with soil bacteria rhizobium and different quantities of struvite were applied, compared to conventionally fertilized bean plants. The bush bean *Phaseolus vulgaris* "Pongo" was used in the experiments due to previous tryouts showing a good production in the perlite substrate as well as for being a highly consumed leguminous crop in Spain. The objective is to show the benefits and costs of this fertilizer alternative when compared to mineral fertilizer and provide knowledge towards reducing resource extraction for Urban Agriculture (UA) as well as avoiding possible emissions into natural ecosystems.

2. Materials & methods

2.1. Description of integrated Rooftop Greenhouse (i-RTG)

The experiments were conducted in the greenhouse laboratories for UA located on the integrated Rooftop Greenhouse Laboratory (i-RTG-Lab) of the Environmental Science and technology building (ICTA-UAB) located in the Universitat Autònoma de Barcelona campus (UTM: 42°29'24" E, 45°94'36" N). The irrigation system is hydroponic

on substrate with primary the use of rainwater. The 900m² rainwater harvesting system (RWHS) is included in the building structure as well as a 100m³ storage tank located underground from which the water for irrigation is pumped to the cropping sites. The building structure and its year-round production have been previously analyzed to identify the environmental impact reduction due to the connectivity and synergy between the greenhouse and the building (Sanjuan-Delmás et al., 2018). This building has two greenhouse laboratories for UA on the fourth floor, where this experiment was conducted. The beans were planted on the South- West facing laboratory (Urban Agriculture Laboratory 2) with a total area of 122.8m² as can be seen on the plant layout shown in Fig. 1.

Several sensors were used to monitor temperature (T107 Campbell Scientific) and relative humidity of the i-RTG cropping areas (Table 1 in the Supplementary information). Irrigation water, water drainage, electric conductivity and pH for each irrigation line were measured three times a week.

2.2. Plant materials and growth conditions

The seedlings were obtained from a nursery, where the seeds were inoculated with the rhizobium mix and transported to the i-RTG 10 days after planting. The production system is soilless with perlite substrate and nutrient solution given through the 2 L/h drip irrigation system. The cropping area was arranged in twelve rows with four 40 L perlite bags each (Fig. 1). Four bean plants were planted in each 1 m long perlite bag, making a total of 192 plants, divided in three treatments (64 per treatment). The plantation frame was 0.125m² within a total cropping area of 84.6m². The irrigation was set 4 times a day for 3 min giving a total amount of 400 mL per day for each plant.

Two experiments were performed. The first experiment took place in 2019, starting on the 16th of January and ending on the 10th of April. The second experiment took place in 2020, and the plants were transplanted on the 13th of February and the experiment was finalized on the 7th of May, lasting each one a total of 84 days. We used different concentrations of struvite during the two experiments to determine the N and P assimilation rates and how the yield was affected. In 2019, the bean plants were treated with 2 g of struvite (0.25 g of P; 0.114 g of N) per plant (SR2) and 5 g of struvite (0.625 g of P; 0.285 g of N) per plant (SR5). In 2020, we incremented the amount of struvite to 10 g

(1.25 g of P; 0.57 g of N) per plant (SR10), and 20 g of struvite (2.5 g of P; 1.14 g of N) per plant (SR20). The inoculation was made prior to their sowing, embedding the bean seeds in the commercial liquid rhizobium mixture before planting. We performed a control experiment both years parallel to the treatments under the same temporal and climatic conditions but fertilized with a full nutrient solution, with zero struvite and without inoculation. These control treatments were also made with the same crop and during the same time, lasting 84 days in the rooftop greenhouse as well.

The struvite granules were placed close to the root area to ensure a better absorption by the plant. To avoid possible runoff of struvite granules a 1 L bag with small holes for water drainage, was placed around the root area to retain the crystalline granules close to the plant. The granulated urine derived struvite was given directly to the plants rhizosphere after transplanting them into the integrated greenhouse.

2.2.1. Commercial inorganic fertilizer

Two nutrient solutions (Table 1) were made for both campaigns, one standard full nutrient solution (NS) with nitrogen, phosphorous and magnesium and a second solution deficient in nitrogen, phosphorous and magnesium with a higher content in K₂SO₄ to avoid potassium as a limiting factor. All nutrients were mixed into a concentrated solution stored in 50 L tanks, further diluted with rainwater when irrigated in a ratio of 1:100 (NS:Rainwater).

2.2.2. Commercial Rhizobium inoculant

The inoculant used for this experiment was obtained through a company based in Karlsruhe, Germany, nadicom GmbH. This 1 L liquid product contained a mixture of two rhizobia strains, *Rhizobium phaseoli* and *Rhizobium giardinii*, that were directly applied on the bean seeds (except for the Control) before planting and again 5 days after transplanting to the ICTA RTG- Lab. The manufacture and transport of this commercial product was not included in the LCA as an input for our alternatively fertilized crops, since the production impact has been considered minimal.

2.2.3. Commercial struvite

The struvite used for the experiment was urine derived, obtained from a wastewater treatment plant (WWTP) in Denmark. The plant recovers struvite from the digestate flow through the addition of reagents to reach stoichiometric levels that trigger struvite precipitation. The




Fig. 1. Experimental layout of the experiment in the i-RTG.

Table 1

Nutrient solution content, Control NS as the full NS for all control plants and the Mg, P, N-free NS for all treated plants with rhizobia inoculation and struvite. *Hortrilon: micronutrients in the form of chelates: 5.0% iron (Fe); 2.5% manganese (Mn); 2.5% copper (Cu); 0.6% zinc (Zn); 0.5% boron (B); 0.5% molybdenum (Mo); **sequestrene: addition of iron in the form of 2,4% - iron (Fe) chelated; 2,4% - iron (Fe), chelated by [o,o] EDDHA; 1,2% p/p to avoid ferric chlorosis.

Nutrients applied	Control NS	Mg, P, N-FREE NS
KPO ₄ H ₂	136 mg/L	–
KNO ₃	101 mg/L	–
K ₂ SO ₄	304.5 mg/L	435 mg/L
Ca(NO ₃) ₂	410 mg/L	–
CaCl ₂	111 mg/L	111 mg/L
Mg(NO ₃) ₂	148.3 mg/L	–
Hortrilon*	0.1 mg/L	0.1 mg/L
Sequestrene**	0.1 mg/L	0.1 mg/L

obtained struvite (Mg(NH₄)PO₄·6H₂O) has a composition conformed by 12.5% w/w phosphorous; 5.7% w/w nitrogen and 9.9% w/w magnesium and a granule size of 1 to 3 mm. The heavy metal content in struvite from different origins and production systems has been analyzed and set at levels under the European threshold, also ranging far below the amount of possible impurities that can be found with the production of phosphate rock as well as untreated sewage sludge from WWTP (Bastida et al., 2019).

2.3. Experimental analyses and nutrient balances

Water samples were taken from each irrigation system as well as the drained water 3 times a week. Production of the bean plants was counted and weighted. The amount of drained leachates were measured daily and sampled three times a week. The concentrations of Cl[−], NO₂[−], NO₃[−], PO₄^{3−}, SO₄^{2−}, Ca²⁺, K⁺ and Mg²⁺ were measured using ionic chromatography. Additionally, the pH and EC were measured daily for both the nutrient solutions and leachate water. To reduce the possible error generated through the irrigation and sampling the generated data was adjusted to a curve. The incoming and outgoing nutrients were quantified as well as the nutrients found in the plant biomass and beans. The plant biomass was collected at the end of the experiment with a sample number of 8 plants per treatment. These samples were dried and weighted before being digested with a Single Reaction Chamber microwave with concentrated HNO₃. The digested samples were then analyzed using Optical Spectrometry (ICP-OES). The same procedure was applied to the obtained production of beans, sampled throughout the experiment. The final balance per plant was assessed with the following equation:

$$Fns + Fs + Ffix = Fp + Fb + Fpl + Fl \quad (1)$$

Equation 1: Fns = g nutrients in nutrient solution, Fs = g in struvite, Ffix = g nutrients, Fp = g nutrients in production, Fb = g nutrients in biomass, Fpl = g nutrients in perlite, Fl = g nutrients in leachates.

To calculate the fraction needed per plant to close the balance, the following equation was used (Eq. (2)):

$$\text{Balance\%} = 100 \cdot \frac{Fp}{(Fns + Fs + Ffix)} + \frac{Fb}{(Fns + Fs + Ffix)} + \frac{Fpl}{(Fns + Fs + Ffix)} + \frac{Fl}{(Fns + Fs + Ffix)} \quad (2)$$

Equation 2: Fp = g nutrients in production, Fb = g nutrients in biomass, Fpl = g nutrients in perlite, Fl = g nutrients in leachates, Fns = g nutrients in nutrient solution, Fs = g in struvite, Ffix = g nutrients obtained through N₂ fixation.

The following results depict the data collected in 2019 for plant biomass, irrigation and leachate nutrient content as well as yield production and nutrient content. The 2020 study was included to provide

further information on the effect of greater struvite quantities to increase the yield. Therefore, the LCA results for 2020 only defer from the 2019 inventory in the amount of struvite used as well as the yield.

Additionally an analysis to calculate the fraction of N in the biomass obtained from N₂ fixation was made, using an elemental analyzer-isotopic ratio mass spectrometer (EA-IRMS; Thermo Fisher Scientific), attaining the δ¹⁵N values (in ‰) for our treatments SR2, SR5 and control as well as our alternative fertilizer struvite which was set in 7.1‰. Contributions from each source (atmospheric or struvite) were then calculated with the following equation (Shearer and Kohl, 1993; Unkovich et al., 2002; Arndt et al., 2004), using the lowest δ¹⁵N value obtained as our 'B' value (−1.16‰) (Shearer and Kohl, 1989; Peoples et al., 2002; Kermah et al., 2018):

$$\%Ndfa = \frac{\delta^{15}N_{\text{Source 2}} - \delta^{15}N_{\text{Sink}}}{\delta^{15}N_{\text{Source 2}} - 'B' \text{ value}} \times 100 \quad (3)$$

Equation 3: %Ndfa (Nitrogen derived from N₂ fixation from the atmosphere), δ¹⁵N Source 2 (‰) corresponds to the δ¹⁵N value of struvite, δ¹⁵N Sink (‰) corresponds to the δ¹⁵N value from the sample, 'B' value set at −1.16‰.

2.4. Life cycle assessment (LCA)

The LCA is a tool with a standardized methodology (ISO 2006) used to determine the environmental performance of goods in all stages of their life cycle of the four proposed treatments (SR2, SR5, SR10 and SR20) and control. The scope of the LCA study is cradle to gate of the bean production system. The functional unit (FU) chosen is 1 kg of fresh beans at the collection point. The cut-off method in the Simapro software was applied which allocates the benefit of the recycled materials to the recycled products. To calculate the life cycle environmental impacts of the treatment, we used the Simapro software and the Ecoinvent 3.5 attributional database. The following impact categories (IC) were selected, all from the ReCiPe (H) Midpoint method: Global warming (GW), Terrestrial acidification (TA), Freshwater eutrophication (FE), Marine Eutrophication (ME), Fossil Resource Scarcity (FRS) and Ecotoxicity (ET), which is the sum of Freshwater, Marine and Terrestrial ecotoxicities.

The system definition is illustrated by Fig. 2, which differentiates between the subsystems infrastructure, operation and end of life. For infrastructure, we considered the production and end of life of the greenhouse, the rainwater harvesting system and the auxiliary equipment such as pumps and fertirrigation installed and all the transportation required. All steps shown in Fig. 2 for raw material extraction, processing, transport to construction site, construction/maintenance, as well as the transport to the landfill or recycling site were considered. On the operational side, the study includes the production, use and end of life, including transport, of all the resources required for the duration of the experiments (perlite substrate, fertilizers, struvite, pesticides, water, and energy). Exceptions to this are the production of nursery plants and the composting of the residual biomass as well as the rhizobium production. For the production of struvite the additional inputs for controlled precipitation were accounted, in this case the chemical inputs can be seen in the LCA inventory, which consisted in MgCl₂, Energy and NaOH for 1 kg of struvite as described by the technology developed by Ostara® (Amann et al., 2018). The used wastewater for the struvite precipitation was not considered within the system boundaries for this study.

For the end of life subsystem of our production several assumptions were made. The remaining biomass generated in the greenhouse goes to composting as well as the used substrate after 5 years of use. The composting of the residual biomass was not considered within the system boundaries. The leachate water was discharged into the urban water cycle entering the wastewater treatment plant. All phosphates and nitrates discharged into the water are therefore considered direct




Fig. 2. Representation of the LCA system boundaries of the present experiment for fresh bean production. Division between the operational system, the infrastructure and the end of life subsystems. LCA inventory represented on the left and the LCA stages on the right. The system boundary defined by the dotted line delimiting within the accounted materials and stages in this study.

emissions to water, in the case of the treatments fertilized with struvite, also as direct emissions to the air. For the system infrastructure it was considered that the RWHS as well as the auxiliary equipment were assumed to be disposed of into the landfill. The distance to the landfill and recycling site were assumed to be 30 km from the greenhouse.

The inventory data for the infrastructure and auxiliary equipment was compiled from Sanyé et al., 2012; Sanyé Mengual, 2015; Sanjuan-Delmás et al., 2018; Rufi-Salís et al., 2020c. For both the rainwater harvesting system (RWHS) and the i-RTG System a lifespan of 50 years was considered while the auxiliary equipment was set at 10 years, taking into account previous work by Sanjuan-Delmás et al., 2018 and Rufi-Salís et al., 2020c. Emission factors for N to air were calculated according to Llorach-Massana et al., 2017, while N and P in water were directly measured. The emissions to air in struvite were calculated taking into account the emission factor of the total nitrogen in the applied quantity of struvite, even when not all struvite was dissolved.

For the transportation of all materials average values for the transport to markets were given. The transport to the i-RTG was then added with a distance of 50 km for all pesticides, fertilizers and auxiliary equipment as well as the struvite and rhizobium applied. Transport distance of 850 km was applied for the substrate bags following the methodology of Sanjuan-Delmás et al., 2018. No transport of the horticultural production was considered since one of the benefits of urban agriculture is the on-site selling of the products, therefore the product procurement by the consumer is located outside our system boundaries.

The data for the operation was collected during the experiment, including the amount of fertilizers, the substrate used as well as the energy used to work the irrigation system. The energy used during the campaign was estimated by the water pumps and amount of water pumped to the greenhouse and crops.

The full inventory is available in Supporting information (Tables 3 to 7 from the Supplementary information).

3. Results

3.1. Yields and nutrient balance

The total and average-per-plant production for both campaigns (2019 and 2020) can be seen in Table 2 for each treatment. The results show that as struvite concentration is increased from SR2 to SR20, production also increases from 1899.2 g to 4821.5 g indicating a significant

assimilation rate of P by the plants. It is crucial to point out that the two controls differ greatly as well from one campaign to the other. Since the campaign of 2020 began in February and the 2019 campaign in January, we can consider the climatic conditions as an explanation for greater productions, taking into account that minimal temperatures were higher in 2020 (by more than 6 °C), as well as the average temperature throughout the experiment (Table 1 Supplementary information). These conditions would enable a greater and faster production of flowers and an earlier bean growth. The first bean pod harvest made in 2019 was 49 days after transplanting the bean plants into the greenhouse while in 2020 the bean pod harvest began 39 days after transplanting (Fig. 1 in the Supplementary information). Due to the different climatic conditions and resulting productions the control treatment for both campaigns has to be considered as a reference. The observed productions do not reach more than 60% of the achieved production in the control treatment, staying between 40 and 60%, in both campaigns.

3.2. Nutrient fluxes

The obtained nutrient balance can be seen in Table 3. Here the nutrient content for nitrogen, phosphorous, magnesium, potassium, sulfate and calcium in the incoming nutrient solution can be observed, as well as the outgoing fluxes of production (bean pods), biomass (leaves, stems, roots), perlite and leachates. In the case of nitrogen the fixated N₂ is also taken into account as seen in Fig. 3. Table 2 in the Supplementary information also provides some incoming (struvite input) and outgoing flows (leachates) for the SR10 and SR20 treatments for P, N and Mg.

Table 2

Production for campaigns 2019 and 2020 for all treatments. *Difference to control = percentage in relation to control as 100%.

Production Treatment	2019			2020		
	SR2	SR5	Control 1	SR10	SR20	Control2
Total	1899.2 g	2375.6 g	4726.7 g	3542.2 g	4821.5 g	8198.4 g
Average per plant	59.3 g	74.2 g	147.7 g	110.7 g	150.6 g	256.2 g
Dif to control*	40.2%	50.3%	100%	43.2%	58.8%	100%

Table 3

Nutrient balance per plant for N, P, Mg, K, S and Ca F_p = g nutrients in production, F_b = g nutrients in biomass, F_{pl} = g nutrients in perlite, F_l = g nutrients in leachates, F_{ns} = g nutrients in nutrient solution, F_s = g in struvite, F_{fix} = g nutrients obtained through N_2 fixation *perlite was obtained from Sanjuan-Delmas.

Nutrient	Treatment	Nutrient solution + struvite (F_{ns} + F_s)	Atm N_2 Fix (F_{fix})	Production (F_p)		Biomass (F_b)		Perlite* (F_{pl})	Leachates (F_l)		Balance
		g	g	g	%	g	%	%	g	%	%
N	SR2	0.114	0.293	0.152	37%	0.204	50%	0%	0.031	8%	95%
	SR5	0.285	0.301	0.159	27%	0.262	45%	0%	0.052	9%	81%
	Control	1.271	0.123	0.376	27%	0.375	27%	6%	0.323	23%	83%
P	SR2	0.25	0	0.021	8%	0.036	14%	0%	0.041	17%	40%
	SR5	0.625	0	0.030	5%	0.057	9%	0%	0.055	9%	23%
	Control	0.740	0	0.081	11%	0.109	15%	6%	0.274	37%	69%
Mg	SR2	0.198	0	0.011	6%	0.020	10%	0%	0.096	49%	64%
	SR5	0.495	0	0.016	3%	0.050	10%	0%	0.133	27%	40%
	Control	0.461	0	0.032	7%	0.060	13%	0%	0.292	63%	83%
K	SR2	6.357	0	0.173	3%	0.270	4%	0%	4.249	67%	74%
	SR5	6.357	0	0.206	3%	0.357	6%	0%	4.128	65%	74%
	Control	4.774	0	0.576	12%	0.739	15%	0%	3.664	77%	104%
S	SR2	1.626	0	0.009	1%	0.030	2%	0%	1.420	87%	90%
	SR5	1.626	0	0.011	1%	0.040	2%	0%	1.405	86%	90%
	Control	1.611	0	0.030	2%	0.060	4%	0%	1.040	65%	70%
Ca	SR2	1.713	0	0.021	1%	0.150	9%	3%	1.126	66%	79%
	SR5	1.713	0	0.025	1%	0.195	11%	3%	1.374	80%	96%
	Control	1.719	0	0.061	4%	0.475	28%	3%	1.127	66%	100%

The obtained results for the fixed g of N where achieved thanks to previous studies with isotopic N_{15} analyses where the percentages of fixation for SR2, SR5 and the 2019 control where obtained. The average percentage of fixed nitrogen for SR2 was 82% while for SR5 it was 72% and 16% for the control of the total N found in production and biomass. This fixed nitrogen was further given as an additional inflow.

The balance for nutrients N, H, K, and Ca is close to 100%, indicating that the inflows can be traced almost entirely in the different outflows. On the other hand, there are losses of P and Mg in the SR2 and SR5 treatments, which are unaccounted for in the mass balance showing percentages under 50% reaching values as low as 23% for the P balance in SR5. We consider that the reason for such low accounting of the balance for P and Mg is the possibility of them remaining undissolved in the perlite bag. While Sanjuan-Delmas provides information of the amounts of these nutrients in fertigation that can be found in the perlite, the remaining P, Mg and N from struvite left in the perlite bag was not determined. This factor can generate uncertainty in our nutrient balance which has to be taken into account. When observing the percentages of out coming P and Mg in treatments SR10 and SR20 (Table 2 in the Supplementary information) we can observe a trend in the leached amount being around 8 to 11% for SR10 and SR20 respectively. These leachate percentages close to the observed in the SR2 and SR5 treatments further support the idea of an uncompleted dissolution in the perlite bag. On the other hand the N balance seems to fit the incoming and out coming flows, while the amount of the given N with the struvite is respectively lower to Mg and P and the atmospheric N_2 fixation also amounts to additional nitrogen in plant biomass and production. When regarding the percentage of discarded P and N into the leachates we can observe a reduction of almost threefold in N when comparing the control (23%) to both treatments (SR2 with 8% and SR5 with 9%) and double (SR2 with 17%) or even fourfold (SR5 with 9%) in P compared to the control (37%). When regarding the quantity of P in the leachate water of the control treatment and the SR20 treatment we can see that it is almost identical, pointing out that the given quantity of struvite has achieved flows to waste water similar to the control treatment. On the other hand, when considering the percentages in regard to the total amount given to the plant the control treatment has leached more than threefold the added P (37%) compared to the SR20 treatment (11%). When observing the N incoming flows the input given for the control treatment in the irrigation is similar to the quantity given, in form of struvite, in the SR20 treatment. We must bear in mind that for the treatment SR20 the quantity of N gained from atmospheric fixation is not known. On the other hand the leached outflow for the

SR20 accounts for about a 13%, almost half of the leached amount in the control (23%). The lower quantities of P and N measured in water, although similar or greater N and P quantities were given to the control treatment, indicate a much slower dissolution of this fertilizer, reinforcing the idea of a remaining undissolved amount of struvite in the perlite bag.

3.3. Environmental performance of the treatments

The LCA impacts per functional unit (FU) are disaggregated into the life cycle stages that resulted in the highest impacts for all four treatments and controls, as shown in Fig. 4. Since the controls resulted in higher yields, consequently the impacts are reduced considerably in all categories.

Within each impact category we can clearly state that the greenhouse structure and the rainwater harvesting system account for most of the generated impact especially in GW, TA, ME and FRS. This can be due to the large transport distances, the processing and construction of larger amounts of materials like aluminum and steel.

While the auxiliary equipment and fertilizers seem to have lower impacts in most categories, the implication of the latter in the ME, FE and TA categories is of great importance for the control treatments where a full nutrient solution was used. Even when the emissions to air and water of struvite were taken into account, the reduction in these categories for treatments SR2 and SR5 is especially clear.

While the higher production in the control treatment reduces impacts of the RTG- infrastructure and RWHS, the impact generated by the fertilizers is still greater in all IC for the control despite the higher yields, only being surpassed in the treatment SR20 for FE. The percentage contribution of the accounted system stages can be further observed in Fig. 5. The reduction of the impact generated by the alternative fertilizer can be seen when comparing the smaller percentages for fertilization in the treatments SR2, SR5 SR10 and SR20 to the control treatments 1 and 2. For the treatment SR20 we can again observe an increase of the impact in the FE category.

When adding the percentages corresponding to the infrastructure (RTG-structure, RWHS, auxiliary equipment) and operation (energy, fertilizers, pesticides, and substrate) stages of production for each impact category, a shift of the weight of impact contribution with the alternative fertilization can be seen (Table 4).

The change in the fertilization mainly generates a shift in the eutrophication impact categories (FE and ME) which reach up to more than 55% of the total impact of the operation in the control treatment while




Fig. 3. Nutrient flow representation for N (left) and P (right). Fp = g nutrients in production, Fb = g nutrients in biomass, Fpl = g nutrients in perlite, Fl = g nutrients in leachates, Fns = g nutrients in nutrient solution, Fs = g in struvite, Ffix = g nutrients obtained through N_2 fixation.

staying under 30% in both treatment SR2 and SR5. It is also worth mentioning that overall the change in the fertilization has an effect on all categories, shifting the weight of the impact from the operational phase to the infrastructure when comparing the control to all other treatments.

Due to the great percentage taken up by the greenhouse structure and rainwater harvesting system, the contribution of the operational side of the bean production is overshadowed. Since the infrastructure remains the same for all treatments and is highly specific to this particular site, it was excluded from consideration in Fig. 6 for a better exploration of the effects of the substituting fertilizer (Fig. 2 Supplementary information for Environmental performance of the operation system in percentage per IC).

When observing Fig. 6 the applied fertilization appears to be the main cause for emissions in all IC in the operation subsystem. While yield is smaller in all four struvite and rhizobium treatments, emissions

remain lower than controls 1 and 2 in categories GW and AT. In the case of FE, ME and ET emissions from SR2 and SR5 remain below the control 1 treatment while being higher for the treatments SR10 and SR20. In the case of fossil resource scarcity it is worth mentioning that the reason for a higher emission in these two categories is not bound to fertilization but due to an increase of the weight of substrate and energy in the operation impact.

While emissions are mostly lower in the four alternative fertilizer treatments, especially in SR20 we can observe an increase with greater amounts of struvite in both categories FE and ME. To further understand the changes in emissions bound to fertilization, Fig. 7 depicts all accounted factors considered in the LCA for the fertilization.

When observing the fertilization emission in Fig. 7, great impact reductions are made due to the reduced emissions to air and water (as seen in TA, GW, FE and ME) and transport of fertilizers (as seen in ET,




Fig. 4. Total System and operation impacts in relation to the functional unit.




Fig. 5. Impact of system components (%).

FRS and again GW). Here we can appreciate that the increase of impact seen for treatment SR20 achieved in categories FE and ME is due to water emissions increase due to greater struvite quantities.

4. Discussion

The life cycle assessment performed on the bean production experiments in soilless substrate fertilized with struvite and rhizobium has shown that there are significant benefits in terms of eutrophication. These findings confirm studies of other authors such as (Sanjuan-Delmás et al., 2018; Rufi-Salís et al., 2020b) in which fertilization has been deemed as a major contributor to the environmental footprint of urban agriculture. However, because the yield is lower than that of the conventional mineral fertilizer, the impacts associated to the infrastructure required for the fertilizer substitution increase. Fertilization has shown to be of great importance in the impacts regarding our bean production. The sole removal of the nitrogen, phosphorous and magnesium

from the fertigation has shifted the weight of the emissions from the operational part to the infrastructure in a drastic manner. These emission reductions not only affected the expected IC (ME and FE) but all due to their transport (for GW, ET and FRS) and the emissions to Air (for TA) as seen in Fig. 7. While this information depicts great flaws in the implementation of such production systems it also gives a great chance for improvement. While the application of struvite has shown to fulfill the entire cycle of the crop with some yield reduction its production and transport do not affect the given IC to a greater extent except for the treatment SR20. The accumulation of all three fertilizers (P, N and Mg) in one and the possibility of its local generation and application can considerably improve the operational footprint of our agricultural systems. While the struvite recovery technology developed by Ostara® and used for this experiment requires inputs of MgCl, energy and NaOH, has been seen to have lower environmental impacts compared to other processes (Rufi-Salís et al., 2020a) further advancements are being made on the use of saltwater, to further reduce the use of chemical

Table 4

Emission origin in our experiment from Infrastructure or Operation of the System in each impact category (IC).

IC	SR2		SR5		Control 1	
	Infrastructure	Operation	Infrastructure	Operation	Infrastructure	Operation
GW	96%	4%	95%	5%	82%	18%
TA	90%	10%	90%	10%	59%	41%
FE	76%	24%	73%	27%	45%	55%
ME	90%	10%	85%	15%	51%	49%
ET	91%	9%	91%	9%	79%	21%
FRS	95%	5%	95%	5%	89%	11%
IC	SR10		SR20		Control 2	
	Infrastructure	Operation	Infrastructure	Operation	Infrastructure	Operation
GW	95%	5%	95%	5%	82%	18%
TA	89%	11%	88%	12%	59%	41%
FE	66%	34%	47%	53%	45%	55%
ME	80%	20%	70%	30%	51%	49%
ET	89%	11%	89%	11%	79%	21%
FRS	95%	5%	95%	5%	89%	11%




Fig. 6. Impact of the production operation in relation to the functional unit.

resources and potentially lowering its environmental footprint (Martínez-Blanco et al., 2014; Hasler et al., 2015; Amann et al., 2018).

On the other hand, as we have seen in the nutrient fluxes for the SR2 and SR5 treatments, the percentages of the balances for P and Mg remain low as well as the SR10 and SR20 leachate fluxes. The previously described slow dissolution of the struvite fertilizer (Bhuiyan et al., 2007; Degryse et al., 2017) has been identified as the reason for the lower balance percentages, leaving struvite in the bag that has still not been diluted. This dissolution could be remedied with a lower pH in

the irrigation as well as an increase of the irrigation points inside the bag. The location of the struvite itself with regard to the root area has also been regarded as relevant for its plant uptake (Degryse et al., 2017). On the other hand the remains of struvite inside the bag can favor the reuse and recycling of the perlite bag for a less P and Mg demanding crop in the case of treatments SR2 and even a second production of beans like in the case of SR5. A second bean production without the addition of struvite would even further reduce the needed inputs and its operational footprint.




Fig. 7. Impact associated to the treatment fertilization in relation to the functional unit.




Fig. 8. Yield reduction (in %) impact on IC (in %) for infrastructure and operation systems.

The use of alternative fertilizers like struvite avoids the consumption of mineral or synthetic fertilizers (Lam et al., 2020) described as fertilizer offset accounting. The environmental benefit of the use of struvite should not only be accounted in the moment of its use (emissions to air and water) and transport but in the avoided production of N, P, and Mg fertilizer. Even further, the environmental benefit of the removal of these nutrients from urban waterbodies should be taken into account as well. As described before, the generation of struvite has requirements but removes potential water and air emissions from WWTP (Ishii and Boyer, 2015; Igos et al., 2017; Lam et al., 2020). While this last benefit has not been taken into account for this study, the further use of struvite as fertilizer and the consequent fertilization offset accounting have, and can be well observed in these results with the emission reductions in almost all IC.

The yield reduction in all treatments compared to the respective controls has a great impact on the production footprint. While a higher production has been reported with a greater application of struvite, a limitation to reach greater yields still remains. Plausible explanations

for these losses are the reduced struvite dissolution (Degryse et al., 2017), the higher P requirement due to the rhizobia symbiosis (Long, 1989; Olivera et al., 2004) or a possible electrochemical imbalance causing a reduced uptake of cations in the root area as described by Kontopoulou et al. (2015).

While the yield reduction remains unclear, its impact on the environmental performance is quick to be identified in the obtained results. The loss of production increases the environmental footprint of production, reaching higher emissions than the control treatments (especially when infrastructure is considered), even with lower total emissions of P and N to water in all treatments (Table 3 in the Supplementary information).

A higher yield without the additional use of these fertilizers would decrease the impact of our production, which begs the question as up to what yield loss percentage we can afford and still remain more sustainable than the control treatment.

To answer this question the control (2019) yield was regarded as our hypothetical 100% yield and therefore a scenario with 0% yield




Fig. 9. Yield reduction (in %) impact on IC (in %) for operation system.

reduction. From here on the emissions for all IC were calculated with a yield loss of 10% to 60%. The values used for the 60% and 50% yield loss were directly taken from the SR2 and SR5 treatments respectively, since the obtained yields corresponded to the simulated yield losses. The emissions were also derived from the SR2 and SR5 treatments, since the emissions to water from treatments SR10 and SR20 were deemed as too similar to the control with no major yield increase (in comparison to the respective control treatment).

Figs. 8 and 9 depict these scenarios for the yield reduction impact in infrastructure and operation and only operation respectively. The control treatment line in both Figs. 8 and 9 is where the baseline from the control treatment was set. Above this control line the emissions are increased with regards to the control (in %), while below this line the impacts are decreased. When considering the infrastructure and operation (Fig. 8) we can observe that no yield can be lost in order to bring all IC under the control treatment line.

While TA, FE and ME are well below the baseline with a yield reduction of 30% other IC like GW and ET start to decrease at 10% yield loss and below. In the case of Fig. 9, when infrastructure is not considered, a 50% yield loss can occur and still decrease all emissions in all IC.

The importance of reducing mineral or synthetic fertilizer to avoid emission in all IC has been regarded throughout the experiment, although the importance of maintaining production levels while using alternative fertilization methods has been laid out clearly. While impact categories like FE and ME are greatly decreased, the capacity to affect other categories in a significant way can only be achieved with low yield reductions. Especially when considering urban agriculture, the production system might entail more complex infrastructure (rooftop greenhouse, indoor agriculture), leaving reduced margins of yield loss.

The slow dissolution of struvite and the feasibility of its use in soil-less agriculture make this fertilizer a good candidate to avoid further P extraction and loss. Due to the uncertainty of the estimation of available mineral P in the next centuries, new ways of nutrient recovery need to be considered in our immediate future (Alewell et al., 2020). This work has demonstrated that the emissions to water, especially for P can be reduced in comparison to conventional fertilization methods. When talking about the extraction of these nutrients to produce struvite further efforts should be made to make this process possible in wastewater treatment plants, reducing transport emissions of agricultural fertilizers to surrounding urban and agricultural areas. Emissions related to transportation of said minerals can be ultimately reduced as well as avoiding emission and loss of nutrients into urban water streams in WWTP (Carpenter and Bennett, 2011; Harder et al., 2019). Ultimately, the capacity to recover P nutrients in local scales can reduce agricultural pressures to obtain said fertilizer in due to its distribution or market instability and increasing prices (Kataki et al., 2016; Alewell et al., 2020).

5. Conclusions

The present work mainly aimed to analyze the environmental impact of a bean production with the use of alternative fertilizing methods of struvite and the inoculation of rhizobium bacteria. It also aimed to study the feasibility of production with the mentioned change in fertilization, exploring different potential yield losses to gain a broader view of the possibilities this methodology presents. To this purpose, two experiments with different struvite quantities were made and a quantification of the environmental impact using life cycle assessment as our tool. Three main conclusions can be drawn from this study:

Firstly, the total reduction of nitrogen and phosphorous mineral and synthetic fertilizers for vegetable production has been shown to be viable with the use of the recycled slow-releasing fertilizer struvite and the bacterial inoculation with rhizobium strains. Although a yield reduction in all cases was observed compromising its efficiency to reduce the environmental impact in all IC except FE and ME.

Secondly, the use of struvite and rhizobium inoculation reduced emissions in all IC mainly due to transport and emissions to air and

water due to a slower dissolution in the substrate. The struvite, being available in all WWTP installations can be obtained with no great environmental cost in its operation while reducing transport and extraction of three separate minerals (N, P and Mg).

Thirdly, the complexity of the infrastructure and operational inputs will increase the environmental impact in all IC, as well as the yield loss. Only the reduction of yield loss up to 0% can equal the environmental impact of the control treatment in all selected categories when the infrastructure is considered. Without the infrastructure the margins for yield loss can range up to 50% staying below the control treatment. Therefore we consider crucial to reduce infrastructure complexity in the prospect of urban agriculture as well as the reduction of mineral and synthetic fertilizers to truly reduce potential environmental impacts.

CRediT authorship contribution statement

Verónica Arcas-Pilz: Conceptualization, Methodology, Formal analysis, Data curation, Writing – original draft. **Martí Rufi-Salís:** Methodology, Data curation, Writing – review & editing. **Felipe Parada:** Methodology, Writing – review & editing. **Xavier Gabarrell:** Conceptualization, Methodology, Writing – review & editing, Supervision. **Gara Villalba:** Conceptualization, Methodology, Writing – review & editing, Supervision, Funding acquisition.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors are grateful to the Spanish Ministry of Economy, Industry and Competitiveness (Spain) for the grant awarded to V. Arcas-Pilz (FPI-MINECO 2018); to the Universitat Autònoma de Barcelona for awarding a research scholarship to M. Rufi-Salís (PIF-UAB 2017) and to the National Commission for Scientific and Technological Research (Chile) for the grant awarded to F. Parada (PFCHA-CONICYT 2018 – Folio 72180248). The research leading to these results has received funding from the European Research Council under the European Horizon2020 research and innovation program under grant agreement No 862663 (FoodE) as well as ERC grant agreement n° 818002 URBAG, awarded to Gara Villalba.

This work was partially supported by the Spanish Ministry of Economy, Industry and Competitiveness (AEU/FEDER) [CTM2016-75772-C3-1-R] and the “María de Maeztu” program for Units of Excellence in R&D (CEX2019-000940-M).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2020.144744>.

References

- Ackerman, J.N., et al., 2013. Evaluation of manure-derived struvite as a phosphorus source for canola. *Can. J. Plant Sci.* 93 (3), 419–424. <https://doi.org/10.4141/cjps2012-207>.
- Ahmed, N., et al., 2018. Struvite recovered from various types of wastewaters: characteristics, soil leaching behaviour, and plant growth. *Land Degrad. Dev.* 29 (9), 2864–2879. <https://doi.org/10.1002/ldr.3010>.
- Alewell, C., et al., 2020. Global phosphorus shortage will be aggravated by soil erosion. *Nat. Commun.* <https://doi.org/10.1038/s41467-020-18326-7>.
- Amann, A., et al., 2018. Environmental impacts of phosphorus recovery from municipal wastewater. *Resour. Conserv. Recycl.* <https://doi.org/10.1016/j.resconrec.2017.11.002>.
- Araujo, J., Urbano, B., González-Andrés, F., 2020. Comparative environmental life cycle and agronomic performance assessments of nitrogen fixing rhizobia and mineral nitrogen fertiliser applications for pulses in the Caribbean region. *J. Clean. Prod.* 267, 122065. <https://doi.org/10.1016/j.jclepro.2020.122065>.

- Arndt, S.K., et al., 2004. Nitrogen fixation and metabolism by groundwater-dependent perennial plants in a hyperarid desert. *Oecologia* 141 (3), 385–394. <https://doi.org/10.1007/s00442-004-1655-7>.
- Bastida, F., et al., 2019. The effects of struvite and sewage sludge on plant yield and the microbial community of a semiarid Mediterranean soil. *Geoderma*. <https://doi.org/10.1016/j.geoderma.2018.10.046>.
- Bhuiyan, M.I.H., Mavinic, D.S., Beckie, R.D., 2007. A solubility and thermodynamic study of struvite. *Environ. Technol.* 28 (9), 1015–1026. <https://doi.org/10.1080/0959332080618857>.
- Carpenter, S.R., Bennett, E.M., 2011. Reconsideration of the planetary boundary for phosphorus. *Environ. Res. Lett.* <https://doi.org/10.1088/1748-9326/6/1/014009>.
- Degryse, F., Baird, R., da Silva, Rodrigo C., et al., 2017. Dissolution rate and agronomic effectiveness of struvite fertilizers – effect of soil pH, granulation and base excess. *Plant Soil* 410 (1), 139–152. <https://doi.org/10.1007/s11104-016-2990-2>.
- Fisher, R.F., Long, S.R., 1992. Rhizobium-plant signal exchange. *Nature* 357 (6380), 655–660. <https://doi.org/10.1038/357655a0>.
- Gopalakrishnan, S. et al. (2015) 'Plant growth promoting rhizobia: challenges and opportunities', 3 Biotech. Springer Berlin Heidelberg, 5(4), pp. 355–377. doi: <https://doi.org/10.1007/s13205-014-0241-x>.
- Harder, R., et al., 2019. 'Recycling nutrients contained in human excreta to agriculture: pathways, processes, and products', Critical Reviews in Environmental Science and Technology. Taylor & Francis 49 (8), 695–743. <https://doi.org/10.1080/10643389.2018.1558889>.
- Hasler, K. et al. (2015) 'Life cycle assessment (LCA) of different fertilizer product types', European Journal of Agronomy. Elsevier B.V., 69, pp. 41–51. doi: <https://doi.org/10.1016/j.eja.2015.06.001>.
- Igos, E., et al., 2017. Assessment of environmental impacts and operational costs of the implementation of an innovative source-separated urine treatment. *Water Res.* 126, 50–59. <https://doi.org/10.1016/j.watres.2017.09.016>.
- Ishii, S.K.L., Boyer, T.H., 2015. 'Life cycle comparison of centralized wastewater treatment and urine source separation with struvite precipitation: focus on urine nutrient management', Water Research. Elsevier Ltd 79, 88–103. <https://doi.org/10.1016/j.watres.2015.04.010>.
- Karak, T. and Bhattacharyya, P. (2011) 'Human urine as a source of alternative natural fertilizer in agriculture: a flight of fancy or an achievable reality', Resources, Conservation and Recycling. Elsevier B.V., 55(4), pp. 400–408. doi: <https://doi.org/10.1016/j.resconrec.2010.12.008>.
- Kataki, S. et al. (2016) 'Phosphorus recovery as struvite: recent concerns for use of seed, alternative Mg source, nitrogen conservation and fertilizer potential', Resources, Conservation and Recycling. Elsevier B.V., 107, pp. 142–156. doi: <https://doi.org/10.1016/j.resconrec.2015.12.009>.
- Kermah, M., et al., 2018. 'N₂-fixation and N contribution by grain legumes under different soil fertility status and cropping systems in the Guinea savanna of northern Ghana', Agriculture, Ecosystems and Environment. Elsevier 261 (September 2017), 201–210. <https://doi.org/10.1016/j.agee.2017.08.028>.
- Kontopoulou, C. K. et al. (2015) 'Responses of hydroponically grown common bean fed with nitrogen-free nutrient solution to root inoculation with N-fixing bacteria', HortScience, 50(4), pp. 597–602. doi: 10.21273/HORTSCI.50.4.597.
- Kontopoulou, C.K., et al., 2017. Impact of rhizobial inoculation and reduced N supply on biomass production and biological N₂ fixation in common bean grown hydroponically. *J. Sci. Food Agric.* 97 (13), 4353–4361. <https://doi.org/10.1002/jsfa.8202>.
- Lam, K.L., Zlatanović, L., van der Hoek, J.P., 2020. Life cycle assessment of nutrient recycling from wastewater: a critical review. *Water Res.* 173, 115519. <https://doi.org/10.1016/j.watres.2020.115519>.
- Lewis, W.M., Wurtsbaugh, W.A., Paerl, H.W., 2011. Rationale for control of anthropogenic nitrogen and phosphorus to reduce eutrophication of inland waters. *Environ. Sci. Technol.* 45 (24), 10300–10305. <https://doi.org/10.1021/es202401p>.
- Linderholm, K., Tillman, A. M. and Mattsson, J. E. (2012) 'Life cycle assessment of phosphorus alternatives for Swedish agriculture', Resources, Conservation and Recycling. Elsevier B.V., 66, pp. 27–39. doi: <https://doi.org/10.1016/j.resconrec.2012.04.006>.
- Llorach-Massana, P., et al., 2017. N₂O emissions from protected soilless crops for more precise food and urban agriculture life cycle assessments. *J. Clean. Prod.* 149, 1118–1126. <https://doi.org/10.1016/j.jclepro.2017.02.191>.
- Long, S.R., 1989. Rhizobium-legume nodulation: life together in the underground. *Cell* 56 (2), 203–214. [https://doi.org/10.1016/0092-8674\(89\)90893-3](https://doi.org/10.1016/0092-8674(89)90893-3).
- Martínez-Blanco, J., et al., 2014. 'Application challenges for the social life cycle assessment of fertilizers within life cycle sustainability assessment', Journal of Cleaner Production. Elsevier Ltd 69, 34–48. <https://doi.org/10.1016/j.jclepro.2014.01.044>.
- Massey, M.S., Davis, J.G., Sheffield, R.E., Ippolito, J. A. (2007) 'Struvite production from dairy wastewater and its potential as a fertilizer for organic production in calcareous soils', International Symposium on Air Quality and Waste Management for Agriculture, (September). Available at: <https://elibrary-asabe.org.ezproxy2.library.arizona.edu/azdez.asp?search=0&IID=1&AID=23823&CID=aqwm2007&v=&i=&t=2>.
- Olivera, M., et al., 2004. Growth, nitrogen fixation and ammonium assimilation in common bean (*Phaseolus vulgaris*): effect of phosphorus. *Physiol. Plant.* 121 (3), 498–505. <https://doi.org/10.1111/j.0031-9317.2004.00355.x>.
- Peoples, M.B., Boddey, R.M., Herridge, D.F., 2002. Quantification of Nitrogen Fixation. Nitrogen Fixation at the Millennium. Elsevier B.V. <https://doi.org/10.1016/b978-044450965-9/50013-6>.
- Rahman, M.M., et al., 2014. 'Production of slow release crystal fertilizer from wastewaters through struvite crystallization – a review', Arabian Journal of Chemistry. King Saud University 7 (1), 139–155. <https://doi.org/10.1016/j.arabjc.2013.10.007>.
- Rufi-Salís, M., Brunnhofer, N., et al. (2020a) 'Can wastewater feed cities? Determining the feasibility and environmental burdens of struvite recovery and reuse for urban regions', Science of the Total Environment. Elsevier B.V., 737, p. 139783. doi: <https://doi.org/10.1016/j.scitotenv.2020.139783>.
- Rufi-Salís, M., Calvo, M. J., et al. (2020b) 'Exploring nutrient recovery from hydroponics in urban agriculture: an environmental assessment', Resources, Conservation and Recycling. Elsevier, 155(November 2019), p. 104683. doi: <https://doi.org/10.1016/j.resconrec.2020.104683>.
- Rufi-Salís, M., Petit-Boix, A., et al., 2020c. Identifying eco-efficient year-round crop combinations for rooftop greenhouse agriculture. *International Journal of Life Cycle Assessment*. The International Journal of Life Cycle Assessment. doi: <https://doi.org/10.1007/s11367-019-01724-5>.
- Ryther, John H., Dunstan, W.M., 1971. Nitrogen, phosphorus, and eutrophication in the coastal marine environment. *Science* 171, 1008–1014.
- Sammauria, R., et al., 2020. Microbial inoculants: potential tool for sustainability of agricultural production systems. *Arch. Microbiol.* 202 (4), 677–693. <https://doi.org/10.1007/s00203-019-01795-w>.
- Sanjuan-Delmás, D., et al., 2018. Environmental assessment of an integrated rooftop greenhouse for food production in cities. *J. Clean. Prod.* 177, 326–337. <https://doi.org/10.1016/j.jclepro.2017.12.147>.
- Sanyal, D., Osorno, J.M., Chatterjee, A., 2020. Influence of rhizobium inoculation on dry bean yield and symbiotic nitrogen fixation potential. *J. Plant Nutr.* 43 (6), 798–810. <https://doi.org/10.1080/01904167.2020.1711946>.
- Sanyé Mengual, E., 2015. 'Sustainability Assessment of Urban Rooftop Farming Using an Interdisciplinary Approach', TDX (Tesis Doctorals en Xarxa). Universitat Autònoma de Barcelona Available at: <http://www.tesisenred.net/handle/10803/308336>. (Accessed 19 February 2019).
- Sanyé, E., et al., 2012. Life cycle assessment of energy flow and packaging use in food purchasing. *J. Clean. Prod.* <https://doi.org/10.1016/j.jclepro.2011.11.067>.
- Savvas, D. et al. (2018) 'Impact of different rhizobial strains and reduced nitrogen supply on growth, yield and nutrient uptake in cowpea grown hydroponically', Acta Horticulturae, (1227), pp. 417–424. doi: 10.17660/ActaHortic.2018.1227.52.
- Shaddel, S., et al., 2020. 'Struvite crystallization by using raw seawater: improving economics and environmental footprint while maintaining phosphorus recovery and product quality', Water Research. Elsevier Ltd 173, 115572. <https://doi.org/10.1016/j.watres.2020.115572>.
- Shearer, G. and Kohl, D. H. (1989) 'Natural ¹⁵N abundance as a method of estimating the contribution of biologically fixed nitrogen to N₂-fixing systems: potential for non-legumes', in Skinner, F. A., Boddey, R. M., and Fendrik, I. (eds) Nitrogen Fixation With Non-Legumes: The Fourth International Symposium on 'Nitrogen Fixation With Non-Legumes', Rio de Janeiro, 23–28 August 1987, Dordrecht: Springer Netherlands, pp. 289–299. doi: https://doi.org/10.1007/978-94-009-0889-5_33.
- Shearer, G., Kohl, D.H., 1993. 'Natural abundance of ¹⁵N: fractional contribution of two sources to a common sink and use of isotope discrimination', Nitrogen Isotope Techniques. Academic Press, pp., 89–125 <https://doi.org/10.1016/B978-0-08-092407-6.50009-2>.
- Siegner, A.B., Acey, C., Sowerwine, J., 2020. 'Producing urban agroecology in the East Bay: from soil health to community empowerment', Agroecology and Sustainable Food Systems. Taylor & Francis 44 (5), 566–593. <https://doi.org/10.1080/21683565.2019.1690615>.
- Talboys, P.J., et al., 2016. Struvite: a slow-release fertiliser for sustainable phosphorus management? *Plant Soil* 401 (1–2), 109–123. <https://doi.org/10.1007/s11104-015-2747-3>.
- Thompson, L., 2013. Field evaluation of the availability for corn and soybean of phosphorus recovered as struvite from corn fiber processing for bioenergy. Graduate Theses and Dissertations Available at: <http://lib.dr.iastate.edu/etd/13173>.
- Tornaghi, C., 2017. Urban agriculture in the food-Disabling City: (re)defining urban food justice, reimagining a politics of empowerment. *Antipode* 49 (3), 781–801. <https://doi.org/10.1111/anti.12291>.
- Unkovich, M. J. et al. (2002) 'Nitrogen isotope fractionation in the fodder tree tagasaste (*Chamaecytisus proliferus*) and assessment of N₂ fixation inputs in deep sandy soils of Western Australia', Functional Plant Biology, 27(10), p. 921. doi: <https://doi.org/10.1071/pp99201>.
- Uysal, A., et al., 2014. Optimization of struvite fertilizer formation from baker's yeast wastewater: growth and nutrition of maize and tomato plants. *Environ. Sci. Pollut. Res.* 21 (5), 3264–3274. <https://doi.org/10.1007/s11356-013-2285-6>.