

RESEARCH

Open Access



Minimal set of periods for continuous self-maps of the eight space

Jaume Llibre^{1*} and Ana Sá²

*Correspondence:

jlilibre@mat.uab.es

¹Departament de Matemàtiques,
Universitat Autònoma de Barcelona,
08193 Bellaterra, Barcelona, Spain
Full list of author information is
available at the end of the article

Abstract

Let G_k be a bouquet of circles, i.e., the quotient space of the interval $[0, k]$ obtained by identifying all points of integer coordinates to a single point, called the branching point of G_k . Thus, G_1 is the circle, G_2 is the eight space, and G_3 is the trefoil. Let $f : G_k \rightarrow G_k$ be a continuous map such that, for $k > 1$, the branching point is fixed.

If $\text{Per}(f)$ denotes the set of periods of f , the minimal set of periods of f , denoted by $\text{MPer}(f)$, is defined as $\bigcap_{g \sim f} \text{Per}(g)$ where $g : G_k \rightarrow G_k$ is homological to f .

The sets $\text{MPer}(f)$ are well known for circle maps. Here, we classify all the sets $\text{MPer}(f)$ for self-maps of the eight space.

MSC: 37E15

Keywords: Continuous maps; Periodic orbit; Period; Minimal period; The space in shape of eight

1 Introduction and statement of the results

In dynamical systems it is often the case that topological information can be used to study qualitative or quantitative properties of the system. This work deals with the problem of determining the set of periods of the periodic orbits of a map given the homology class of the map.

A *finite graph* (simply a *graph*) G is a topological space formed by a finite set of points V (points of V are called *vertices*) and a finite set of open arcs (called *edges*) in such a way that each open arc is attached by its endpoints to vertices. An open arc is a subset of G homeomorphic to the open interval $(0, 1)$. Note that a finite graph is compact since it is the union of a finite number of compact subsets (the closed edges and the vertices). Notice that a closed edge is homeomorphic either to the closed interval $[0, 1]$ or to the circle. It may be either connected or disconnected, and it may have isolated vertices.

The *valence* of a vertex is the number of edges with the vertex as an endpoint (where the closed edges homeomorphic to a circle are counted twice). The vertices with valence 1 of a connected graph are *endpoints* of the graph and the vertices with valence larger than 2 are *branching points*.

Suppose that $f : G \rightarrow G$ is a continuous map, in what follows a *graph map*. A *fixed point* of f is a point x in G such that $f(x) = x$. We will call x a *periodic point of period n* if x is a

© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

fixed point of f^n but it is not fixed by any f^k for $1 \leq k < n$. We denote by $\text{Per}(f)$ the set of natural numbers corresponding to periods of the periodic points of f .

Let G be a connected graph, and let f be a graph map. Then f induces endomorphisms $f_{*k} : H_k(G) \rightarrow H_k(G)$ for $k = 0, 1$ on the integral homology groups of G , where $H_0(G) \approx \mathbb{Z}$ (because G is connected), and $H_1(G) \approx \mathbb{Z} \oplus \cdots \oplus \mathbb{Z}$, where k is the number of independent circuits or loops of G as elements of $H_1(G)$. A *circuit* of G is a subset of G homeomorphic to the circle. The endomorphisms f_{*0} and f_{*1} are represented by integer matrices. Furthermore, since G is connected, f_{*0} is the identity.

The endomorphism f_{*1} will play the main role in our analysis of the minimal sets of periods for graph maps on G . In what follows f_{*1} will be denoted by f_* . For example, if $H_1(G) \approx \mathbb{Z} \oplus \mathbb{Z}$ and

$$f_* = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

this means that the graph G has two independent oriented circuits. Moreover, if the first circuit covers itself exactly a_1 times following the same orientation (not necessarily in a consecutive way) and exactly a_2 times following the converse orientation (not necessarily in a consecutive way), then $a = a_1 - a_2$. Similarly, if the first circuit covers the second one exactly b_1 times following the same orientation (not necessarily in a consecutive way) and exactly b_2 times following the converse orientation (not necessarily in a consecutive way), then $b = b_1 - b_2$. An analogous explanation can be given with the second independent circuit and with b and d instead of a and c , respectively.

Let G_k be a *bouquet of k circles*, that is, the quotient space of $[0, k]$ obtained by identifying all points of integer coordinates to a single point. Notice that G_1 is the *circle* and that G_2 is usually called the *eight space*. For the G_k graph, we have $H_0(G_k) \approx \mathbb{Z}$, $H_1(G_k) \approx \mathbb{Z} \oplus \cdots \oplus \mathbb{Z}$, $f_{*0} \approx \text{id}$, and $f_{*1} = f_* = A$, where A is a $k \times k$ integral matrix. For more details on graph maps, see [16] or [18].

Our main goal is to study the set $\text{Per}(f)$ for graph maps. More explicitly, we want to provide a description of the minimal set of periods (see below) attained within the homology class of a given graph map. When the map $g : G \rightarrow G$ is homological to f (i.e., g induces the same endomorphisms as f on the homology groups of G), we shall write $g \simeq f$. We define the *minimal set of periods* of f to be the set

$$\text{MPer}(f) = \bigcap_{g \simeq f} \text{Per}(g).$$

From its definition $\text{MPer}(f)$ is the maximal subset of periods contained in $\text{Per}(g)$ for all $g \simeq f$.

Our main objective is to *characterize the minimal sets of periods* $\text{MPer}(f)$ for graph maps $f : G_i \rightarrow G_i$ with the branching point a fixed point for $i = 2, 3$. So, always $1 \in \text{MPer}(f)$. Even for circle maps $f : G_1 \rightarrow G_1$ the characterization of all minimal sets of periods $\text{MPer}(f)$ is interesting and nontrivial, see Theorem A. This result was stated by Efremova [12] and Block et al. [9] without giving a complete proof. As far as we know, the first complete proof was given in [4].

We denote by $\mathbb{N}$ the set of all natural numbers and by $k\mathbb{N}$ the set $\{kl : l \in \mathbb{N}\}$.

Theorem A Let $f : G_1 \rightarrow G_1$ be a circle map such that the endomorphism induced by f on the first homology group is $f_* = (d)$ (i.e., d is the degree of f). Then the following statements hold:

- (a) If $d \notin \{-2, -1, 0, 1\}$, then $\text{MPer}(f) = \mathbb{N}$.
- (b) If $d = -2$, then $\text{MPer}(f) = \mathbb{N} \setminus \{2\}$.
- (c) If $d \in \{-1, 0\}$, then $\text{MPer}(f) = \{1\}$.
- (d) If $d = 1$, then $\text{MPer}(f) = \emptyset$.

In the next theorem we characterize the minimal sets of periods for eight maps, i.e., for continuous maps $f : G_2 \rightarrow G_2$.

Theorem B Let $f : G_2 \rightarrow G_2$ be an eight map such that

$$f_* = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

Suppose that the branching point is a fixed point. Then the following statements hold:

- (a) If $\{a, d\} \not\subset \{-2, -1, 0, 1\}$, then $\text{MPer}(f) = \mathbb{N}$.
- (b) If $-2 \in \{a, d\}$ and $\{a, d\} \subset \{-2, -1, 0, 1\}$, then

$$\text{MPer}(f) = \begin{cases} \mathbb{N} \setminus \{2\} & \text{if } bc = 0, \\ \mathbb{N} & \text{if } bc \neq 0. \end{cases}$$

- (c) Assume that $\{a, d\} \subset \{-1, 0, 1\}$.

- (c1) If $|a| + |d| = 2$, then

$$\text{MPer}(f) = \begin{cases} \{1\} & \text{if } bc = 0, \\ \mathbb{N} \setminus \{2\} & \text{if } bc = 1, \\ \mathbb{N} & \text{if } bc = -1 \text{ or } |bc| > 1. \end{cases}$$

- (c2) If $|a| + |d| = 1$ and

- (c21) $a = 1, d = 0$, then

$$\text{MPer}(f) = \begin{cases} \{1\} & \text{if } bc = 0, \\ \mathbb{N} \setminus \{2\} & \text{if } (b, c) \in R, \\ \mathbb{N} & \text{otherwise;} \end{cases}$$

where $R = \{(1, 1), (-1, -1), (1, 2), (-1, -2)\}$.

- (c22) $a = 0, d = 1$, then it follows (c21) interchanging b and c .

- (c23) $a = -1, d = 0$, then

$$\text{MPer}(f) = \begin{cases} \{1\} & \text{if } bc = 0, \\ \mathbb{N} \setminus \{2\} & \text{if } (b, c) \in R, \\ \mathbb{N} \setminus \{3\} & \text{if } bc = -1, \\ \mathbb{N} & \text{otherwise.} \end{cases}$$

- (c24) $a = 0, d = -1$, then it follows (c23) interchanging b and c .

(c3) If $|a| + |d| = 0$, then

$$\text{MPer}(f) = \begin{cases} \{1\} & \text{if } bc = 0 \text{ or } bc = 1, \\ \{1, 2\} & \text{if } bc = -1, \\ \{1\} \cup (2\mathbb{N} \setminus \{2\}) & \text{if } bc = 2, \\ \{1\} \cup (2\mathbb{N} \setminus \{4\}) & \text{if } bc = -2, \\ \{1\} \cup 2\mathbb{N} & \text{if } |bc| > 2. \end{cases}$$

We remark that Theorem B implies Theorem A if f has a fixed point by choosing, for instance, $a = b = c = 0$.

The study of the minimal set of periods of a homotopy class of maps instead of its homology class is the main objective of the fixed point theory, see for instance the books of Brown [10], Jiang [13], and Kiang [15]. Other extensions from circle maps to n -dimensional torus have been done in [2] and [14], and from circle maps to transversal n -sphere maps in [11]. Some different results on the periods of graph maps have been given in [1, 3, 5–8, 16, 17].

Finally, we recall that the classification of the dynamics on the graph maps helps the classification of the homeomorphisms on the surfaces, see for instance [19] and the references quoted therein.

This work is organized as follows. How to obtain a given period for a graph map by using the notion of f -covering is described in Sect. 2. The proof of Theorem B is given in Sect. 3.

2 Periods and f -covering

Let $f : G \rightarrow G$ be a graph map and $x \in G$ be a periodic point of period n . The set $\{x, f(x), \dots, f^{n-1}(x)\}$ is called the *periodic orbit* of x .

A set $I \subset G$ will be called an *interval* if there is a homeomorphism $h : J \rightarrow I$, where J is $[0, 1]$, $(0, 1]$, $[0, 1)$, or $(0, 1)$. The set $h((0, 1))$ will be called the *interior* of I . If $J = [0, 1]$, the interval I will be called *closed*; if $J = (0, 1)$, it will be called *open*. Notice that it may happen that the above terminology does not coincide with the one used when we think about I as a subset of G (the same applies to the edges of G). For example, if $G = I = [0, 1]$ and $h = \text{identity}$, then for I regarded as a subset of the topological space G , I is both open and closed, and the interior of I is I .

Let C_i and C_j be two circuits of G_k . A closed interval $I = [u, v]$ is *basic* if $I \subset C_i$, $f(I) = C_j$, where $\{i, j\} \subset \{1, 2, \dots, k\}$, $f(u) = f(v) = p$, where p is the branching point of G_k , and there is no other closed interval $K \subsetneq I$ such that $f(K) = C_j$. Of course the definition of basic interval also applies to the particular case that $[u, v] = C_i$. Let I and J be two basic intervals, $K \subset I$, $L \subset J$ be two subintervals. Then we say that K *f -covers* L , and we write $K \rightarrow L$ if there exists a closed subinterval M of K such that $f(M) = L$. If $L = J = C_j$, we say that $K = I$ *f -covers* L because either $f(K) = L$ or $K = I = C_i$ and $f(K) = L$ by the definition of basic intervals.

Lemma 2.1 *Suppose that $I_1, I_2, \dots, I_n$ are intervals such that $I_1 \rightarrow I_2 \rightarrow \dots \rightarrow I_n \rightarrow I_1$ with I_1 different from a circuit. Then there is a fixed point z of f^n such that $z \in I_1, f(z) \in I_2, \dots, f^{n-1}(z) \in I_n$.*

Proof Since $I_n \rightarrow I_1$ and I_1 is not a circuit, there is a closed interval $J_n \subset I_n$ such that $f(J_n) = I_1$. Similarly, there are closed intervals or circuits $J_1, \dots, J_{n-1}$ such that, for each $k = 1, \dots, n -$

$1, J_k \subset I_k$ and $f(J_k) = J_{k+1}$. It follows that $f^n(J_1) = I_1$ and since $J_1 \subset I_1$ and I_1 is not a circuit, by Bolzano's theorem f^n has a fixed point $z \in J_1$. Clearly, $z \in I_1, f(z) \in I_2, \dots, f^{n-1}(z) \in I_n$. $\square$

A sequence of the form $I_1 \rightarrow I_2 \rightarrow \dots \rightarrow I_n \rightarrow I_1$ is called a *loop of length n* . Let $I_1 \rightarrow I_2 \rightarrow \dots \rightarrow I_n \rightarrow I_1$ and $J_1 \rightarrow J_2 \rightarrow \dots \rightarrow J_m \rightarrow J_1$ be two loops such that $I_1 = J_1$. We define the *concatenation* of these two loops as the loop $I_1 \rightarrow I_2 \rightarrow \dots \rightarrow I_n \rightarrow I_1 \rightarrow J_2 \rightarrow \dots \rightarrow J_m \rightarrow I_1$. We say that a loop is an m -repetition, $m \geq 2$, of a given loop if it is the concatenation of that loop with itself m times. We say that a loop is *non-repetitive* if it is not an m -repetition of any of its subloops with $m \geq 2$.

In what follows, a G_k -map f is a continuous map $f: G_k \rightarrow G_k$ such that $f(p) = p$, where p is the branching point of G_k .

Proposition 2.2 *Let f be a G_k -map. Suppose that f has two intervals I_1 and I_2 such that $\text{Int}(I_1) \cap \text{Int}(I_2) = \emptyset$ and $I_1 \cap I_2$ has no fixed points. If f has the subgraph $\odot I_1 \rightleftharpoons I_2 \odot$, then $\text{Per}(f) = \mathbb{N}$.*

Proof Clearly, since $p \notin I_1 \cap I_2$, at least one of the intervals, I_1 or I_2 , is not a circuit. Without loss of generality, we assume that I_1 is not a circuit. We consider the non-repetitive loop $I_1 \rightarrow I_2 \rightarrow I_1 \rightarrow \dots \rightarrow I_1$ of length $n \geq 2$. Since $\text{Int}(I_1) \cap \text{Int}(I_2) = \emptyset$ and $I_1 \cap I_2$ has no fixed points, by Lemma 2.1, there is a periodic point z of f with period $n \geq 2$. That is, $\text{Per}(f) = \mathbb{N}$. $\square$

In what follows when we say “we have two intervals A and B ” we are really saying that we have two different intervals A and B . We remark that if we have two basic intervals I_1 and I_2 such that $p \notin I_1 \cap I_2$, then they satisfy the assumptions of Proposition 2.2.

Proposition 2.3 *Let f be a G_k -map. Suppose that f has three intervals I_1, I_2 , and I_3 such that $\text{Int}(I_i) \cap \text{Int}(I_j) = \emptyset$ for all $i \neq j$ and $I_i \cap I_j$ has no fixed points for some $i \neq j$. If f has the subgraph $\odot I_1 \rightarrow I_2 \rightarrow I_3 \rightarrow I_1$, then $\text{Per}(f) \supset \mathbb{N} \setminus \{2\}$. Moreover, if $I_2 \cap I_3 = \emptyset$ and $I_3 \rightarrow I_2$, then $2 \in \text{Per}(f)$.*

Proof We consider the non-repetitive loop $I_1 \rightarrow I_2 \rightarrow I_3 \rightarrow I_1 \rightarrow \dots \rightarrow I_1$ of length $n \geq 3$. Since $\text{Int}(I_i) \cap \text{Int}(I_j) = \emptyset$ for all $i \neq j$ and $I_i \cap I_j$ has no fixed points for some $i \neq j$, by Lemma 2.1, there is a periodic point z of f with period $n \geq 3$. Therefore, $\text{Per}(f) \supset \mathbb{N} \setminus \{2\}$.

We suppose now that $I_2 \cap I_3 = \emptyset$ and $I_3 \rightarrow I_2$. We consider the non-repetitive loop $I_2 \rightarrow I_3 \rightarrow I_2$ of length 2. By Lemma 2.1 there is a periodic point z of f with period 2. $\square$

We remark that if we have three basic intervals I_1, I_2 , and I_3 such that $p \notin I_i$ for some $i \in \{1, 2, 3\}$, then we are in the assumptions of Proposition 2.3.

3 The eight

In this section we shall prove Theorem B. The two circuits of G_2 are denoted by C_1 and C_2 . If f_* is given as in Theorem B, we consider that the circuit C_1 covers itself $|a|$ times and it covers C_2 $|c|$ times. Similarly for the circuit C_2 .

Proof of statement (a) of Theorem B Suppose that $\{a, d\} \not\subset \{-2, -1, 0, 1\}$.

Case 1: Assume that $\{a, d\} \not\subset \{-2, -1, 0, 1, 2\}$. Without loss of generality, we may assume that $|a| \geq 3$. From the graph of f (see for instance Fig. 1), it is clear that there are two basic





Figure 1 Examples of maps with $\{a, d\} \not\subset \{-2, -1, 0, 1, 2\}$

Figure 2 $I_1 = [p, a_1]$, $I_2 = [b_0, b_1]$, and $I_3 = [a_1, a_2]$



intervals I_1 and I_2 in C_1 such that $p \notin I_1 \cap I_2$ and f has the subgraph of Proposition 2.2, so $\text{Per}(f) = \mathbb{N}$. That is, $\text{MPer}(f) = \mathbb{N}$.


Case 2: Suppose that $2 \in \{a, d\}$ and $\{a, d\} \subset \{-2, -1, 0, 1, 2\}$. Without loss of generality, we may assume that $a = 2$.

Since $a = 2$, this means that f has at least two basic intervals I_1 and I_2 in C_1 such that f has the subgraph of Proposition 2.2. If $p \notin I_1 \cap I_2$, then, by Proposition 2.2, $\text{Per}(f) = \mathbb{N}$. But not always I_1 and I_2 satisfy that $p \notin I_1 \cap I_2$. In this case let p and a_0 be the endpoints of I_1 , b_0 and p be the endpoints of I_2 (see for instance Fig. 2).

We establish an ordering in the intervals I_1 and I_2 in such a way that p is the smallest element of I_1 and the greatest of I_2 . Set $I_1 = [p, a_0]$ and $I_2 = [b_0, p]$. Notice that we may have $a_0 = b_0$. Consider the subset $(f|_{I_1})^{-1}(a_0)$ of C_1 . Let a_1 be the infimum of the points in $(f|_{I_1})^{-1}(a_0)$. Consider the subset $(f|_{I_2})^{-1}(a_0)$ of C_1 and choose b_1 to be the infimum of the points in $(f|_{I_2})^{-1}(a_0)$. Set $I_{11} = [p, a_1]$, $I_{12} = [a_1, a_0]$, and $I_{21} = [b_0, b_1]$. Now we take the interval $I_{13} = [a_1, a_2]$, where a_2 denotes the infimum of the points in the subset $(f|_{I_{12}})^{-1}(b_1)$ of C_1 . Then f has the subgraph $\odot I_{11} \rightarrow I_{13} \rightrightarrows I_{21} \rightarrow I_{11}$. Since $I_{21} \cap I_{13} = \emptyset$, by Proposition 2.3, $n \in \text{Per}(f)$ for all $n \geq 1$. Therefore, $\text{MPer}(f) = \mathbb{N}$. This proves statement (a). $\square$

Proof of statement (b) of Theorem B Suppose that $-2 \in \{a, d\}$ and $\{a, d\} \subset \{-2, -1, 0, 1\}$. Without loss of generality, we may assume that $a = -2$.

First we suppose that $bc \neq 0$. We always have four basic intervals I_1, I_2, I_3 , and I_4 , $I_1, I_2, I_3 \subset C_1$, and $I_4 \subset C_2$ such that either $p \notin I_1 \cap I_3$ or $I_2 \cap I_4 = \emptyset$ and f has the subgraph



(see for instance Fig. 3).

If $p \notin I_1 \cap I_3$, by Proposition 2.2, $\text{Per}(f) = \mathbb{N}$. If $I_2 \cap I_4 = \emptyset$, by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$. Therefore, if $bc \neq 0$, $\text{MPer}(f) = \mathbb{N}$.

We suppose now that $bc = 0$. As it can be deduced from the examples of Fig. 4, $2 \notin \text{MPer}(f)$.

Since $a = -2$, this means that f has at least two basic intervals I_1 and I_2 in C_1 such that f has the subgraph of Proposition 2.2. If $p \notin I_1 \cap I_2$, then by Proposition 2.2 $\text{Per}(f) = \mathbb{N}$. But not always $p \notin I_1 \cap I_2$. In this case let p and a_0 be the endpoints of I_1 , b_0 and p be the endpoints of I_2 (see for instance Fig. 5). We consider an ordering in the intervals I_1 and I_2 in such a way that p is the smallest element of I_1 and the greatest of I_2 . Write $I_1 = [p, a_0]$ and $I_2 = [b_0, p]$. Notice that we may have $a_0 = b_0$. Consider the subsets $(f|I_1)^{-1}(a_0)$ and $(f|I_2)^{-1}(a_0)$ of C_1 . Let a_1 be the infimum of the points in $(f|I_1)^{-1}(a_0)$ and b_1 be the infimum

(a) $a = -2, d = 0, bc = 1$ (b) $a = -2, d = 1, bc = 1$ (c) $a = -2, d = -1, bc = -1$




Figure 3 Examples of maps with $a = -2$ and $bc \neq 0$






Figure 4 Examples of maps with $a = -2, d \in \{-2, -1, 0, 1\}, bc = 0$, and $2 \notin \text{Per}(f)$

Figure 5 $I_{11} = [p, a_1]$, $I_{12} = [a_1, a_0]$, and $I_{21} = [b_1, p]$ **Figure 6** Examples of maps with $\{a, d\} \subset \{-1, 0, 1\}$, $|a| + |d| = 2$, and $bc = 0$ 

of the points in $(f|_{I_2})^{-1}(a_0)$. Set $I_{11} = [p, a_1]$, $I_{12} = [a_1, a_0]$, and $I_{21} = [b_1, p]$. Then f has the subgraph $\odot_{I_{12}} \rightarrow I_{11} \rightarrow I_{21} \rightarrow I_{12}$. Since we are in the assumptions of Proposition 2.3, $n \in \text{Per}(f)$ for all $n \neq 2$. Therefore, $\text{MPer}(f) = \mathbb{N} \setminus \{2\}$. This proves statement (b). $\square$

Proof of statement (c1) of Theorem B Suppose that $\{a, d\} \subset \{-1, 0, 1\}$ and $|a| + |d| = 2$. We consider first the case $bc = 0$. Without loss of generality, we may assume that $c = 0$. From the examples of Fig. 6 it is clear that $n \notin \text{MPer}(f)$ for any $n \in \mathbb{N}$ larger than 1, so $\text{MPer}(f) = \{1\}$ since the branching is fixed.

We assume now that $|bc| > 1$. From the graph of f (see for instance Fig. 7) it is easy to see that we always have three basic intervals I_1, I_2 , and I_3 , with $I_1, I_2 \subset C_1$ and $I_3 \subset C_2$ such that $p \notin I_i$ for some $i \in \{1, 2, 3\}$ and f has the subgraph of Proposition 2.3, so $\text{Per}(f) \supset \mathbb{N} \setminus \{2\}$. Now we will prove that $2 \in \text{MPer}(f)$.

If $\{b, c\} \not\subset \{-2, -1, 1, 2\}$, that is, if either $|b| \geq 3$ or $|c| \geq 3$, we can choose I_2 in one circuit and I_3 in the other circuit in such a way that $I_2 \cap I_3 = \emptyset$ (see (a), (b), and (c) of Fig. 7) and $I_3 \rightarrow I_2$. By Proposition 2.3, $2 \in \text{Per}(f)$. If $\{b, c\} \subset \{-2, -1, 1, 2\}$, in general there do not exist two basic intervals I_i and I_j , $I_i \neq I_j$, such that $p \notin I_i \cap I_j$ and $I_i \rightrightarrows I_j$ (see (e) and (f) of Fig. 7). If they exist, then by Lemma 2.1, considering the non-repetitive loop $I_i \rightarrow I_j \rightarrow I_i$, there is a periodic point z of f with period 2. If they do not exist, we shall find two intervals with empty intersection such that one f -covers the other.

We suppose first that $|bc| = 2$. We may assume, without loss of generality, that $|b| = 1$ and $|c| = 2$. We know that f has five basic intervals I_1, I_2, I_3, I_4 , and I_5 , the first three in C_1 and the other two in C_2 , such that $f(I_2) = f(I_3) = f(I_5) = C_2$ and $f(I_1) = f(I_4) = C_1$. Let p and a_0 be the endpoints of I_2 , a_0 and a_1 be the endpoints of I_1 , a_1 and p be the endpoints of I_3 (see for instance Fig. 8).

We consider an ordering in the intervals I_1, I_2 , and I_3 in such a way that p is the smallest element of I_2 and the greatest of I_3 . Set $I_2 = [p, a_0]$, $I_1 = [a_0, a_1]$, and $I_3 = [b_0, p]$. We have two possibilities for the interval I_4 : either $I_4 = [p, b_0]$ or $I_4 = [b_0, p]$. If $I_4 = [p, b_0]$ and $b = 1$, let b_1 be the supremum of the points in $(f|_{I_4})^{-1}(a_1)$ and $I_{42} = [b_1, b_0]$. We have $I_{42} \rightrightarrows I_3$ and




Figure 7 Examples of maps with $\{a, d\} \subset \{-1, 0, 1\}$, $|a| + |d| = 2$, and $|bc| > 1$




Figure 8 Examples of maps with $\{a, d\} \subset \{-1, 0, 1\}$, $|a| + |d| = 2$ and $|bc| = 2$

$I_{4_2} \cap I_3 = \emptyset$, so, by Lemma 2.1, $2 \in \text{Per}(f)$. If $I_4 = [p, b_0]$ and $b = -1$, set $b_1 = \sup\{(f|_{I_4})^{-1}(a_0)\}$ and $I_{4_2} = [b_1, b_0]$. Then $I_{4_2} \rightleftharpoons I_2$ and $I_{4_2} \cap I_2 = \emptyset$ so, by Lemma 2.1, $2 \in \text{Per}(f)$. If $I_4 = [b_0, p]$ and $b = 1$, write $b_1 = \inf\{(f|_{I_4})^{-1}(a_0)\}$ and $I_{4_1} = [b_0, b_1]$. Then $I_{4_1} \rightleftharpoons I_2$ and $I_{4_1} \cap I_2 = \emptyset$, so, by




Figure 9 Examples of maps with $\{a, d\} \subset \{-1, 0, 1\}$, $|a| + |d| = 2$, and $|bc| = 4$




Figure 10 Examples of maps with $\{a, d\} \subset \{-1, 0, 1\}$, $|a| + |d| = 2$, $bc = 1$, and $2 \notin \text{Per}(f)$

Lemma 2.1, $2 \in \text{Per}(f)$). If $I_4 = [b_0, p]$ and $b = -1$, take $b_1 = \inf\{(f|_{I_4})^{-1}(a_1)\}$ and $I_{4_1} = [b_0, b_1]$. Then $I_{4_1} \rightleftharpoons I_3$ and $I_{4_1} \cap I_3 = \emptyset$, so, by **Lemma 2.1**, $2 \in \text{Per}(f)$.

Suppose now that $|bc| = 4$. We know that f has six basic intervals I_1, I_2, I_3, I_4, I_5 , and I_6 , the first three in C_1 and the other three in C_2 , such that $f(I_2) = f(I_3) = f(I_5) = C_2$ and $f(I_1) = f(I_4) = f(I_6) = C_1$ (see for instance Fig. 9). Using the same ordering as the above set $I_2 = [p, a_0]$, $I_1 = [a_0, a_1]$, $I_3 = [b_0, p]$, $I_4 = [p, b_0]$, $I_5 = [b_0, b_1]$, and $I_6 = [b_1, p]$. If $b = 2$, set $b_2 = \inf\{(f|_{I_6})^{-1}(a_0)\}$ and $I_{6_1} = [b_1, b_2]$. Then $I_{6_1} \rightleftharpoons I_2$ and $I_{6_1} \cap I_2 = \emptyset$ so, by **Lemma 2.1**, $2 \in \text{Per}(f)$. If $b = -2$, write $b_2 = \inf\{(f|_{I_6})^{-1}(a_1)\}$ and $I_{6_1} = [b_1, b_2]$. Then $I_{6_1} \rightleftharpoons I_3$ and $I_{6_1} \cap I_3 = \emptyset$ so, by **Lemma 2.1**, $2 \in \text{Per}(f)$. Therefore, if $|bc| > 1$, $\text{MPer}(f) = \mathbb{N}$.

We suppose that $|bc| = 1$. We assume that $\mathbf{b} = \mathbf{c} = \mathbf{1}$. As it can be seen from examples (a), (c), and (e) of Fig. 10, $2 \notin \text{MPer}(f)$. Now we will prove that $\text{Per}(f) = \mathbb{N} \setminus \{2\}$.

We know that f has four basic intervals I_1, I_2, I_3 , and I_4 , the first two in C_1 and the other two in C_2 , such that $f(I_1) = f(I_3) = C_1$ and $f(I_2) = f(I_4) = C_2$. We have four possibilities for these intervals. Let $a_0 \in I_1 \cap I_2$ and $b_0 \in I_3 \cap I_4$ (see for instance Fig. 11). First, we take




Figure 11 Examples of maps with $\{a, d\} \subset \{-1, 0, 1\}$, $|a| + |d| = 2$, and $b = c = 1$

the interval I_3 to be $[p, b_0]$. Set $I_{3_2} = [b_1, b_0]$ where $b_1 = \sup\{(f|_{I_3})^{-1}(a_0)\}$. If $I_1 = [p, a_0]$, then f has the subgraph $\odot I_4 \rightarrow I_{3_2} \rightarrow I_2 \rightarrow I_4$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N} \setminus \{2\}$. If $I_1 = [a_0, p]$, then f has the subgraph $\odot I_1 \rightarrow I_2 \rightarrow I_{3_2} \rightarrow I_1$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N} \setminus \{2\}$.

Now we take the interval I_3 to be $[b_0, p]$. Set $b_1 = \inf\{(f|_{I_3})^{-1}(a_0)\}$ and $I_{3_1} = [b_0, b_1]$. If $I_1 = [p, a_0]$, then f has the subgraph $\odot I_1 \rightarrow I_2 \rightarrow I_{3_1} \rightarrow I_1$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N} \setminus \{2\}$. If $I_1 = [a_0, p]$, then f has the subgraph $\odot I_4 \rightarrow I_{3_1} \rightarrow I_2 \rightarrow I_4$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N} \setminus \{2\}$. Therefore, if $|a| = |d| = 1$ and $b = c = 1$, then $\text{MPer}(f) = \mathbb{N} \setminus \{2\}$.

We assume now that $\mathbf{b} = \mathbf{c} = -1$. As it can be seen from examples (b), (d), and (f) of Fig. 10, $2 \notin \text{MPer}(f)$. Now we will prove that $\text{Per}(f) = \mathbb{N} \setminus \{2\}$.

We know that f has four basic intervals I_1, I_2, I_3 , and I_4 , the first two in C_1 and the other two in C_2 , such that $f(I_1) = f(I_3) = C_1$ and $f(I_2) = f(I_4) = C_2$. We have four possibilities for these intervals. Let $a_0 \in I_1 \cap I_2$ and $b_0 \in I_3 \cap I_4$ (see for instance Fig. 12). First we take I_3 to be the interval $[p, b_0]$. Consider $b_1 = \sup\{(f|_{I_3})^{-1}(a_0)\}$ and $I_{3_2} = [b_1, b_0]$. If $I_1 = [p, a_0]$, then f has the subgraph $\odot I_1 \rightarrow I_2 \rightarrow I_{3_2} \rightarrow I_1$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N} \setminus \{2\}$. If $I_1 = [a_0, p]$, then f has the subgraph $\odot I_4 \rightarrow I_{3_2} \rightarrow I_2 \rightarrow I_4$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N} \setminus \{2\}$.

If $I_3 = [b_0, p]$, consider $b_1 = \inf\{(f|_{I_3})^{-1}(a_0)\}$ and $I_{3_1} = [b_0, b_1]$. If $I_1 = [p, a_0]$, then f has the subgraph $\odot I_4 \rightarrow I_{3_1} \rightarrow I_2 \rightarrow I_4$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N} \setminus \{2\}$. If $I_1 = [a_0, p]$, then f has the subgraph $\odot I_1 \rightarrow I_2 \rightarrow I_{3_1} \rightarrow I_1$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N} \setminus \{2\}$. Therefore, if $b = c = -1$, then $\text{MPer}(f) = \mathbb{N} \setminus \{2\}$. Hence, if $|a| + |d| = 2$ and $bc = 1$, $\text{MPer}(f) = \mathbb{N} \setminus \{2\}$.




Figure 12 Examples of maps with $\{a, d\} \subset \{-1, 0, 1\}$, $|a| + |d| = 2$, and $b = c = -1$

We consider now the case $\mathbf{b} = -1$ and $\mathbf{c} = 1$. We know that f has four basic intervals I_1 , I_2 , I_3 , and I_4 , the first two in C_1 and the other two in C_2 , such that $f(I_1) = f(I_3) = C_1$ and $f(I_2) = f(I_4) = C_2$. We have four possibilities for these intervals. Let $a_0 \in I_1 \cap I_2$ and $b_0 \in I_3 \cap I_4$ (see for instance Fig. 13). We suppose first that $I_2 = [a_0, p]$. If $I_3 = [p, b_0]$, choose $a_1 = \inf\{(f|_{I_2})^{-1}(b_0)\}$ and set $I_{2_1} = [a_0, a_1]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_1} \rightleftharpoons I_3 \rightarrow I_1$ with $I_3 \cap I_{2_1} = \emptyset$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$. If $I_3 = [b_0, p]$, denote $b_1 = \inf\{(f|_{I_3})^{-1}(a_0)\}$ and $I_{3_1} = [b_0, b_1]$. Then f has the subgraph $\odot I_4 \rightarrow I_{3_1} \rightleftharpoons I_2 \rightarrow I_4$ with $I_2 \cap I_{3_1} = \emptyset$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$.

We consider now $I_2 = [p, a_0]$. If $I_3 = [p, b_0]$, set $b_1 = \sup\{(f|_{I_3})^{-1}(a_0)\}$ and $I_{3_2} = [b_1, b_0]$. Then f has the subgraph $\odot I_4 \rightarrow I_{3_2} \rightleftharpoons I_2 \rightarrow I_4$ with $I_2 \cap I_{3_2} = \emptyset$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$. If $I_3 = [b_0, p]$, write $a_1 = \sup\{(f|_{I_2})^{-1}(b_0)\}$ and $I_{2_2} = [a_1, a_0]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_2} \rightleftharpoons I_3 \rightarrow I_1$ with $I_3 \cap I_{2_2} = \emptyset$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$. Therefore, if $b = -1$ and $c = 1$, then $\text{MPer}(f) = \mathbb{N}$.

We consider now the case $\mathbf{b} = 1$ and $\mathbf{c} = -1$. We know that f has four basic intervals I_1 , I_2 , I_3 , and I_4 , the first two in C_1 and the other two in C_2 , such that $f(I_1) = f(I_3) = C_1$ and $f(I_2) = f(I_4) = C_2$. We have again four possibilities for these intervals. Let $a_0 \in I_1 \cap I_2$ and $b_0 \in I_3 \cap I_4$ (see for instance Fig. 14). We take the interval I_2 to be $[a_0, p]$. If $I_3 = [p, b_0]$, define $b_1 = \sup\{(f|_{I_3})^{-1}(a_0)\}$ and $I_{3_2} = [b_1, b_0]$. It follows that f has the subgraph $\odot I_4 \rightarrow I_{3_2} \rightleftharpoons I_2 \rightarrow I_4$ with $I_2 \cap I_{3_2} = \emptyset$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$. If $I_3 = [b_0, p]$, consider $a_1 = \inf\{(f|_{I_2})^{-1}(b_0)\}$ and $I_{2_1} = [a_0, a_1]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_1} \rightleftharpoons I_3 \rightarrow I_1$ with $I_3 \cap I_{2_1} = \emptyset$, and we get, by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$.

Suppose that $I_2 = [p, a_0]$. If $I_3 = [p, b_0]$, set $a_1 = \sup\{(f|_{I_2})^{-1}(b_0)\}$ and $I_{2_2} = [a_1, a_0]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_2} \rightleftharpoons I_3 \rightarrow I_1$ with $I_3 \cap I_{2_2} = \emptyset$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$.




Figure 13 Examples of maps with $\{a, d\} \subset \{-1, 0, 1\}$, $|a| + |d| = 2$, $b = -1$, and $c = 1$




Figure 14 Examples of maps with $\{a, d\} \subset \{-1, 0, 1\}$, $|a| + |d| = 2$, $b = 1$, and $c = -1$

If $I_3 = [b_0, p]$, consider $b_1 = \inf\{(f|_{I_3})^{-1}(a_0)\}$ and $I_{3_1} = [b_0, b_1]$. Then f has the subgraph $\odot I_4 \rightarrow I_{3_1} \rightleftharpoons I_2 \rightarrow I_4$ with $I_2 \cap I_{3_1} = \emptyset$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$. Therefore, if $b = 1$ and $c = -1$, then $\text{MPer}(f) = \mathbb{N}$. Hence, if $|a| + |d| = 2$ and $bc = -1$, then $\text{MPer}(f) = \mathbb{N}$. This completes the proof of statement (c1). $\square$

Proof of statement (c21) of Theorem B We assume now that $\mathbf{a} = \mathbf{1}$ and $\mathbf{d} = \mathbf{0}$. If $\mathbf{bc} = \mathbf{0}$, then $\text{MPer}(f) = \{1\}$ as it can be deduced from the examples of Fig. 15. We suppose that b and c are such that $|\mathbf{bc}| > 1$ and $(\mathbf{b}, \mathbf{c}) \notin \{(2, 1), (2, -1), (-2, 1), (-2, -1)\}$. From the graph of f (see for instance Fig. 16) it follows that there are three basic intervals I_1, I_2 , and I_3 , $I_1, I_2 \subset C_1$, $I_3 \subset C_2$, such that either $p \notin I_1 \cap I_2$ or $p \notin I_1 \cap I_3$ and f has the subgraph of Proposition 2.3, so $\text{Per}(f) \supset \mathbb{N} \setminus \{2\}$.

If $\{\mathbf{b}, \mathbf{c}\} \not\subset \{-2, -1, 1, 2\}$, then we can choose I_2 and I_3 such that $I_2 \cap I_3 = \emptyset$, and by Proposition 2.3, $2 \in \text{Per}(f)$. If $\{\mathbf{b}, \mathbf{c}\} \subset \{-2, -1, 1, 2\}$, in general there do not exist two basic intervals I_i and I_j , $I_i \neq I_j$, such that $p \notin I_i \cap I_j$ and $I_i \rightleftharpoons I_j$. If they exist, then by Lemma 2.1, considering the non-repetitive loop $I_i \rightarrow I_j \rightarrow I_i$, there is a periodic point z of f with period 2. If they do not exist (see for instance (c) and (d) of Fig. 16) and $(b, c) \in \{(1, 2), (-1, -2)\}$, $2 \notin \text{Per}(f)$ as we can see from the examples of Fig. 17. Now we will prove that if $(b, c) \in \{(1, -2), (-1, 2)\}$ or $|b| = |c| = 2$, then $2 \in \text{Per}(f)$.

We suppose first that $(b, c) \in \{(1, -2), (-1, 2)\}$. We know that f has four basic intervals I_1, I_2, I_3 , and I_4 , the first three in C_1 and $I_4 = C_2$, such that $f(I_1) = f(I_4) = C_1$ and $f(I_2) = f(I_3) = C_2$. Let p and a_0 be the endpoints of I_2 , a_0 and a_1 be the endpoints of I_1 , a_1 and p be the endpoints of I_3 (see for instance Fig. 18). We consider an ordering in the intervals I_1, I_2 , and I_3 in such a way that p is the smallest element of I_2 and the greatest of I_3 . Under these

Figure 15 Examples of maps with $a = 1, d = 0$, and $bc = 0$





Figure 16 Examples of maps with $a = 1, d = 0$, $(b, c) \notin \{(2, 1), (2, -1), (-2, 1), (-2, -1)\}$, and $|\mathbf{bc}| > 1$




Figure 17 Examples of maps with $a = 1, d = 0$, $(b, c) \in \{(1, 2), (-1, -2)\}$, and $2 \notin \text{Per}(f)$




Figure 18 Examples of maps with $a = 1$, $d = 0$, and $(b, c) \in \{(1, -2), (-1, 2)\}$




Figure 19 Examples of maps with $a = 1$, $d = 0$, and $|b| = |c| = 2$

assumptions, set $I_2 = [p, a_0]$, $I_1 = [a_0, a_1]$, and $I_3 = [a_1, p]$. Define $b_0 = \sup\{(f|_{I_4})^{-1}(a_0)\}$, $I_{4_1} = [p, b_0]$, and $I_{4_2} = [b_0, p]$. Set $a_2 = \inf\{(f|_{I_3})^{-1}(b_0)\}$ and $I_{3_1} = [a_1, a_2]$. If $(b, c) = (1, -2)$, we have $I_{4_2} \rightleftharpoons I_{3_1}$ and $I_{4_2} \cap I_{3_1} = \emptyset$. If $(b, c) = (-1, 2)$, we get $I_{4_1} \rightleftharpoons I_{3_1}$ and $I_{4_1} \cap I_{3_1} = \emptyset$. So, by Lemma 2.1, $2 \in \text{Per}(f)$.

Suppose now that $|b| = |c| = 2$. We know that f has five basic intervals I_1, I_2, I_3, I_4 , and I_5 , the first three in C_1 and the other two in C_2 , such that $f(I_2) = f(I_3) = C_2$ and $f(I_1) = f(I_4) = f(I_5) = C_1$. Taking an ordering similar to the previous case, define the intervals $I_2 = [p, a_0]$, $I_1 = [a_0, a_1]$, $I_3 = [a_1, p]$, $I_4 = [p, b_0]$, and $I_5 = [b_0, p]$ (see for instance Fig. 19). Set $a_2 = \sup\{(f|_{I_2})^{-1}(b_0)\}$ and $I_{2_2} = [a_2, a_0]$. If $c = 2$, we have $I_{2_2} \rightleftharpoons I_5$ and $I_{2_2} \cap I_5 = \emptyset$. If $c = -2$, we have $I_{2_2} \rightleftharpoons I_4$ and $I_{2_2} \cap I_4 = \emptyset$. So, by Lemma 2.1, $2 \in \text{Per}(f)$. Therefore, if $|bc| > 1$ and




Figure 20 Examples of maps with $a = 1$, $d = 0$, and $(b, c) \in \{(2, 1), (-2, 1)\}$

$(b, c) \notin \{(2, 1), (2, -1), (-2, 1), (-2, -1)\}$, we have $\text{MPer}(f) = \mathbb{N} \setminus \{2\}$ if $(b, c) \in \{(1, 2), (-1, -2)\}$ and $\text{MPer}(f) = \mathbb{N}$ otherwise.

We assume that $|\mathbf{bc}| > 1$ and $(\mathbf{b}, \mathbf{c}) \in \{(2, 1), (2, -1), (-2, 1), (-2, -1)\}$. We know that f has four basic intervals I_1, I_2, I_3 , and I_4 , the first two in C_1 and the others in C_2 , such that $f(I_1) = f(I_3) = f(I_4) = C_1$ and $f(I_2) = C_2$. Let p and a_0 be the endpoints of I_1 and I_2 , and b_0 and p be the endpoints of I_3 and I_4 (see for instance Figs. 20 and 21). For each pair (b, c) , we have two possibilities for the intervals I_1 and I_2 . If $(b, c) \in \{(2, 1), (-2, 1)\}$ and $I_2 = [a_0, p]$, write $a_1 = \inf\{(f|_{I_2})^{-1}(b_0)\}$ and $I_{2_1} = [a_0, a_1]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_1} \rightrightarrows I_3 \rightarrow I_1$ with $I_3 \cap I_{2_1} = \emptyset$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$. If $I_2 = [p, a_0]$, consider $a_1 = \sup\{(f|_{I_2})^{-1}(b_0)\}$ and $I_{2_2} = [a_1, a_0]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_2} \rightrightarrows I_4 \rightarrow I_1$ with $I_4 \cap I_{2_2} = \emptyset$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$.

If $(b, c) \in \{(-2, -1), (2, -1)\}$ and $I_2 = [a_0, p]$, set $a_1 = \inf\{(f|_{I_2})^{-1}(b_0)\}$ and $I_{2_1} = [a_0, a_1]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_1} \rightrightarrows I_4 \rightarrow I_1$ with $I_4 \cap I_{2_1} = \emptyset$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$. If $I_2 = [p, a_0]$, consider $a_1 = \sup\{(f|_{I_2})^{-1}(b_0)\}$ and $I_{2_2} = [a_1, a_0]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_2} \rightrightarrows I_3 \rightarrow I_1$ with $I_3 \cap I_{2_2} = \emptyset$, and by Proposition 2.3, $\text{Per}(f) = \mathbb{N}$. Therefore, if $|\mathbf{bc}| > 1$ and $(\mathbf{b}, \mathbf{c}) \in \{(2, 1), (2, -1), (-2, 1), (-2, -1)\}$, $\text{MPer}(f) = \mathbb{N}$.

We consider the case $|\mathbf{bc}| = 1$. First assume that $\mathbf{bc} = 1$. As we can see from the examples of Fig. 22, $2 \notin \text{MPer}(f)$. Now we will prove that $\text{Per}(f) = \mathbb{N} \setminus \{2\}$.

We know that f has three basic intervals I_1, I_2 , and I_3 , the first two in C_1 and $I_3 = C_2$, such that $f(I_1) = f(I_3) = C_1$ and $f(I_2) = C_2$. We have two possibilities for the intervals I_1 and I_2 : either p is the smallest element of I_1 and the greatest of I_2 or p is the smallest element of I_2 and the greatest of I_1 (see for instance Fig. 23). In the assumption that $b = c = 1$, if $I_1 = [p, a_0]$, write $b_0 = \inf\{(f|_{I_3})^{-1}(a_0)\}$, $I_{3_1} = [p, b_0]$, $a_1 = \inf\{(f|_{I_2})^{-1}(b_0)\}$, and $I_{2_1} =$




Figure 21 Examples of maps with $a = 1$, $d = 0$, and $(b, c) \in \{(-2, -1), (2, -1)\}$




Figure 22 Examples of maps with $a = 1$, $d = 0$, $bc = 1$, and $2 \notin \text{Per}(f)$

$[a_0, a_1]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_1} \rightarrow I_{3_1} \rightarrow I_1$ and by Proposition 2.3, $\text{Per}(f) \supset \mathbb{N} \setminus \{2\}$. If $I_1 = [a_0, p]$, define $b_0 = \sup\{(f|_{I_3})^{-1}(a_0)\}$, $I_{3_2} = [b_0, p]$, $a_1 = \sup\{(f|_{I_2})^{-1}(b_0)\}$, and $I_{2_2} = [a_1, a_0]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_2} \rightarrow I_{3_2} \rightarrow I_1$, and by Proposition 2.3, $\text{Per}(f) \supset \mathbb{N} \setminus \{2\}$.

If $b = c = -1$, we consider first the case $I_1 = [p, a_0]$. Set $b_0 = \sup\{(f|_{I_3})^{-1}(a_0)\}$, $I_{3_2} = [b_0, p]$, $a_1 = \inf\{(f|_{I_2})^{-1}(b_0)\}$, and $I_{2_1} = [a_0, a_1]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_1} \rightarrow I_{3_2} \rightarrow I_1$, and by Proposition 2.3, $\text{Per}(f) \supset \mathbb{N} \setminus \{2\}$. If $I_1 = [a_0, p]$, write $b_0 = \inf\{(f|_{I_3})^{-1}(a_0)\}$, $I_{3_1} = [p, b_0]$, $a_1 = \sup\{(f|_{I_2})^{-1}(b_0)\}$, and $I_{2_2} = [a_1, a_0]$. Then f has the subgraph $\odot I_1 \rightarrow I_{2_2} \rightarrow I_{3_1} \rightarrow I_1$, and by Proposition 2.3, $\text{Per}(f) \supset \mathbb{N} \setminus \{2\}$. Therefore, if $a = 1$, $d = 0$, and $bc = 1$, $\text{MPer}(f) = \mathbb{N} \setminus \{2\}$.

Assume now that $bc = -1$. We know that f has three basic intervals I_1 , I_2 , and I_3 , the first two in C_1 and $I_3 = C_2$, such that $f(I_1) = f(I_3) = C_1$ and $f(I_2) = C_2$. We have two possibilities for the intervals I_1 and I_2 : either p is the smallest element of I_1 and the greatest of I_2 or p is the smallest element of I_2 and the greatest of I_1 (see for instance Fig. 24). Define $b_0 = \inf\{(f|_{I_3})^{-1}(a_0)\}$, $I_{3_1} = [p, b_0]$, $I_{3_2} = [b_0, p]$, and $a_1 = \inf\{(f|_{I_2})^{-1}(b_0)\}$.




Figure 23 Examples of maps with $a = 1$, $d = 0$, and $bc = 1$






Figure 24 Examples of maps with $a = 1$, $d = 0$, and $bc = -1$

If $I_1 = [a_0, p]$, let $I_{2_1} = [p, a_1]$ and $I_{2_2} = [a_1, a_0]$. Consider $a_2 = \inf\{(f|_{I_1})^{-1}(a_1)\}$. We write $I_{1_1} = [a_0, a_2]$ and $I_{1_2} = [a_2, p]$. If $b = 1$ and $c = -1$ (see (a) of Fig. 24), f has the subgraph




We consider the non-repetitive loops $I_{1_1} \rightarrow I_{2_1} \rightarrow I_{3_2} \rightarrow I_{1_1}$ and $I_{1_2} \rightarrow I_{1_1} \rightarrow I_{2_1} \rightarrow I_{3_2} \rightarrow I_{1_2} \rightarrow \dots \rightarrow I_{1_2}$ of lengths 3 and $n \geq 4$, respectively. From the first loop and by Lemma 2.1, there is a periodic point z of f with period 3; from the second loop and by Lemma 2.1, there is a periodic point z of f with period $n \geq 4$. Moreover, $I_{3_1} \rightleftharpoons I_{2_2}$ and $I_{3_1} \cap I_{2_2} = \emptyset$, so, by Lemma 2.1, $2 \in \text{Per}(f)$. Hence, $\text{Per}(f) = \mathbb{N}$. If $b = -1$ and $c = 1$ (see (b) of Fig. 24), f has the subgraph




Now from the non-repetitive loops $I_{1_1} \rightarrow I_{2_1} \rightarrow I_{3_1} \rightarrow I_{1_1}$ and $I_{1_2} \rightarrow I_{1_1} \rightarrow I_{2_1} \rightarrow I_{3_1} \rightarrow I_{1_2} \rightarrow \dots \rightarrow I_{1_2}$ of lengths 3 and $n \geq 4$, respectively, and $I_{3_2} \rightleftharpoons I_{2_2}$ and $I_{3_2} \cap I_{2_2} = \emptyset$, it follows that $\text{Per}(f) = \mathbb{N}$.

If $I_1 = [p, a_0]$, let $I_{2_1} = [a_0, a_1]$, $I_{2_2} = [a_1, p]$. Define $a_2 = \sup\{(f|_{I_1})^{-1}(a_1)\}$, $I_{1_1} = [a_0, a_2]$, and $I_{1_2} = [a_2, p]$. If $b = 1$ and $c = -1$ (see (c) of Fig. 24), f has the subgraph



Again from the non-repetitive loops $I_{1_2} \rightarrow I_{2_2} \rightarrow I_{3_1} \rightarrow I_{1_2}$ and $I_{1_1} \rightarrow I_{1_2} \rightarrow I_{2_2} \rightarrow I_{3_1} \rightarrow I_{1_1} \rightarrow \dots \rightarrow I_{1_1}$ of lengths 3 and $n \geq 4$, respectively, $I_{3_2} \rightleftharpoons I_{2_1}$ and $I_{3_2} \cap I_{2_1} = \emptyset$, $\text{Per}(f) = \mathbb{N}$. If $b = -1$ and $c = 1$ (see (d) of Fig. 24), f has the subgraph



We consider the non-repetitive loops $I_{1_2} \rightarrow I_{2_2} \rightarrow I_{3_2} \rightarrow I_{1_2}$ and $I_{1_1} \rightarrow I_{1_2} \rightarrow I_{2_2} \rightarrow I_{3_2} \rightarrow I_{1_1} \rightarrow \dots \rightarrow I_{1_1}$ of lengths 3 and $n \geq 4$, respectively, $I_{3_1} \rightleftharpoons I_{2_1}$ and $I_{3_1} \cap I_{2_1} = \emptyset$. We obtain

that $\text{Per}(f) = \mathbb{N}$. Therefore, if $a = 1$, $d = 0$ and $bc = -1$, $\text{MPer}(f) = \mathbb{N}$. This completes the proof of statement (c21). $\square$

Proof of statement (c22) of Theorem B If $\mathbf{a} = \mathbf{0}$ and $\mathbf{d} = \mathbf{1}$, by using the same kind of arguments as those in the case $a = 1$ and $d = 0$, and interchanging b and c , we obtain statement (c22). $\square$

Proof of statement (c23) of Theorem B We suppose that $\mathbf{a} = -\mathbf{1}$ and $\mathbf{d} = \mathbf{0}$. If $\mathbf{bc} = \mathbf{0}$, then $\text{MPer}(f) = \{1\}$ as it can be seen from the examples of Fig. 25. The cases in which $\text{MPer}(f)$ is either $\mathbb{N} \setminus \{2\}$ or $\mathbb{N}$ can be proved following exactly the same kind of arguments as those in the proof of statement (c21).

Assume now that $\mathbf{bc} = -\mathbf{1}$. From the examples of Fig. 26 we can see that $3 \notin \text{MPer}(f)$.

We know that f has three basic intervals I_1 , I_2 , and I_3 , the first two in C_1 and $I_3 = C_2$, such that $f(I_1) = f(I_3) = C_1$ and $f(I_2) = C_2$. We have two possibilities for the intervals I_1 and I_2 : either p is the smallest element of I_1 and the greatest of I_2 or p is the smallest element of I_2 and the greatest of I_1 (see for instance Fig. 27). Denote $b_0 = \inf\{(f|_{I_3})^{-1}(a_0)\}$, $I_{3_1} = [p, b_0]$, $I_{3_2} = [b_0, p]$, and $a_1 = \inf\{(f|_{I_2})^{-1}(b_0)\}$.

If $I_1 = [a_0, p]$, let $I_{2_1} = [p, a_1]$ and $I_{2_2} = [a_1, a_0]$. Consider $a_2 = \inf\{(f|_{I_1})^{-1}(a_1)\}$. Write $I_{1_1} = [a_0, a_2]$ and $I_{1_2} = [a_2, p]$. If $b = 1$ and $c = -1$ (see (a) of Fig. 27), f has the subgraph $\odot I_{1_1} \rightarrow I_{1_2} \rightarrow I_{2_1} \rightarrow I_{3_2} \rightarrow I_{1_1}$. We consider the non-repetitive loop $I_{1_1} \rightarrow I_{1_2} \rightarrow I_{2_1} \rightarrow I_{3_2} \rightarrow I_{1_1} \rightarrow \dots \rightarrow I_{1_1}$ of length $n \geq 4$. By Lemma 2.1 there is a periodic point z of f with period $n \geq 4$. Moreover, $I_{3_1} \rightleftharpoons I_{2_2}$ and $I_{3_1} \cap I_{2_2} = \emptyset$, so, by Lemma 2.1, $2 \in \text{Per}(f)$. Hence, $\text{Per}(f) = \mathbb{N} \setminus \{3\}$. If $b = -1$ and $c = 1$ (see (b) of Fig. 27), f has the subgraph $\odot I_{1_1} \rightarrow I_{1_2} \rightarrow I_{2_1} \rightarrow I_{3_1} \rightarrow I_{1_1}$. We consider the non-repetitive loop $I_{1_1} \rightarrow I_{1_2} \rightarrow I_{2_1} \rightarrow I_{3_1} \rightarrow I_{1_1} \rightarrow \dots \rightarrow I_{1_1}$ of length $n \geq 4$. By Lemma 2.1 there is a periodic point z of f with

Figure 25 Examples of maps with $a = -1$, $d = 0$, and $bc = 0$





Figure 26 Examples of maps with $a = -1$, $d = 0$, $bc = -1$, and $3 \notin \text{Per}(f)$




Figure 27 Examples of maps with $a = 1$, $d = 0$ and $bc = -1$

period $n \geq 4$. Moreover, $I_{3_2} \rightleftharpoons I_{2_2}$ and $I_{3_2} \cap I_{2_2} = \emptyset$, so, by Lemma 2.1, $2 \in \text{Per}(f)$. Hence, $\text{Per}(f) = \mathbb{N} \setminus \{3\}$.

If $I_1 = [p, a_0]$, let $I_{2_1} = [a_0, a_1]$ and $I_{2_2} = [a_1, p]$. Consider $a_2 = \sup\{(f|_{I_1})^{-1}(a_1)\}$. Write $I_{1_1} = [p, a_2]$ and $I_{1_2} = [a_2, a_0]$. If $b = 1$ and $c = -1$ (see (c) of Fig. 27), f has the subgraph $\odot I_{1_2} \rightarrow I_{1_1} \rightarrow I_{2_2} \rightarrow I_{3_1} \rightarrow I_{1_2}$. From the non-repetitive loop $I_{1_2} \rightarrow I_{1_1} \rightarrow I_{2_2} \rightarrow I_{3_1} \rightarrow I_{1_2} \rightarrow \dots \rightarrow I_{1_2}$ of length $n \geq 4$, $I_{3_2} \rightleftharpoons I_{2_1}$, and $I_{3_2} \cap I_{2_1} = \emptyset$, we obtain that $\text{Per}(f) = \mathbb{N} \setminus \{3\}$. If $b = -1$ and $c = 1$ (see (d) of Fig. 27), f has the subgraph $\odot I_{1_2} \rightarrow I_{1_1} \rightarrow I_{2_2} \rightarrow I_{3_2} \rightarrow I_{1_2}$. Using the non-repetitive loop $I_{1_2} \rightarrow I_{1_1} \rightarrow I_{2_2} \rightarrow I_{3_2} \rightarrow I_{1_2} \rightarrow \dots \rightarrow I_{1_2}$ of length $n \geq 4$, $I_{3_1} \rightleftharpoons I_{2_1}$ and $I_{3_1} \cap I_{2_1} = \emptyset$, we get that $\text{Per}(f) = \mathbb{N} \setminus \{3\}$. Therefore, if $a = -1$, $d = 0$, and $bc = -1$, $\text{MPer}(f) = \mathbb{N} \setminus \{3\}$. This completes the proof of statement (c23). $\square$

Proof of statement (c24) of Theorem B If $\mathbf{a} = \mathbf{0}$ and $\mathbf{d} = -\mathbf{1}$, by using the same kind of arguments as those in the case $a = -1$ and $d = 0$, and interchanging b and c , we obtain statement (c24). $\square$

Proof of statement (c3) of Theorem B We suppose that $\mathbf{a} = \mathbf{d} = \mathbf{0}$. If $\mathbf{bc} = \mathbf{0}$ or $\mathbf{bc} = \mathbf{1}$, we can deduce from the examples of Fig. 28 that $\text{MPer}(f) = \{1\}$. If $\mathbf{bc} = -\mathbf{1}$, then $\text{MPer}(f) = \{1, 2\}$ (see for instance Fig. 29).

We assume now that $|\mathbf{bc}| = 2$. Since $a = d = 0$, we may assume without loss of generality that $|b| = 1$ and $|c| = 2$. We consider first the case $\mathbf{bc} = -2$. Clearly, $\{1, 2\} \subset \text{Per}(f)$, no other odd number belongs to $\text{MPer}(f)$ and $4 \notin \text{MPer}(f)$ as it can be deduced from Fig. 30. Now we will prove that $n \in \text{Per}(f)$ for any n even larger than 4.

We know that f has three basic intervals I_1 , I_2 , and I_3 , the first two in C_1 and $I_3 = C_2$, such that $f(I_1) = f(I_2) = C_2$ and $f(I_3) = C_1$ (see for instance Fig. 31). Consider




Figure 28 Examples of maps with $a = d = 0$ and either $bc = 0$ or $bc = 1$




Figure 29 Examples of maps with $a = d = 0$ and $bc = -1$




Figure 30 Examples of maps with $a = d = 0$, $bc = -2$, and $4 \notin \text{Per}(f)$




Figure 31 Examples of maps with $\sigma = d = 0$ and $bc = -2$





Figure 32 Examples of maps with $\sigma = d = 0$, $bc = 2$, and $2 \notin \text{Per}(f)$



$b_0 = \inf\{(f|_{I_3})^{-1}(a_0)\}$, $a_1 = \inf\{(f|_{I_1})^{-1}(b_0)\}$, $b_1 = \inf\{(f|_{I_3})^{-1}(a_1)\}$. Set $I_{11} = [p, a_1]$, $I_{12} = [a_1, a_0]$, I_{31} the interval with endpoints b_1 and p , I_{32} the interval with endpoints b_1 and b_0 , and I_{33} the interval with endpoints b_0 and p . Then f has the subgraph



We consider the non-repetitive loops $I_{32} \rightarrow I_{12} \rightarrow I_{32}$ and $I_2 \rightarrow I_{32} \rightarrow I_{12} \rightarrow I_{31} \rightarrow I_{11} \rightarrow I_{33} \rightarrow I_2 \rightarrow \dots \rightarrow I_{33} \rightarrow I_2$ of lengths 2 and n even, $n \geq 6$, respectively. We have $I_{32} \cap I_{12} = \emptyset$, so, from the first loop and by Lemma 2.1, there is a periodic point z of f with period 2; from the second loop and by Lemma 2.1, there is a periodic point z of f with period n even $n \geq 6$. Therefore, if $bc = -2$, then $\text{MPer}(f) = \{1\} \cup (2\mathbb{N} \setminus \{4\})$.

We suppose that $bc = 2$. No odd number other than 1 belongs to $\text{MPer}(f)$, as it can be seen from the examples of Fig. 32. Also from Fig. 32 we can deduce that $2 \notin \text{MPer}(f)$. Now we will prove that $n \in \text{Per}(f)$ for any n even larger than 2.

We know that f has three basic intervals I_1 , I_2 , and I_3 , the first two in C_1 and $I_3 = C_2$, such that $f(I_1) = f(I_2) = C_2$ and $f(I_3) = C_1$ (see for instance Fig. 33). Denote $b_0 = \inf\{(f|_{I_3})^{-1}(a_0)\}$, $a_1 = \inf\{(f|_{I_1})^{-1}(b_0)\}$, and $b_1 = \inf\{(f|_{I_3})^{-1}(a_1)\}$. Write $I_{11} = [p, a_1]$, $I_{12} = [a_1, a_0]$, I_{32} the in-



terval with endpoints b_1 and b_0 , and I_{33} the interval with endpoints b_0 and p . Then f has the subgraph $I_{32} \rightarrow I_{12} \rightarrow I_{33} \rightarrow I_2 \rightarrow I_{32}$. We take the non-repetitive loop $I_2 \rightarrow I_{32} \rightarrow I_{12} \rightarrow I_{33} \rightarrow I_2 \rightarrow \dots \rightarrow I_{33} \rightarrow I_2$ of length n even, $n \geq 4$. By Lemma 2.1, there is a periodic point z of f with period n even $n \geq 4$. Therefore, if $bc = 2$, then $\text{MPer}(f) = \{1\} \cup (2\mathbb{N} \setminus \{2\})$.

We consider now the case $|bc| > 2$. We must separate the case $|b| = |c| = 2$ from the others. If $|b| > 2$ or $|c| > 2$, then there are three basic intervals I_1 , I_2 , and I_3 such that $I_2 \cap I_3 = \emptyset$ and $I_1 \rightarrow I_3 \rightarrow I_2$ (see for instance Fig. 34). By Lemma 2.1, the non-repetitive loop $I_2 \rightarrow I_3 \rightarrow I_2$ gives a periodic point z of f with period 2, and the non-repetitive loop $I_1 \rightarrow I_3 \rightarrow I_2 \rightarrow I_3 \rightarrow \dots \rightarrow I_2 \rightarrow I_3 \rightarrow I_1$ of length n even larger than 2 gives a periodic point z of f with period n even. No odd number other than 1 belongs to $\text{MPer}(f)$. Therefore, if $|b| > 2$ or $|c| > 2$, then $\text{MPer}(f) = \{1\} \cup 2\mathbb{N}$.

We suppose that $|b| = |c| = 2$. Clearly, no odd number other than 1 belongs to $\text{MPer}(f)$. Now we will prove that $n \in \text{Per}(f)$ for any n even.

We know that f has four basic intervals I_1 , I_2 , I_3 , and I_4 , the first two in C_1 and the others in C_2 , such that $f(I_1) = f(I_2) = C_2$ and $f(I_3) = f(I_4) = C_1$ (see for instance Fig. 35). Consider $b_1 = \inf\{(f|_{I_3})^{-1}(a_0)\}$ and $a_1 = \inf\{(f|_{I_1})^{-1}(b_1)\}$. Denote $I_{11} = [p, a_1]$, $I_{12} = [a_1, a_0]$, $I_2 = [a_0, p]$, $I_{31} = [p, b_1]$, $I_{32} = [b_1, b_0]$, and $I_4 = [b_0, p]$. If $(b, c) \in \{(2, 2), (-2, 2)\}$, then f has the subgraph $I_2 \rightarrow I_4 \rightarrow I_{12}$. We take the non-repetitive loops $I_4 \rightarrow I_{12} \rightarrow I_4$ and $I_2 \rightarrow I_4 \rightarrow I_{12} \rightarrow I_4 \rightarrow \dots \rightarrow I_{12} \rightarrow I_4 \rightarrow I_2$ of lengths 2 and n even larger than 2, respectively. By Lemma 2.1, the first loop gives a periodic point z of f with period 2, and the second loop gives a periodic point z of f with period n even larger than 2. Hence, if $(b, c) \in \{(2, 2), (-2, 2)\}$, $\text{Per}(f) = \{1\} \cup 2\mathbb{N}$.

If $(b, c) = (-2, -2)$, then f has the subgraph $I_4 \rightarrow I_{11} \rightarrow I_{32}$. We consider the non-repetitive loops $I_{32} \rightarrow I_{11} \rightarrow I_{32}$ and $I_4 \rightarrow I_{11} \rightarrow I_{32} \rightarrow I_{11} \rightarrow \dots \rightarrow I_{32} \rightarrow I_{11} \rightarrow I_4$ of lengths 2 and n even larger than 2, respectively. By Lemma 2.1, the first loop gives a periodic point z of f




Figure 35 Examples of maps with $a = d = 0$ and $|b| = |c| = 2$

with period 2, and the second loop gives a periodic point z of f with period n even larger than 2. Hence, if $(b, c) = (-2, -2)$, $\text{Per}(f) = \{1\} \cup 2\mathbb{N}$.

If $(b, c) = (2, -2)$, then f has the subgraph $I_4 \rightleftharpoons I_2 \rightleftharpoons I_{3_2}$. We consider the non-repetitive loops $I_2 \rightarrow I_{3_2} \rightarrow I_2$ and $I_4 \rightarrow I_2 \rightarrow I_{3_2} \rightarrow I_2 \rightarrow \dots \rightarrow I_{3_2} \rightarrow I_2 \rightarrow I_4$ of lengths 2 and n even larger than 2, respectively. By Lemma 2.1, the first loop gives a periodic point z of f with period 2, and the second loop gives a periodic point z of f with period n even larger than 2. Hence, if $(b, c) = (-2, -2)$, $\text{Per}(f) = \{1\} \cup 2\mathbb{N}$. Therefore, if $|b| = |c| = 2$, then $\text{MPer}(f) = \{1\} \cup 2\mathbb{N}$. This completes the proof of statement (c3). $\square$

Acknowledgements

We thank our universities for the facilities that they gave us during the elaboration of this work.

Funding

The first author has been partially supported by the Ministerio de Ciencia, Innovación y Universidades, Agencia Estatal de Investigación grants MTM2016-77278-P (FEDER), the Agència de Gestió d'Ajuts Universitaris i de Recerca grant 2017SGR1617, and the H2020 European Research Council grant MSCA-RISE-2017-777911.

Availability of data and materials

No materials are necessary.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Both authors have contributed approximately in one half to the paper. Both authors read and approved the final manuscript.

Author details

¹Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain. ²Departamento de Matemática, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 11 December 2019 Accepted: 2 December 2020 Published online: 18 January 2021

References

1. Abdulla, U.G., Abdulla, R.U., Abdulla, M.U., Iqbal, N.H.: Second minimal orbits, Sharkovski ordering and universality in chaos. *Int. J. Bifurc. Chaos Appl. Sci. Eng.* **27**, 1–24 (2017)
2. Alsedà, L., Baldwin, S., Llibre, J., Swanson, R., Szlenk, W.: Minimal sets of periods for torus maps via Nielsen numbers. *Pac. J. Math.* **169**, 1–32 (1995)
3. Alsedà, L., Juher, D., Mumbú, P.: On the preservation of combinatorial types for maps on trees. *Ann. Inst. Fourier (Grenoble)* **55**(7), 2375–2398 (2005)
4. Alsedà, L., Llibre, J., Misiurewicz, M.: *Combinatorial Dynamics and Entropy in Dimension One*, 2nd edn. Advanced Series in Nonlinear Dynamics, vol. 5. World Scientific, Singapore (2000)
5. Alsedà, L., Ruette, S.: Rotation sets for graph maps of degree 1. *Ann. Inst. Fourier (Grenoble)* **58**(4), 1233–1294 (2008)
6. Arai, T.: The structure of dendrites constructed by pointwise P-expansive maps on the unit interval. *Discrete Contin. Dyn. Syst.* **36**, 43–61 (2016)
7. Bernhardt, C.: Vertex maps for trees: algebra and periods of periodic orbits. *Discrete Contin. Dyn. Syst.* **14**(3), 399–408 (2006)
8. Bernhardt, C.: Vertex maps on graphs-trace theorems. *Fixed Point Theory Appl.* **2011**, 8 (2011)
9. Block, L., Guckenheimer, J., Misiurewicz, M., Young, L.S.: Periodic points and topological entropy of one dimensional maps. In: *Global Theory of Dynamical Systems. Lecture Notes in Math.*, vol. 819, pp. 18–34. Springer, Berlin (1980)
10. Brown, R.F.: *The Lefschetz Fixed Point Theorem*. Scott, Foresman, Chicago (1971)
11. Casasayas, J., Llibre, J., Nunes, A.: Periodic orbits of transversal maps. *Math. Proc. Camb. Philos. Soc.* **118**, 161–181 (1995)
12. Efremova, L.S.: Periodic orbits and the degree of a continuous map of a circle. *Differ. Integer Equ. (Gor'kii)* **2**, 109–115 (1978) (in Russian)
13. Jiang, B.: *Lectures on Nielsen Fixed Point Theory*. Contemporary Mathematics, vol. 14. Am. Math. Soc., Providence (1983)
14. Jiang, B., Llibre, J.: Minimal sets of periods for torus maps. *Discrete Contin. Dyn. Syst.* **4**, 301–320 (1998)
15. Kiang, T.H.: *The Theory of Fixed Point Classes*. Springer, Berlin (1989)
16. Llibre, J.: Periodic points of one dimensional maps. In: Mira, C., Netzer, N., Simó, C., Targonski, G. (eds.) *European Conference on Iteration Theory, ECIT 89*, pp. 194–198. World Scientific, Singapore (1991)
17. Llibre, J., Misiurewicz, M.: Negative periodic orbits for graph maps. *Nonlinearity* **19**(3), 741–746 (2006)
18. Llibre, J., Sá, A.: Periods for continuous self-maps of a bouquet of circles. *C. R. Acad. Sci. Paris Sér. I Math.* **318**, 1035–1040 (1994)
19. Los, J.E., Nitecki, Z.H.: Edge-transitive graph automorphisms and periodic surface homeomorphisms. *Discrete dynamical systems. Int. J. Bifurc. Chaos Appl. Sci. Eng.* **9**(9), 1803–1813 (1999)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)