DOI: 10.1002/clt2.12010

REVIEW

Non-specific lipid-transfer proteins: Allergen structure and function, cross-reactivity, sensitization, and epidemiology

Isabel J. Skypala¹ Ricardo Asero² I Domingo Barber^{3,4} I Lorenzo Cecchi⁵ Arazeli Diaz Perales⁶ Karin Hoffmann-Sommergruber⁷ Helide A. Pastorello⁸ I Ines Swoboda⁹ Joan Bartra¹⁰ I Didier G. Ebo¹¹ Arazeli Diaz Perales¹² I Joan Bartra¹⁰ I Didier G. Ebo¹¹ Arazeli Margaretha A. Faber¹¹ I Montserrat Fernández-Rivas¹² I Francesca Gomez¹³ I Anastasios. P. Konstantinopoulos¹⁴ I Olga Luengo¹⁵ I Ronald van Ree¹⁶ I Enrico Scala¹⁷ I Stephen J. Till^{18,19} I European Academy of Allergy Clinical Immunology (EAACI) Task Force: Non-specific Lipid Transfer Protein Allergy Across Europe

- ⁷Department of Pathophysiology and Allergy Research, Medical University of Vienna, Austria
- ⁸Unit of Allergology and Immunology, ASST Grande Ospedale Metropolitano Niguarda, University of Milan, Milan, Italy
- ⁹Biotechnology Section, FH Campus Wien, University of Applied Sciences, Vienna, Austria
- ¹⁰Hospital Clinic de Barcelona, IDIBAPS, Universitat de Barcelona, ARADyAL, Barcelona, Spain
- ¹¹Department of Immunology, Allergology, Rheumatology and Infla-Med Centre of Excellence, Faculty of Medicine and Health Sciences, University of Antwerp and Antwerp University Hospital, Ghent, Belgium
- ¹²Department of Allergy, Hospital Clínico San Carlos, Universidad Complutense de Madrid, IdISSC, ARADyAL, Madrid, Spain
- ¹³Allergy Unit, IBIMA- Hospital Regional Universitario de Malaga, Malaga, and Spanish Network for Allergy RETICS de Asma, Reacciones adversas y Alérgicas (ARADyAL), Madrid, Spain
- ¹⁴Allergy Department, 2nd Paediatric Clinic, University of Athens, Athens, Greece
- ¹⁵Allergy Unit, Internal Medicine Department, Vall d'Hebron University Hospital, Universitat Autònoma de Barcelona, ARADyAL, Barcelona, Spain
- ¹⁶Department of Experimental Immunology and Department of Otorhinolaryngology, Amsterdam University Medical Centers, location AMC, Amsterdam, The Netherlands
- ¹⁷Experimental Allergy Unit, Istituto Dermopatico Dell'immacolata, IRCCS, FLMM, Rome, Italy
- ¹⁸Peter Gorer Department of Immunobiology, King's College London, London, UK
- ¹⁹Department of Allergy, Guy's & St Thomas' NHS Foundation Trust, London, UK

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2021 The Authors. Clinical and Translational Allergy published by John Wiley and Sons Ltd on behalf of European Academy of Allergy and Clinical Immunology.

¹Department of Allergy & Clinical Immunology, Royal Brompton & Harefield NHS Foundation Trust, Imperial College, London, UK

²Ambulatorio di Allergologia, Clinica San Carlo, Milan, Italy

³IMMA, School of Medicine, Universidad San Pablo CEU, CEU Universities, Madrid, Spain

⁴RETIC ARADYAL RD16/0006/0015, Instituto de Salud Carlos III, Madrid, Spain

⁵SOS Allergy and Clinical Immunology, USL Toscana Centro, Prato, Italy

⁶Departamento de Biotecnología-Biología Vegetal, Centro de Biotecnología y Genómica de Plantas (CBGP, UPM-INIA), Universidad Politécnica de Madrid, Madrid, Spain



Correspondence

Isabel Skypala, Department of Allergy & Clinical Immunology, Royal Brompton Hospital, Sydney Street, London SW3 6NP, UK. Email:

Abstract

Background: Discovered and described 40 years ago, non-specific lipid transfer proteins (nsLTP) are present in many plant species and play an important role protecting plants from stressors such as heat or drought. In the last 20 years, sensitization to nsLTP and consequent reactions to plant foods has become an increasing concern. **Aim:** The aim of this paper is to review the evidence for the structure and function of nsLTP allergens, and cross-reactivity, sensitization, and epidemiology of nsLTP allergy. **Materials and Methods:** A Task Force, supported by the European Academy of Allergy & Clinical Immunology (EAACI), reviewed current evidence and provide a signpost for future research. The search terms for this paper were "Non-specific Lipid Transfer Proteins", "LTP syndrome", "Prup 3", "plant food allergy", "pollen-food syndrome".

Results: Most nsLTP allergens have a highly conserved structure stabilised by 4disulphide bridges. Studies on the peach nsLTP, Pru p 3, demonstrate that nsLTPs are very cross-reactive, with the four major IgE epitopes of Pru p 3 being shared by nsLTP from other botanically related fruits. These nsLTP allergens are to varying degrees resistant to heat and digestion, and sensitization may occur through the oral, inhaled or cutaneous routes. In some populations, Pru p 3 is the primary and sole sensitizing allergen, but many are poly-sensitised both to botanically un-related nsLTP in foods, and non-food sources of nsLTP such as Cannabis sativa, Platanus acerifolia, (plane tree), Ambrosia artemisiifolia (ragweed) and Artemisia vulgaris (mugwort). Initially, nsLTP sensitization appeared to be limited to Mediterranean countries, however more recent studies suggest clinically relevant sensitization occurs in North Atlantic regions and also countries in Northern Europe, with nsLTP sensitisation profiles being broadly similar.

Discussion: These robust allergens have the potential to sensitize and provoke symptoms to a large number of plant foods, including those which are raw, cooked or processed. It is unknown why some sensitized individuals develop clinical symptoms to foods whereas others do not, or indeed what other allergens besides Pru p 3 may be primary sensitising allergens. It is clear that these allergens are also relevant in non-Mediterranean populations and there needs to be more recognition of this.

Conclusion: Non-specific LTP allergens, present in a wide variety of plant foods and pollens, are structurally robust and so may be present in both raw and cooked foods. More studies are needed to understand routes of sensitization and the world-wide prevalence of clinical symptoms associated with sensitization to these complex allergens.

KEYWORDS allergy, epidemiology, food, lipid transfer protein, LTP, sensitization

1 | BACKGROUND

Non-specific lipid-transfer proteins (nsLTPs) are a ubiquitous multigene family of molecules expressed in the Plant kingdom, starting life 470 million years ago in Marchantiophyta, the first plants grown on land after red and green algae.¹ Despite their longevity, nsLTPs were first discovered in plants only about 40 years ago, by Jean-Claude Kader, whose numerous and fundamental studies advanced the botanical knowledge regarding this family of proteins.² The biological longevity of nsLTPs throughout time is probably also due to their important role in defence of plants against biotic and abiotic stress and their expression in different plant tissues under different environmental conditions.³ Their immunoglobulin E (IgE)-binding capacity was first demonstrated in peach fruits and Parietaria pollen, and then

was detected by IgE immunoblotting in peach, cherry, apricot and plum. It was found not to be birch (*Betula verrucosa*) cross-reactive, unlike the Bet v 1 homologues, which at the time were considered to be the main trigger of allergic symptoms to fruits. The novel nsLTP allergens seemed initially to be an important cause of symptoms only in southern Europe.⁴ The peach nsLTP, Pru p 3, was first fully sequenced and characterized simultaneously in Italy and Spain, identifying a 9 kDa protein, of 91–93 amino acids long, with a high p (isoelectric point) and a conserved 8 cysteine motif, now considered as the major structural characteristic shared by all members of the nsLTP protein family.^{5,6}

Many further studies have demonstrated that nsLTPs are relevant allergens of several pollens and plant foods belonging to both the widespread Rosaceae family and many other distantly related species and can elicit systemic reactions in nsLTP sensitized subjects.⁷ Unlike Bet v 1-related or profilin-related food allergy, reactions involving LTP allergens have also been demonstrated to occur with cereals.⁸ It has also been observed that nsLTP sensitization could induce other food-related allergic conditions, including 'Baker's' asthma and food-dependent, exercise-induced anaphylaxis (FDEIA).^{9,10} Sensitization to nsLTP and the elicited clinical reactions were initially mostly reported to affect subjects living in the Mediterranean area, with those who were not pollen-sensitized typically experiencing more severe reactions.^{11,12}

In some countries (notably Italy and Spain), y Pru p 3, the nsLTP from peach (*Prunus persica*), is the first sensitizing allergen in children who may subsequently develop sensitization to a larger number of nsLTP-containing foods.¹³ According to the data published from a number of different countries, Pru p 3 is also a marker allergen in adult-onset symptoms attributed to nsLTP allergens. However, nsLTPs present in different routes of exposure and contact and different climate zones.^{14,15} The mechanism underlying the induction of sensitization to Pru p 3 has recently been demonstrated, identifying the role of the ligand in the allergic immune response which may lead to further insights into how nsLTP sensitization may develop.¹⁶

Over the last 30 years, many researchers have contributed to our understanding of the structure, function and IgE-binding properties of nsLTPs, providing evidence of the clinical entity of nsLTP syndrome(s), now well recognized by the entire International Scientific Community. However, the challenging nature of this condition, the characterization of an increasing number of nsLTP proteins in different foods and the complexities surrounding sensitization mechanisms led to the formation of a Task Force, supported by the European Academy of Allergy & Clinical Immunology (EAACI). The Task Force's aim was to produce a summary of the current evidence and provide a signpost for future research needs to ensure that this fascinating condition can be more effectively managed in the future. This paper is the first of the two papers, with the second one reviewing the evidence for the diagnosis and management of symptoms linked to nsLTP sensitization. The search terms for this paper were 'Non-specific Lipid Transfer

Proteins', 'LTP syndrome', 'Pru p 3', 'plant food allergy', 'pollen-food syndrome'.

2 | MOLECULAR STRUCTURE AND BIOLOGICAL FUNCTION OF nsLTPs

nsLTPs form a large protein family, known to occur in land plants, where they are abundantly expressed in most tissues.^{1,17} These nsLTPs are small, non-glycosylated proteins of 6–9 kDa and basic isoelectric points of 9.² They are encoded by multigene families and occur in many flowering plants.¹⁸ Currently, 46 nsLTPs have been listed by the WHO/IUIS Allergen Nomenclature Sub-Committee (Table 1). Most nsLTPs are synthesized with an N-terminal signal peptide, which directs the proteins to the extracellular, apoplastic space. Several nsLTPs also contain a C-terminal sequence motif for post-translational addition of a glycosylphosphatidylinositol (GPI)-anchor, such GPI-anchors that target the nsLTPs to the plasma membrane surfaces.^{19,20} Based on their molecular mass, nsLTPs are grouped into two types, LTP1 (9–10 kDa; around 90 amino acids) and LTP2 (6-7 kDa; around 70 amino acids).²¹ However, the majority of allergenic nsLTPs belong to the nsLTP1 type.

Structural studies on nsLTPs from wheat, maize, barley and rice were performed by x-ray and / or nuclear magnetic resonance analyses^{22–26} and the crystal structure of Prup 3 was described in 2005 by Pasquato and colleagues²⁷ (see Figure 1). These studies confirmed the overall highly conserved structure of nsLTPs, consisting of four alphahelices, which are stabilized by four highly conserved disulphide bridges and connected by flexible loops. The characteristic of the nsLTP structure is an internal, tunnel-like, hydrophobic cavity, which allows binding and transportation of various lipids^{2,17} (see Figure 2). The major biological role of nsLTPs seems to be the transfer and deposition of lipids for the assembly of complex barrier polymers on the surface of plant tissues and organs, such as waxes on leaves²⁸, suberin in seeds and roots, and sporopollenin in pollen grains.²⁹ Such barriers are required to control the fluxes of gases and liquids, but are also important for the protection of the plants from biotic and abiotic stress. Indeed, the involvement of nsLTPs in signalling during pathogen attack^{30,31} and in the defence against temperature extremes or drought has been shown.³²

Structural studies of liganded nsLTPs provided evidence that the hydrophobic internal cavity can bind to a range of different ligands including fatty acids, acyl-coenzyme A and phospholipids, thus showing great flexibility of the cavity. Furthermore, ligand binding affects the orientation of certain conserved amino acid residues at the C-terminal region and induces a conformational change as shown for Pru p 3, Jug r 3 and Mal d 3.^{33–35} These structural modifications can also modify the IgE-binding capacity of nsLTPs.^{33–35} Moreover, evidence has accumulated that the lipid-ligand bound to Pru p 3 can act as an adjuvant to promote sensitization to Pru p 3 via CD1d-mediated activation of invariant natural killer T-cells (iNKTs).¹⁶ These findings show that nsLTP-lipid interaction can increase the allergenic properties of nsLTPs and might promote sensitization to these proteins.³⁶

⁴└─WILEY-ॐ

TABLE 1 Allergen nomenclature as agreed by the WHO/IUIS Allergen Nomenclature Sub Committee

	Species	Allergen	Biochemical name	MW	UniProt
1	Ambrosia artemisiifolia (Short ragweed)	Amb a 6	nsLTP 1	10 kDa	O04004
2	Apium graveolens (Celery)	Api g 2	nsLTP 1	9 kDa	E6Y8S8
3	Apium graveolens (Celery)	Api g 6	nsLTP 2	7 kDa	P86809
4	Arachis hypogaea (Peanut, groundnut)	Ara h 9	nsLTP 1	9.8 kDa	B6CEX8
5	Arachis hypogaea (Peanut, groundnut)	Ara h 16	nsLTP 2	8.5 kDa	-
6	Arachis hypogaea (Peanut, groundnut)	Ara h 17	nsLTP 1	11 kDa	-
7	Artemisia vulgaris (Mugwort, wormwood)	Art v 3	nsLTP 1	12 kDa	P0C088
8	Asparagus officinalis (Asparagus)	Aspa o 1	nsLTP 1	9 kDa	-
9	Brassica oleracea (Cabbage and others)	Bra o 3	nsLTP 1	9 kDa	-
10	Cannabis sativa (Indian hemp)	Can s 3	nsLTP 1	9 kDa	W0U0V5
11	Castanea sativa (Chestnut)	Cas s 8	nsLTP 1	9 kDa (red)	-
12	Citrus limon (Lemon)	Cit I 3	nsLTP 1	9.6 kDa	P84160
13	Citrus reticulata (Tangerine)	Cit r 3	nsLTP 1	9 kDa	P84161
14	Citrus sinensis (Sweet orange)	Cit s 3	nsLTP 1	9.46 kDa	P84161
15	Corylus avellana (Hazelnut)	Cor a 8	nsLTP 1	9 kDa	Q9ATH2
16	Fragaria ananassa (Strawberry)	Fra a 3	nsLTP 1	9 kDa	Q8VX12
17	Helianthus annuus (Sunflower)	Hel a 3	nsLTP 1	9 kDa	Q7X9Q5
18	Hevea brasiliensis (Para rubber tree (latex))	Hev b 12	nsLTP 1	9 kDa	Q8RYA8
19	Juglans regia (English walnut)	Jug r 3	nsLTP 1	9 kDa	C5H617
20	Lactuca sativa (Cultivated lettuce)	Lac s 1	nsLTP 1	9 kDa	-
21	Lens culinaris (Lentil)	Len c 3	nsLTP 1	9 kDa	A0AT29
22	Lupinus angustifolius (Narrow-leaved blue lupin)	Lup an 3	nsLTP 1	11 kDa	-
23	Malus domestica (Apple)	Mal d 3	nsLTP 1	9 kDa	Q5J026
24	Morus nigra (Mulberry)	Mor n 3	nsLTP 1	10 kDa	P85894
25	Musa acuminata (Banana)	Mus a 3	nsLTP 1	9 kDa	P86333
26	Olea europaea (Olive)	Ole e 7	nsLTP 1	9.5 kDa	P81430
27	Parietaria judaica (Wall Pellitory)	Par j 1	PhosphoLTP	15 kDa	P43217
28	Parietaria judaica (Wall Pellitory)	Par j 2	PhosphoLTP	10-14 kDa	P55958
29	Parietaria officinalis (Pellitory)	Par o 1	PhosphoLTP	15 kDa	-
30	Phaseolus vulgaris (Green bean, French bean)	Pha v 3	nsLTP 1	8.8-9.0 kDa	D3W146
31	Platanus acerifolia (London plane tree)	Pla a 3	nsLTP 1	10 kDa	-
32	Prunus armeniaca (Apricot)	Pru ar 3	nsLTP 1	9 kDa	P81651
33	Prunus avium (Sweet cherry)	Pru av 3	nsLTP 1	10 kDa	Q9M5X8
34	Prunus domestica (European plum)	Pru d 3	nsLTP 1	9 kDa	P82534
35	Prunus dulcis (Almond)	Pru du 3	nsLTP 1	9 kDa	C0L0I5
36	Prunus persica (Peach)	Pru p 3	nsLTP 1	10 kDa	P81402
37	Punica granatum (Pomegranate)	Pun g 1	nsLTP 1	9 kDa	A0A059STC4
38	Pyrus communis (Pear)	Pyr c 3	nsLTP 1	9 kDa	Q9M5X6
39	Rubus idaeus (Red raspberry)	Rub i 3	nsLTP 1	11 kDa	Q0Z8V0
40	Sinapis alba (Yellow mustard)	Sin a 3	nsLTP 1	12.3 kDa	E6Y2L9





FIGURE 1 Structure of lipid-transfer protein allergens

So far, it has been shown that four major (immunodominant) IgE epitopes of Pru p 3 are shared by LTPs of fruits from the Rosaceae family.^{37,38} In a cohort of Spanish and Italian peach allergic patients, a Th2-dominated response was identified, and among several T-cell epitopes, immunodominant T-cell-activating regions of Pru p 3 were detected.³⁹ In contrast, other nsLTPs from more distantly related species, such as Cor a 8 in hazelnut (Corylus avellane) and Hel a 3s in sunflower seed (Helianthus annuus), do not contain these conserved epitope sequences. In a study on patients allergic to hazelnut and peach, the immunodominant role of Pru p 3 was confirmed, but no Cor a 8-specific T-cell epitopes were detected suggesting cross-reactivity rather than genuine primary sensitization to Cor a 8.40 These molecular data suggest that Pru p 3 could be the primary sensitizer behind nsLTP-driven food allergy in some populations.

The disulphide bridges contribute to the folding of nsLTPs into a compact structure with high thermal and proteolytic stability, which might explain the high allergenicity of nsLTPs.^{41,42} In vitro experiments confirmed the stability of Pru p 3 and its resistance to digestion and heat treatment in an acidic environment by circular dichroism (CD) spectroscopy . However, when performing these experiments under neutral pH conditions, the stability of these proteins is reduced as shown for Prup 3 and Cor a 8 which might be due to the cleavage of disulphide bonds at neutral pH and high temperature.⁴³⁻⁴⁵ Interestingly, stability to endolvsosomal degradation also seems to differ within the nsLTP protein family, since Pru p 3 showed higher resistance as compared to Cor a 8.⁴⁰ These data point towards a selective resistance of individual nsLTPs. It remains to be shown, whether the

BOX 1. Definitions

nsLTP sensitization-sensitization to one or more nsLTPs nsLTP allergy-clinically relevant sensitization to one or more nsLTPs

💐_WILEY___5

nsLTP syndrome-multiple plant food allergies attributed to nsl TP sensitization

binding of lipids or the structural differences among individual members of the nsLTP family might cause this selective resistance ⁴⁴

The abundancy of allergenic nsLTPs in the culprit food can also contribute to the allergenic risk. While the concentration of Pru p 3 in peach peel is comparatively high (approx. 6 mg/g peach peel; it is less in apple peel (approx. 66 µg/g tissue).^{45,46} Regarding tree nuts, nsLTPs from walnut and hazelnut are accumulated in the brown skin of the embryo and the concentration per nut is comparatively low compared to peach.47

HIERARCHY OF CROSS-REACTIVITY 3

Patients suffering from an 'nsLTP syndrome' frequently display symptoms with multiple plant-derived foods due to the ubiquitous distribution of these pan-allergens.^{48,49} The clinical relevance of these allergens, as well as the primary sensitizer, varies greatly depending on the patient's age and geographic area.⁵⁰ Initial epidemiological studies on nsLTP sensitization have mainly been conducted in southern Europe on adult or adolescent populations.⁴⁹ In this region, most allergists agree that peach is the primary driving source behind nsLTP sensitization, although nsLTPs of some pollen, particularly plane tree and mugwort, have been proposed as contributing to sensitization.⁵¹ According to Palacin's study, where the IgE-binding capacity of a number of purified LTPs was analysed using sera from nsLTP syndrome patients; Pru p 3 (peach) was the most frequently recognized, followed by Mal d 3 (apple), Cit r 3 (orange), Bra o 3 (cabbage) Sin a 3 (mustard) and the tree nut LTPs, Jug r 3 (walnut) and Cas s 8 (chestnut). In contrast, Tri a 14, the wheat nsLTP, was seldom recognized in the population studied.52

Furthermore, IgE binding frequently does not predict clinical cross-reactivity; most patients did not report any clinical reaction to lentil, soybean, maize and sesame, despite specific IgE levels to the nsLTPs in these foods. This might suggest either that in these foods, nsLTPs are less abundant or thermal processing reduces allergenicity.^{53,54} So, patients with an 'nsLTP syndrome' are usually polysensitized to multiple foods, to some extent, influenced by the amount of allergen present in the sources, the route of sensitization, geographical patterns and the sensitizing agent. Primary sensitization through Can s 3 exposure might, for example, result in a different array of hierarchical order of food nsLTP reactivity.⁵⁵

4 | ROUTES OF SENSITIZATION TO nsLTP

Sensitization to nsLTP can occur in several different ways. The characterization of lipid ligands bound to nsLTPs has shed light on the potential innate stimulatory properties of nsLTPs, which might account for their high sensitization potential.^{16,56} In early life, sensitization to foods such as milk, egg and peanut may occur through cutaneous exposure due to skin barrier disruption and exposure via the epi-cutaneous route, with IgG1+ B-cell immunity preceding the initiation of IgE responses.^{57,58} Sensitization to nsLTP may also occur through the cutaneous route; is reported that peach-induced contact urticaria has been linked to nsLTP sensitization.⁵⁹ Sensitization to foods via the gastrointestinal tract is a common cause of food-allergic reactions, and this is also true of nsLTP allergens. Tordesillas et al. demonstrated that Pru p 3 has a high capacity to cross the gastric barrier, a factor that could be relevant to the capacity for sensitization by the gastric route.⁶⁰

Just as reactions to plant foods also frequently occur through sensitization to inhaled pollen aeroallergens provoking cross-reactivity to plant foods, sensitization to nsLTP in foods via inhalation may also occur.⁶¹ Case reports describe a patient who developed respiratory symptoms when exposed to peach tree crops, and a second patient who developed rhinitis after working in a wholesale fruit warehouse, both of whom subsequently developed severe reactions to eating peaches and other foods.^{62,63} Garcia and colleagues subsequently showed that peach leaves contain Pru p 3, which might explain up to 10% of residents in an area with extensive orchard-tree crops were mono-sensitized to Pru p 3, and the finding that Pru p 3 is an occupational respiratory allergen in peach crop workers.^{19,64} nsLTP in a variety of foods has also been shown to be relevant in occupational allergy in those involved in processing other plant food crops including asparagus, wheat and maize.^{9,65-69}

Rodriguez and colleagues generated animal models of anaphylaxis to Pru p 3 following sensitization via the nasal route in combination with lipopolysaccharide (LPS).⁷⁰ Recently, Rosace et al. have described a progressive remodelling process of the oral mucosa, along with respiratory allergy progression, using profilin as an allergen.⁷¹ Co-localization of CD11+ and CD4+ cells demonstrates that the oral mucosa is an immunocompetent organ and can contribute to the progression from respiratory to food allergy in Bet v 1 homologues, profilin and nsLTP allergic syndromes.

Respiratory exposure and subsequent clinical reactivity to nsLTP can also occur due to inhalation of Cannabis sativa. The first description of a case of a patient sensitized to Pru p 3, with nsLTP sensitization mediated by smoking marijuana, was published by Gamboa in 2007 and has been followed by other publications supporting the relevance of C. sativa as a potential source of nsLTP sensitization and cannabis-related allergy.^{14,72-74} Research from northern/central Europe suggests marijuana inhalation is an effective entry route to nsLTP sensitization in the absence of Pru p 3-mediated allergy.⁷⁵⁻⁷⁷ Decuyper and colleagues reported that 45% of 120 patients allergic to C. sativa and sensitized to Can s 3, and the nsLTP in C. sativa had severe and generalized allergy to plant foods.78 Passive inhalation of smoke from C. sativa can also lead to sensitization to Can s 3, and other nsLTP allergens.⁷⁹ However, sensitization to Can s 3 may not be universal; a publication from the United States, reporting on the sequencing of allergens from 23 patients with a positive skin prick test to a crude extract of C. sativa, did not find anything sensitized to Can s 3.80

5 | SENSITIZATION TO POLLEN nsLTP ALLERGENS

Sensitization to nsLTP in plant foods via the inhaled route may also involve nsLTP in pollens.

The existence of pollen food syndrome mediated by nsLTPs was first described by García-Sellés in 2002.⁸¹ Pollen nsLTPs such as Art v 3 from Artemisia vulgaris (mugwort) and Pla a 3 from Platanus acerifolia (plane tree) share partial cross-reactivity with Rosaceae nsLTPs, and could thus initiate the sensitization process and/or contribute to the progressive recognition of multiple nsLTPs.⁸² In some areas, high exposure to nsLTP-containing pollens could result in the development of allergic respiratory symptoms, with a proportion of individuals also affected by nsLTP-driven food allergy.^{15,83-85} For example, the pollen allergen Pla a 3 was first noted to play a clinically relevant role by clinicians in Barcelona, an area with high Platanus pollen exposure where a significant percentage of pollen-allergic patients (12%) are mono-sensitized to Platanus.^{86,87} Thus, Pla a 3 sensitization is associated with nsLTP food sensitization, and in inhibition assays plane pollen extracts inhibited IgE binding to food nsLTPs by between 50% and 100%.88

Likewise, some with nsLTP sensitization to foods are also affected by respiratory symptoms due to nsLTP allergens in the pollen, with cross-sectional studies suggesting that in some cases the nsLTP within the pollen is the primary sensitizer.^{84–86} Studies suggest that Art v 3 could also act as a primary sensitizer, demonstrating the clinical relevance of pollen nsLTPs in areas of high exposure to Artemisia pollen.^{85,89} For example, *Artemisia* is the dominant allergenic pollen in China, and studies have demonstrated that this might be a mechanism of sensitization to Pru p 3, and the cause of peach allergy in the Chinese population.⁸³ In areas with high exposure to either Artemisia pollen as Gran Canaria Island⁴ or to Platanus pollen as Barcelona,





FIGURE 2 Structure of Pru p 3: (A) surface model, (B) ribbon model, (C) internal cavity with ligand binding

epidemiological studies have also demonstrated a broader recognition pattern to nsLTPs in sensitized and symptomatic patients.⁵²

It has also been demonstrated that patients sensitized to Pru p 3 can show sensitization to Art v 3 and/or Pla a 3 even in geographical areas lacking Artemisia/plane tree pollen, presumably due to IgE cross-reactivity.^{84,85} Despite a lack of Artemisia pollen, a recent UK study reported a strong correlation between Pru p 3 and both Pla a 3 and Art v 3, with the *r*-value being greater than that for a matched group of

Italian subjects.⁹⁰. A study performed in patients with a clinical history of peach anaphylaxis demonstrated dual sensitization to both Pru p 3 and Art v 3, and therefore, even in the absence of atmospheric mugwort pollen, there is a possibility that these patients could still develop a respiratory allergy if exposed to mugwort.^{88,91} The presence of nsLTP sensitization in the United Kingdom, and other parts of more northerly European latitudes suggests the primary sensitizing nsLTP might be from a pollen rather than from a food.^{90,92}

Allergenic pollen nsLTPs have also been described from other sources including Parietaria judaica (Par j 2) and olive tree (Oleum Europea) (Ole e 7). Sensitization to Parietaria has been reported in subjects from northern European countries free from this weed, suggesting this is another expression of cross-reactivity and probably to relevant.^{90,92} A study by Tordesillas and colleagues in 2011 reported that there was no cross-reactivity between nsLTP allergens from olive and parietaria, or between the pollen nsLTP and nsLTP allergens from foods.⁹³ However, a more recent study from Ciprandi and colleagues reported that in a group of children with allergic rhinitis due to Parietaria pollen allergy and sensitization to Pru p 3, 50% reported a food allergy or oral allergy symptoms, and 25% reported anaphylaxis after ingesting nsLTP-containing foods.⁹⁴ It is also unclear how pertinent Ole e 7 is in terms of diagnostic relevance. Although it has less than 20% amino acid sequence identity with Pru p 3, recognition of Ole e 7 is associated with reactions to plantderived food (mainly peach, walnut and peanut), and 80% of Ole e 7 reactors are sensitized to food nsLTPs versus only 30% of Ole e 1 reactors.⁹⁵ A recent study reported that common IgG and IgE epitopes were identified in Ole e 7 and Pru p 3, with inhibition studies suggesting that primary sensitization to Ole e 7 could lead to sensitization to Pru p 3.96

There has been some suggestion that the greater the atmospheric burden of birch pollen, the lower the incidence of nsLTP sensitization, leading to a hypothesis that high levels of birch pollen protect against nsLTP allergy.⁹⁷ It is suggested one potential explanation could be lower reactivity of effector cells exposed to the nsLTPs, as described by Christensen et al., on the basis of interference by other IgE molecules bound to the high-affinity IgE receptors on effector cells.⁹⁸ However, a recent study from Belgium demonstrate that many subjects are sensitized to both nsLTP allergens in foods and pollens.⁹² A study on nsLTP sensitized adults in the UK study reported that co-sensitization to birch pollen allergens did not seem to be protective against severe reactions, with the use of adrenaline and subsequent hospital attendance being significantly reported.⁹⁰

6 | EPIDEMIOLOGY

Epidemiological data have most often involved the reporting of the rates of sensitization to the peach nsLTP Pru p 3. Using ISAC microarray, Scala et al. reported that 9% of 23,000 patients living in central and southern Italy were sensitized to Pru p 3.⁹⁹ In Spain, Barber et al. detected Pru p 3 sensitization in 12% of 2000 Spanish



FIGURE 3 World-wide reported lipid-transfer protein sensitization

patients with pollen allergy.^{100,101} This proportion was confirmed by Gonzalez-Mancebo et al. who reported that 53/430 (12.3%) of outpatients seen in their allergy unit for any reason were sensitized to Pru p 3.¹⁰² Those sensitized to Pru p 3 may not always be symptomatic and the symptom severity may vary. A study on nsLTP syndrome reported that in a group of 87 subjects sensitized to Pru p 3, 44% experienced anaphylaxis, 43% skin or oropharyngeal reactions, and 13% were asymptomatic.¹⁰³ It is generally accepted that nsLTP sensitization often occurs early in life.^{104,105} Pru p 3 sensitization among children in a Spanish cohort was twice as high as that in adults.¹⁰⁰ Another study reported that the level of Pru p 3 sensitization in a large population of Italian subjects peaks around the start of the third decade of life.¹⁰⁶ Further, a study in LTP-allergic adults suggested that the younger the age of onset, the higher the titres of specific IgE to Pru p 3 that are detected in patients.¹³

In addition to those who are mono-sensitized to Pru p 3, studies have also investigated subjects with sensitization to multiple nsLTP and usually have clinically relevant allergy to different plant foods. These individuals have the so-called LTP syndrome.¹⁰⁷ Rosaceae fruits and nuts are the main plant food allergens involved in this syndrome.¹⁰⁸ In a nationwide review of 17 Italian allergy clinics, 210 subjects sensitized to nsLTP reported reactions to multiple food triggers including peach (85%), apple (35%) and tree nuts (20%). Offending foods included also other Rosaceae (apricot, plum, pear, almond, strawberry), cereals (maize, beer, rice), kiwifruit, grapes, lettuce, cauliflower and broccoli.¹²

Following initial identification around 20 years ago, nsLTP sensitization appears to be most prevalent in the 'Mediterranean basin', a climate area which is defined by a southern European latitude. Although precise epidemiologic data from the general population are lacking, there is little doubt that sensitization to nsLTPs is a significant cause of primary food allergy in adults living in the Mediterranean basin^{12,109} where it accounts for a large-if not the largest-proportion of food-induced anaphylaxis.⁸ Interestingly, there are marked geographic variations in the prevalence of LTP hypersensitivity within single countries. This has been best shown for Italy due to its very different climate zones, that is, northern (continental climate) and central southern (Mediterranean climate), with nsLTP sensitization dominating in the latter.^{12,110} Similar observations have been seen in two Spanish studies, where nsLTP-sensitization prevalence among patients with seasonal allergy in different areas was 2% and 40%, respectively.^{52,101} However, a Mediterranean climate on its own is not the only factor in the development of sensitization to nsLTP. A high prevalence of nsLTP sensitization is also found in Portugal, which, despite being part of the Iberian Peninsula, is an Atlantic country.¹¹¹ In addition, those living in areas of Spain and Italy characterized either by a continental or Atlantic climate may also be sensitized to nsLTP.^{12,112}

The geographic variability of nsLTP sensitization can be best appreciated in multinational studies, where the differences between northern and southern European countries are evident.¹¹³ However, the presence of nsLTP sensitization in European countries other than the traditional 'endemic' Mediterranean areas is also emerging-see Figure 3. A study from Belgium reported nsLTP sensitization rates in selected pollen and/or plant food allergic children and adults to exceed 20% (albeit clinically irrelevant in most cases).⁹² Other reports have been published from Austria.^{114,115} France.⁷⁵ Belgium.¹¹⁶ Germany,^{117,118} Poland¹¹⁹ and the United Kingdom.^{90,120}. Outside Europe, nsLTP hypersensitivity and allergy seem relevant in certain areas of China.^{83,121} A single report has been published from

Australia.¹²² Remarkably, nsLTP sensitization has not yet been reported in the Americas or Africa.

7 | FUTURE RESEARCH NEEDS

Although experimental and clinical studies of nsLTP sensitization have increased significantly in the last decade, there are still several issues not fully understood. Although some definitions have been agreed by the Task Force, it is important to develop internationally agreed criteria and definitions for nsLTP sensitization and nsLTP syndromes, especially when determining the prevalence of nsLTP sensitization. In paediatric populations, nsLTP sensitization has some peculiarities: the lower risk of severe reactions compared to adults and the common occurrence of subclinical sensitization. Continent-wide epidemiological prevalence studies in both adults and children are urgently needed to evaluate the prevalence of nsLTP sensitization, its geographical distribution, risk/protective factors, genetic predisposition, clinical features and natural history.

The variation in the severity of symptoms also warrants investigation. It has already been demonstrated in one study that there is a link between multiple nsLTP sensitizations and systemic reactions.⁹² It is possible that patients sensitized to multiple nsLTPs, with strong concomitant exposure to pollen-derived nsLTP allergens, could display a broad T-cell stimulatory repertoire, similar to that reported for profilin allergy.^{123,124} The reports of more severe reactions in those with nsLTP sensitization but without pollinosis confirms the complex nature of food-allergic mechanisms and identifies a need for

Key Messages

- nsLTPs are a large group of allergens found in land plants; the majority of allergenic nsLTPs belong to LTP (type)1 (9–10 kDa; around 90 amino acids).
- 2. nsLTPs are highly cross-reactive with particular hierarchies of sensitization depending on the primary sensitizer.
- 3. In Italy and Spain, sensitization frequently occurs in childhood, with peach nsLTP (Pru p 3) often being the sole sensitizing nsLTP.
- Sensitization may be both via the oral, inhaled or cutaneous route, and may be initiated by a pollen nsLTP.
- Although first described in Italy and Spain, nsLTP sensitization occurs in many European countries and has also been reported in China.
- When an allergy to a plant food is suspected, nsLTP allergens should not be ruled out as a trigger, wherever the geographical location

studies into such mechanisms that may ameliorate severe nsLTP reactions. Evaluation of sensitization profiles to pollen and food in different regions of Europe might contribute to the identification of primary sensitizers, whether they be from plant foods, pollens or other inhalants such as cannabis, thus paving the way for prevention and control measures.

8 | CONCLUSION

Investigating the structure and function of nsLTP has led to an understanding of their allergenic potential, which is exacerbated due to the high degree of cross-reactivity between nsLTP from different plant species. These allergens are present in fruits, vegetables, nuts, seeds, legumes and cereals, and food nsLTPs have been shown to be able to sensitize through the oral, respiratory and cutaneous routes. Exposure to peaches has been linked to the development of primary sensitization to the peach nsLTP, Pru p 3, which dominates in the Mediterranean areas of Europe. However, the sensitization process might depend on both individual (dietary/ occupational) and/or geographical factors. Although data suggest it is an important cause of primary food allergy in the southern European countries, the extent of nsLTP sensitization could also be significant in the rest of Europe. In this context, the role of exposure to pollen needs further study, with some pollens, such as those of mugwort and plane tree, potentially playing a major role in sensitization to food nsLTPs. Despite high levels of sensitization to Pru p 3, many children remain asymptomatic or experience only mild skin symptoms. Moderate or severe clinical symptoms are much more likely to present in teenage years or in adults, despite levels of Prup 3 falling. Sensitization to nsLTP is common and likely to be much more widespread than has been previously thought. It is therefore likely to be an important cause of plant food allergy in many adults in Europe and beyond.

ACKNOWLEDGEMENT

Not applicable.

CONFLICTS OF INTEREST

The following authors declare that they have competing interests: K. Hoffmann-Somergruber has received support from the Austrian Science funds (FWF) by SFB Subproject F-4603, L. Cecchi has received honoraria from Malseci, Menarini, Mylan and Thermofisher, D Barber is a scientific advisor to ALK and AIMMUNE, E.Scala has received consultant arrangements and speakers' bureau participation from Stallergenes and Thermo Fisher Scientific. M. Fernández-Rivas declares grants from Spanish Government (ISCIII; MINECO), Aimmune Therapeutics and Diater to her organization; consultancy fees from Aimmune, DBV, Novartis, SPRIM; lecture fees from Aimmune, ALK, Diater, GSK, HAL Allergy, Thermofisher Scientific, outside the submitted work. R. van Ree undertakes consultancies for HAL Allergy BV, Citeq BV and Angany Inc, has received honoraria from HAL 10 |___WILEY_

Allergy BV, ThermoFisher Scientific and funding support from the European Commission, Dutch Science Foundation, Health Holland. I. Skypala, D. Ebo, M. Faber, A Diaz-Perales, I Swoboda, F Gomez, R. Asero, E. Pastorello, O. Luengo, S Till, J Bartra, A Konstantinopoulos have no conflict of interests to declare

AUTHOR CONTRIBUTIONS

Isabel Skypala established the Task Force, set up and chaired the meetings, determined the sections of the manuscript, merged all of the sections from individual contributors, added key points, wrote the abstract, made amendments to the manuscript following both internal and external reviews and submitted the manuscript in its final format. Ricardo Asero, Domingo Barber, Lorenzo Cecchi, Arazeli Diaz Perales, Karin Hoffmann-Sommergruber, Elide A. Pastorello, and Ines Swoboda attended and contributed to the meetings, wrote individual sections of the manuscript, reviewed it once all the sections had been put together and following internal and external review. Joan Bartra, Didier G. Ebo, Margaretha A. Faber, Montserrat Fernández-Rivas, Francesca Gomez, A. P. Konstantinopoulos, Olga Luengo, Ronald van Ree, Enrico Scala and Stephen J. Till attended and contributed to the meetings and reviewed the manuscript following review.

ORCID

Isabel J. Skypala b https://orcid.org/0000-0003-3629-4293 Ricardo Asero b https://orcid.org/0000-0002-8277-1700 Domingo Barber b https://orcid.org/0000-0002-5488-5700 Lorenzo Cecchi b https://orcid.org/0000-0002-0658-2449 Arazeli Diaz Perales b https://orcid.org/0000-0002-1093-3627 Karin Hoffmann-Sommergruber b https://orcid.org/0000-0002-8830-058X

Elide A. Pastorello b https://orcid.org/0000-0002-3753-1783 Ines Swoboda b https://orcid.org/0000-0002-9164-1721 Joan Bartra b https://orcid.org/0000-0001-7767-4730 Didier G. Ebo b https://orcid.org/0000-0003-0672-7529 Margaretha A. Faber b https://orcid.org/0000-0002-1277-5052 Montserrat Fernández-Rivas b https://orcid.org/0000-0003-1748-2328

Francesca Gomez https://orcid.org/0000-0003-1607-6686 Anastasios. P. Konstantinopoulos https://orcid.org/0000-0002-7060-653X

Olga Luengo https://orcid.org/0000-0002-8905-2776 Enrico Scala https://orcid.org/0000-0002-9391-9168 Stephen J. Till https://orcid.org/0000-0003-4518-3093

REFERENCES

- Edstam MM, Viitanen L, Salminen TA, Edqvist J. Evolutionary history of the non-specific lipid transfer proteins. *Mol Plant*. 2011;4:947-964.
- Kader JC. Lipid transfer proteins in plants. Ann Rev Plant Physiol Mol Biol. 1996;47:627-654.
- 3. Liu F, Zhang X, Lu C, et al. Non-specific lipid transfer proteins in plants: presenting new advances and an integrated functional analysis. *J Exp Bot.* 2015;66:56635681.

- 4. Pastorello EA, Ortolani C, Farioli L, et al. C. Allergenic crossreactivity among peach, apricot, plum, and cherry in patients with oral allergy syndrome: an in vivo and in vitro study. J Allergy Clin Immunol. 1994;94:699-707.
- Pastorello EA, Farioli L, Pravettoni V, et al. The major allergen of peach (Prunus persica) is a lipid transfer protein. J Allergy Clin Immunol. 1999;103:520-526.
- Sánchez-Monge R, Lombardero M, García-Sellés FJ, Barber D, Salcedo G. Lipid-transfer proteins are relevant allergens in fruit allergy. J Allergy Clin Immunol. 1999;103:514-519.
- Asero R, Antonicelli L, Arena A, et al. Causes of food-induced anaphylaxis in Italian adults: a multi-centre study. Int Arch Allergy Immunol. 2009;150:271-277.
- Pastorello EA, Farioli L, Stafylaraki C, et al. Wheat-dependent exercise-induced anaphylaxis caused by a lipid transfer protein and not by ω-5 gliadin. Ann Allergy Asthma Immunol. 2014;112:386-387.
- 9. Palacin A, Quirce S, Armentia A, et al. Wheat lipid transfer protein is a major allergen associated with baker's asthma. J Allergy Clin Immunol. 2007;120:1132-1138.
- 10. Romano A, Scala E, Rumi G, et al. Lipid transfer proteins: the most frequent sensitizer in Italian subjects with food-dependent exercise-induced anaphylaxis. *Clin Exp Allergy*. 2012;42:1643-1653.
- Pastorello EA, Farioli L, Pravettoni V, et al. Pru p 3-sensitised Italian peach-allergic patients are less likely to develop severe symptoms when also presenting IgE antibodies to Pru p 1 and Pru p 4. Int Arch Allergy Immunol. 2011;156:362-372.
- 12. Asero R, Antonicelli L, Arena A, et al. EpidemAAITO: features of food allergy in Italian adults attending allergy clinics: a multi-centre study. *Clin Exp Allergy*. 2009;39:547-555.
- 13. Pastorello EA, Farioli L, Stafylaraki C, et al. Anti-rPru p 3 IgE levels are inversely related to the age at onset of peach-induced severe symptoms reported by peach-allergic adults. *Int Arch Allergy Immunol.* 2013;162:45-49.
- Gamboa P, Sanchez-Monge R, Sanz ML, Palacín A, Salcedo G, Diaz-Perales A. Sensitization to Cannabis sativa caused by a novel allergenic lipid transfer protein, Can s 3. J Allergy Clin Immunol. 2007;120:1459-1460.
- Deng S, Yin J. Mugwort Pollen-related food allergy: lipid transfer protein sensitization and correlation with the severity of allergic reactions in a Chinese population. *Allergy Asthma Immunol Res.* 2019;11:116-128.
- Tordesillas L, Cubells-Baeza N, Gómez-Casado C, et al. Mechanisms underlying induction of allergic sensitization by Pru p3. *Clin Exp Allergy*. 2017;47:1398-1408.
- Salminen TA, Blomqvist K, Edqvist J. Lipid transfer proteins: classification, nomenclature, structure, and function. *Planta*. 2016;244:971-997.
- Boutrot F, Chantret N, Gautier MF. Genome-wide analysis of the rice and Arabidopsis non-specific lipid transfer protein (nsLTPp) gene families and identification of wheat nsLTP genes by EST data mining. *BMC Genomics.* 2008;9:86.
- Debono A, Yeats TH, Rose JK, et al. Arabidopsis LTPG is a glycosylphosphatidylinositol-anchored lipid transfer protein required for export of lipids to the plant surface. *Plant Cell.* 2009;21:1230-1238.
- Lee SB, Go YS, Bae HJ, et al. Disruption of glycosylphosphatidylinositol-anchored lipid transfer protein gene altered cuticular lipid composition, increased plastoglobules, and enhanced susceptibility to infection by the fungal pathogen Alternaria brassicicola. *Plant Physiol.* 2009;150:42-54.
- Lin CH, Li L, Lyu PC, Chang JY. Distinct unfolding and refolding pathways of lipid transfer proteins LTP1 and LTP2. *Protein J*. 2004;23:553-566.
- 22. Gincel E, Simorre JP, Caille A, Marion D, Ptak M, Vovelle F. Threedimensional structure in solution of a wheat lipid-transfer protein

from multidimensional 1H-NMR data. A new folding for lipid carriers. *Eur J Biochem*. 1994;226:413-422.

- Shin DH, Lee JY, Hwang KY, Kim KK, Suh SW. High-resolution crystal structure of the non-specific lipid-transfer protein from maize seedlings. *Structure*. 1995;3:189-199.
- Gomar J, Petit MC, Sodano P, et al. Solution structure and lipid binding of a nonspecific lipid transfer protein extracted from maize seeds. *Protein Sci.* 1996;5:565-577.
- Lerche MH, Kragelund BB, Bech LM, Poulsen FM. Barley lipidtransfer protein complexed with palmitoyl CoA: the structure reveals a hydrophobic binding site that can expand to fit both large and small lipid-like ligands. *Structure*. 1997;5:291-306.
- Lee JY, Min K, Cha H, Shin DH, Hwang KY, Suh SW. Rice nonspecific lipid transfer protein: the 1.6 A crystal structure in the unliganded state reveals a small hydrophobic cavity. J Mol Biol. 1998;276:437-448.
- Pasquato N, Berni R, Folli C, et al. Crystal structure of peach Pru p 3, the prototypic member of the family of plant non-specific lipid transfer protein pan-allergens. J Mol Biol. 2006;356:684-694.
- Liu F, Xiong X, Wu L, et al. BraLTP1, a lipid transfer protein gene involved in epicuticular wax deposition, cell proliferation and flower development in Brassica napus. *PLoS One*. 2014;9(10):e110272.
- Edstam MM, Blomqvist K, Eklöf A, Wennergren U, Edqvist J. Coexpression patterns indicate that GPI-anchored non-specific lipid transfer proteins are involved in accumulation of cuticular wax, suberin and sporopollenin. *Plant Mol Biol.* 2013;83:625-649.
- García-Olmedo F, Molina A, Segura A, Moreno M. The defensive role of nonspecific lipid-transfer proteins in plants. *Trends Microbiol*. 1995;3:72-74.
- Sun JY, Gaudet DA, Lu ZX, Frick M, Puchalski B, Laroche A. Characterization and antifungal properties of wheat nonspecific lipid transfer proteins. *Mol Plant Microbe Interact.* 2008;21: 346-360.
- Guo L, Yang H, Zhang X, Yang S. Lipid transfer protein 3 as a target of MYB96 mediates freezing and drought stress in Arabidopsis. *J Exp Bot.* 2013;64:1755-1767.
- Dubiela P, Aina R, Polak D, et al. Enhanced Pru p 3 IgE-binding activity by selective free fatty acid-interaction. J Allergy Clin Immunol. 2017;140:1728-1731.
- Dubiela P, Del Conte R, Cantini F, et al. Impact of lipid binding on the tertiary structure and allergenic potential of Jug r 3, the non-specific lipid transfer protein from walnut. *Sci Rep.* 2019;9:2007.
- Aina R, Dubiela P, Geiselhart S, et al. Distinct Lipid Transfer Proteins display different IgE-binding activities that are affected by fatty acid binding. *Allergy*. 2019;74:827-831.
- Scheurer S, Schülke S. Interaction of non-specific lipid-transfer proteins with plant-derived lipids and its impact on allergic sensitization. Front Immunol. 2018;9:1389.
- Borges JP, Barre A, Culerrier R, Granier C, Didier A, Rougé P. Lipid transfer proteins from Rosaceae fruits share consensus epitopes responsible for their IgE-binding cross-reactivity. *Biochem Biophys Res Commun.* 2008;365:685-690.
- García-Casado G, Pacios LF, Díaz-Perales A, et al. Identification of IgE-binding epitopes of the major peach allergen Pru p 3. J Allergy Clin Immunol. 2003;112:599-605.
- Schulten V, Radakovics A, Hartz C, et al. Characterization of the allergic T-cell response to Pru p 3, the nonspecific lipid transfer protein in peach. J Allergy Clin Immunol. 2009;124:100-107.
- Schulten V, Nagl B, Scala E, et al. Pru p 3, the nonspecific lipid transfer protein from peach, dominates the immune response to its homolog in hazelnut. *Allergy*. 2011;66:1005-1013.
- Asero R, Mistrello G, Roncarolo D, et al. Lipid transfer protein: a pan-allergen in plant-derived foods that is highly resistant to pepsin digestion. *Int Arch Allergy Immunol.* 2000;122:20-32.

- 42. Lindorff-Larsen K, Winther JR. Surprisingly high stability of barley lipid transfer protein, LTP1, towards denaturant, heat and proteases. *FEBS Lett.* 2001;488:145-148.
- Offermann LR, Bublin M, Perdue ML, et al. Structural and functional characterization of the hazelnut allergen Cor a 8. J Agric Food Chem. 2015;63:9150-9158.
- Bublin M, Eiwegger T, Breiteneder H. Do lipids influence the allergic sensitization process? J Allergy Clin Immunol. 2014;134:521-529.
- Gaier S, Marsh J, Oberhuber C, et al. Purification and structural stability of the peach allergens Pru p 1 and Pru p 3. *Mol Nutr Food Res.* 2008;52(Suppl 2):S220-S229.
- Sancho AI, van Ree R, van Leeuwen A. Measurement of lipid transfer protein in 88 apple cultivars. Int Arch Allergy Immunol. 2008;146:19-26.
- 47. Pastorello EA, Farioli L, Pravettoni V, et al. Lipid transfer protein and vicilin are important walnut allergens in patients not allergic to pollen. J Allergy Clin Immunol. 2004;114:908-914.
- Van Winkle RC, Chang C. The biochemical basis and clinical evidence of food allergy due to lipid transfer proteins: a comprehensive review. *Clin Rev Allergy Immunol.* 2014;46:211-224.
- 49. Salcedo G, Sanchez-Monge R, Diaz-Perales A, Garcia-Casado G, Barber D. Plant non-specific lipid transfer proteins as food and pollen allergens. *Clin Exp Allergy*. 2004;34:1336-1341.
- Asero R, Mistrello G, Roncarolo D, et al. Immunological crossreactivity between lipid transfer proteins from botanically unrelated plant-derived foods: a clinical study. *Allergy*. 2002;57: 900-906.
- Pascal M, Muñoz-Cano R, Reina Z, et al. Lipid transfer protein syndrome: clinical pattern, cofactor effect and profile of molecular sensitization to plant-foods and pollens. *Clin Exp Allergy*. 2012; 42:1529-1539.
- Palacín A, Gómez-Casado C, Rivas LA, et al. Graph based study of allergen cross-reactivity of plant lipid transfer proteins (LTPs) using microarray in a multicenter study. *PLoS One*. 2012;7(12):e50799.
- 53. Asero R. In patients with LTP syndrome food-specific IgE show a predictable hierarchical order. *Eur Ann Allergy Clin Immunol.* 2014;46:142-146.
- 54. Asero R, Mistrello G, Roncarolo D, Amato S, Falagiani P. Why do lipid transfer protein-hypersensitive patients tolerate bean (and other legumes)? *Int Arch Allergy Immunol.* 2005;137:236-240.
- 55. Decuyper, II, Faber MA, Lapeere H, et al. Cannabis allergy: A diagnostic challenge. *Allergy*. 2018;73:1911-1914.
- Cubells-Baeza N, Gómez-Casado C, Tordesillas L, et al. Identification of the ligand of Pru p 3, a peach LTP. *Plant Mol Biol.* 2017;94:33-44.
- 57. Kong J, Chalcraft K, Mandur TS, et al. Comprehensive metabolomics identifies the alarmin uric acid as a critical signal for the induction of peanut allergy. *Allergy*. 2015;70:495-505.
- Jiménez-Saiz R, Ellenbogen Y, Koenig JFE, et al. IgG1(+) B-cell immunity predates IgE responses in epicutaneous sensitization to foods. *Allergy*. 2019;74:165-175.
- Asero R. Peach-induced contact urticaria is associated with lipid transfer protein sensitization. Int Arch Allergy Immunol. 2011;154:345-348.
- Tordesillas L, Gómez-Casado C, Garrido-Arandia M, et al. Transport of Pru p 3 across gastrointestinal epithelium an essential step towards the induction of food allergy? *Clin Exp Allergy*. 2013;43:1374-83.
- Werfel T, Asero R, Ballmer-Weber BK, et al. Position paper of the EAACI: food allergy due to immunological cross-reactions with common inhalant allergens. *Allergy*. 2015;70:1079-1090.
- García BE, Lombardero M, Echechipía S, et al. Respiratory allergy to peach leaves and lipid-transfer proteins. *Clin Exp Allergy*. 2004;34:291-295.

¹² WILEY-

- Borghesan F, Mistrello G, Roncarolo D, Amato S, Plebani M, Asero R. Respiratory allergy to lipid transfer protein. Int Arch Allergy Immunol. 2008;147:161-165.
- Pérez-Calderón R, Gonzalo-Garijo MÁ, Rodríguez-Velasco FJ, Sánchez-Vega S, Bartolomé-Zavala B. Occupational respiratory allergy in peach crop workers. *Allergy*. 2017;72:1556-1564.
- Tabar AI, Alvarez-Puebla MJ, Gomez B, et al. Diversity of asparagus allergy: clinical and immunological features. *Clin Exp Allergy*. 2004;34:131-136.
- Pastorello EA, Farioli L, Robino AM, Trambaioli C, Conti A, Pravettoni V. A lipid transfer protein involved in occupational sensitization to spelt. J Allergy Clin Immunol. 2001;108:145-146.
- Guillen D, Barranco P, Palacín A, Quirce S. Occupational rhinoconjunctivitis due to maize in a snack processor: a cross-reactivity study between lipid transfer proteins from different cereals and peach. Allergy Asthma Immunol Res. 2014;6:470-473.
- Safi H, Wangorsch A, Lidholm J, et al. Identification and molecular characterization of allergenic non-specific lipid-transfer protein from durum wheat (Triticum turgidum). *Clin Exp Allergy*. 2019;49:120-129.
- Díaz-Perales A, Tabar AI, Sánchez-Monge R, et al. Characterization of asparagus allergens: A relevant role of lipid transfer proteins. J Allergy Clin Immunol. 2002;110:790-796.
- Rodriguez MJ, Aranda A, Fernandez TD, et al. LPS promotes Th2 dependent sensitisation leading to anaphylaxis in a Pru p 3 mouse model. *Sci Rep.* 2017;7:40449.
- Rosace D, Gomez-Casado C, Fernandez P, et al. Profilin-mediated food-induced allergic reactions are associated with oral epithelial remodeling. J Allergy Clin Immunol. 2019;143:681-690.
- Rojas Pérez-Ezquerra P, Sánchez-Morillas L, Davila-Ferandez G, et al. Contact urticaria to Cannabis sativa due to a lipid transfer protein (LTP). Allergol Immunopathol. 2015;43:231-233.
- Larramendi CH, López-Matas MÁ, Ferrer A, et al. Prevalence of sensitization to Cannabis sativa. Lipid-transfer and thaumatin-like proteins are relevant allergens. *Int Arch Allergy Immunol.* 2013;162:115-22.
- 74. Decuyper, II, Rihs HP, Van Gasse AL, et al. Cannabis allergy: what the clinician needs to know in 2019. *Expert Rev Clin Immunol.* 2019;15:599-606.
- Metz-Favre C, Pauli G, Bessot JC, De Blay F. Molecular allergology in practice: an unusual case of LTP sensitization. *Eur Ann Allergy Clin Immunol.* 2011;43:193-195.
- Drouet M, Hoppe A, Moreau AS, Bonneau JC, Leclere JM, Le Sellin J. Cannabis and crossed allergy with food. *Rev Pneumol Clin.* 2017;73:290-293.
- Ebo DG, Swerts S, Sabato V, et al. New food allergies in a European non-Mediterranean region: is Cannabis sativa to blame? *Int Arch Allergy Immunol.* 2013;161:220-228.
- Decuyper, II, Van Gasse AL, Faber MA, et al. Exploring the diagnosis and profile of cannabis allergy. *J Clin Immunol. Pract.* 2019;7: 983-989.
- Decuyper, II, Faber MA, Sabato V, et al. Where there's smoke, there's fire: cannabis allergy through passive exposure. J Allergy Clin Immunol Pract. 2017;5:864-865.
- Nayak AP, Green BJ, Sussman G, et al. Characterization of Cannabis sativa allergens. Ann Allergy Asthma Immunol. 2013;111:32-37.
- García-Sellés FJ, Díaz-Perales A, Sánchez-Monge R, et al. Patterns of Reactivity to lipid transfer proteins of plant foods and Artemisia pollen: an in vivo study. *Int Arch of Allergy Immunol.* 2002;128. 115-22.
- Enrique E, Ahrazem O, Bartra J, et al. Lipid transfer protein is involved in rhinoconjunctivitis and asthma produced by rice inhalation. J Allergy Clin Immunol. 2005;116:926-928.
- Gao ZS, Yang ZW, Wu SD. Peach allergy in China: a dominant role for mugwort pollen lipid transfer protein as a primary sensitizer. J Allergy Clin Immunol. 2013;131:224-225.

- Lauer I, Miguel-Moncin MS, Abel T, et al. Identification of a plane pollen lipid transfer protein (Pla a 3) and its immunological relation to the peach lipid-transfer protein, Pru p 3. *Clin Exp Allergy*. 2007;37:261-269.
- Lombardero M, García-Sellés FJ, Polo F, et al. Prevalence of sensitization to Artemisia allergens Art v 1, Art v 3 and Art v 60kDa. Cross-reactivity among Art v 3 and other relevant lipidtransfer allergens. *Clin Exp Allergy*. 2004;34:1415-1421.
- Wangorsch A, Larsson H, Messmer M, et al. Molecular cloning of plane pollen allergen Pla a 3 and its utility as diagnostic marker for peach associated plane pollen allergy. *Clin Exp Allergy*. 2016;46:764-774.
- Barber D, Díaz-Perales A, Villalba M, Chivato T. Challenges for allergy diagnosis in regions with complex pollen exposures. *Curr Allergy Asthma Rep.* 2015;15:496.
- Scala E, Till SJ, Asero R, et al. Lipid transfer protein sensitization: reactivity profiles and clinical risk assessment in an Italian cohort. *Allergy*. 2015;70:933-943.
- 89. Scala E, Cecchi L, Abeni D, et al. Pla a 2 and Pla a 3-reactivity identify plane tree-allergic patients with respiratory symptoms or food allergy. *Allergy*. 2017;72:671-674.
- Skypala IJ, Cecchi L, Shamji MH, Scala E, Till S. Lipid Transfer Protein allergy in the United Kingdom: Characterization and comparison with a matched Italian cohort. *Allergy*. 2019;74:1340-1351.
- Sánchez-López J, Tordesillas L, Pascal M, et al. J. Role of Art v 3 in pollinosis of patients allergic to Pru p 3. J Allergy Clin Immunol. 2014;133:1018-1025.
- Faber MA, Van Gasse AL, Decuyper, II, et al. IgE-reactivity profiles to nonspecific lipid transfer proteins in a north western European country. J Allergy Clin Immunol. 2017;139:679-682.
- Tordesillas L, Sirvent S, Díaz-Perales A, et al. Plant lipid transfer protein allergens: no cross-reactivity between those from foods and olive and Parietaria pollen. *Int Arch Allergy Immunol.* 2011;156:291-296.
- Ciprandi G, Del Barba P, Silvestri M, Barberi S, Tosca MA. Pru p 3 sensitization in children with allergy to Parietaria pollens. *Acta Biomed*. 2019;90:265268.
- Scala E, Abeni D, Pomponi D, et al. Ole e 1, Ole e 7, and Ole e 9: Identifying distinct clinical subsets of olive tree-allergic patients. J Allergy Clin Immunol. 2016;137:629-631.
- Oeo-Santos C, Navas A, Benedé S, et al. New insights into the sensitization to nonrelated nsLTPs from pollen and food: New role of the allergen Ole e 7. Allergy. 2019.
- Rial MJ, Sastre J. Food allergies caused by allergenic lipid transfer proteins: what is behind the geographic restriction? *Curr Allergy Asthma Rep.* 2018;18:56.
- Christensen LH, Holm J, Lund G, Riise E, Lund K. Several distinct properties of the IgE repertoire determine effector cell degranulation in response to allergen challenge. J Allergy Clin Immunol. 2008;122:298-304.
- Scala E, Alessandri C, Bernardi ML, et al. Cross-sectional survey on immunoglobulin E reactivity in 23,077 subjects using an allergenic molecule-based microarray detection system. *Clin Exp Allergy*. 2010;40:911-921.
- 100. Barber D, de la Torre F, Feo F, et al. Understanding patient sensitization profiles in complex pollen areas: a molecular epidemiological study. *Allergy*. 2008;63:1550-1558.
- Barber D, de la Torre F, Lombardero M, et al. Component-resolved diagnosis of pollen allergy based on skin testing with profilin, polcalcin and lipid transfer protein pan-allergens. *Clin Exp Allergy*. 2009;39:1764-1773.
- 102. González-Mancebo E, González-de-Olano D, Trujillo MJ, et al. Prevalence of sensitization to lipid transfer proteins and profilins in a population of 430 patients in the south of Madrid. *J Investig Allergol Clin Immunol.* 2011;21:278-82.

- Basagaña M, Elduque C, Teniente-Serra A, Casas I, Roger A. Clinical profile of lipid transfer protein syndrome in a Mediterranean area. J Investig Allergol Clin Immunol. 2018;28:58-60.
- Pascal M, Vazquez-Ortiz M, Folque MM, et al. Asymptomatic LTP sensitisation is common in plant-food allergic children from the Northeast of Spain. *Allergol Immunopathol (Madr)*. 2016;44: 351-358.
- 105. Mastrorilli C, Tripodi S, Caffarelli C, et al. Endotypes of pollen-food syndrome in children with seasonal allergic rhinoconjunctivitis: a molecular classification. *Allergy*. 2016;71:1181-1191.
- Ciprandi G, De Amici M, Di Martino ML, Barocci F, Comite P. The impact of age on Pru p 3 IgE production in Italy. *Asia Pac Allergy*. 2017;7:42-47.
- 107. Pastorello EA, Robino AM. Clinical role of lipid transfer proteins in food allergy. *Mol Nutr Food Res.* 2004;48:356-362.
- 108. Asero R, Pravettoni V. Anaphylaxis to plant-foods and pollen allergens in patients with lipid transfer protein syndrome. *Curr Opin Allergy Clin Immunol.* 2013;13:379-385.
- Cuesta-Herranz J, Barber D, Blanco C, et al. Differences among pollen-allergic patients with and without plant food allergy. *Int Arch Allergy Immunol.* 2010;153:182-192.
- 110. Scala E, Villalta D, Uasuf CG, et al. An atlas of IgE sensitization patterns in different Italian areas. A multicenter, cross-sectional study. *Eur Ann Allergy Clin Immunol.* 2018;50:217-225.
- 111. Rodrigues-Alves R, Lopez A, Pereira-Santos MC, et al. Clinical, anamnestic and serological features of peach allergy in Portugal. Int Arch Allergy Immunol. 2009;149:65-73.
- 112. Azofra J, Berroa F, Gastaminza G, et al. Lipid transfer protein syndrome in a non-Mediterranean area. *Int Arch Allergy Immunol.* 2016;169:181-188.
- 113. Fernández-Rivas M, Bolhaar S, González-Mancebo E, et al. Apple allergy across Europe: how allergen sensitization profiles determine the clinical expression of allergies to plant foods. J Allergy Clin Immunol. 2006;118:481-488.
- 114. Gaier S, Oberhuber C, Hemmer W, et al. Pru p 3 as a marker for symptom severity for patients with peach allergy in a birch pollen environment. J Allergy Clin Immunol. 2009;124:166-167.
- 115. Mothes-Luksch N, Raith M, Stingl G, et al. Pru p 3, a marker allergen for lipid transfer protein sensitization also in Central Europe. *Allergy*. 2017;72:1415-1418.

- 116. Ebo DG, Ahrazem O, Lopez-Torrejon G, Bridts CH, Salcedo G, Stevens WJ. Anaphylaxis from mandarin (Citrus reticulata): identification of potential responsible allergens. *Int Arch Allergy Immunol.* 2007;144:39-43.
- 117. Schad SG, Trcka J, Vieths S, et al. Wine anaphylaxis in a German patient: IgE-mediated allergy against a lipid transfer protein of grapes. Int Arch Allergy Immunol. 2005;136:159-164.
- 118. Gülsen A, Jappe U. Lipid transfer protein sensitization in an appleallergic patient: a case report from northern Europe. *Eur Ann Allergy Clin Immunol.* 2019;51:80-83.
- Ukleja-Sokołowska N, Gawrońska-Ukleja E, Żbikowska-Gotz M, Sokołowski Ł, Bartuzi Z. Recurrent anaphylaxis in patient allergic to eggplant - a Lipid transfer protein (LTP) syndrome. Asian Pac J Allergy Immunol. 2018;36:109-112.
- 120. Anantharachagan A, Sammour R, Vijayadurai P. Non-specific lipid transfer protein allergy in United Kingdom. *Ann Allergy Asthma Immunol.* 2019;123:618-620.
- Ma S, Nie L, Li H, Wang R, Yin J. Component-resolved diagnosis of peanut allergy and its possible origins of sensitization in China. Int Arch Allergy Immunol. 2016;169:241-248.
- 122. Murad A, Katelaris CH, Baumgart K. A case study of apple seed and grape allergy with sensitization to nonspecific lipid transfer protein. *Asia Pac Allergy*. 2016;6:129-132.
- 123. Obeso D, Mera-Berriatua L, Rodríguez-Coira J, et al. Multi-omics analysis points to altered platelet functions in severe food-associated respiratory allergy. *Allergy*. 2018;73:2137-2149.
- 124. Lund G, Brand S, Ramos T, et al. Strong and frequent T-cell responses to the minor allergen PhI p 12 in Spanish patients IgEsensitized to profilins. *Allergy*. 2018;73:1013-1021.

How to cite this article: Skypala IJ, Asero R, Barber D, et al. Non-specific lipid-transfer proteins: Allergen structure and function, cross-reactivity, sensitization, and epidemiology. *Clin Transl Allergy*. 2021;1–13. https://doi.org/10.1002/clt2.12010