

PHASE PORTRAITS OF SEPARABLE QUADRATIC SYSTEMS AND A BIBLIOGRAPHICAL SURVEY ON QUADRATIC SYSTEMS

TAO LI¹ AND JAUME LLIBRE²

ABSTRACT. Although planar quadratic differential systems and their applications have been studied in more than one thousand papers, we still have no complete understanding of these systems. In this paper we have two objectives.

First we provide a brief bibliographical survey on the main results about quadratic systems. Here we do not consider the applications of these systems to many areas as in Physics, Chemist, Economics, Biology, ...

Second we characterize the new class of planar separable quadratic polynomial differential systems. For such class of systems we provide the normal forms which contain one parameter, and using the Poincaré compactification and the blow up technique, we prove that there exist 10 non-equivalent topological phase portraits in the Poincaré disc for the separable quadratic polynomial differential systems.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

Let $P(x, y)$ and $Q(x, y)$ be two real polynomials of degree 2. Then the differential system

$$(1) \quad \dot{x} = P(x, y), \quad \dot{y} = Q(x, y),$$

is called a *planar quadratic polynomial differential system*, or in what follows simply a *quadratic system*. As usual the dot denotes derivative with respect to an independent variable t , called the time.

Quadratic systems began to be studied at the beginning of the twentieth century. According to Coppel [76] it seems that the first work on quadratic systems was written in 1904 by Büchel [41]. In 1966 Coppel [76] published a short survey on quadratic systems, another short survey on these systems appeared in 1982 by Chicone and Tian [63].

Quadratic systems have been intensively studied in the past several decades and a large number of valuable results were obtained, see the books [22, 189, 226] dedicated completely to quadratic systems and references therein, and

2010 *Mathematics Subject Classification.* Primary 34C05.

Key words and phrases. separable system, quadratic system, phase portrait, Poincaré compactification.

also the references in [188]. But the investigations on quadratic systems are still very far from being completed.

It is well known that one of the main subjects in the dynamics of planar polynomial differential systems is to characterize their global phase portraits in the Poincaré disc, see subsection 2.2 for the definition of the Poincaré compactification and the Poincaré disc. For the quadratic systems a complete characterization of the global phase portraits is an extremely difficult task due to the fact that these systems depend on 12 parameters, see for instance [22]. Instead many particular subfamilies of quadratic systems have been analyzed, for instance:

- structurally stable quadratic systems modulo limit cycles [5, 192],
- structurally unstable quadratic systems of codimension one [12],
- all the configurations of singularities of quadratic systems [13, 16, 17, 18, 19, 20, 21, 22, 25, 27, 32, 33, 36, 68],
- quadratic systems with a center [35, 83, 92, 117, 118, 119, 128, 141, 175, 176, 177, 178, 179, 191, 193, 208, 214, 234],
- quadratic systems with an isochronous center [56, 75, 164],
- Hopf bifurcation in quadratic systems [35],
- quadratic systems with a weak focus of third order [9, 145],
- quadratic systems with a weak focus of second order [14],
- quadratic systems with a weak focus and an invariant straight line [15],
- quadratic systems with weak singularities [215],
- Lotka–Volterra quadratic systems [47, 203, 204, 222, 50, 45],
- Bernoulli quadratic systems [142],
- quadratic systems without finite singularities, also called quadratic foliations or chordal systems [95, 96, 113, 212],
- every quadratic system has finitely many limit cycles [34],
- Abel quadratic systems [155],
- weak Hilbert’s 16th problem for quadratic systems: there are hundreds of papers see the references in the book of Christopher and Li [64] and [57, 58, 59, 74, 98, 99, 102, 103, 104, 105, 107, 108, 109, 121, 122, 123, 124, 125, 126, 129, 130, 206, 223, 225, 231],
- limit cycles in quadratic systems [11, 60, 61, 62, 69, 77, 85, 86, 87, 101, 106, 110, 116, 120, 131, 146, 163, 180, 181, 182, 183, 184, 190, 207, 209, 227, 228, 229, 233],
- quadratic-linear systems, i.e. one of the two equations of the system is defined by a polynomial of degree one [46, 162],
- integrability of quadratic–linear systems with [151, 152, 153],
- quadratic systems with a unique finite singularity [73],
- quadratic systems with a focus and one anti-saddle modulo limit cycles [8],
- quadratic systems with all points at infinity as singularities [3, 97, 174, 196, 198, 202],

- quadratic systems with a higher order singularity with two zero eigenvalues [112],
- quadratic systems with a finite and an infinite saddle-node [30, 31],
- quadratic systems with a semi-elemental triple node [29],
- singular points determine quadratic systems [24],
- restricted version of the Hilbert's 16th problem for quadratic [111],
- Darboux integrability for quadratic systems of [44, 47, 48, 55, 66, 78, 133, 140, 165, 171, 194, 195],
- Darboux invariants in quadratic systems [38, 138, 139],
- quadratic systems with a rational first integral [23, 49, 135, 136, 137],
- quadratic systems with a polynomial first integral [26, 51, 93],
- quadratic Hamiltonian systems [6, 7, 115, 175],
- quadratic systems of Darboux type [216],
- bounded quadratic systems [72, 80, 82, 127],
- phase portraits of quadratic systems in applications [186, 187],
- quadratic systems with a polynomial inverse integrating factor [70, 71],
- homogeneous quadratic systems [40, 79, 167, 169, 220, 221],
- quadratic systems with an integrable saddle [28, 39],
- quadratic systems with a weak saddle [43, 114],
- integrability of quadratic systems [159],
- quadratic systems with a symmetric center and simple infinite singular points [166, 217],
- singularly perturbed quadratic systems [147],
- algebraic limit cycles in quadratic systems [1, 52, 54, 67, 88, 89, 90, 91, 132, 143, 144, 148, 149, 154, 156, 157, 185, 224, 230],
- quadratic systems with a unique finite singular point of multiplicity two, possessing two zero eigenvalues [173],
- quadratic systems with a single finite singularity which in addition is simple [213],
- quadratic systems with a finite singular point of multiplicity four [211, 219],
- statistical measure of quadratic systems [10],
- quadratic systems with a singular point of multiplicity three [218],
- quadratic systems with invariant algebraic curves [94, 134, 150, 158, 160, 161],
- quadratic systems with invariant straight lines of total multiplicity greater than or equal to four [197, 199, 200, 201],
- quadratic systems with a semi-elemental triple node [4],
- quadratic systems with two parallel invariant straight lines [42],
- quadratic systems with complex conjugate invariant lines meeting at a finite point [205, 210],
- quadratic systems with invariant algebraic curves of arbitrarily high degree without rational first integrals [53, 65, 170].

A differential equation

$$(2) \quad \frac{dy}{dx} = F(x, y)$$

is said to be *separable* if $F(x, y)$ can be written as the form $F_1(x)F_2(y)$. In parallel we say that the planar differential system (1) is separable if written into the form

$$\frac{dy}{dx} = \frac{Q(x, y)}{P(x, y)} = F(x, y),$$

this differential equation is separable. The nonlinear separable differential systems are special because they can be solved by the separation method of variables. Although the solutions of a large number of separable differential systems may be complex and even can provide integrals which cannot be computed. Thus it is also necessary to study the separable differential systems using the qualitative theory of the differential systems.

In this paper we shall study the phase portraits in the Poincaré disc of the following family of separable quadratic polynomial differential systems

$$(3) \quad \dot{x} = a_1x^2 + a_2x + a_3, \quad \dot{y} = b_1y^2 + b_2y + b_3,$$

where

$$(4) \quad a_1^2 + b_1^2 \neq 0, \quad a_1^2 + a_2^2 + a_3^2 \neq 0, \quad b_1^2 + b_2^2 + b_3^2 \neq 0.$$

The next theorem is our main result.

Theorem 1. *The global phase portraits of the separable quadratic polynomial differential systems (3) satisfying (4) are topologically equivalent to one of the 10 phase portraits in the Poincaré disc described in Figure 1.*

The paper is organized as follows. In section 2 we recall some basic notions and tools for studying the phase portraits of polynomial differential systems in the Poincaré disc. In particular, subsection 2.1 contains some definitions and results about finite and infinite singular points, and subsection 2.2 reviews the Poincaré compactification. In section 3 we provide the normal forms of systems (3) satisfying (4). In sections 4 and 5 the infinite and finite singular points are studied respectively. Finally, Theorem 1 is proved in section 6.

2. PRELIMINARIES

In this section we shortly review some basic notions, results and tools which are involved in the investigation of the phase portraits of planar polynomial differential systems in the Poincaré disc, see [84, 232] for more details.




Figure 1. Phase portraits in the Poincaré disc of systems (3) satisfying (4). Here the pair (S,R) which appears in each phase portrait in the Poincaré disc denotes the number of separatrices S and the number of canonical regions R of the corresponding phase portrait.

2.1. Singular points. Let $Z(x, y) = (X(x, y), Y(x, y))$ be a planar smooth vector field. A point $p \in \mathbb{R}^2$ is said to be a *singular point* of Z if $X(p) = Y(p) = 0$. Let J be the Jacobian matrix of Z at the singular point p , i.e.,

$$J = \begin{pmatrix} X_x(p) & X_y(p) \\ Y_x(p) & Y_y(p) \end{pmatrix},$$

where the subscripts x and y denote the partial derivative with respect to x and y , respectively. Depending on the character of the matrix J , an isolated singular point p is called

- (i) a *hyperbolic* singular point if J has two eigenvalues with nonzero real part,
- (ii) a *semi-hyperbolic* singular point if J has a unique zero eigenvalue,
- (iii) a *nilpotent* singular point if J has two zero eigenvalues and the matrix J is not identically zero,
- (iv) a *linearly zero* singular point if the matrix J is identically zero.




Figure 2. Phase portraits of saddle-nodes.

For a hyperbolic, semi-hyperbolic or nilpotent singular point p , the local phase portrait of Z at p has been thoroughly studied, see for instance Theorems 2.15, 2.19 and 3.5 of [84], respectively. In particular, a hyperbolic singular point p is a *saddle* (resp. *node*) if J at p has two real eigenvalues with opposite (resp. same) sign, and a *strong focus* if the eigenvalues are complex with nonzero real part. If J at p has a pair purely imaginary eigenvalues, then it may be a *weak focus* or a *center*, see [84, 232]. Since Theorems 2.19 of [84] will be used repeatedly later on, we summarize it in what follows.

Theorem 2. *Let the origin O be an isolated singular point of the vector field $Z = (X(x, y), Y(x, y)) = (\tilde{X}(x, y), \lambda y + \tilde{Y}(x, y))$, where \tilde{X} and \tilde{Y} are analytic in a neighborhood of the origin with $\tilde{X}(0, 0) = \tilde{Y}(0, 0) = D\tilde{X}(0, 0) = D\tilde{Y}(0, 0) = 0$ and $\lambda > 0$. Let $y = f(x)$ be the solution of the equation $\lambda y + \tilde{Y}(x, y) = 0$ in a neighborhood of O , and suppose that the function $g(x) = \tilde{X}(x, f(x))$ has the expression $g(x) = a_m x^m + o(x^m)$, where $m \geq 2$ and $a_m \neq 0$.*

- (a) *If m is odd and $a_m < 0$, then O is a topological saddle. Moreover, the two stable separatrices are tangent to the x -axis at O .*
- (b) *If m is odd and $a_m > 0$, then O is an unstable topological node.*
- (c) *If m is even, then O is a saddle-node, i.e., a point whose neighborhood is separated into two parts by one unstable separatrix that is tangent to the positive y -axis at O and one unstable separatrix that is tangent to the negative y -axis at O . When $a_m < 0$ (resp. $a_m > 0$), the left (resp. right) part is a parabolic sector with unstable invariant manifolds of O and the right (resp. left) part is two hyperbolic sectors separated by one stable separatrix that is tangent to the positive (resp. negative) x -axis at O , see Figure 2(1) (resp. (2)).*

Remark 3. The case $\lambda < 0$ can be reduced to $\lambda > 0$ reversing the time. Moreover the case $Z = (\lambda x + \tilde{X}, \tilde{Y})$ can be reduced to $Z = (\tilde{X}, \lambda y + \tilde{Y})$ by the change of variables $(x, y) \rightarrow (y, x)$. As we will see the two changes will

be repeatedly used to study the local phase portrait at a semi-hyperbolic singular point.

The local phase portrait of a linearly zero singular point p can be determined by using the *blow up technique*, see [2, 81, 84] for more details.

2.2. Poincaré compactification. It is well known that an important tool for studying the phase portraits of a quadratic polynomial vector field $Z = (X, Y)$ is the *Poincaré compactification*. For completeness we shortly review it in this subsection, for more details see Chapter 5 of [84], or [232].

Consider the 2-dimensional sphere $\mathbb{S}^2 := \{(s_1, s_2, s_3) \in \mathbb{R}^3 : s_1^2 + s_2^2 + s_3^2 = 1\}$, called the *Poincaré sphere*, and its tangent plane at the point $(0, 0, 1)$, this plane is identified with the plane \mathbb{R}^2 where the vector field Z is defined. According with [84] the vector field Z in \mathbb{R}^2 can be extended analytically to a vector field $p(Z)$ on \mathbb{S}^2 by using the central projection f which maps each point Q in \mathbb{R}^2 onto two points on \mathbb{S}^2 using the straight line through Q and the origin $(0, 0, 0)$. Notice that we obtain two copies of Z on \mathbb{S}^2 by the central projection f , one in the northern hemisphere and the other in the southern hemisphere. So we have a vector field Z^* defined on $\mathbb{S}^2 \setminus \mathbb{S}^1$, where $\mathbb{S}^1 := \{(s_1, s_2, s_3) \in \mathbb{S}^2 : s_3 = 0\}$ is the equator of \mathbb{S}^2 . Doing the rescaling $s_3^2 Z^*$ we extend the quadratic polynomial vector field Z^* to the whole sphere, and as usually we denote the vector field $y_3^3 Z^*$ by $p(Z)$, called the *Poincaré compactification* of the vector field Z . The dynamics of $p(Z)$ near the equator \mathbb{S}^1 corresponds to the dynamics of Z in a neighborhood of the infinity of \mathbb{R}^2 .

It is sufficient to consider the Poincaré compactification restricted to the northern hemisphere $\{(s_1, s_2, s_3) \in \mathbb{S}^2 : s_3 > 0\}$ union the equator \mathbb{S}^1 in order to study the phase portrait of the vector field Z . Moreover the phase portrait is drawn on the so-called *Poincaré disc* $\mathbb{D}^2 := \{(s_1, s_2) \in \mathbb{R}^2 : s_1^2 + s_2^2 \leq 1\}$, obtained projecting the northern hemisphere union the equator onto \mathbb{D}^2 using the projection $\pi(s_1, s_2, s_3) = (s_1, s_2)$.

For working with $p(Z)$ on \mathbb{S}^2 we need the local charts (U_i, ϕ_i) and (V_i, ψ_i) for $i = 1, 2, 3$, where

$$U_i := \{(s_1, s_2, s_3) \in \mathbb{S}^2 : s_i > 0\}, \quad V_i := \{(s_1, s_2, s_3) \in \mathbb{S}^2 : s_i < 0\},$$

$\phi_i : U_i \rightarrow \mathbb{R}^2$ and $\psi_i : V_i \rightarrow \mathbb{R}^2$ are defined by

$$\phi_i(s_1, s_2, s_3) = -\psi_i(s_1, s_2, s_3) = (s_m/s_i, s_n/s_i) = (u, v),$$

where $m < n$ and $m, n \neq i$. Then the expression of $p(Z)$ in the local chart U_1 is given by

$$(5) \quad \dot{u} = z^2 \left(-uX \left(\frac{1}{z}, \frac{u}{z} \right) + Y \left(\frac{1}{z}, \frac{u}{z} \right) \right), \quad \dot{z} = -z^3 X \left(\frac{1}{z}, \frac{u}{z} \right).$$

In the local chart U_2 the vector field $p(Z)$ is given by

$$(6) \quad \dot{v} = z^2 \left(X \left(\frac{v}{z}, \frac{1}{z} \right) - vY \left(\frac{v}{z}, \frac{1}{z} \right) \right), \quad \dot{z} = -z^3 Y \left(\frac{v}{z}, \frac{1}{z} \right).$$

The compactified vector field $p(Z)$ in the local chart U_3 is just Z . The expression of $p(Z)$ in V_i is the expression in U_i multiplied by -1 for $i = 1, 2, 3$.

The singular points of $p(Z)$ in $\mathbb{S}^2 \setminus \mathbb{S}^1$ correspond to the finite singular points of Z , and they can be obtained by using the local chart U_3 . The infinite singular points of Z are the singular points of systems (5) and (6) lying in \mathbb{S}^1 , we remark that their coordinate $z = 0$. It is worth mentioning that we only need to look at $U_1|_{z=0}$ and $U_2|_{(0,0)}$ to study the infinite singular points of Z .

Two Poincaré compactified vector fields on \mathbb{D}^2 are *topologically equivalent* if there is a homeomorphism from one vector field to the other sending orbits to orbits, and preserving or reversing the direction of all the orbits.

According to [168, 172] a *separatrix* of $\pi(p(Z))$ on \mathbb{D}^2 is one of following orbits: finite singular points, limit cycles, all orbits at the infinity, and the two boundaries of a hyperbolic sector at a finite or infinite singular point. We denote the set formed by all separatrices by Σ . Then Σ is a closed set as it is proved in [172]. An open connected component of $\mathbb{D}^2 \setminus \Sigma$ is called a *canonical region* of $\pi(p(Z))$. Then the *separatrix configuration* Σ^* of $\pi(p(Z))$ is the union of the set Σ with an orbit for each canonical region. Two separatrix configurations Σ_1^* and Σ_2^* are *topologically equivalent* if there exists a homeomorphism from Σ_1^* to Σ_2^* sending the orbits of one separatrix configuration to the orbits of the other, and preserving or reversing the direction of all the orbits.

Finally we recall the Neumann's Theorem [172], which characterizes the topological equivalence of two global phase portraits in the Poincaré disc \mathbb{D}^2 .

Theorem 4. *Two Poincaré compactified polynomial vector fields $\pi(p(Z_1))$ and $\pi(p(Z_2))$ with finitely many separatrices are topologically equivalent if and only if their separatrix configurations are topological equivalent.*

We shall use this theorem for proving Theorem 1.

3. NORMAL FORMS

In this section we study the normal forms of system (3) satisfying (4). The next lemma is the main result of this section.

Lemma 5. *System (3) satisfying (4), after a linear change of variables and a scaling of its independent variable t , can be written in one of the following normal forms:*

- (I) $\dot{x} = 1, \quad \dot{y} = y^2 + \xi;$
- (II) $\dot{x} = x, \quad \dot{y} = y^2 + \xi;$
- (III) $\dot{x} = x^2, \quad \dot{y} = y^2 + \xi;$
- (IV) $\dot{x} = x^2 + 1, \quad \dot{y} = y^2 + \xi;$
- (V) $\dot{x} = x^2 - 1, \quad \dot{y} = y^2 + \xi.$

Proof. Since $a_1^2 + b_1^2 \neq 0$ by condition (4), we only need to consider the case of $b_1 \neq 0$ by the change of variables $(x, y) \rightarrow (y, x)$.

If $a_1 = a_2 = 0$, we have $a_3 \neq 0$ from (4), and then the change of variables

$$(x, y) \rightarrow (a_3x, y/b_1 - b_2/(2b_1))$$

transforms system (3) into system (I) with $\xi = b_1b_3 - b_2^2/4$.

If $a_1 = 0$ and $a_2 \neq 0$, then the change

$$(x, y, t) \rightarrow (x - a_3/a_2, a_2y/b_1 - b_2/(2b_1), t/a_2)$$

transforms system (3) into system (II) with $\xi = b_1b_3/a_2^2 - b_2^2/(4a_2^2)$.

If $a_1 \neq 0$ doing the change

$$(x, y, t) \rightarrow ((\Delta x - a_2)/(2a_1), \Delta y/(2b_1) - b_2/(2b_1), 2t/\Delta),$$

where

$$\Delta := \begin{cases} 1 & \text{if } a_2^2 - 4a_1a_3 = 0, \\ \sqrt{|a_2^2 - 4a_1a_3|} & \text{if } a_2^2 - 4a_1a_3 \neq 0, \end{cases}$$

system (3) becomes system (III) (resp. (IV), (V)) when $a_2^2 - 4a_1a_3 = 0$ (resp. $< 0, > 0$). Here $\xi = 4b_1b_3/\Delta^2 - b_2^2/\Delta^2$. \square

From Lemma 5 we only need to study the phase portraits of systems (I)–(V) in order to obtain all phase portraits of system (3) satisfying (4) in the Poincaré disc.

4. INFINITE SINGULAR POINTS

In this section we study the infinite singular points of systems (I)–(V).

Lemma 6. *In the local chart U_1 system (I) has a unique infinite singular point, namely the origin, which is linearly zero having the local phase portrait presented in*

- (a) Figure 3(2) if $\xi > 0$,
- (b) Figure 3(3) if $\xi = 0$,
- (c) Figure 4(2) if $\xi < 0$.




Figure 3. Blow up of system (7) with $\xi \geq 0$ at O .

Proof. In the local chart U_1 system (I) writes

$$(7) \quad \dot{u} = u^2 + \xi z^2 - u z^2, \quad \dot{z} = -z^3.$$

When $z = 0$ the origin O is the unique singular point of system (7) and it is linearly zero.

To determine the local phase portrait of system (7) at O we use the blow up technique as follows. Since the characteristic polynomial of the linear part of system (7) at O is $\mathcal{F}(u, z) = z(u^2 + \xi z^2)$, $z = 0$ is a characteristic direction. Thus, using the u -directional blow up $(u, z) = (\bar{u}, \bar{u}\bar{z})$ we change system (7) into

$$(8) \quad \dot{\bar{u}} = \bar{u}(1 + \xi \bar{z}^2 - \bar{u} \bar{z}^2), \quad \dot{\bar{z}} = -\bar{z}(1 + \xi \bar{z}^2),$$

after cancelling the common factor \bar{u} doing the change of time $t \rightarrow t/\bar{u}$. For $\bar{u} = 0$ the origin O is always a singular point of system (8) and it is a hyperbolic saddle. Since $\bar{u} = 0$ and $\bar{z} = 0$ are two invariant straight lines of (8), the two stable (resp. unstable) invariant manifolds of the saddle are contained in the \bar{z} -axis (resp. \bar{u} -axis), see Figure 3(1).

If $\xi \geq 0$, then O is the unique singular point of system (8) when $\bar{u} = 0$. Moreover $z = 0$ is the unique characteristic direction of system (7) at O if $\xi > 0$, while if $\xi = 0$, system (7) has exactly two characteristic directions at O , $z = 0$ and $u = 0$. Note that these characteristic directions are invariant straight lines of system (7). Hence going back through the used blow up in the last paragraph and using the direction of vector field on these invariant straight lines, we get the local phase portrait of system (7) at O as shown in Figure 3(2) if $\xi > 0$, and in Figure 3(3) if $\xi = 0$, i.e. statements (a) and (b) hold.

If $\xi < 0$ then system (8) for $\bar{z} = 0$ has two additional singular points $(0, p^\pm)$ with $p^\pm = \pm 1/\sqrt{-\xi}$. They are semi-hyperbolic because the two eigenvalues at $(0, p^\pm)$ are 0 and 2. We next apply Theorem 2 to determine the local phase portraits at these two points. Translating $(0, p^+)$ to the




Figure 4. Blow up of system (7) with $\xi < 0$ at O .

origin we see that the quantities λ, m, a_m in Theorem 2 associated to system (8) are $\lambda = 2, m = 2, a_m = 1/\xi < 0$, respectively. Hence a straightway application of Theorem 2(c) yields that $(0, p^+)$ is a saddle-node with two unstable separatrices in the \bar{z} -axis and one stable separatrix in $\{(\bar{u}, \bar{z}) : \bar{u} > 0, \bar{z} = p^+\}$, combining that $\bar{u} = 0$ and $\bar{z} = p^+$ are two invariant straight lines of system (8). Similarly, translating $(0, p^-)$ to the origin we get $\lambda = 2, m = 2, a_m = 1/\xi < 0$, and thus $(0, p^-)$ is a saddle-node with two unstable separatrices in the \bar{z} -axis and one stable separatrix in $\{(\bar{u}, \bar{z}) : \bar{u} > 0, \bar{z} = p^-\}$ by Theorem 2(c) again. This concludes the local phase portraits of system (8) at $(0, p^\pm)$, see Figure 4(1) where the local phase portrait at O is also drawn. Consequently, the local phase portrait of system (7) at O is as shown in Figure 4(2), going back to system (7) through the blow up $(u, z) = (\bar{u}, \bar{u}\bar{z})$. This completes the proof of statement (c). \square

Lemma 7. *In the local chart U_1 system (II) has a unique infinite singular point, namely the origin, which is linearly zero having the local phase portrait presented in*

- (a) *Figure 5(2) when $\xi > 0$,*
- (b) *Figure 5(3) when $\xi = 0$,*
- (c) *Figure 6(2) when $\xi < 0$.*

Proof. In the local chart U_1 system (II) becomes

$$(9) \quad \dot{u} = u^2 - uz + \xi z^2, \quad \dot{z} = -z^2.$$

For $z = 0$ system (9) has a unique singular point, namely the origin O . Clearly, it is linearly zero.

Next we use the blow up technique to study the local phase portrait of system (9) at O . Since the characteristic polynomial of the linear part of




Figure 5. Blow up of system (9) with $\xi \geq 0$ at O .

system (9) at the origin O is $\mathcal{F}(u, z) = z(u^2 + \xi z^2)$, $z = 0$ is a characteristic direction. Thus using the blow up technique with the change $(u, z) = (\bar{u}, \bar{u}\bar{z})$ we obtain

$$(10) \quad \dot{\bar{u}} = \bar{u}(1 - \bar{z} + \xi \bar{z}^2), \quad \dot{\bar{z}} = -\bar{z}(1 + \xi \bar{z}^2),$$

after cancelling the common factor \bar{u} doing the change $t \rightarrow t/\bar{u}$. For $\bar{u} = 0$ the origin O is always a singular point of system (10), particularly it is a hyperbolic saddle with two unstable (resp. stable) manifolds contained in the \bar{u} -axis (resp. \bar{z} -axis), see Figure 5(1).

If $\xi \geq 0$, then O is the unique singular point of system (10) for $\bar{u} = 0$. Moreover $z = 0$ is the unique characteristic direction of system (9) at O if $\xi > 0$, while if $\xi = 0$, system (9) has exactly two characteristic directions at O , $z = 0$ and $u = 0$. Since these characteristic directions are invariant straight lines of system (9), going back to system (9) through the last blow up and using the direction of vector fields on these lines, we get the local phase portrait of system (9) at O as it is shown in Figure 5(2) if $\xi > 0$, and in Figure 5(3) if $\xi = 0$, i.e. statements (a) and (b) hold.

If $\xi < 0$, system (10) for $\bar{u} = 0$ has two additional singular points, $(0, p^\pm)$ with $p^\pm = \pm 1/\sqrt{-\xi}$ where the eigenvalues are $-p^\pm$ and 2. Thus $(0, p^-)$ is an unstable hyperbolic node if $p^- < 0$, and $(0, p^+)$ is a hyperbolic saddle if $p^+ > 0$. This concludes the local phase portraits of system (10) at O and $(0, p^\pm)$ as shown in Figure 6(1). Going back to system (9) we get the local phase portrait of system (9) at O as it is shown in Figure 6(2), consequently statement (c) holds. \square

Lemma 8. *In the local chart U_1 each one of the systems (III), (IV) and (V) has two infinite singular points: the origin and $(1, 0)$. Moreover the origin is a stable hyperbolic node, and $(1, 0)$ is a hyperbolic saddle with two unstable separatrices contained in the u -axis.*




Figure 6. Blow up of system (9) with $\xi < 0$ at the origin.

Proof. In the local chart U_1 systems (III), (IV) and (V) are

$$(11) \quad \begin{aligned} \dot{u} &= -u + u^2 + \xi z^2, & \dot{z} &= -z, \\ \dot{u} &= -u + u^2 + \xi z^2 - uz^2, & \dot{z} &= -z - z^3, \\ \dot{u} &= -u + u^2 + \xi z^2 + uz^2, & \dot{z} &= -z + z^3, \end{aligned}$$

respectively. Thus each system in (11) has two singular points O and $(1, 0)$ for $z = 0$. For each system the eigenvalues at O (resp. $(1, 0)$) are always -1 (resp. 1) and -1 . Thus both O and $(1, 0)$ are hyperbolic, particularly O is a stable node and $(1, 0)$ is a saddle. Since $z = 0$ is an invariant straight line of each system in (11), the two unstable manifolds of $(1, 0)$ are contained in the u -axis. This ends the proof of Lemma 8. \square

The last four lemmas studied the infinite singular points of systems (I)-(V) in the local chart U_1 . The next lemma verifies whether the origin is an infinite singular point of these systems in the local chart U_2 .

Lemma 9. *In the local chart U_2 the origin is an infinite singular point for each one of systems (I)-(V). Moreover it is a stable hyperbolic node.*

Proof. We observe that systems (I)-(V) can be written in a unified form as

$$(12) \quad \dot{x} = \kappa_1 x^2 + \kappa_2 x + \kappa_3, \quad \dot{y} = y^2 + \xi,$$

where $\kappa_i \in \{-1, 0, 1\}$, $i = 1, 2, 3$. In the local chart U_2 system (12) writes

$$\begin{aligned} \dot{v} &= -v + \kappa_1 v^2 + \kappa_2 vz + \kappa_3 z^2 - \xi vz^2, \\ \dot{z} &= -z - \xi z^3, \end{aligned}$$

which obviously has the origin as a singular point. Since all the two eigenvalues at the origin are -1 , it is a stable hyperbolic node. \square

5. FINITE SINGULAR POINTS

This section is devoted to study the finite singular points of systems (I)-(V).

Lemma 10. *Systems (I) and (IV) have no finite singular points.*

Proof. Lemma 10 follows from $\dot{x} > 0$ systems (I) and (IV). \square

Lemma 11. *On the finite singular points of system (II) we have the following statements.*

- (a) *If $\xi > 0$, there is no finite singular points.*
- (b) *If $\xi = 0$, there is a unique finite singular point, the origin. Moreover it is a semi-hyperbolic saddle-node with two unstable separatrices contained in the x -axis and one stable separatrix contained in the negative y -axis.*
- (c) *If $\xi < 0$, there are two finite singular points, $(0, \pm\sqrt{-\xi})$. Moreover $(0, \sqrt{-\xi})$ is an unstable hyperbolic node and $(0, -\sqrt{-\xi})$ is a hyperbolic saddle with two stable separatrices contained in the y -axis.*

Proof. The number of finite singular points of system (II) equals the number of roots of the equation $y^2 + \xi = 0$. Thus statement (a) holds directly if $\xi > 0$.

If $\xi = 0$, then O is the unique finite singular point, where the eigenvalues are 1 and 0, so that it is semi-hyperbolic. To determine the local phase portrait we use the change $(x, y) \rightarrow (y, x)$ to transform system (II) into the system of Theorem 2, i.e. the system

$$(13) \quad \dot{x} = x^2, \quad \dot{y} = y.$$

A direct application of Theorem 2(c) concludes that O is a saddle-node of system (13) with two unstable separatrices contained in the y -axis and one stable separatrix contained in the negative x -axis. Here the x -axis and y -axis are invariant straight lines of (13). Therefore statement (b) holds going back to system (II).

If $\xi < 0$, system (II) has two finite singular points $(0, \pm\sqrt{-\xi})$, where the eigenvalues are 1 and $\pm 2\sqrt{-\xi}$. This yields statement (c) joining the fact that $x = 0$ is an invariable straight line. \square

Lemma 12. *On the finite singular points of system (III) we have the following statements.*

- (a) *If $\xi > 0$, there is no finite singular points.*
- (b) *If $\xi = 0$, there is a unique finite singular point, the origin. Moreover, it is linearly zero with the local phase portrait presented in Figure 7(2).*

- (c) If $\xi < 0$, there are two finite singular points, $(0, \pm\sqrt{-\xi})$. Moreover $(0, \sqrt{-\xi})$ is a semi-hyperbolic saddle-node with two unstable separatrices contained in the y -axis and one stable separatrix contained in $\{(x, y) : x < 0, y = \sqrt{-\xi}\}$, $(0, -\sqrt{-\xi})$ is a semi-hyperbolic saddle-node with two stable separatrices contained in the y -axis and one unstable separatrix contained in $\{(x, y) : x > 0, y = -\sqrt{-\xi}\}$.

Proof. As in system (II) the number of finite singular points of system (III) is equal to the number of roots of the equation $y^2 + \xi = 0$. Thus statement (a) holds directly if $\xi > 0$.

If $\xi = 0$, then O is the unique finite singular point of system (III) and it is linearly zero. Next we use the blow up technique to study the local phase portrait at O . Since the characteristic polynomial of system (III) is $\mathcal{F}(x, y) = xy(x - y)$, $y = 0$ is a characteristic direction. Thus doing the x -directional blow up $(x, y) = (\bar{x}, \bar{x}\bar{y})$ we get the system

$$(14) \quad \dot{\bar{x}} = \bar{x}, \quad \dot{\bar{y}} = -\bar{y} + \bar{y}^2,$$

after cancelling the common factor \bar{x} using the change $t \rightarrow t/\bar{x}$. For $\bar{x} = 0$ system (14) has two singular points, $(0, 1)$ and O . Since the eigenvalues at $(0, 1)$ (resp. O) are 1 and 1 (resp. -1), $(0, 1)$ is an unstable hyperbolic node and O is a hyperbolic saddle with two stable (resp. unstable) separatrices contained in the \bar{y} -axis (resp. \bar{x} -axis), see Figure 7(1). Finally we get the local phase portrait of system (III) at O as shown in Figure 7(2), going back to system (III) through the used blow up. This yields statement (b).




Figure 7. Blow up of system (III) with $\xi = 0$ at O .

If $\xi < 0$, system (III) has two exactly singular points, $(0, \pm\sqrt{-\xi})$, where the eigenvalues are 0 and $\pm 2\sqrt{-\xi}$. So they are semi-hyperbolic. To determine the local phase portrait at $(0, \sqrt{-\xi})$ we use the change $(x, y) \rightarrow (x, y + \sqrt{-\xi})$ to transform system (III) into the normal form of system of Theorem 2, i.e.

$$(15) \quad \dot{x} = x^2, \quad \dot{y} = 2\sqrt{-\xi}y + y^2.$$

Then the quantities λ, m, a_m in Theorem 2 associated to system (15) are $\lambda = 2\sqrt{-\xi} > 0, m = 2, a_m = 1$. Thus together with the invariant lines $x = 0$ and $y = 0$, O is a saddle-node of system (15) with two unstable separatrices contained in the y -axis and one stable separatrix contained in the negative x -axis. Coming back to system (III) we finally get that $(0, \sqrt{-\xi})$ is a saddle-node with two unstable separatrices contained in the y -axis and one stable separatrix contained in $\{(x, y) : x < 0, y = \sqrt{-\xi}\}$. Regarding the local phase portrait at $(0, -\sqrt{-\xi})$, using the change $(x, y, t) \rightarrow (x, y - \sqrt{-\xi}, -t)$ we can carry out a similar analysis, and thus $(0, -\sqrt{-\xi})$ is a saddle-node with two stable separatrices contained in the y -axis and one unstable separatrix contained in $\{(x, y) : x > 0, y = -\sqrt{-\xi}\}$. That is, statement (c) holds. \square

Lemma 13. *For system (V) all finite singular points must lie in the invariant straight lines $x = 1$ and $x = -1$.*

- (a) *In the line $x = 1$ system (V) has*
 - (a1) *no singular points if $\xi > 0$;*
 - (a2) *a unique singular point $(1, 0)$ if $\xi = 0$, and it is a semi-hyperbolic saddle-node with two unstable separatrices contained in the x -axis and one stable separatrix contained in $\{(x, y) : x = 1, y < 0\}$;*
 - (a3) *two singular points $(1, \pm\sqrt{-\xi})$ if $\xi < 0$, and $(1, \sqrt{-\xi})$ is an unstable hyperbolic node and $(1, -\sqrt{-\xi})$ is a hyperbolic saddle with two stable separatrices contained in the line $x = 1$.*
- (b) *In the line $x = -1$ system (V) has*
 - (b1) *no singular points if $\xi > 0$;*
 - (b2) *a unique singular point $(-1, 0)$ if $\xi = 0$, and it is a semi-hyperbolic saddle-node with two stable separatrices contained in the x -axis and one unstable separatrix contained in $\{(x, y) : x = -1, y > 0\}$;*
 - (b3) *two singular points $(-1, \pm\sqrt{-\xi})$ if $\xi < 0$, and $(-1, -\sqrt{-\xi})$ is a stable hyperbolic node and $(-1, \sqrt{-\xi})$ is a hyperbolic saddle with two unstable separatrices contained in the line $x = -1$.*

Proof. We neglect the proof of this lemma because it is completely similar to the proof of Lemma 11 by using Theorem 2 and the fact that $x = \pm 1$ are invariant straight lines. \square

6. PROOF OF THEOREM 1

To prove Theorem 1 we can equivalently study all global phase portraits for each one of the systems (I)-(V) by Lemma 5. We see that all finite singular points of each one of systems (I)-(VI) must lie on invariant straight lines. This means that there are no periodic orbits, and consequently no limit cycles. Combining the investigation of infinite singular points in section 4

with the one of finite singular points in section 5, we conclude that all separatrices can be determined in a natural and unique way except for the ones in system (IV) with $\xi > 0$, and particularly the following statements hold.

- (i) The global phase portrait of system **(I)** is topologically equivalent to the one of
 - (i.a) Figure 1(I.1) if $\xi > 0$ from Lemmas 6(a), 9 and 10;
 - (i.b) Figure 1(I.2) if $\xi = 0$ from Lemmas 6(b), 9 and 10;
 - (i.c) Figure 1(I.3) if $\xi < 0$ from Lemmas 6(c), 9 and 10.
- (ii) The global phase portrait of system **(II)** is topologically equivalent to the one of
 - (ii.a) Figure 1(II.1) if $\xi > 0$ from Lemmas 7(a), 9 and 11(a);
 - (ii.b) Figure 1(II.1) if $\xi = 0$ from Lemmas 7(b), 9 and 11(b);
 - (ii.c) Figure 1(II.2) if $\xi < 0$ from Lemmas 7(c), 9 and 11(c).
- (iii) The global phase portrait of system **(III)** is topologically equivalent to the one of
 - (iii.a) Figure 1(III.1) if $\xi > 0$ from Lemmas 8, 9 and 12(a);
 - (iii.b) Figure 1(III.2) if $\xi = 0$ from Lemmas 8, 9 and 12(b);
 - (iii.c) Figure 1(III.3) if $\xi < 0$ from Lemmas 8, 9 and 12(c).
- (iv) Since $y = x$ is a solution of system (IV) with $\xi = 1$, it follows from Lemmas 8, 9 and 10 that the global phase portrait of system **(IV)** is topologically equivalent to the one of
 - (iv.a) Figure 1(III.1) if $\xi > 1$;
 - (iv.b) Figure 1(IV) if $\xi = 1$;
 - (iv.c) Figure 8(a) if $\xi < 1$.
- (v) The global phase portrait of system **(V)** is topologically equivalent to the one of
 - (v.a) Figure 1(III.1) if $\xi > 0$ from Lemmas 8, 9 and 13(a1)(b1);
 - (v.b) Figure 8(b) if $\xi = 0$ from Lemmas 8, 9 and 13(a2)(b2);
 - (v.c) Figure 1(V) if $\xi < 0$ from Lemmas 8, 9 and 13(a3)(b3).

From Theorem 4 we easily observe that the global phase portrait of Figure 8(a) is topologically equivalent to the one of Figure 1(III.1), and the global phase portraits of Figure 8(b) is topologically equivalent to the one of Figure 1(III.3). Consequently, according to the items (i), \dots , (v), and Theorem 4, we finally obtain the 10 non-topologically equivalent global phase portraits in Figure 1. This ends the proof of Theorem 1.




Figure 8. Two global phase portraits.

ACKNOWLEDGEMENTS

The first author is partially supported by the grant CSC #201906240094 from the P.R. China. The second author is partially supported by the Ministerio de Ciencia, Innovación y Universidades, Agencia Estatal de Investigación grants MTM2016-77278-P (FEDER), the Agència de Gestió d'Ajuts Universitaris i de Recerca grant 2017SGR1617, and the H2020 European Research Council grant MSCA-RISE-2017-777911.

REFERENCES

- [1] M. ALBERICH-CARRAMIÑANA, A. FERRAGUT AND J. LLIBRE, *Quadratic planar differential systems having algebraic limit cycles*, preprint 2019.
- [2] M. J. ÁLVAREZ, A. FERRAGUT AND X. JARQUE, *A survey on the blow up technique*, Internat. J. Bifur. Chaos **21** (2011), 3103–3118.
- [3] A.A. ANDRONOV, E. A. LEONTOVICH, I. I. GORDON AND A. G. MAIER, *Theory of bifurcations of dynamic systems on a plane*, Translated from the Russian. Halsted Press [A division of John Wiley and Sons], New York-Toronto, Ont.; Israel Program for Scientific Translations, Jerusalem-London, 1973.
- [4] J.C. ARTÉS, *Global phase portraits of quadratic polynomial differential systems with a semi-elemental triple node*, Int. J. Bifur. Chaos **23** (2013), 1350140.
- [5] J.C. ARTÉS, R. KOIJ AND J. LLIBRE, *Structurally stable quadratic vector fields*, Memoires Amer. Math. Soc. **134** (639), 1998.
- [6] J.C. ARTÉS AND J. LLIBRE, *Quadratic Hamiltonian vector fields*, Differential Equations **107** (1994), 80–95.
- [7] J.C. ARTÉS AND J. LLIBRE, *Corrigendum: Quadratic Hamiltonian vector fields*, J. Differential Equations **129** (1996), 559–560.
- [8] J.C. ARTÉS AND J. LLIBRE, *Phase portraits for quadratic systems having a focus and one anti-saddle*, Rocky Mountain J. Math. **24** (1994), 875–889.
- [9] J.C. ARTÉS AND J. LLIBRE, *Quadratic vector fields with a weak focus of third order*, Publ. Mat. **41** (1997), 7–39.
- [10] J.C. ARTÉS AND J. LLIBRE, *Statistical measure of quadratic vector fields*, Resenhas do Instituto de Matematica e Estatistica da Universidade de Sao Paulo **6** (2003), 85–97.
- [11] J.C. ARTÉS AND J. LLIBRE, *Nonexistence of limit cycles for a class of structurally stable quadratic vector fields*, Disc. Cont. Dyn. Syst. **17** (2007), 259–270.
- [12] J.C. ARTÉS, J. LLIBRE AND A. C. REZENDE, *Structurally unstable quadratic vector fields of codimension one*, Birkhäuser/Springer, 2018.

- [13] J.C. ARTÉS, J. LLIBRE, A.C. REZENDE, D. SCHLOMIUK AND N. VULPE, *Global configurations of singularities for quadratic differential systems with exactly two finite singularities of total multiplicity four*, Electron. J. Qual. Theory Differ. Equ. **2014** (2014), 1–43.
- [14] J.C. ARTÉS, J. LLIBRE AND D. SCHLOMIUK, *The geometry of quadratic differential systems with a weak focus of second order*, Internat. J. Bifur. Chaos **16** (2006), 3127–3194.
- [15] J. C. ARTÉS, J. LLIBRE AND D. SCHLOMIUK, *The geometry of quadratic polynomial differential systems with a weak focus and an invariant straight line*, Internat. J. Bifur. Chaos **20** (2010), 3627–3662.
- [16] J. C. ARTÉS, J. LLIBRE, D. SCHLOMIUK AND N. VULPE, *Configurations of singularities for quadratic differential systems with total finite multiplicity lower than 2*, Bul. Acad. Stiinte Repub. Mold. Mat. (2013), 72–124.
- [17] J. C. ARTÉS, J. LLIBRE, D. SCHLOMIUK AND N. VULPE, *Geometric configurations of singularities for quadratic differential systems with three distinct real simple finite singularities*, J. Fixed Point Theory Appl. **14** (2013), 555–618.
- [18] J.C. ARTÉS, J. LLIBRE, D. SCHLOMIUK AND N. VULPE, *Geometric configurations of singularities for quadratic differential systems with total finite multiplicity $m_f = 2$* , Electron. J. Differential Equations **2014** (2014), 1–79.
- [19] J. C. ARTÉS, J. LLIBRE, D. SCHLOMIUK AND N. VULPE, *Global configurations of singularities for quadratic differential systems with total finite multiplicity three and at most two real singularities*, Qual. Theory Dyn. Syst. **13** (2014), 305–351.
- [20] J. C. ARTÉS, J. LLIBRE, D. SCHLOMIUK AND N. VULPE, *Global configurations of singularities for quadratic differential systems with exactly three finite singularities of total multiplicity four*, Electron. J. Qual. Theory Differ. Equ. **2015** (2015), 1–60.
- [21] J. C. ARTÉS, J. LLIBRE, D. SCHLOMIUK AND N. VULPE, *From topological to geometric equivalence in the classification of singularities at infinity for quadratic vector fields*, Rocky Mountain J. of Math. **45** (2015), 29–113.
- [22] J.C. ARTÉS, J. LLIBRE, D. SCHLOMIUK AND N. VULPE, *Geometric configurations of singularities of planar polynomial differential systems. A global classification on the quadratic case*, to appear in Birkhäuser, 2020.
- [23] J.C. ARTÉS, J. LLIBRE AND N. VULPE, *Quadratic systems with a rational first integral of degree 2: a complete classification in the coefficient space \mathbb{R}^{12}* , Rend. Circ. Mat. Palermo (2) **56** (2007), 417–444.
- [24] J.C. ARTÉS, J. LLIBRE AND N. VULPE, *When singular points determine quadratic systems*, Electron. J. Differential Equations **2008** (2008), 1–37.
- [25] J.C. ARTÉS, J. LLIBRE AND N. VULPE, *Singular points of quadratic systems: A complete classification in the coefficient space \mathbb{R}^{12}* , Internat. J. Bifur. Chaos **18** (2008), 313–362.
- [26] J.C. ARTÉS, J. LLIBRE AND N. VULPE, *Quadratic systems with a polynomial first integral: a complete classification in the coefficient space \mathbb{R}^{12}* , J. Differential Equations **246** (2009), 3535–3558.
- [27] J.C. ARTÉS, J. LLIBRE AND N. VULPE, *Complete geometric invariant study of two classes of quadratic systems*, Electron. J. Differential Equations **2012** (2012), 1–35.
- [28] J.C. ARTÉS, J. LLIBRE AND N. VULPE, *Quadratic systems with an integrable saddle: A complete classification in the coefficient space \mathbb{R}^{12}* , Nonlinear Anal. **75** (2012), 5416–5447.
- [29] J.C. ARTÉS, A. C. REZENDE AND R. OLIVEIRA, *Global phase portraits of quadratic polynomial differential systems with a semi-elemental triple node*, Internat. J. Bifur. Chaos **23** (2013), 1350140.
- [30] J.C. ARTÉS, A. C. REZENDE AND R. OLIVEIRA, *The geometry of quadratic polynomial differential systems with a finite and an infinite saddle-node (A, B)* , Internat. J. Bifur. Chaos **24** (2014), 1450044.

- [31] J.C. ARTÉS, A. C. REZENDE AND R. D. S. OLIVEIRA, *The geometry of quadratic polynomial differential systems with a finite and an infinite saddle-node (C)*, Internat. J. Bifur. Chaos **25** (2015), 1530009.
- [32] V. BALTAG AND N. VULPE, *The number and multiplicity of singular points of a quadratic differential system*, (Russian) Dokl. Akad. Nauk. **323** (1992), 9–12; translation in Russian Acad. Sci. Dokl. Math. **45** (1992), 235–238.
- [33] V.A. BALTAG AND N.I. VULPE, *Affine-invariant conditions for determining the number and multiplicity of singular points of quadratic differential systems*, Izv. Akad. Nauk Respub. Moldova Mat. **1** (1993), 39–48.
- [34] R. BAMON, *Quadratic vector fields in the plane have a finite number of limit cycles*, Inst. Hautes Études Sci. Publ. Math., No. 64 (1986), 111–142.
- [35] N.N. BAUTIN, *On the number of limit cycles which appear with the variation of coefficients from an equilibrium position of focus or center type*, Mat. Sb. (N.S) **30** (7)2 (1952), 181–196 (Russian), Transl. Amer. Math. Soc. Ser. (1), **100** (1954), 397–413.
- [36] A. BERLINSKII, *On the coexistence of singular points of different types*, (Russian), Izv. Vyss. Ucebn. Zaved. Matematika **18** (1960), 27–32.
- [37] J. BERNOULLI, *Explicationes, Annotationes Additiones ad ea, quae in Actis sup. de Curva Elastica, Isochrone Paracentrica, Velaria, hinc inde memorata, paratim controversa leguntur; ubi de Linea mediarum directionum, aliisque novis*, Acta Eruditorum, 1965 (Cited [100]).
- [38] Y. BOLAÑOS, J. LLIBRE AND C. VALLS, *Phase portraits of quadratic Lotka–Volterra systems with a Darboux invariant in the Poincaré disc*, Comm. in Contemporary Math. **16** (2014), 1350041.
- [39] Y. BOLAÑOS, J. LLIBRE AND C. VALLS, *Liouvillian first integrals for quadratic systems with an integrable saddle*, Rocky Mountain J. Math. **45** (2015), 1765–1779.
- [40] D. BOULARAS, *A new classification of planar homogeneous quadratic systems*, (English summary) Qual. Theory Dyn. Syst. **2** (2001), 93–110.
- [41] W. BÜCHEL, *Zur topologie der durch eine gewöhnliche differentialgleichung erster ordnung und ersten grades definierten kurvenschar*, Mitteil. der Math. Gesellsch. in Hamburg **4** (1904), 33–68.
- [42] C. A. BUZZI AND D. J. TONON, *Quadratic planar systems with two parallel invariant straight lines*, Qual. Theory Dyn. Syst. **7** (2009), 295–316.
- [43] SUI LIN CAI, *The weak saddle and separatrix cycle of a quadratic system*, Acta Math. Sinica **30** (1987), 553–559 (Chinese).
- [44] L. CAIRÓ, M.R. FEIX AND J. LLIBRE, *Darboux method and invariants for the Lotka–Volterra and complex quadratic systems*, J. Math. Physics **40** (1999), 2074–2091.
- [45] L. CAIRÓ, H. GIACOMINI AND J. LLIBRE, *Liouvillian first integrals for the planar Lotka–Volterra systems*, Rend. Circ. Mat. Palermo (2) **52** (2003), 389–418.
- [46] L. CAIRÓ AND J. LLIBRE, *Phase portraits of planar semi-homogeneous vector fields*, Nonlinear Anal. **29** (1997), 783–811.
- [47] L. CAIRÓ AND J. LLIBRE, *Integrability of the 2D Lotka–Volterra system via polynomial first integrals and polynomial inverse integrating factors*, J. Phys. A **33** (2000), 2407–2417.
- [48] L. CAIRÓ AND J. LLIBRE, *Darbouxian first integrals and invariants for real quadratic systems having an invariant conic*, J. Physics A: Math. Gen. **35** (2002), 589–608.
- [49] L. CAIRÓ AND J. LLIBRE, *Phase portraits of quadratic polynomial vector fields having a rational first integral of degree 2*, Nonlin. Anal. **67** (2007), 327–348.
- [50] F. CAO AND J. JIANG, *The classification on the global phase portraits of two-dimensional Lotka–Volterra system*, J. Dyn. Diff. Equat. **20** (2008), 797–830.
- [51] J. CHAVARRIGA, B. GARCÍA, J. LLIBRE, J.S. PÉREZ DEL RÍO AND J.A. RODRÍGUEZ, *Polynomial first integrals of quadratic vector fields*, J. Differential Equations **230** (2006), 393–421.

- [52] J. CHAVARRIGA, H. GIACOMINI AND J. LLIBRE, *Uniqueness of algebraic limit cycles for quadratic systems*, J. Math. Anal. Appl. **261** (2001), 85–99.
- [53] J. CHAVARRIGA AND M. GRAU, *Invariant algebraic curves linear in one variable for planar real quadratic systems*, Appl. Math. Comput. **138** (2003), 291–308.
- [54] J. CHAVARRIGA, J. LLIBRE AND J. SOROLLA, *Algebraic limit cycles of quadratic systems*, J. Differential Equations, **200** (2003), 206–244.
- [55] J. CHAVARRIGA, J. LLIBRE AND J. SOTOMAYOR, *Algebraic solutions for polynomial systems with emphasis in the quadratic case*, Exposition. Math. **15** (1997), 161–173.
- [56] J. CHAVARRIGA AND M. SABATINI, *A survey of isochronous centers*, Qual. Theory Dyn. Syst. **1** (1999), 1–70.
- [57] F. CHEN, C. LI, J. LLIBRE AND Z. ZHANG, *A unified proof on the weak Hilbert’s 16th problem for $n = 2$* , J. Differential Equations **221** (2006), 309–342.
- [58] G. CHEN, C. LI, C. LIU AND J. LLIBRE, *The cyclicity of period annuli of some classes of reversible quadratic systems*, Disc. Cont. Dyn. Syst. Series A **16** (2006), 157–177.
- [59] L. CHEN, X. MA, J. ZHANG AND C. LI, *Cyclicity of several quadratic reversible systems with center of genus one*, J. Appl. Anal. Comput. **1** (2011), 439–437.
- [60] L. S. CHEN AND M. S. WANG, *The relative position and number of limit cycles of a quadratic differential system*, Acta Math. Sinica **22** (1979), 751–768.
- [61] L.A. CHERKAS, J.C. ARTÉS AND J. LLIBRE, *Quadratic systems of limit cycles of normal size*, Bulletin of Academy of Sciences of Moldova (Matematica) **1**(41) (2003), 31–46.
- [62] C. CHICONE AND D. SHAFER, *Separatrix and limit cycles of quadratic systems and Dulac’s theorem*, Trans. Amer. Math. Soc. **278** (1983), 585–612.
- [63] C. CHICONE AND J. TIAN, *On general properties of quadratic systems*, The American Mathematical Monthly, Vol. **89**, No. 3 (Mar., 1982), pp. 167–178.
- [64] C. CHRISTOPHER AND C. LI, *Limit cycles of differential equations*, Advanced Courses in Mathematics, CRM Barcelona, Birkhäuser Verlag, Basel, 2007.
- [65] C. CHRISTOPHER AND J. LLIBRE, *A family of quadratic polynomial differential systems with invariant algebraic curves of arbitrarily high degree without rational first integrals*, Proc. Amer. Math. Soc. **130** (2002), 2025–2030.
- [66] C. CHRISTOPHER, J. LLIBRE AND J. PEREIRA, *Multiplicity of invariant algebraic curves in polynomial vector fields*, Pacific J. Math., Vol. **29** (2007), 63–117.
- [67] C. CHRISTOPHER, J. LLIBRE AND G. ŚWIRSZCZ, *Invariant algebraic curves of large degree for quadratic systems*, J. Math. Anal. Appl. **303** (2005), 450–461.
- [68] A. CIMA, A. GASULL AND F. MAÑOSAS, *Some applications of the Euler–Jacobi formula to differential equations*, Proc. Amer. Math. Soc. **118** (1993), 151–163.
- [69] B. COLL AND J. LLIBRE, *Limit cycles for a quadratic system with an invariant straight line and some evolution of phase portraits*, Colloquia Mathematica Societatis Janos Bolyai, Vol. **53**, Szeget, Hungary, (1988), 111–123.
- [70] B. COLL, A. FERRAGUT AND J. LLIBRE, *Phase portraits of the quadratic systems with a polynomial inverse integrating factor*, Internat. J. Bifur. Chaos **19** (2009), 765–783.
- [71] B. COLL, A. FERRAGUT AND J. LLIBRE, *Polynomial inverse integrating factors for quadratic differential systems*, Nonlinear Anal. **73** (2010), 881–914.
- [72] B. COLL, A. GASULL AND J. LLIBRE, *Some theorems on the existence, uniqueness and non existence of limit cycles for quadratic systems*, J. Differential Equations **67** (1987), 372–399.
- [73] B. COLL, A. GASULL AND J. LLIBRE, *Quadratic systems with a unique finite rest point*, Publ. Mat. **32** (1988), 199–259.
- [74] B. COCLSWLL, C. LI AND R. PROHENS, *Quadratic perturbations of a class of quadratic reversible systems with two centers*, Disc. Cont. Dyn. Syst. **24** (2009), 699–729.
- [75] R. CONTI, *Uniformly isochronous centers of polynomial systems in \mathbb{R}^2* , Lecture Notes in Pure and Appl. Math. **152** (1994), 21–31.

- [76] W.A. COPPEL, *A Survey of Quadratic Systems*, J. Differential Equations **2** (1966), 293–304.
- [77] W.A. COPPEL, *Some quadratic systems with at most one limit cycle*, Dynamics reported, Vol. **2**, 61–88, Dynam. Report. Ser. Dynam. Systems Appl. **2**, Wiley, Chichester, 1989.
- [78] G. DARBOUX, *Mémoire sur les équations différentielles du premier ordre et du premier degré*, Bull. Sci. Math. 2me série **2**(1) (1878), 60–96; 123–144; 151–200.
- [79] T. DATE, *Classification and analysis of two-dimensional real homogeneous quadratic differential equation systems*, J. Differential equations **32** (1979), 311–334.
- [80] R.J. DICKSON AND L.M. PERKO, *Bounded quadratic systems in the plane*, J. Differential Equations **7** (1970), 251–273.
- [81] F. DUMORTIER, *Techniques in the theory of local bifurcations: Blow-up, normal forms, nilpotent bifurcations, singular perturbations*, in Bifurcations and Periodic Orbits of Vector Fields, Springer, 1993, 19–73.
- [82] F. DUMORTIER, C. HERSSENS AND L. PERKO, *Local bifurcations and a survey of bounded quadratic systems*, J. Differential Equations **165** (2000), 430–467.
- [83] F. DUMORTIER, C. LI AND Z. ZHANG, *Unfolding of a quadratic integrable system with two centers and two unbounded heteroclinic loops*, J. Differential Equations **139** (1997), 146–193.
- [84] F. DUMORTIER, J. LLIBRE AND J. C. ARTÉS, *Qualitative Theory of Planar Differential Systems*, Springer-Verlag, Berlin, 2006.
- [85] F. DUMORTIER, J. LLIBRE AND J.C. ARTÉS, *Limit cycles near hyperbolas in quadratic systems*, J. Differential Equations **246** (2009), 235–260.
- [86] F. DUMORTIER, R. ROUSSARIE AND C. ROUSSEAU, *Hilbert’s 16th problem for quadratic vector fields*, J. Differential Equations **110** (1994), 86–133.
- [87] J. ECALLE, *Introduction aux fonctions analysables et preuve constructive de la conjecture du Dulac*, Actualités Math., Paris, Hermann, Paris, 1992.
- [88] R.M. EVDOKIMENCO, *Construction of algebraic paths and the qualitative investigation in the large of the properties of integral curves of a system of differential equations*, Differential Equations **6** (1970), 1349–1358.
- [89] R.M. EVDOKIMENCO, *Behavior of integral curves of a dynamic system*, Differential Equations **9** (1974), 1095–1103.
- [90] R.M. EVDOKIMENCO, *Investigation in the large of a dynamic system*, Differential Equations **15** (1979), 215–221.
- [91] V.F. FILIPTOV, *Algebraic limit cycles*, Differential Equations **9** (1973), 983–986.
- [92] M. FROMMER, *Über das Auftreten von Wirbeln und Strudeln (geschlossener und spiraliger Integralcurven) in der Umgebung rationaler Unbestimmtheitsstellen*, Math. Ann. **109** (1934), 395–424 .
- [93] B. GARCIA, J. LLIBRE AND J. PÉREZ DEL RIO, *Phase portraits of the quadratic vector fields with a polynomial first integral*, Rend. Circ. Mat. Palermo (2) **55** (2006), 420–440.
- [94] I. GARCÍA AND J. LLIBRE, *Classical planar algebraic curves realizable by quadratic polynomial differential systems*, Internat. J. Bifur. Chaos **27** (2017), 1750141.
- [95] A. GASULL, S. LI REN AND J. LLIBRE, *Chordal quadratic systems*, Rocky Mountain J. Math. **16** (1986), 751–782.
- [96] A. GASULL AND J. LLIBRE, *On the nonsingular quadratic differential equations in the plane*, Proc. Amer. Math. Soc. **104** (1988), 793–794.
- [97] A. GASULL AND R. PROHENS, *Quadratic and cubic systems with degenerate infinity*, J. Math. Anal. Appl. **198** (1996), 25–34.
- [98] L. GAVRILOV, *The infinitesimal 16th Hilbert problem in the quadratic case*, Invent. Math. **143** (2001), 449–497.
- [99] L. GAVRILOV AND E. HOROZOV, *Limit cycles of perturbations of quadratic Hamiltonian vector fields*, J. Math. Pures Appl. **72** (1993), 2013–238.

- [100] E. HAIRER, S.P. NORSETT AND G. WANNER, *Solving ordinary differential equations I: Nonstiff problems*, Berlin, New York, Springer-Verlag, 1993.
- [101] D. HILBERT, *Mathematical problems*, reprinted from Bull. Amer. Mat. Soc. **8** (1902), 437–479, in Bull. Amer. Math. **37** (2000), 407–436.
- [102] C. HONG AND L. ZOU, *Abelian integrals and critical periods for a class of reversible quadratic systems*, Sichuan Daxue Xuebao **49** (2012), 960–964.
- [103] X. HONG, R. MA AND B. WANG, *Estimation of the number of zeros of Abelian integrals of a quadratic reversible system*, Math. Pract. Theory **45** (2015), 270–276.
- [104] X. HONG, S. XIE AND L. CHEN, *Estimating the number of zeros for Abelian integrals of quadratic reversible centers with orbits formed by higher order curves*, Internat. J. Bifur. Chaos **26** (2016), 1650020.
- [105] X. HONG, S. XIE AND R. MA, *On the Abelian integrals of quadratic reversible centers with orbits formed by genus 1 curves of higher degree*, J. Math. Anal. Appl. **429** (2015), 924–941.
- [106] X. HUANG AND J. W. REYN, *On the limit cycle distribution over two nests in quadratic systems*, Bull. Austral. Math. Soc. **52** (1995), 461–474.
- [107] I.D. ILIEV, *The cyclicity of the period annulus of the quadratic Hamiltonian triangle*, J. Differential Equations **128** (1996), 309–326.
- [108] I.D. ILIEV, *Perturbations of quadratic centers*, Bull. Sci. Math. **122** (1998), 107–161.
- [109] I.D. ILIEV, C. LI AND J. YU, *Bifurcations of limit cycles in a reversible quadratic system with a center, a saddle and two nodes*, Commun. Pure Appl. Anal. **9** (2010), 583–610.
- [110] YU. ILYASHENKO, *Finiteness theorems for limit cycles*, Translation of Mathematical Monographs **94**, 1991.
- [111] YU. ILYASHENKO AND J. LLIBRE, *A restricted version of the Hilbert’s 16th problem for quadratic vector fields*, Moscou Math. J. **10** (2010), 317–335.
- [112] P. DE JAGER, *Phase portraits for quadratic systems with a higher order singularity with two zero eigenvalues*, J. Differential Equations **87** (1990), 169–204.
- [113] X. JARQUE, J. LLIBRE AND D. S. SHAFER, *Structurally stable quadratic foliations*, Rocky Mountain J. Math. **38** (2008), 489–530.
- [114] P. JOYAL AND C. ROUSSEAU, *Saddle quantities and applications*, J. Differential Equations **78** (1989), 374–399.
- [115] Y.F. KALIN AND N. I. VULPE, *Affine invariant conditions for the topological discrimination of quadratic Hamiltonian differential systems*, (Russian) Differ. Uravn. **34** (1998), 298–302; translation in Differential equations **34** (1998), 297–301.
- [116] V. KALOSHIN, *The existential Hilbert 16-th problem and an estimate for cyclicity of elementary polycycles*, Invent. Mat. **151** (2003), 451–512.
- [117] W. KAPTEYN, *On the midpoints of integral curves of differential equations of the first order and the first degree*, (Dutch) Nederl. Acad. Wetensch. Verslagen, Afd. Natuurkunde Koninkl. Nederland. **XIX**(2), (1911), 1446–1457.
- [118] W. KAPTEYN, *New investigations on the midpoints of integrals of differential equations of the first order and the first degree*, (Dutch) Nederl. Acad. Wetensch. Verslagen, Afd. Natuurkunde **XX**(2), (1912), 1354–1365.
- [119] E.F. KIRNICKAJA AND K.S. SIBIRSCHI, *Conditions for two centers for a quadratic differential system (Russian)*, Differentsial’nye Uravneniya **14** (1978), 1589–1593; translation in Differential Equations **14** (1978), 1129–1133.
- [120] C. LI, *Non-existence of limit cycles around a weak focus of order three for any quadratic system*, Chinese Ann. Math. Series B **7** (1986), 174–190.
- [121] C. LI, *Abelian integrals and limit cycles*, Qual. Theory Dyn. Syst. **11** (2012), 111–128.
- [122] C. LI, W. LI, J. LLIBRE AND Z. ZHANG, *Linear estimate for the number of zeros of Abelian integrals for quadratic isochronous centers*, Nonlinearity **13** (2000), 1775–1800.

- [123] C. LI AND J. LLIBRE, *A Unified study on the cyclicity of period annulus of the reversible quadratic Hamiltonian systems*, J. Dyn. Diff. Equat. **16** (2004), 271–295.
- [124] C. LI AND J. LLIBRE, *The cyclicity of period annulus of a quadratic reversible Lotka–Volterra system*, Nonlinearity **22** (2009), 2971–2979.
- [125] C. LI AND J. LLIBRE, *Quadratic perturbations of a quadratic reversible Lotka–Volterra system*, Qual. Theory Dyn. Syst. **9** (2010), 235–239.
- [126] C. LI, J. LLIBRE AND Z. ZHANG, *Abelian integrals of quadratic Hamiltonian vector fields with an invariant straight line*, Publ. Math. **39** (1995), 355–366.
- [127] C. LI, J. LLIBRE AND Z. ZHANG, *Weak focus, limit cycles and bifurcations for bounded quadratic systems*, J. Differential Equations **115** (1995), 193–223.
- [128] C. LI AND R. ROUSSARIE, *The cyclicity of the elliptic segment loops of the reversible quadratic Hamiltonian systems under quadratic perturbations*, J. Differential Equations **205** (2004), 488–520.
- [129] C. LIU, *Limit cycles bifurcated from some reversible quadratic centers with a non-algebraic first integral*, Nonlinearity **25** (2012), 1653–1660.
- [130] C. LIU, *The cyclicity of period annuli of a class of quadratic reversible systems with two centers*, J. Differential Equations **252** (2012), 5260–5273.
- [131] J. LLIBRE, *Averaging theory and limit cycles for quadratic systems*, Radovi Matematički **11** (2002), 215–228.
- [132] J. LLIBRE, *Open problems on the algebraic limit cycles of planar polynomial vector fields*, Bull. Academy of Sciences of Moldova (Matematica) **56** (2008), 19–26.
- [133] J. LLIBRE AND J. C. MEDRADO, *Darboux integrability and reversible quadratic vector fields*, Rocky Mountain J. Math. **35** (2005), 1999–2057.
- [134] J. LLIBRE, M. MESSIAS AND A. C. REINOL, *Global phase portraits of quadratic and cubic systems with two nonconcentric circles as invariant algebraic curves*, Dyn. Syst. **32** (2017), 374–390.
- [135] J. LLIBRE AND R. OLIVEIRA, *Phase portraits of quadratic polynomial vector fields having a rational first integral of degree 3*, Nonlinear Anal. **70** (2009), 3549–3560.
- [136] J. LLIBRE AND R. OLIVEIRA, *Erratum to “Phase portraits of quadratic polynomial vector fields having a rational first integral of degree 3”*, Nonlinear Anal. **70** (2009), 3549–3560, Nonlinear Anal., **71** (2009), 6378–6379.
- [137] J. LLIBRE AND R.D.S. OLIVEIRA, *Phase portraits of quadratic polynomial vector fields having a rational first integral of degree 3*, Nonlin. Anal. **70** (2009), 3549–3560.
- [138] J. LLIBRE AND R. OLIVEIRA, *Quadratic systems with invariant straight lines of total multiplicity two having Darboux invariants*, Comm. Contemporary Math. **17** (2015), 1450018.
- [139] J. LLIBRE AND R. OLIVEIRA, *Quadratic systems with an invariant conic having Darboux invariants*, Comm. Contemporary Math. **20** (2018), 1750033.
- [140] J. LLIBRE, J.S. PÉREZ DEL RÍO AND J.A. RODRÍGUEZ, *Phase portraits of a new class of integrable quadratic vector fields*, Dyn. Cont. Disc. Impulsive Systems **7** (2000), 595–616.
- [141] J. LLIBRE, J.S. PÉREZ DEL RÍO AND J.A. RODRÍGUEZ, *Averaging analysis of a perturbed quadratic center*, Nonlinear Analysis **46** (2001), 45–51.
- [142] J. LLIBRE, W.F. PEREIRA AND C. PESSOA, *Phase portraits of Bernoulli quadratic polynomial differential systems*, Electron. J. Differential Equations **2020** (2020), 1–19.
- [143] J. LLIBRE, R. RAMIREZ AND A. SADOVSKAIA, *On the 16th Hilbert problem for algebraic limit cycles*, J. Differential Equations **248** (2010), 1401–1409.
- [144] J. LLIBRE, R. RAMIREZ AND A. SADOVSKAIA, *On the 16th Hilbert problem for limit cycles on nonsingular algebraic curves*, J. Differential Equations **250** (2011), 983–999.
- [145] J. LLIBRE AND D. SCHLOMIUK, *The geometry of differential quadratic systems with a weak focus of third order*, Canadian J. Math. **56** (2004), 310–343.
- [146] J. LLIBRE AND D. SCHLOMIUK, *On the limit cycles bifurcating from an ellipse of a quadratic center*, Disc. Cont. Dyn. Syst. Series A **35** (2015), 1091–1102.

- [147] J. LLIBRE AND P.R. DA SILVA, *Singularly perturbed quadratic polynomial differential systems on the plane*, Brazilian J. Math. and Stat. **25** (2007), 137–150.
- [148] J. LLIBRE AND G. ŚWIRSZCZ, *Relationships between limit cycles and algebraic invariant curves for quadratic systems*, J. Differential Equations **229** (2006), 529–537.
- [149] J. LLIBRE AND G. ŚWIRSZCZ, *Classification of quadratic systems admitting the existence of an algebraic limit cycle*, Bull. Sci. Math. **131** (2007), 405–421.
- [150] J. LLIBRE AND Y. TIAN, *Algebraic limit cycles bifurcating from algebraic ovals of quadratic centers*, Internat. J. Bifur. Chaos **28** (2018), 1850145.
- [151] J. LLIBRE AND C. VALLS, *Analytic integrability of quadratic-linear polynomial differential systems*, Ergodic Theory Dyn. Syst. **31** (2011), 245–258.
- [152] J. LLIBRE AND C. VALLS, *Liouvillian first integrals of quadratic-linear polynomial differential systems*, J. Math. Anal. Appl. **379** (2011), 188–199.
- [153] J. LLIBRE AND C. VALLS, *Liouvillian and analytic integrability of the quadratic vector fields having an invariant ellipse*, Acta Math. Sinica **30** (2014), 453–466.
- [154] J. LLIBRE AND C. VALLS, *On the uniqueness of algebraic limit cycles for quadratic polynomial differential systems with two pairs of equilibrium points at infinity*, Geometriae Dedicata **191** (2017), 37–52.
- [155] J. LLIBRE AND C. VALLS, *Global phase portraits for the Abel quadratic polynomial differential equations of second kind with \mathbb{Z}_2 -symmetries*, Canadian Math. Bull. **61** (2018), 149–165.
- [156] J. LLIBRE AND C. VALLS, *Algebraic limit cycles for quadratic polynomial differential systems*, Disc. Cont. Dyn. Syst. Series B **23** (2018), 2475–2485.
- [157] J. LLIBRE AND C. VALLS, *Algebraic limit cycles on quadratic polynomial differential systems*, Proc. Edinb. Math. Soc. **61** (2018), 499–512.
- [158] J. LLIBRE AND C. VALLS, *Global phase portraits of quadratic systems with a complex ellipse as invariant algebraic curve*, Acta Math. Sinica **34** (2018), 801–811.
- [159] J. LLIBRE AND C. VALLS, *Integrability of reversible and equivariant quadratic polynomial differential systems in the plane*, Rocky Mountain J. Math. **49** (2019), 579–591.
- [160] J. LLIBRE AND J. YU, *Global phase portraits of quadratic systems with an ellipse and a straight line as invariant algebraic curves*, Electron. J. Differential Equations **2015**, 1–14.
- [161] J. LLIBRE AND J. YU, *Global phase portraits of quadratic systems with a hyperbola and a straight line as invariant algebraic curves*, Electron. J. Differential Equations **2018**, 1–19.
- [162] J. LLIBRE AND X. ZHANG, *Topological phase portraits of planar semi-linear quadratic vector fields*, Houston J. Math. **27** (2001), 247–296.
- [163] J. LLIBRE AND X. ZHANG, *Non-existence, existence and uniqueness of limit cycles for quadratic polynomial differential systems*, Proc. Royal Soc. Edinburgh-Section A Mathematics **149** (2019), 1–14.
- [164] W.S. LOUD, *Behavior of the period of solutions of certain plane autonomous systems near centers*, Contributions to Differential Equations **3** (1964), 21–36.
- [165] V.A. LUNKEVICH AND K. S. SIBIRSCHI, *Integrals of general quadratic systems in cases of a center*, (Russian), Differentsialnye Uravneniya **8** (1982), 786–792; translation in Differential Equations **8** (1982), 563–568.
- [166] M. LUPAN AND N. VULPE, *Classification of quadratic systems with a symmetry center and simple infinite singular points*, Bul. Acad. Stiinte Repub. Mold. Mat. (2003), 102–119.
- [167] L.S. LYAGINA, *Integralnye Krivye Uravneniya $dy/dx = (ax^2 + bxy + cy^2)/(dx^2 + exy + fy^2)$* , Uspehi Matem. Nauk (N.S.) **6** (1951), 171–183.
- [168] L. MARKUS, *Global structure of ordinary differential equations in the plane*, Trans. Amer. Math. Soc. **76** (1954), 127–148.

- [169] L. MARKUS, *Quadratic differential equations and non-associative algebras*, in “Contributions to the theory of nonlinear oscillations”, Vol. **V**, 185–213, Annals of Mathematics Studies, No. **45**, Princeton University Press, Princeton, N.J. 1960.
- [170] J. MOULIN OLLAGNIER, *About a conjecture on quadratic vector fields*, J. Pure Appl. Algebra **165** (2001), 227–234.
- [171] J. MOULIN OLLAGNIER, *Liouvillian integration of the Lotka-Volterra system*, Qual. Theory Dyn. Syst. **2** (2001), 307–358.
- [172] D. NEUMANN, *Classification of continuous flows on 2-manifolds*, Proc. Amer. Math. Soc. **48** (1975), 73–81.
- [173] I.V. NIKOLAEV AND N.I. VULPE, *Topological classification of quadratic systems with a unique finite second order singularity with two zero eigenvalues*, Izv. Akad. Nauk Respub. Moldova Mat. (1993), 3–8.
- [174] I.V. NIKOLAEV AND N.I. VULPE, *Topological classification of quadratic systems at infinity*, J. London Math. Soc. (2) **55** (1997), 473–488.
- [175] J. PAL AND D. SCHLOMIUK, *Summing up the dynamics of quadratic Hamiltonian systems with a center*, Canad. J. Math. **49** (1997), 583–599.
- [176] L. PENG, *Quadratic perturbations of a quadratic reversible center of genus one*, Front. Math. China **6** (2011), 911–930.
- [177] L. PENG, Z. FENG AND C. LIU, *Quadratic perturbations of a quadratic reversible Lotka-Volterra system with two centers*, Disc. Cont. Dyn. Syst. **34** (2014), 4807–4826.
- [178] L. PENG AND Y. LEI, *Bifurcations of limit cycles from a quadratic reversible center with the unbounded elliptic separatrix*, Bull. Iranian Math. Soc. **39** (2013), 1225–1248.
- [179] L. PENG AND Y. LI, *On the limit cycles bifurcating from a quadratic reversible center of genus one*, Mediter. J. Math. **11** (2014), 373–392.
- [180] I. PETROVSKI AND E. LANDIS, *On the number of limit cycles of the equation $dy/dx = P(x, y)/Q(x, y)$, where P and Q are polynomials of 2nd degree*, (Russian) Mat. Sb. N.S. **37**(79) (1955), 209–250.
- [181] I. PETROVSKI AND E. LANDIS, *On the number of limit cycles of the equation $dy/dx = M(x, y)/N(x, y)$, where the M and N are polynomials of second degree*, (Russian) Dokl. Akad. Nauk SSSR (N.S.) **102** (1955), 29–32.
- [182] I. PETROVSKI AND E. LANDIS, *On the number of limit cycles of the equation $dy/dx = P(x, y)/Q(x, y)$, where P and Q are polynomials of the second degree*, American Mathematical Society Translations, Ser. 2 **10** (1958), 177–221.
- [183] I. PETROVSKI AND E. LANDIS, *Corrections to the articles “On the number of limit cycles of the equations $dy/dx = P(x, y)/Q(x, y)$, where P and Q are polynomials of 2nd degree” and “On the number of limit cycles of the equation $dy/dx = P(x, y)/Q(x, y)$, where P and Q are polynomials”*, (Russian) Mat. Sb. (N.S.) **48**(90) (1959), 253–255.
- [184] I. PETROVSKI AND E. LANDIS, *A letter to the editors*, (Russian) Mat. Sb. (N.S.) **73** (115) (1967), 160.
- [185] QIN YUAN-XUN, *On the algebraic limit cycles of second degree of the differential equation $dy/dx = \sum_{0 \leq i+j \leq 2} a_{ij}x^i y^j / \sum_{0 \leq i+j \leq 2} b_{ij}x^i y^j$* , Acta Math. Sinica **8** (1958), 23–35.
- [186] J.W. REYN, *Phase portraits of a quadratic system of differential equations occurring frequently in applications*, Nieuw Arch. Wisk. (4) **5** (1987), 107–151.
- [187] J. W. REYN, *Classes of quadratic systems of differential equations in the plane*, Dynamical systems (Tianjin, 1990/1991), 146–180; translation in Nankai Ser. Pure Appl. Math. Theoret. Phys. **4**, World Sci. Publ., River Edge, NJ, 1993.
- [188] J. W. REYN, *A bibliography of the qualitative theory of quadratic systems of differential equations in the plane*, Third edition, Delft University of Technology, Faculty of Technical Mathematics and Informatics 2, 1994.
- [189] J. REYN, *Phase Portraits of Planar Quadratic Systems*, Mathematics and Its Applications vol. **583**, Springer, 2007.

- [190] C. ROUSSEAU, C. SHAN AND H. ZHU, *Finite cyclicity of some graphics through a nilpotent point of saddle type inside quadratic systems*, Qual. Theory Dyn. Syst. **15** (2016), 237–256.
- [191] N.A. SAKHARNIKOV, *On Frommer's conditions for the existence of a center*, (Russian) Prikl. Mat. Mek. **12** (1948), 669–670.
- [192] G.T. DOS SANTOS, *Classification of generic quadratic vector fields without limit cycles*, Lecture Notes in Mathematics, Vol. **597**, 605–640, Springer, New York/Berlin, 1977.
- [193] D. SCHLOMIUK, *Algebraic particular integrals, integrability and the problem of the center*, Trans. Amer. Math. Soc. **338** (1993), 799–841.
- [194] D. SCHLOMIUK, *Elementary first integrals and algebraic invariant curves of differential equations*, Exposition. Math. **11** (1993), 433–454.
- [195] D. SCHLOMIUK, J. GUCKENHEIMER AND R. RAND, *Integrability of plane quadratic vector fields*, Exposition. Math. **8** (1990), 3–25.
- [196] D. SCHLOMIUK AND J. PAL, *On the geometry in the neighborhood of infinity of quadratic differential systems with a weak focus*, Qual. Theory Dyn. Syst. **2** (2001), 1–43.
- [197] D. SCHLOMIUK AND N. VULPE, *Planar quadratic differential systems with invariant straight lines of at least five total multiplicity*, Qual. Theory Dyn. Syst. **5** (2004), 135–194.
- [198] D. SCHLOMIUK AND N. VULPE, *Geometry of quadratic differential systems in the neighborhood of infinity*, J. Differential Equations **215** (2005), 357–400.
- [199] D. SCHLOMIUK AND N. VULPE, *Integrals and phase portraits of planar quadratic differential systems with invariant lines of at least five total multiplicity*, Rocky Mountain J. Math. **38** (2008), 2015–2075.
- [200] D. SCHLOMIUK AND N. VULPE, *Integrals and phase portraits of planar quadratic differential systems with invariant lines of total multiplicity four*, Bul. Acad. Stiinte Repub. Mold. Mat. (2008), 27–83.
- [201] D. SCHLOMIUK AND N. VULPE, *Planar quadratic differential systems with invariant straight lines of total multiplicity four*, Nonlinear Anal. **68** (2008), 681–715.
- [202] D. SCHLOMIUK AND N. VULPE, *The full study of planar quadratic differential systems possessing a line of singularities at infinity*, J. Dyn. Diff. Equat. **20** (2008), 737–775.
- [203] D. SCHLOMIUK AND N. VULPE, *Global classification of the planar Lotka–Volterra differential systems according to their configurations of invariant straight lines*, J. Fixed Point Theory Appl. **8** (2010), 177–245.
- [204] D. SCHLOMIUK AND N. VULPE, *Global topological classification of Lotka–Volterra quadratic differential systems*, Electron. J. Differential Equations **2012** (2012), 1–69.
- [205] D. SCHLOMIUK AND X. ZHANG, *Quadratic differential systems with complex conjugate invariant lines meeting at a finite point*, J. Differential Equations **265** (2018), 3650–3684.
- [206] Y. SHAO AND A. CHUNXIANG, *Quadratic perturbations of a class of quadratic reversible Lotka–Volterra systems*, Internat. J. Bifur. Chaos. **23** (2013), 1350137.
- [207] S. SHI, *A concrete example of the existence of four limit cycles for plane quadratic systems*, Sci. Sinica **23** (1980), 153–158.
- [208] K.S. SIBIRSKI, *On conditions for the presence of a center and a focus*, (Russian), Kisinev. Gos. Univ. Uch. Zap. **11** (1954), 115–117.
- [209] J. SOTOMAYOR AND R. PATERLINI, *Quadratic vector fields with finitely many periodic orbits*, Lecture Notes in Math. **1007** (1983), 753–766.
- [210] G. SUO AND Y. CHEN, *The real quadratic systems with two conjugate imaginary straight line solutions*, Ann. Differential Equations **2** (1986), 197–207.

- [211] M. VOLDMAN, I.U. T. CALIN AND N. I. VULPE, *Affine invariant conditions for the topological distinction of quadratic systems with a critical point of the 4th multiplicity*, Publ. Mat. **40** (1996), 431–441.
- [212] A. VOLDMAN AND N. VULPE, *Affine invariant conditions for topologically distinguishing quadratic systems without finite critical points*, Izv. Akad. Nauk Respub. Moldova Mat. (1995), 100–112.
- [213] A. VOLDMAN AND N. VULPE, *Affine invariant conditions for topologically distinguishing quadratic systems with $m_f = 1$* , Nonlinear Anal. **31** (1998), 171–179.
- [214] N.I. VULPE, *Affine invariant conditions for the topological distinction of quadratic systems with a center*, (Russian) Differentsialnye Uravneniya **19** (1983), 371–379; translation in Differential equations **19** (1983), 273–280.
- [215] N. VULPE, *Characterization of the finite weak singularities of quadratic systems via invariant theory*, Nonlinear Anal. **74** (2011), 6553–6582.
- [216] N.I. VULPE AND A.YU. LIKHOVETSKI, *Coefficient conditions for the topological discrimination of quadratic systems of Darboux type*, (Russian) Mat. Issled., No. **106**, Differ. Uravneniya i Mat. Fizika, (1989), 34–49.
- [217] N. VULPE AND M. LUPAN, *Quadratic systems with dicritical points*, Izv. Akad. Nauk Respub. Moldova Mat. (1994), 52–60.
- [218] N.I. VULPE AND I.V. NIKOLAEV, *Topological classification of QS with a unique third order singular point*, Izv. Akad. Nauk Respub. Moldova Mat. (1992), 37–44.
- [219] N.I. VULPE AND I.V. NIKOLAEV, *Topological classification of a quadratic system with a four-fold singular point*, (Russian) Differentsialnye Uravneniya **29** (1993), 1669–1674; translation in Differential equations **29** (1993), 1449–1453.
- [220] N.I. VULPE AND K.S. SIBIRSCHI, *Affinely invariant coefficient conditions for the topological distinctness of quadratic systems*, Mat. Issled. **10** (1975), 15–28.
- [221] N.I. VULPE AND K.S. SIBIRSCHI, *Geometric classification of a quadratic differential system*, (Russian) Differentsialnye Uravneniya **13** (1977), 803–814; translation in Differential equations **13** (1977), 548–556.
- [222] A. WÖRZ-BUSERKROS, *A complete classification of all two-dimensional Lotka–Volterra systems*, Differential Equations Dynam. Systems **1** (1993), 101–118.
- [223] J. WU, L. PENG AND C. LI, *On the number of limit cycles in perturbations of quadratic reversible center*, J. Aust. Math. Soc. **92** (2012), 409–423.
- [224] A.I. YABLONSKII, *Limit cycles of a certain differential equations*, Differential Equations **2** (1966), 335–344 (In Russian).
- [225] S. YAKOVENKO, *Quantitative theory of ordinary differential equations and the tangential Hilbert 16th problem*, In: On Finiteness in Differential Equations and Diophantine Geometry. CRM Monogr. Ser., vol. **24**, 41–10, Amer. Math. Soc., Providence, 2005.
- [226] Y. YE, *Theory of limit cycles*, Transl. Math. Monogr., vol. **66**, Amer. Math. Soc., Providence, RI, 1986.
- [227] A. ZEGELING AND R. E. KOOLJ, *The distribution of limit cycles in quadratic systems with four finite singularities*, J. Differential Equations **151**, (1999), 373–385.
- [228] P. ZHANG, *On the distribution and number of limit cycles for quadratic system with two foci*, (Chinese) Acta Math. Sinica (Chin. Ser.) **44** (2001), 37–44.
- [229] P. ZHANG, *On the distribution and number of limit cycles for quadratic systems with two foci*, Qual. Theory Dyn. Syst. **3** (2002), 437–463.
- [230] X. ZHANG, *The 16th Hilbert problem on algebraic limit cycles*, J. Differential Equations **251** (2011), 1778–1789.
- [231] Y.K. ZHANG, C.P. LI AND B.Y. LI, *The number of zeros of Abelian integrals for a class of plane quadratic integrable systems with one center*, J. Systems Sci. Math. Sci. **32** (2012), 626–640.

- [232] Z. ZHANG, T. DING, W. HUANG AND Z. DONG, *Qualitative Theory of Differential Equations*, Science Publisher, 1985 (in Chinese); Transl. Math. Monogr., vol. **101**, Amer. Math. Soc., Providence, RI, 1992.
- [233] Y. ZHAO AND H. ZHU, *Bifurcations of limit cycles from a non-Hamiltonian quadratic integrable system with homoclinic loop*, Infinite dimensional dynamical systems, Fields Inst. Commun. **64**, Springer, New York, (2013), 445–479.
- [234] H. ŻOŁĄDEK, *Quadratic systems with center and their perturbations*, J. Differential Equations 109 (1994), 223–273.

¹ DEPARTMENT OF MATHEMATICS, SICHUAN UNIVERSITY, 610064 CHENGDU, SICHUAN, P.R. CHINA

Email address: t.l.scumath@gmail.com

² DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BELLATERRA, BARCELONA, CATALONIA, SPAIN

Email address: jllibre@mat.uab.cat