

Research Report

Neural signatures of predictive language processing in Parkinson's disease with and without mild cognitive impairment



Patricia León-Cabrera^a, Javier Pagonabarraga^{b,c,d,e,**}, Joaquín Morís^f,
 Saül Martínez-Horta^{b,c,d,e}, Juan Marín-Lahoz^{b,c,d,e},
 Andrea Horta-Barba^{b,c,d,e}, Helena Bejr-Kasem^{b,c,d,e},
 Jaime Kulisevsky^{b,c,d,e} and Antoni Rodríguez-Fornells^{a,g,*}

^a Cognition and Brain Plasticity Unit (CBPU), Department of Cognition, Development and Educational Psychology, Institute of Neuroscience, University of Barcelona and Bellvitge Biomedical Research Institute (IDIBELL), Barcelona, Spain

^b Movement Disorders Unit, Neurology Department, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain

^c Biomedical Research Institute (IIB-Sant Pau), Barcelona, Spain

^d Universitat Autònoma de Barcelona (UAB), Spain

^e Centro de Investigación en Red-Enfermedades Neurodegenerativas (CIBERNED), Spain

^f Department of Educational and Evolutive Psychology, University of Granada, Granada, Spain

^g Catalan Institution for Research and Advanced Studies (ICREA), Barcelona, Spain

ARTICLE INFO

Article history:

Received 8 December 2020

Reviewed 07 February 2021

Revised 4 March 2021

Accepted 31 March 2021

Action editor Michael Schwarze

Published online 30 April 2021

Keywords:

Semantic prediction

Semantic processing

Parkinson's disease

Mild cognitive impairment

ABSTRACT

Cognitive deficits are common in Parkinson's disease (PD), with some PD patients meeting criteria for mild cognitive impairment (MCI). An unaddressed question is whether linguistic prediction is preserved in PD. This ability is nowadays deemed crucial for achieving fast and efficient comprehension, and it may be negatively impacted by cognitive deterioration in PD. To fill this gap of knowledge, we used event-related potentials (ERPs) to evaluate mechanisms of linguistic prediction in a sample of PD patients (on dopamine compensation) with and without MCI. To this end, participants read sentence contexts that were predictive or not about a sentence-final word. The final word appeared after one sec, matching or mismatching the prediction. The introduction of the interval allowed to capture neural responses both before and after sentence-final words, reflecting semantic anticipation and semantic processing. PD patients with normal cognition (N = 58) showed ERP responses comparable to those of matched controls. Specifically, in predictive contexts, a slow negative potential developed prior to sentence-final words, reflecting semantic anticipation. Later, expected words elicited reduced N400 responses (compared to unexpected words), indicating facilitated semantic processing. PD patients with MCI (N = 20) showed, in addition, a prolongation of the N400 congruency effect (compared to matched PD patients without MCI), indicating that further cognitive decline impacts

* Corresponding author. Cognition and Brain Plasticity Unit Department of Cognition, Development and Educational Psychology Faculty of Psychology, University of Barcelona, L'Hospitalet de Llobregat, Barcelona 08097, Spain. Fax: +34 934024268.

** Corresponding author. Movement Disorders Unit, Neurology Department, Hospital de la Santa Creu i Sant Pau, Barcelona, 08041, Spain.

E-mail addresses: jpagonabarraga@santpau.cat (J. Pagonabarraga), Antoni.rodriiguez@icrea.cat (A. Rodríguez-Fornells).

<https://doi.org/10.1016/j.cortex.2021.03.032>

0010-9452/© 2021 Published by Elsevier Ltd.

semantic processing. Finally, lower verbal fluency scores correlated with prolonged N400 congruency effects and with reduced pre-word differences in all PD patients ($N = 78$). This relevantly points to a role of deficits in temporal-dependent mechanisms in PD, besides prototypical frontal dysfunction, in altered semantic anticipation and semantic processing during sentence comprehension.

© 2021 Published by Elsevier Ltd.

1. Introduction

Parkinson's disease (PD) is a chronic, neurodegenerative disorder that, in addition to motor defects, involves difficulties in a variety of cognitive domains, including executive functions, visuospatial skills, memory or language (Kudlicka, Clare, & Hindle, 2011; Muslimovic, Post, Speelman, & Schmand, 2005; Pell & Monetta, 2008). These deficits tend to progress in severity and meet criteria for mild cognitive impairment (PD-MCI) in up to 40% of patients within the first 5 years of disease (Aarsland et al., 2010; Litvan et al., 2011). The development of PD-MCI is a robust predictor of further conversion to dementia (PDD). However, the pattern of progression of PD-MCI and the rate of conversion to PDD is highly variable between patients. Among the several cognitive phenotypes characterizing PD-MCI, changes in visuo-perceptive skills, language production and comprehension, and semantic and episodic memory have been highlighted to better predict conversion to PDD than deficits mostly ascribed to frontal-executive dysfunction (Horta-Barba et al., 2020; Martinez-Horta & Kulisevsky, 2019; Lang et al., 2019). Yet, the mechanisms underlying deficits in language comprehension in PD and PD-MCI have been scarcely studied.

An aspect of language comprehension that has not been evaluated in PD is predictive language processing, which plays an important role in achieving fast and efficient language processing (Kutas & Federmeier, 2011). Accordingly, readers and listeners probabilistically infer and pre-activate different aspects of upcoming words (e.g., van Berkum, Brown, Zwitserlood, Kooijman & Hagoort, 2005; Wicha, Bates, Moreno, & Kutas, 2003). Much evidence of language prediction has been derived from the N400 event-related potential (ERP) component, an index of semantic processing (Kutas & Hillyard, 1980). The N400, a negativity peaking ~ 400 msec after word onset, is reduced for words that are more predictable in a given context (N400 context effect) (Kutas & Hillyard, 1984), which is considered to reflect facilitated semantic processing owing to the prediction of the word or some of its semantic features (Federmeier & Kutas, 1999). Most relevantly, recent research has shown that processing differences arise even before the word is presented. In particular, prediction negative potentials (PNP) – sustained, slow cortical potentials consistent with semantic anticipation – precede sentence-final words in predictive contexts, eliciting larger amplitudes before words that are strongly expected in a given context (e.g., before the word “ball” in “The goalkeeper managed to catch the... ball”) (León-Cabrera, Rodríguez-Fornells, & Morís, 2017; León-Cabrera, Flores, Rodríguez-Fornells, & Morís, 2019;

Grisoni, Miller, & Pulvermüller, 2017; for a review, see Pulvermüller & Grisoni, 2020).

Importantly, prior research suggests that linguistic prediction is reduced in populations with limited cognitive resources (Federmeier, McLennan, De Ochoa, & Kutas, 2002; Federmeier & Kutas, 2005; Federmeier, Kutas, & Schul, 2010; Payne & Federmeier, 2018; Wlotko & Federmeier, 2012; for reviews, see; Payne & Silcox, 2019; Huettig, 2015). For instance, older adults seem to take less advantage of semantically rich contexts to facilitate subsequent processing (Federmeier & Kutas, 2005) and those with lower verbal fluency scores exhibit diminished or absent ERP effects associated with prediction, suggesting the adoption of a ‘wait-and-see’, incremental processing strategy instead (Federmeier et al., 2002). As previously stated, PD involves significant impairments in cognitive capacities including executive functions and working memory (WM), which may secondarily impact language comprehension in these patients (Grossman et al., 2002). Thus, we hypothesized that predictive language processing could be impaired in PD, especially in those individuals with worse cognitive status, as in the case of those individuals with PD-MCI. In addition, there is behavioral evidence suggesting that PD patients have difficulties using lexical and sentential information to selectively activate appropriate meanings (Copland, Chenery & Murdoch, 2000a, 2000b, 2001; Copland, Seife, Ashley, Hudson, & Chenery, 2009). Also, several studies have pointed to slower or delayed lexical and semantic activation in PD (Angwin, Chenery, Copland, Murdoch, & Silburn, 2005; Arnott, Chenery, Murdoch, & Silburn, 2001), including research focused on N400 effects (Angwin, Dissanayaka, & Moorcroft et al., 2017; but see; Friederici, Kotz, Werheid, Hein, & von Cramon, 2003; Angwin, Dissanayaka, McMahon, Silburn, Copland, 2017). These deficits in lexical and semantic processing could also hinder the ability to appropriately use sentence contexts to guide and activate representations ahead of time during sentence comprehension.

To address this, the current study investigated ERP signatures of semantic prediction and semantic processing in PD with normal cognition (PD-NC) and in PD-MCI, as well as in a group of healthy control participants matched in terms of age, gender and education. We presented sentence contexts that were predictive or not of a sentence-final word that was delayed by one sec (see Table 2), thus allowing to capture the above-mentioned ERP signatures in the anticipatory and processing stages of prediction. In addition, in predictive contexts, half of the final words were semantically incongruent, thus representing a prediction mismatch.

Following previous studies in healthy adult population (León-Cabrera et al., 2017, 2019), we hypothesized that

controls would anticipate the final word after predictive contexts, as indexed by the development of a PNP in the anticipatory stage. In PD-NC and PD-MCI, potential failure to anticipate would be reflected in a comparatively reduced or absent PNP. In the processing stage, after the presentation of the final word, we expected controls to exhibit canonical N400 context effects, consistent with facilitated processing of congruent words as a result of prediction (Kutas & Federmeier, 2011). On the other hand, if PD patients adopt a more incremental strategy for comprehension, they might still exhibit N400 context effects because, even in the absence of prediction, congruent words are easier to integrate than incongruent words. However, compared to controls, the magnitude of the effect or its onset latency might be delayed, reflecting more difficult or slower semantic processing compared to controls, who would benefit from prediction instead. Lastly, following prior findings in older adults (Federmeier et al., 2002), we explored whether the targeted ERP signatures of linguistic prediction were reduced or diminished in PD patients with lower verbal fluency scores.

2. Materials and methods

No part of the study procedures or analyses was pre-registered prior to the research being conducted. In the following sections, we report how we determined our sample size, all data exclusions, all inclusion/exclusion criteria, whether inclusion/exclusion criteria were established prior to data analysis, all manipulations, and all measures in the study.

2.1. Participants

Participants were prospectively recruited from a sample of outpatients regularly attending the Movement Disorders Clinic at Hospital de la Santa Creu i Sant Pau (Barcelona, Spain). They were invited to participate in a longitudinal study involving comprehensive neuropsychological testing and two EEG recording sessions (one at baseline and one in a follow-up session after 1 year). Healthy adults were also invited to participate in the study to serve as controls. The current study focused on the data collected in the baseline session. The project was approved by the research ethics committee of the Hospital de la Santa Creu i Sant Pau. Before inclusion, written informed consent was obtained from all the participants.

The conditions of our ethics approval do not permit public archiving of anonymized study data. Readers seeking access to the data should contact the corresponding authors. Access can be granted only to qualified researchers under a collaboration agreement with the corresponding authors and in accordance with ethical procedures governing the reuse of clinical data, including the completion of a formal data sharing agreement and approval of the local ethics committee.

2.2. Clinical and cognitive testing

PD patients had been diagnosed by a neurologist with expertise in movement disorders. All patients accomplished steps 1 and 2 of the UKPDSBB criteria, and three or more of the four first supportive positive criteria of step 3 (Hughes, Daniel,

Kilford, & Lees, 1992). Motor status and stage of illness were assessed by the MDS-UPDRS-III and Hoehn & Yahr scales (Goetz et al., 2008). Demographic variables including age, gender, years of formal education, disease onset, medication history, as well as the levodopa equivalent daily dosage (LDD) (Tomlinson et al., 2010) were collected for all patients (Table 1). All participants were classified as having either normal cognition or MCI based on their score in the Parkinson's Disease - Cognitive Rating Scale (PD-CRS) (Pagonabarraga et al., 2008). A cut-off score of ≤ 83 was used to classify patients as PD-MCI (Fernández de Bobadilla et al., 2013). Participants with dementia were excluded according to the MDS diagnostic criteria for PD with dementia (Emre et al., 2007). In PD patients, cognition was examined during the 'on' state, and all participants were on stable doses of dopaminergic drugs during the 4 weeks before inclusion.

In addition, all participants completed standardized semantic and phonological verbal fluency tests (Peña-Casanova et al., 2009). In these tests, they were asked to either name as many animals (semantic verbal fluency) or words beginning with the letter P (phonological verbal fluency) as possible in 1 min. For each participant, total verbal fluency scores were obtained by averaging the direct scores in the semantic and phonological fluency tests (for another study using total verbal fluency scores, see Federmeier et al., 2002).

Legal copyright restrictions prevent public archiving of the neuropsychological tests used, which can be obtained from the copyright holders in the cited references.

2.3. Final samples

A total of 135 participants completed the baseline EEG recording session. Of those participants, 102 were PD patients and 33 were healthy controls. The data of some participants were excluded because the files were corrupt ($n = 3$) or their recordings were excessively noisy ($n = 4$) as determined by visual inspection of the raw data, leaving 128 participants available for analyses. From the remaining participants, 97 were PD patients and 31 were controls. Controls were excluded if they had mild cognitive impairment ($n = 3$) or their cognitive status was not available ($n = 2$). Lastly, participants with less than 20 available trials in any condition after EEG pre-processing ($n = 9$) were also excluded from the analyses (see EEG pre-processing section for more information about the artefact detection and rejection procedure).

The final pool of data consisted of 88 PD patients and 26 controls, from which matched samples were subsequently obtained to perform the following comparisons according to the goals of the present study: 1) PD patients with normal cognition (PD-NC) ($N = 58$) with an age-, gender- and education-matched Control group ($N = 24$), and 2) a subsample of Parkinson's disease patients with normal cognition (sPD-NC) ($N = 19$) with an age-, gender-, education- and years of disease evolution-matched PD patients with mild cognitive impairment (PD-MCI) ($N = 20$). To avoid the comparison of patients classified as sPD-NC and PD-MCI but having similar PD-CRS total score, we selected participants with PD-CRS scores of < 75 for the PD-MCI group and of > 85 for the sPD-NC.

The demographic and clinical features of the final samples are reported in Table 1. Controls and PD-NC patients did not

Table 1 – Demographics and clinical features of the patient and control samples.

	Control	PD-NC	P	sPD-NC	PD-MCI	P
N	24	58		19	20	
Age (years)	62.85 (6.82) [59.9, 65.7]	64.47 (7.61) [62.4, 66.4]	.37	70.75 (2.70) [69.4, 72]	72.22 (4.48) [70.1, 74.3]	.22
Education (years)	14.5 (4.04) [12.7, 16.2]	13.2 (4.57) [12, 14.4]	.23	12.42 (5.08) [9.9, 14.8]	10.4 (5.16) [7.9, 12.8]	.22
Women (%)	41.6	31	.35	42.1	45	.85
Disease duration (years)	n/a	5.26 (3.00) [4.4, 6]	–	5.68 (3.06) [4.2, 7.1]	6.28 (3.8) [4.4, 8]	.59
Levodopa dose (mg/d)	n/a	265.25 (249.58)	–	252.87 (293.74)	384.4 (215.04)	.12
LEDD (mg/d)	n/a	456.04 (230.68)	–	397.18 (236.08)	532.83 (209.70)	.09
PD-CRS total score	106.16 (8.74)	99.72 (10.86)	.01	98.2 (9.17)	67.15 (5.39)	<.001
[≤ 83 cutoff]	[102.4, 109.85]	[96.8, 102.5]		[93.7, 102.6]	[64.6, 69.6]	
Total verbal fluency score (DS)	19.66 (3.12)	18.01 (4.18)	.08	17.63 (4.32)	10.52 (2.08)	<.001

The values correspond to means, standard deviations (in round brackets) and 95% mean confidence intervals (CI) (in square brackets). For simplicity, CIs are shown only for the relevant variables in the group comparisons.
p-values were determined with independent samples *t*-tests between groups for continuous and ordinal variables and with Pearson Chi-Square for nominal variables (Gender).
 n/a = not applicable; LEDD = Total levodopa equivalent daily dosage; PD-CRS = Parkinson's Disease Cognitive Rating Scale; DS = Direct score.

differ in years of age ($t(1,80) = -.90, p = .37, 95\% \text{ CI } [-5.19, 1.95]$), years of education ($t(1,80) = 1.20, p = .23, 95\% \text{ CI } [-.84, 3.43]$), gender ($X^2(1, 82) = .853, p = .35$), or verbal fluency scores ($t(1,80) = 1.73, p = .08, 95\% \text{ CI } [-.23, 3.53]$). They differed in PD-CRS total score ($t(1,80) = -2.57, p = .01, 95\% \text{ CI } [-11.41, -1.46]$), but both groups were above the established score cut-off of 83 for PD-MCI diagnosis. Likewise, sPD-NC and PD-MCI patients did not differ in years of age ($t(1,37) = -1.23, p = .22, 95\% \text{ CI } [-3.89, .94]$), years of education ($t(1,37) = -1.30, p = .22, 95\% \text{ CI } [-1.30, 5.34]$), gender ($X^2(1, 39) = .033, p = .855$), disease duration ($t(1,37) = -.53, p = .59, 95\% \text{ CI } [-2.84, 1.65]$) or levodopa equivalent daily dose ($t(1,37) = -1.74, p = .09, 95\% \text{ CI } [-293.94, 22.64]$), but differed in PD-CRS total score ($t(1,37) = 12.96, p < .001, 95\% \text{ CI } [26.2, 35.91]$) and total verbal fluency scores ($t(1,37) = 6.59, p < .001, 95\% \text{ CI } [4.92, 9.29]$). For all comparisons, equal variances were assumed based on null results in Levene's tests.

2.4. Materials

We used the materials from León-Cabrera et al. (2019) consisting of high (HC) or low (LC) constraining sentence contexts, establishing a stronger or weaker expectation for an upcoming word (see Table 2 for sentence examples). A total of 312 sentences were included, of which 208 had HC sentence contexts (66.6%) and 104 had LC sentence contexts (33.3%). The sentences were originally created and categorized as either HC (mean cloze probability = 76%, $SD \pm 17.7\%$) or LC (mean cloze probability = 6.1%, $SD \pm 10.3\%$) by Mestres-Missé, Rodríguez-Fornells, & Münte, 2007. Within the HC condition, half of the

sentences ended with a congruent word and half with a semantically incongruent word. For the sake of task brevity to avoid fatigue, we decided to include the congruency contrast (congruent vs incongruent contrast) only in the HC condition. Sentences were further divided in two lists of 156 sentences, each list containing half of the sentences of each condition, and the use of one or another list was pseudo-randomized across participants. In sum, each participant read the following sentences: 52 HC with congruent endings (HC congruent), 52 HC with incongruent endings (HC incongruent), and 52 LC with congruent endings (LC congruent) (Table 2). All final words were nouns. Congruent ones were the best completions – the word with the highest cloze-probability (i.e., percentage of individuals that supply that word as a continuation for that sentence) (Taylor, 1953) – and incongruent words were selected from the ESPAL database (Duchon, Perea, Sebastián-Gallés, Martí, & Carreiras, 2013) so that they matched the congruent endings in the variables of mean word length, mean number of syllables, word frequency, familiarity, imaginability and concreteness. The stimuli and task presentation code can be downloaded at <https://osf.io/3FSP9/>.

2.5. Procedure

Participants were tested individually. After the external electrodes and the EEG cap were placed, electrode impedances were checked. Participants were instructed to keep their gaze fixated on the center of the screen and to blink only during the blinking interval. After completing a short training block (4 sentences), the proper experiment started. They were

Table 2 – Example sentences of each condition (original sentences in Spanish in brackets).

Condition	Sentence context	Final word
HC congruent	The goalkeeper managed to catch the (El portero fue capaz de atrapar la)	ball (pelota)
HC incongruent	The goalkeeper managed to catch the (El portero fue capaz de atrapar la)	shore (orilla)
LC congruent	As a present she gave her son a (Le ha regalado a su hijo una)	ball (pelota)

instructed to read the words that would appear at the center of the screen attentively and silently.

The experiment consisted of 26 blocks of 6 trials each (2 HC congruent, 2 HC incongruent, 2 LC congruent). The order of the sentences within each block was randomized. Words were white on black background (font type: Arial; font size: 24 points). On each trial, a fixation point (a cross) appeared at the center of the screen (800 msec). Then, a sentence was presented word-by-word (300 msec per word, followed by a 200 msec blank screen). Between the penultimate and the final word, the blank screen remained for a period of 1 sec (pre-word interval). After the final word (300 msec duration), a blank screen was presented (1000 msec) before the blinking signal (1 sec; depiction of an eye at the center of the screen). The next trial began after the offset of the blinking signal (see Fig. 1 for a simplified display of the trial structure).

To ensure that participants were reading attentively, they were asked to judge whether the previous sentence was congruent or not (“did the previous sentence make sense?”) every two blocks. The question remained on screen until they answered (‘yes’ or ‘no’) using the keyboard. The next block started immediately after the test except for even-numbered blocks, which were followed by a resting period that could be resumed anytime by the participant. All groups performed above chance in the congruency probe task (see Supplementary Materials for the accuracy of each group).

2.6. EEG recording

2.6.1. EEG acquisition

Continuous electroencephalogram (EEG; sampling rate = 250 Hz) was recorded from 19 electrodes (FP1/2, F3/4, F7/8, Fz, C3/4, Cz, P3/4, Pz, T3/4, T5/6, O1/2) mounted on a cap following the international 10–20 system positions. The EEG signal was amplified on-line (band-pass filter = .016–1000 Hz) with a Brain Vision amplifier. To monitor ocular artefacts, electrooculogram (EOG) electrodes were recorded with electrodes at bipolar vertical (supraorbital and suborbital ridge of the right eye) and horizontal placements (external ocular canthus of the left and right eyes). All electrode impedances were kept below 5 k Ω . An on-line notch filter (50 Hz) was applied to attenuate high-frequency electrical noise and data were referenced to the mean activity of the left and right mastoid electrodes.

2.6.2. EEG preprocessing

EEG data were preprocessed using ERPlab v7.0.0 (López-Calderon & Luck, 2014) of the EEGLab toolbox v14.0.0 (Delorme & Makeig, 2004), FieldTrip version 20181231 (Oostenweld, Fries,

Maris & Schoffelen, 2011), and custom scripts programmed in MATLAB R2017b. For each participant and condition, the continuous data were segmented into the following epochs of interest: 1) from –100 to 1300 msec from the onset of the penultimate word (pre-word interval), and 2) from –100 to 1000 msec from the target word onset (word interval). Of note, in a previous study with healthy adult population (León-Cabrera et al., 2019), we used a larger pre-word interval that also included the sentence context (from the onset of the sentence until the offset of the delay), which importantly revealed that differences accrued early and gradually during sentence processing. However, we discarded this possibility in the current study because the EEG signal in the sentence context intervals contained many artifacts, as preliminarily assessed through visual inspection. It was not unlikely that participants, especially PD patients and older adults, had greater difficulties in refraining from blinking or from moving for a relatively long interval (4.3 sec). We therefore opted to restrict the analysis of the anticipatory stage to the 1 sec delay interval following previous reports (León-Cabrera et al., 2017, 2019).

Artefact detection and rejection were applied on the epoched data. To facilitate artefact rejection, we computed the vertical (vEOG) and horizontal (hEOG) bipolar EOG channels by subtracting the inferior from the superior vertical EOG channels and the right from the left horizontal EOG channels, respectively. For the pre-word interval, all epochs with activity $\pm 85 \mu\text{V}$ in the ocular channels or $\pm 200 \mu\text{V}$ in any other channel were automatically removed and the remaining epochs were visually inspected and excluded if they contained blinks, muscle activity or large drifts. For the word interval, the concatenated epochs were first subjected to independent component analysis (ICA) decomposition to correct blinking artifacts because preliminary visual inspection revealed that many participants had difficulties refraining from blinking until the blinking signal, leading to a large loss of trials in this interval. ICA was run using the “runica” algorithm as implemented in EEGLab toolbox v14.0.0, excluding the ocular channels from the decomposition. The ocular components were detected and rejected based on visual inspection. After artifact-correction, epochs with activity $\pm 200 \mu\text{V}$ in any but ocular channels were automatically rejected, and those that remained were visually screened for any previously unnoticed artefacts (e.g., muscle activity or large drifts).

For the Control with PD-NC comparison, an average of 89.8% trials per condition were available for analysis (HC = 85.7 trials; LC = 41.9 trials; HC congruent = 49.6 trials; HC incongruent = 49.6 trials; LC congruent = 49.7 trials; overall average = 46 trials per condition), with no significant differences in the percentage of available trials between groups (all




Fig. 1 – Depiction of the structure of a trial. Sentences were presented one word a time. Each word of the sentence context (w1 to w7) was displayed for 300 msec followed by a 200 msec inter-word interval, except for the last word (w7), which was followed by a 1000 msec interval (pre-word interval) leading up to the presentation of the final word (w8).

p -values $> .3$). For the sPD-NC with PD-MCI comparison, an average of 87.4% trials per condition were included (HC = 79.2 trials; LC = 39.2 trials; HC congruent = 49.3 trials; HC incongruent = 49.5 trials; LC congruent = 49.6 trials; overall average = 45 trials per condition), with no significant differences in the percentage of remaining trials between groups (all p -values $> .1$).

2.7. Statistical analyses

2.7.1. Event-related potential analyses

Before performing statistical analyses, data were baseline-corrected to the 100 msec pre-stimulus period and low-pass filtered (as implemented in ERPLab v7.0.0). For the pre-word interval, a 5 Hz low-pass Butterworth filter (12 dB/oct roll-off) was applied to the data to focus exclusively on slow activity (Brunia, Hackley, van Boxtel, Kotani, & Ohgami, 2011). For the word interval, in which the N400 component was targeted, a 30 Hz low-pass Butterworth filter (12 dB/oct roll-off) was used. For the figures presented, the data were filtered using a 12 Hz low-pass Butterworth filter (12 dB/oct roll-off).

2.7.2. Cluster-based permutation analyses

Non-parametrical cluster-based permutation tests were used to assess the differences (Maris & Oostenveld, 2007) (as implemented in Fieldtrip version 20181231 under MATLAB R2017b), a method that exerts proper control for the increased probability of false positives in the context of many comparisons. Broadly, it identifies adjacent time points and channels with similar differences between conditions. The test worked as follows in the present study. For within-subject comparisons, every sample (channel \times time) was compared between two conditions (e.g., HC vs LC) through a dependent t -test. For between-subject comparisons, an independent t -test compared the effect (e.g., HC minus LC) between groups at each sample (channel \times time). Next, adjacent samples were clustered based on a t -value threshold (pre-determined from a one-tailed t -distribution with an alpha level of .05 and $N-1$ degrees of freedom) and the cluster with the largest sum of t -values was selected (cluster-level t -value). To determine whether effects were significant, the Monte Carlo method was used to construct a null distribution from the cluster-level t -values of random partitions obtained by randomly swapping the samples (between conditions and within participants for within-subject comparisons; and between conditions and between groups for between-subject comparisons) (5000 randomizations). Only observed clusters with cluster-level t -values within the 2.5th percentiles (alpha level of .05) of the null distribution were considered significant.

Choosing one-tailed testing is an optimal methodological decision when there are a priori hypotheses about the direction of the differences (Lakens, 2016). It allows to improve statistical sensitivity, which is particularly important when working with clinical samples. Following this, we chose one-tailed testing for the within-subject comparisons, as we had a priori hypotheses both for the pre- and post-word interval. In the pre-word interval, we expected more negative amplitudes for HC than LC (León-Cabrera et al., 2017, 2019). In the post-word interval, in which N400 context effects were targeted, we expected more negative amplitudes (larger N400 responses) for unexpected than for expected words, that is, for HC incongruent than for HC

congruent, as well as for LC congruent relative to HC congruent (Kutas & Federmeier, 2011). In turn, we chose two-tailed testing for the between-subject comparisons, as we did not have a priori hypotheses of the direction of the effects in this case.

2.7.3. Correlations between ERP effects and verbal fluency

After examining ERP effects, we conducted non-parametrical Spearman–Brown correlations to evaluate a potential linear relationship between the ERP effects and verbal fluency performance. All PD patients were included in this analysis ($N = 78$), that is, PD patients with normal cognition ($N = 58$) as well as PD patients with MCI ($N = 20$). For each participant, the total verbal fluency score was obtained by averaging the direct scores in the semantic and phonological fluency subtests (see *Supplementary Materials* for correlations with each subtest individually). ERP measures were labelled and quantified as follows: 1) PNP (LC minus HC mean amplitude difference in the 600 msec before pre-word onset at a left anterior cluster including electrodes FP1, F3, Fz, F7); 2) N400 congruency effect (HC congruent minus HC incongruent mean amplitude difference from 300 to 500 msec post-word onset); 3) N400 constraint effect (LC congruent minus HC congruent mean amplitude difference from 300 to 500 msec post-word onset); and, based on the ERP results, 4) prolonged N400 congruency effect (HC congruent minus HC incongruent mean amplitude difference from 600 to 800 msec post-word onset). All effects tied to the N400 were tested on a centrally-distributed cluster (averaged electrodes: Cz, C3, C4, Pz, P3, P4). All values were normalized before performing correlations. All p -values were corrected with the Holm-Bonferroni correction (Holm, 1979).

3. Results

3.1. Event-related potentials

3.1.1. Controls versus PD patients with normal cognition

Fig. 2A and B shows the grand-average ERPs at the F7 electrode during the pre-word interval in the Control group and the group of PD patients with normal cognition, respectively. The ERPs were computed in the interval from the onset of the penultimate word to the onset of the final word (1300 msec duration, the first 300 msec correspond to the penultimate word, and the last 1000 msec to the delay interval), with a 100 msec pre-stimulus baseline.

Both groups exhibited similar patterns, namely, an N100 associated with the processing of the penultimate word, followed by a positivity, and the later development of a negativity that became progressively larger as the presentation of the final word approached. As expected, based on previous findings in healthy adult population (León-Cabrera et al., 2019), the negativity was more prominent in the HC condition (relative to the LC condition), that is, when the final word could be strongly predicted from the prior context. This difference was confirmed by a significant cluster in the Control group ($p = .01$) (Fig. 2A) as well as in PD individuals with normal cognition ($p = .01$) (Fig. 2B). The cluster showed a similar temporal profile in both groups. It started approximately 800 msec before the final word and continued until the word appeared. In regard to




Fig. 2 – ERP results for the Control and the PD-NC group. **A and B)** Grand-averaged ERPs at the F7 electrode during the preword interval for both conditions for **(A)** Controls and **(B)** PD-NC patients. ERPs are time-locked to the presentation of the penultimate word (100 pre-stimulus baseline). The horizontal bar highlights (grey) the time interval in which significant differences were obtained. Below the ERPs, topographical maps indicating the electrodes (white asterisks) that showed a significant effect of constraint (HC minus LC) in the 600 msec prior to the critical word (in steps of 200 msec). **C and D)** Grand-averaged ERPs at Cz electrode after the presentation of the sentence-final word (100 pre-stimulus baseline) for **(C)** Controls and **(D)** PD-NC patients. The horizontal bars highlight the time interval of the N400 effects of Congruency (green) and Constraint (purple). Topographical maps with electrodes (white asterisks) that showed effects in the time interval in which N400 effects were expected to be maximal (300–500 msec post-word onset).

the spatial distribution, the cluster encompassed mainly frontal electrodes, with a slight left-lateralization. There were no significant differences between groups.

We then turned to the interval after word presentation. As can be observed in Fig. 2C and D, for both groups, semantically unexpected words (HC incongruent and LC congruent) elicited a larger negativity peaking about 400–600 msec (relative to expected words; HC congruent) that is consistent with the canonical features of the N400. These differences were reflected in the significant clusters described below.

The effect of constraint (HC congruent vs LC congruent) was captured by significant clusters in the Control group ($p = .001$) (Fig. 2C) and in the PD group ($p < .001$) (Fig. 2D), whereby words presented in less predictive contexts (LC congruent) showed more negative amplitudes than words in strongly predictive contexts (HC congruent). There were no differences between groups. The cluster started about 250–300 msec after word onset and resolved at about 700 msec, and it exhibited a widespread distribution over the scalp. In fact, the cluster was significant at all sites. On the

other hand, the effect of congruency (HC congruent vs HC incongruent) was reflected in a shorter-lived cluster in the Control group ($p = .007$) (Fig. 2C) and in the PD group ($p < .001$) (Fig. 2D), in this case confirming more negative amplitudes for incongruent than congruent words in strongly predictive contexts. Again, there were no significant differences between groups. The effect was widespread with a seemingly slight rightward focus in magnitude.

3.1.2. Subset of PD patients with normal cognition versus PD with MCI

The mean voltage pattern elicited during the pre-word interval (Fig. 3A and Fig. 3B) exhibited the normative N100 component, followed by a transient positivity and a subsequent negativity in the last part of the interval that evidenced the hypothesized tendency towards more negative amplitudes for HC than LC. However, in this case, there were no statistically significant




Fig. 3 – ERP results for the subgroup of PD patients with normal cognition (PD-NC) and the group of PD-MCI patients. **A and B)** Grand-averaged ERPs at the F7 electrode during the pre-word interval for (A) sPD-NC and (B) PD-MCI patients. ERPs are time-locked to the presentation of the penultimate word (100 pre-stimulus baseline). The grey horizontal bar highlights the time interval in which significant differences were obtained. Below the ERPs, the topographical maps indicate the electrodes (white asterisks) that showed a significant effect of constraint (HC minus LC) in the 600 msec prior to the critical word (in steps of 200 msec). **C and D)** Grand-averaged ERPs at the Cz electrode after the presentation of the sentence-final word (100 pre-stimulus baseline) for (C) sPD-NC and (D) PD-MCI patients. The horizontal bars highlight the time interval of the N400 effect of Congruency (green) and Constraint (purple). The topographical maps show which electrodes (white asterisks) showed effects in the time interval in which N400 effects were expected to be maximal (300–500 msec post-word onset).

differences between conditions, nor any differences between the two groups.

After the presentation of the final word (Fig. 3C and D) there was an observable larger negativity for unexpected (HC incongruent and LC congruent) relative to expected words (HC congruent) in both groups, in line with classical N400 context effects. These observable differences were confirmed by significant clusters as specified next.

There were significant effects of constraint (HC congruent vs LC congruent) both for the group of PD patients without MCI ($p < .001$) (Fig. 3C) and the group of PD patients with MCI ($p < .001$) (Fig. 3D). That is, as expected, words in less

predictive contexts elicited a larger N400 than words in strongly predictive contexts, a difference that encompassed the interval between about 300 and 700 msec post word onset. The scalp distribution of the effect covered a wide set of electrodes over frontal and central regions of the scalp, with a slight frontward amplitude maximum. No differences between groups were found.

As for the effect of congruency in predictive contexts (HC congruent vs HC incongruent), again as expected, significant clusters reflected that incongruent words had more negative amplitudes than congruent words in patients without MCI ($p = .005$) (Fig. 3C), and also in PD patients with MCI ($p = .001$)




Fig. 4 – A) Topographical maps showing the temporal evolution and scalp distribution of the congruency contrast (HC incongruent minus HC congruent) during final word processing, showing a prolonged N400 congruency effect in the group of individuals with PD and MCI. B) Grand-averaged ERPs of the difference waveforms (HC incongruent and HC congruent) of both groups, averaged over the set of electrodes that were part of the significant cluster of differences between the groups (highlighted in white in the topographical map). The grey-colored area shows the timepoints included in the cluster. The topographical map shows the scalp distribution of the averaged t-values within the cluster (cluster t-value \pm 2.02 for an alpha level of .05).

(Fig. 3D). Relevantly, in this case, there were significant differences between the groups (Fig. 4). The difference waveforms (HC incongruent minus HC congruent) of each group were contrasted in a between-group cluster-based permutation test including all electrodes and time points (from 0 to 1000 msec). The test yielded a significant cluster ($p = .008$) that revealed a longer-lasting effect of congruency for the group of patients with MCI, which can be visualized in Fig. 4A. More specifically, the cluster spanned approximately from 630 to 760 msec after word onset at centro-parietal sites (significant electrodes: C3, Cz, C4, P3, Pz, P4, T5, T4, T6, O2) (see the scalp distribution of t -values in Fig. 4B). This result suggests a prolonged N400 congruency effect in the group of PD patients with MCI relative to the group of PD patients without MCI.

3.2. Correlations between ERP effects and verbal fluency scores

After establishing the ERP effects, we examined the association between ERP measures and verbal fluency scores in all PD patients ($N = 78$) (for further specification of this analysis, see Section 2.7.3.) by means of non-parametric Spearman correlations between direct scores in verbal fluency tests, and four ERP measures, namely, 1) PNP (LC minus HC), 2) N400 constraint effect (LC congruent minus HC congruent), 3) N400 congruency effect (HC congruent minus HC incongruent), and 4) prolonged N400 congruency effect (HC congruent minus HC incongruent). Note that, in these analyses, the direction of the condition subtraction is reversed, such that positive values represent differences in the expected direction of the effect (e.g., in the case of the PNP, positive values indicate larger amplitudes for HC than for LC). All correlations were corrected using Holm-Bonferroni correction (indicated as p_{HB} in the text).

The results of the correlational analyses are presented in Table 3. After correcting for multiple comparisons, two of the ERP measures showed a significant correlation with verbal fluency scores: the PNP and the prolonged N400 congruency effect (Fig. 5). In particular, the PNP showed a significant positive correlation with verbal fluency ($r(77) = .326$, $p_{HB} = .010$). Note that we excluded one participant who had outlying scores in the PNP (z -score = -3.91), although the correlation was unaffected by its inclusion ($r(78) = .326$, $p_{HB} = .01$), which is normal when using non-parametric Spearman correlations which are very

Table 3 – Results of correlational analyses between ERP measures and verbal fluency scores in PD patients.

Test	ERP effect	r	p -value (corrected)
Verbal fluency	PNP	.326	.010*
	N400 constraint	.157	.336
	N400 congruency	-.073	.525
	Prolonged N400 congruency	-.358	.005*
PNP effect (LC minus LC); N400 constraint effect (LC congruent minus HC congruent); N400 congruency effect (HC congruent minus HC incongruent); Prolonged N400 congruency effect (HC congruent minus HC incongruent).			
Holm-Bonferroni correction was applied to adjust p -values. Statistically significant correlations are highlighted with asterisks.			

robust to the presence of outliers. This indicates that a larger PNP (more negative amplitudes prior to words in predictive than in unpredictable contexts) is associated with better verbal fluency performance in patients with PD. In turn, the prolongation of the N400 congruency effect (more negative amplitudes for incongruent than for congruent words 600–800 msec after word onset) correlates with worse mean verbal fluency scores ($r(78) = -.358$, $p_{HB} = .005$). No other correlations were statistically significant.

4. Discussion

Motivated by previous evidence of altered predictive language processing in populations with low cognitive resources (Federmeier et al., 2002, 2005, 2010), the present study assessed linguistic prediction in PD with normal cognition and PD with MCI. To this end, ERP modulations were examined before and after encountering words that could or could not be predicted from context, thus capturing both the postulated anticipatory and processing stage of prediction. Specifically, we focused on PNPs as signatures of semantic anticipation, and post-word N400 context effects as indices of processing facilitation by pre-activation. Relative to controls, PD patients with normal cognitive status exhibited the expected ERP signatures, pointing to preserved predictive language processing. In turn, PD patients with MCI showed absent PNPs and a significant prolongation of N400 congruency effects in predictive contexts, suggesting altered mechanisms tied to linguistic prediction. Interestingly, correlational analyses revealed that worse verbal fluency performance was associated with the presence of the ERP pattern observed in the group of PD patients with MCI.

4.1. Neural correlates of predictive sentence processing are preserved in PD-NC

Consistent with prediction, PD patients with normal cognition exhibited the canonical reduction of N400 amplitude for expected (HC congruent) compared to unexpected words (LC congruent and HC incongruent) (Kutas & Hillyard, 1984). This effect is taken to reflect facilitated lexical and semantic activation of expected words as a result of their pre-activation (Federmeier & Kutas, 1999; Lau, Phillips, & Poeppel, 2008). In line with this interpretation, in predictive contexts (HC), PD patients with normal cognition exhibited a frontally-distributed negative PNP that developed progressively prior to the presentation of the final word (León-Cabrera et al., 2019; for similar results, see, Grisoni, Miller & Pullvermüller, 2017; Li, Zhang, Xia, & Swaab, 2017). Studies in healthy population suggest that such PNPs might capture anticipation of semantic aspects of the upcoming word (for a recent review, see Pulvermüller & Grisoni, 2020). More specifically, they might reflect retrieval or maintenance of pre-activated lexical and/or semantic representations (León-Cabrera et al., 2019; Li et al., 2017). Altogether, PD patients with normal cognition seem to make proper use of sentence contexts to predict upcoming linguistic information during comprehension.

More broadly, the present findings are in line with previous studies showing intact N400 effects in sentence processing in




Fig. 5 – Scatterplots and scalp maps of the significant correlations between ERP measures and verbal fluency scores in all PD patients (PD with and without MCI) ($N = 78$). **A)** Scatterplot of the positive correlation ($r = .326$, $p_{HB} = .010$) between mean verbal fluency scores and the PNP (more negative amplitudes prior to words in HC than in LC contexts) and **B)** of the negative correlation ($r = -.358$, $p_{HB} = .005$) between verbal fluency scores and the prolongation of the N400 congruency effect (more negative amplitudes for incongruent than for congruent words 600–800 msec post-word). All values were normalized. Below the scatter diagrams, the r -values (Spearman rho) for all 19 electrodes are plotted on an idealized head. Darker shading indicates larger r -values in the direction of the correlation. Electrodes included in the ROI used to perform the reported correlations with verbal fluency (shown in the scatter diagrams) are highlighted in white. R -values between electrodes were estimated by spherical spline interpolation.

PD patients with normal cognition (Angwin, Dissanayaka, & McMahon, et al., 2017; Friederici et al., 2003), and, in turn, contrasts with behavioral and electrophysiological evidence pointing to altered lexical and semantic processing in these patients (Angwin, Chenery, Copland & Arnott et al., 2004, Angwin, Chenery, Copland & Murdoch et al., 2004, 2006, 2009, Angwin, Dissanayaka, & Moorcroft et al., 2017; Arnott et al., 2001, 2010; Copland et al., 2009; Kutas et al., 2013). Of note, most studies showing abnormal lexical and semantic processing employed tasks that consisted of single-word contexts, instead of sentence contexts. Importantly, sentence processing involves the construction of a high-level meaning representation (Graesser, Singer, & Trabasso, 1994), which serves to constrain in a top-down manner towards the appropriate lexical and semantic representation, perhaps compensating for activation problems within semantic networks in these individuals. Nonetheless, note that all patients in the current study were on dopaminergic medication. PD patients on dopaminergic medication perform better in semantic priming tasks than PD patients off medication (with normal cognition) (Angwin, Chenery, Copland & Arnott et al., 2004, 2009), suggesting that dopamine (DA) mediates semantic activation, arguably through the calibration of the spread and focus of activation within semantic networks (Kischka et al., 1996). DA

has also been linked with anticipatory processes in PD. Specifically, PD patients with normal cognition off medication show reduced amplitudes of the *stimulus-preceding negativity*, suggesting that the DA system is implicated in the anticipation of motivationally salient and rewarding stimuli (Mattox Valle-Inclán & Hackley, 2006). Future studies testing patients off medication would help untangle to what extent dopaminergic compensation contributed to the preservation of otherwise altered mechanisms.

4.2. PD-MCI impacts late semantic processing in predictive contexts

To further investigate the impact of cognitive impairment, we also examined neural responses in a smaller sample of PD-MCI patients. Compared to a matched subgroup of PD patients with normal cognition, PD patients with MCI also exhibited expected N400 amplitude modulations within the normal onset latency (i.e., about 300–400 msec post-word onset). However, most remarkably, PD patients with MCI showed a significantly prolonged N400 congruency effect in predictive contexts, extending up to 700 msec after word onset. Critically, this was not observed in the N400 constraint contrast (LC congruent vs HC congruent), which suggests that the prolongation

stemmed from the ERP response to incongruent words, rather than that to congruent words (i.e., from HC incongruent, instead of HC congruent). With that in mind, it is important to note that N400 context effects do not solely capture prediction effects, but also integration demands (Van Berkum, Hagoort, & Brown, 1999). Interestingly, a recent study showed that prediction effects dominate the earlier portion of the N400 context effect (starting as early as 200 post-word onset), whereas integration effects began later and continued until about 650 msec after word onset (Nieuwland et al., 2020; see also, Brothers, Swaab, & Traxler, 2015; Lau, Namyst, Fogel, & Delgado, 2016). As such, prolonged N400 congruency effects may reflect abnormally sustained efforts to integrate the incongruent word with the previous context (Nieuwland et al., 2020). Curiously, a previous study found that patients with bilateral basal ganglia lesions exhibited similarly prolonged N400 effects in sentence processing (up to 700 msec post-word) (Kotz, Frisch, Von Cramon, & Friederici, 2003). In PD, the worsening of the condition from PD with normal cognition to PD with MCI may lead to altered late integrational semantic processing as well.

More recently, N400 effects have postulated to reflect the ‘prediction error’ (albeit with nuances in their definition, see Willems, Frank, Nijhof, Hagoort, & Van den Bosch, 2016; Rabovsky, Hansen, & McClelland, 2018; Kuperberg, Brothers, & Wlotko, 2020). In predictive processing frameworks (Clark, 2013), predictions are sent top-down from higher to lower level regions to ‘explain away’ the incoming sensory input and, then, the ‘prediction error’ – the portion of information that remains unexplained – is sent back and used as a learning signal to update internal models and improve future predictions. Thus, prolonged N400 congruency effects may indicate difficulties in this ‘learning’ phase, leading to less efficient internal model updating.

Furthermore, also in line with less efficient predictive language processing in PD patients with MCI, there were no effects in the anticipatory stage in this group, pointing to absent semantic anticipation. Note, however, that there were no effects in the matched subgroup of PD patients with normal cognition either. In fact, there were no differences between the two groups. Therefore, the absence or extreme reduction of the PNP in these two groups might be associated with factors other than MCI, such as the older age of the patients in these samples (compared to the whole sample of PD with normal cognition), or insufficient statistical power to detect differences, as sample size was smaller in this case.

4.3. *Worse verbal fluency performance is associated with altered signatures of predictive language processing in PD*

Interestingly, correlational analyses revealed that lower verbal fluency scores were associated with reduced PNP and prolonged N400 congruency effects. Similarly, previous evidence has shown that verbal fluency scores are a good predictor of the status of predictive language processing in healthy older population (Federmeier et al., 2002). Worse performance in verbal fluency is generally associated with executive dysfunction, such as difficulties in rule switching or inhibition. Executive

dysfunction is common with the progression of fronto-striatal deterioration in PD with and without MCI (Aarsland et al., 2010; Kudlicka et al., 2011; Monchi et al., 2004) and has been proposed to indirectly hinder language processing in PD (Grossman et al., 2002, 2003). However, verbal fluency performance depends not only on frontal lobe function, but also on lexical and semantic retrieval processes that are dependent on temporo-parietal structures (Unsworth, Spillers, & Brewer, 2011). In fact, recent findings point to a greater weight of language-related processes than executive function in verbal fluency (Whiteside et al., 2016). Recent meta-analyses and lesion-based studies have also shown that verbal fluency is supported by standard language networks and underlying white-matter connectivity (Baldo, Schwartz, Wilkins, & Dronkers, 2006; Costafreda et al., 2006; Griffis, Nenert, Allendorfer, & Szaflarski, 2017). Therefore, the ERP pattern associated with lower verbal fluency – suggestive of less efficient predictive processing mechanisms, as previously discussed – may reflect not only executive dysfunction, but also disruption in semantic networks, in line with recent findings of difficulties in temporal-dependent functions in PD with and without MCI (Horta-Barba et al., 2020; Martinez-Horta & Kulisevsky, 2019; Lang et al., 2019).

4.4. *Concluding remarks and future directions*

Overall, the current results suggest preserved predictive language processing in PD patients with normal cognitive status. In turn, in PD patients with MCI, greater cognitive limitations might hinder mechanisms associated with semantic prediction in normal circumstances. While these limitations may not prevent the pre-activation of relevant representations, as indexed by canonical N400 amplitude modulations, they might negatively affect sentence processing due to poorer semantic anticipation and less efficient semantic processing when contexts are predictive. Furthermore, both executive dysfunction and damage within semantic networks may underlie these difficulties, as suggested by their association with low verbal fluency scores. Along with recent research (Horta-Barba et al., 2020; Martinez-Horta & Kulisevsky, 2019; Lang et al., 2019), these findings emphasize the value of examining cognitive changes in PD patients with and without MCI in other domains beyond executive functions, like the language domain. Interestingly, altered N400 effects are associated with a higher risk of transiting to dementia in individuals with amnesic MCI (Olichney et al., 2008). Therefore, future longitudinal studies in PD could evaluate changes in N400 congruency effects as potential markers of the transition to PD with MCI and to PD with dementia (for interested readers, see Supplementary Materials for a hierarchical regression analysis further attesting that the N400 congruency effect is a good predictor of global cognitive decline in PD). Finally, as previously highlighted, future studies testing patients off medication could assess the exact role of DA in predictive sentence comprehension. Yet, the clinical value of evaluating PD patients on stable doses of dopaminergic medication must be emphasized, as it is the most representative of their habitual cognitive status and thus has the potential to uncover limitations that may impact the everyday functioning of individuals with PD.

Funding

This study was funded by grants from the Fundació la Marató de TV3 2014/U/477 and 20142910, and by financial support from the Center for Biomedical Research and Neurodegenerative Resources (CIBERNED). PLC was funded with a pre-doctoral grant FPU15/05554 (FPU "Ayudas para la Formación de Profesorado Universitario") of the Spanish Ministry of Education, Culture and Sport. JP was funded by PERIS (expedient number: SLT008/18/00088) of Generalitat de Catalunya. JK and HB-K were funded by FIS PI18/01717. HB -K was funded by Río Hortega CM17/00209 of Instituto de Salud Carlos III (ISCIII) (Spain). JM was funded by Río Hortega CM15/00071 of Instituto de Salud Carlos III (ISCIII) (Spain). None of these funding sources had any involvement in the conduct of the research or the preparation of the article.

CRediT authorship contribution statement

PLC: Conceptualization, Formal analysis, Visualization, Writing - Original Draft, and Writing - Review & Editing; JP: Conceptualization, Supervision, Methodology, Writing - Review & Editing, Resources, and Funding acquisition; JM: Conceptualization, Supervision, Methodology and Writing - Review & Editing; SMH: Investigation and Writing - Review & Editing; JML: Investigation and Writing - Review & Editing; AHB: Investigation; HBK: Investigation; JK: Resources, Writing - Review & Editing, and Funding acquisition; ARF: Conceptualization, Supervision, Methodology, and Writing - Review & Editing.

Declaration of competing interests

The authors declare no disclosure of financial interests and potential conflict of interest.

Acknowledgments

In honor and loving memory of Dr. Jordi Riba, who passed away in the process of preparing this manuscript. We would like to thank Dr. Ion Iarritu, Dr. Laura Ferreri, Miriam Cornellà and Fernando Martínez for their help with data collection and technical support.

Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.cortex.2021.03.032>.

REFERENCES

Aarsland, D., Bronnick, K., Williams-Gray, C., Weintraub, D., Marder, K., Kulisevsky, J., et al. (2010). Mild cognitive

- impairment in Parkinson disease: A multicenter pooled analysis. *Neurology*, 75, 1062–1069. <https://doi.org/10.1212/WNL.0b013e3181f39d0e>
- Angwin, A. J., Arnott, W. L., Copland, D. A., Haire, M. P. L., Murdoch, B. E., Silburn, P. A., et al. (2009). Semantic activation in Parkinson's disease patients on and off levodopa. *Cortex*, 45, 950–959. <https://doi.org/10.1016/j.cortex.2009.02.012>
- Angwin, A. J., Chenery, H. J., Copland, D. A., Arnott, W. L., Grattan, R., Murdoch, B. E., et al. (2006). Priming of semantic features in Parkinson's disease. *Brain and Language*, 99(1–2), 102–103. <https://doi.org/10.1016/j.bandl.2006.06.060>
- Angwin, A. J., Chenery, H. J., Copland, D. A., Arnott, W. L., Murdoch, B. E., & Silburn, P. A. (2004). Dopamine and semantic activation: An investigation of masked direct and indirect priming. *Journal of the International Neuropsychological Society: JINS*, 10(1), 15–25. <https://doi.org/10.1017/S1355617704101033>
- Angwin, A. J., Chenery, H. J., Copland, D. A., Murdoch, B. E., & Silburn, P. A. (2004). The time course of semantic activation in Parkinson's disease. *Brain and Language*, 91, 145–146. <https://doi.org/10.1016/j.bandl.2004.06.076>
- Angwin, A. J., Chenery, H. J., Copland, D. A., Murdoch, B. E., & Silburn, P. A. (2005). Summation of semantic priming and complex sentence comprehension in Parkinson's disease. *Cognitive Brain Research*, 25(1), 78–89. <https://doi.org/10.1016/j.cogbrainres.2005.04.008>
- Angwin, A. J., Dissanayaka, N. N., McMahon, K. L., Silburn, P. A., & Copland, D. A. (2017). Lexical ambiguity resolution during sentence processing in Parkinson's disease: An event-related potential study. *Plos One*, 12(5), Article e0176281. <https://doi.org/10.1371/journal.pone.0176281>
- Angwin, A. J., Dissanayaka, N. N., Moorcroft, A., McMahon, K. L., Silburn, P. A., & Copland, D. A. (2017). A neurophysiological study of semantic processing in Parkinson's disease. *Journal of the International Neuropsychological Society: JINS*, 23(1), 78–89. <https://doi.org/10.1017/S1355617716000953>
- Arnott, W. L., Chenery, H. J., Angwin, A. J., Murdoch, B. E., Silburn, P. A., & Copland, D. A. (2010). Decreased semantic competitive inhibition in Parkinson's disease: Evidence from an investigation of word search performance. *International journal of speech-language pathology*, 12(5), 437–445. <https://doi.org/10.3109/17549507.2010.492875>
- Arnott, W. L., Chenery, H. J., Murdoch, B. E., & Silburn, P. A. (2001). Semantic priming in Parkinson's disease: Evidence for delayed spreading activation. *Journal of Clinical and Experimental Neuropsychology*, 23(4), 502–519. <https://doi.org/10.1076/jcen.23.4.502.1224>
- Baldo, J. V., Schwartz, S., Wilkins, D., & Dronkers, N. F. (2006). Role of frontal versus temporal cortex in verbal fluency as revealed by voxel-based lesion symptom mapping. *Journal of the International Neuropsychological Society: JINS*, 12(6), 896–900. <https://doi.org/10.1017/S1355617706061078>
- Brothers, T., Swaab, T. Y., & Traxler, M. J. (2015). Effects of prediction and contextual support on lexical processing: Prediction takes precedence. *Cognition*, 136, 135–149. <https://doi.org/10.1016/j.cognition.2014.10.017>
- Brunia, C. H., Hackley, S. A., van Boxtel, G. J., Kotani, Y., & Ohgami, Y. (2011). Waiting to perceive: Reward or punishment? *Clinical Neurophysiology*, 122(5), 858–868. <https://doi.org/10.1016/j.clinph.2010.12.039>
- Clark, A. (2013). Whatever next? Predictive brains, situated agents, and the future of cognitive science. *The Behavioral and Brain Sciences*, 36(3), 181–204. <https://doi.org/10.1017/S0140525X12000477>
- Copland, D. A., Chenery, H. J., & Murdoch, B. E. (2000a). Impaired meaning selection for polysemous words in individuals with nonthalamic subcortical lesions and Parkinson's disease. *Brain and Language*, 74(3), 360–363.

- Copland, D. A., Chenery, H. J., & Murdoch, B. E. (2000b). Understanding ambiguous words in biased sentences: Evidence of transient contextual effects in individuals with nonthalamic subcortical lesions and Parkinson's disease. *Cortex*, 36(5), 601–622. [https://doi.org/10.1016/S0010-9452\(08\)70541-0](https://doi.org/10.1016/S0010-9452(08)70541-0)
- Copland, D. A., Chenery, H. J., & Murdoch, B. E. (2001). Discourse priming of homophones in individuals with dominant nonthalamic subcortical lesions, cortical lesions and Parkinson's disease. *Journal of Clinical and Experimental Neuropsychology*, 23(4), 538–556. <https://doi.org/10.1076/jcen.23.4.538.1233>
- Copland, D. A., Sefe, G., Ashley, J., Hudson, C., & Chenery, H. J. (2009). Impaired semantic inhibition during lexical ambiguity repetition in Parkinson's disease. *Cortex*, 45(8), 943–949. <https://doi.org/10.1016/j.cortex.2009.02.023>
- Costafreda, S. G., Fu, C. H., Lee, L., Everitt, B., Brammer, M. J., & David, A. S. (2006). A systematic review and quantitative appraisal of fMRI studies of verbal fluency: Role of the left inferior frontal gyrus. *Human Brain Mapping*, 27, 799–810. <https://doi.org/10.1002/hbm.20221>
- Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of neuroscience methods*, 134(1), 9–21. <https://doi.org/10.1016/j.jneumeth.2003.10.009>
- Duchon, A., Perea, M., Sebastián-Gallés, N., Martí, A., & Carreiras, M. (2013). Espal: One-stop shopping for Spanish word properties. *Behavior Research Methods*, 45(4), 1246–1258. <https://doi.org/10.3758/s13428-013-0326-1>
- Emre, M., Aarsland, D., Brown, R., Burn, D. J., Duyckaerts, C., & Mizuno, Y. (2007). Clinical diagnostic criteria for dementia associated with Parkinson's disease. *Movement Disorders*, 22, 1689–1707. <https://doi.org/10.1002/mds.21507>, 2007.
- Federmeier, K. D., & Kutas, M. (1999). A rose by any other name: Long-term memory structure and sentence processing. *Journal of Memory and Language*, 41(4), 469–495. <https://doi.org/10.1006/jmla.1999.2660>
- Federmeier, K. D., & Kutas, M. (2005). Aging in context: Age-related changes in context use during language comprehension. *Psychophysiology*, 42(2), 133–141. <https://doi.org/10.1111/j.1469-8986.2005.00274.x>
- Federmeier, K. D., Kutas, M., & Schul, R. (2010). Age-related and individual differences in the use of prediction during language comprehension. *Brain and Language*, 115(3), 149–161. <https://doi.org/10.1016/j.bandl.2010.07.006>
- Federmeier, K. D., McLennan, D. B., De Ochoa, E., & Kutas, M. (2002). The impact of semantic memory organization and sentence context information on spoken language processing by younger and older adults: An ERP study. *Psychophysiology*, 39(2), 133–146. <https://doi.org/10.1111/1469-8986.3920133>
- Fernández de Bobadilla, R., Pagonabarraga, J., Martínez-Horta, S., Pascual-Sedano, B., Campolongo, A., & Kulisevsky, J. (2013). Parkinson's disease-cognitive rating scale: Psychometrics for mild cognitive impairment. *Movement Disorders*, 28(10), 1376–1383. <https://doi.org/10.1002/mds.25568>
- Friederici, A. D., Kotz, S. A., Werheid, K., Hein, G., & von Cramon, D. Y. (2003). Syntactic processes in Parkinson's disease: A dissociation between early automatic and late integrational processes. *Neuropsychology*, 17(1), 133–142. <https://doi.org/10.1037/0894-4105.17.1.133>
- Goetz, C. G., Tilley, B. C., Shaftman, S. R., Stebbins, G. T., Fahn, S., Martinez-Martin, P., et al. (2008). Movement Disorder Society-Sponsored Revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): Scale presentation and clinimetric testing results. *Movement Disorders*, 23, 2129–2170. <https://doi.org/10.1002/mds.22340>
- Graesser, A. C., Singer, M., & Trabasso, T. (1994). Constructing inferences during narrative text comprehension. *Psychological Reviews*, 101, 371–395. <https://doi.org/10.1037/0033-295X.101.3.371>
- Griffis, J. C., Nenert, R., Allendorfer, J. B., & Szaflarski, J. P. (2017). Damage to white matter bottlenecks contributes to language impairments after left hemispheric stroke. *NeuroImage: Clinical*, 14, 552–565. <https://doi.org/10.1016/j.nicl.2017.02.019>
- Grisoni, L., Miller, T. M., & Pulvermüller, F. (2017). Neural correlates of semantic prediction and resolution in sentence processing. *Journal of Neuroscience*, 2800–2816. <https://doi.org/10.1523/JNEUROSCI.2800-16.2017>
- Grossman, M., Cooke, A. C., Devita, C., Alsop, D., Detre, J., Chen, W., et al. (2002). Age-related changes in working memory during sentence comprehension: An fMRI study. *Neuroimage*, 15, 302–317. <https://doi.org/10.1006/nimg.2001.0971>
- Grossman, M., Cooke, A., Devita, C., Lee, C., Alsop, D., Detre, J., et al. (2003). Grammatical and resource components of sentence processing in Parkinson's disease: An fMRI study. *Neurology*, 60(5), 775–781. <https://doi.org/10.1212/01.WNL.0000044398.73241.13>
- Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6, 65–70.
- Horta-Barba, A., Pagonabarraga, J., Martínez-Horta, S., Marín-Lahoz, J., Sampedro, F., Fernández-Bobadilla, R., et al. (2020). The free and cued selective reminding test in Parkinson's disease mild cognitive impairment: Discriminative accuracy and neural correlates. *Frontiers in Neurology*, 11, 240. <https://doi.org/10.3389/fneur.2020.00240>
- Huetting, F. (2015). Four central questions about prediction in language processing. *Brain research*, 1626, 118–135. <https://doi.org/10.1016/j.brainres.2015.02.014>
- Hughes, A. J., Daniel, S. E., Kilford, L., & Lees, A. J. (1992). Accuracy of clinical diagnosis of idiopathic Parkinson's disease: A clinico-pathological study of 100 cases. *Journal of Neurological and Neurosurgical Psychiatry*, 55, 181–184. <https://doi.org/10.1136/jnnp.55.3.181>
- Kischka, U., Kammer, T. H., Maier, S., Weisbrod, M., Thimm, M., & Spitzer, M. (1996). Dopaminergic modulation of semantic network activation. *Neuropsychologia*, 34(11), 1107–1113. [https://doi.org/10.1016/0028-3932\(96\)00024-3](https://doi.org/10.1016/0028-3932(96)00024-3)
- Kotz, S. A., Frisch, S., Von Cramon, D. Y., & Friederici, A. D. (2003). Syntactic Language Processing: ERP lesion data on the role of the basal ganglia. *Journal of the International Neuropsychological Society: JINS*, 9(7), 1053–1060. <https://doi.org/10.1017/S1355617703970093>
- Kudlicka, A., Clare, L., & Hindle, J. V. (2011). Executive functions in Parkinson's disease: Systematic review and meta-analysis. *Movement disorders*, 26(13), 2305–2315. <https://doi.org/10.1002/mds.23868>
- Kuperberg, G. R., Brothers, T., & Wlotko, E. W. (2020). A Tale of Two Positivities and the N400: Distinct neural signatures are evoked by confirmed and violated predictions at different levels of representation. *Journal of Cognitive Neuroscience*, 32(1), 12–35. https://doi.org/10.1162/jocn_a_01465
- Kutas, M., & Federmeier, K. D. (2011). Thirty years and counting: Finding meaning in the N400 component of the event-related brain potential (ERP). *Annual Review of Psychology*, 62, 621–647. <https://doi.org/10.1146/annurev.psych.093008.131123>
- Kutas, M., & Hillyard, S. A. (1980). Reading senseless sentences: Brain potentials reflect semantic incongruity. *Science*, 207(4427), 203–205. <https://doi.org/10.1126/science.7350657>
- Kutas, M., & Hillyard, S. A. (1984). Brain potentials during reading reflect word expectancy and semantic association. *Nature*, 307(5947), 161. <https://doi.org/10.1038/307161a0>
- Kutas, M., Iragui, V. J., Niu, Y. Q., D'Avanzo, T. J., Yang, J. C., Salmon, D. P., et al. (2013). Altered N400 congruity effects in Parkinson's disease without dementia. *Cognitive Electrophysiology of Attention: Signals of the Mind*, 254–267. <https://doi.org/10.1016/B978-0-12-398451-7.00020-8>

- Lakens, D. (2016). Will knowledge about more efficient study designs increase the willingness to pre-register?. <https://doi.org/10.31222/osf.io/svzyc>. Retrieved from osf.io/uy8x5.
- Lang, S., Hanganu, A., Gan, L. S., Kibreab, M., Auclair-Ouellet, N., Alrazi, T., et al. (2019). Network basis of the dysexecutive and posterior cortical cognitive profiles in Parkinson's disease. *Movement Disorders*, 34(6), 893–902. <https://doi.org/10.1002/mds.27674>. Jun.
- Lau, E. F., Namyst, A., Fogel, A., & Delgado, T. (2016). A direct comparison of N400 effects of predictability and incongruity in adjective-noun combination. *Collabra*, 2(1). <https://doi.org/10.1525/collabra.40>
- Lau, E. F., Phillips, C., & Poeppel, D. (2008). A cortical network for semantics:(de) constructing the N400. *Nature Reviews Neuroscience*, 9(12), 920–933. <https://doi.org/10.1038/nrn2532>
- León-Cabrera, P., Flores, A., Rodríguez-Fornells, A., & Morís, J. (2019). Ahead of time: Early sentence slow cortical modulations associated to semantic prediction. *Neuroimage*, 189, 192–201. <https://doi.org/10.1016/j.neuroimage.2019.01.005>
- León-Cabrera, P., Rodríguez-Fornells, A., & Morís, J. (2017). Electrophysiological correlates of semantic anticipation during speech comprehension. *Neuropsychologia*, 99, 326–334. <https://doi.org/10.1016/j.neuropsychologia.2017.02.026>
- Litvan, I., Aarsland, D., Adler, C. H., Goldman, J. G., Kulisevsky, J., Mollenhauer, B., et al. (2011). MDS task force on mild cognitive impairment in Parkinson's disease: Critical review of PD-MCI. *Movement Disorders*, 26(10), 1814–1824. <https://doi.org/10.1002/mds.23823>
- Li, X., Zhang, Y., Xia, J., & Swaab, T. Y. (2017). Internal mechanisms underlying anticipatory language processing: Evidence from event-related-potentials and neural oscillations. *Neuropsychologia*, 102, 70–81. <https://doi.org/10.1016/j.neuropsychologia.2017.05.017>
- López-Calderon, J., & Luck, S. J. (2014). ERPLAB: An open-source toolbox for the analysis of event-related potentials. *Frontiers in Human Neuroscience*, 8, 213. <https://doi.org/10.3389/fnhum.2014.00213>
- Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG-and MEG-data. *Journal of Neuroscience Methods*, 164(1), 177–190. <https://doi.org/10.1016/j.jneumeth.2007.03.024>
- Martínez-Horta, & S., & Kulisevsky, J. (2019). Mild cognitive impairment in Parkinson's disease. *Journal of Neural Transmission*, 126(7), 897–904. <https://doi.org/10.1007/s00702-019-02003-1>
- Mattox, S. T., Valle-Inclán, F., & Hackley, S. A. (2006). Psychophysiological evidence for impaired reward anticipation in Parkinson's disease. *Clinical Neurophysiology*, 117(10), 2144–2153. <https://doi.org/10.1016/j.clinph.2006.05.026>
- Mestres-Missé, A., Rodríguez-Fornells, A., & Münte, T. F. (2007). Watching the brain during meaning acquisition. *Cerebral Cortex*, 17(8), 1858–1866. <https://doi.org/10.1093/cercor/bhl094>
- Monchi, O., Petrides, M., Doyon, J., Postuma, R. B., Worsley, K., & Dagher, A. (2004). Neural bases of set-shifting deficits in Parkinson's disease. *Journal of Neuroscience*, 24, 702–710. <https://doi.org/10.1523/JNEUROSCI.4860-03.2004>
- Muslimović, D., Post, B., Speelman, J. D., & Schmand, B. (2005). Cognitive profile of patients with newly diagnosed Parkinson disease. *Neurology*, 65(8), 1239–1245. <https://doi.org/10.1212/01.wnl.0000180516.69442.95>
- Nieuwland, M. S., Barr, D. J., Bartolozzi, F., Busch-Moreno, S., Darley, E., Donaldson, D. I., et al. (2020). Dissociable effects of prediction and integration during language comprehension: Evidence from a large-scale study using brain potentials. *Philosophical Transactions of the Royal Society*. <https://doi.org/10.1098/rstb.2018.0522>. B37520180522.
- Olichney, J. M., Taylor, J. R., Gatherwright, J., Salmon, D. P., Bressler, A. J., Kutas, M., et al. (2008). Patients with MCI and N400 or P600 abnormalities are at very high risk for conversion to dementia. *Neurology*, 70(19 Part 2), 1763–1770. <https://doi.org/10.1212/01.wnl.0000281689.28759.ab>
- Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). Fieldtrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, 1. <https://doi.org/10.1155/2011/156869>, 2011.
- Pagonabarraga, J., Kulisevsky, J., Llebaria, G., García-Sánchez, C., Pascual-Sedano, B., & Gironell, A. (2008). Parkinson's disease-cognitive rating scale: A new cognitive scale specific for Parkinson's disease. *Movement Disorders: Official Journal of the Movement Disorder Society*, 23(7), 998–1005. <https://doi.org/10.1002/mds.22007>
- Payne, B. R., & Federmeier, K. D. (2018). Contextual constraints on lexico-semantic processing in aging: Evidence from single-word event-related brain potentials. *Brain Research*, 1687, 117–128. <https://doi.org/10.1016/j.brainres.2018.02.021>
- Payne, B. R., & Silcox, J. W. (2019). Aging, context processing, and comprehension. In , Vol. 71. *Psychology of learning and motivation* (pp. 215–264). Academic Press. <https://doi.org/10.1016/bs.plm.2019.07.001>
- Pell, M., & Monetta, L. (2008). How Parkinson's disease affects non-verbal communication and language processing. *Language and Linguistics Compass*, 2(5), 739–759. <https://doi.org/10.1111/j.1749-818X.2008.00074.x>
- Peña-Casanova, J., Quiñones-Ubeda, S., Gramunt-Fombuena, N., Quintana-Aparicio, M., Aguilar, M., Badenes, D., et al. (2009). Spanish multicenter normative studies (NEURONORMA project): Norms for verbal fluency tests. *Archives of Clinical Neuropsychology*, 24, 395–411. <https://doi.org/10.1093/arclin/acp042>
- Pulvermüller, F., & Grisoni, L. (2020). Semantic prediction in brain and mind. *Trends in Cognitive Sciences*, 24(10), 781–784. <https://doi.org/10.1093/arclin/acp042>
- Rabovsky, M., Hansen, S. S., & McClelland, J. L. (2018). Modelling the N400 brain potential as change in a probabilistic representation of meaning. *Nature Human Behaviour*, 2(9), 693–705. <https://doi.org/10.1038/s41562-018-0406-4>
- Taylor, W. I. (1953). Cloze procedure: A new tool for measuring readability. *Journal and Mass Communication Quarterly*, 30, 415. <https://doi.org/10.1177/107769905303000401>
- Tomlinson, C. L., Stowe, R., Patel, S., Rick, C., Gray, R., & Clarke, C. E. (2010). Systematic review of levodopa dose equivalency reporting in Parkinson's disease. *Movement Disorders*, 25, 2649–2653. <https://doi.org/10.1002/mds.23429>
- Unsworth, N., Spillers, G. J., & Brewer, G. A. (2011). Variation in verbal fluency: A latent variable analysis of clustering, switching, and overall performance. *Quarterly Journal of Experimental Psychology*, 64(3), 447–466. <https://doi.org/10.1080/17470218.2010.505292>
- Van Berkum, J. J., Brown, C. M., Zwitserlood, P., Kooijman, V., & Hagoort, P. (2005). Anticipating upcoming words in discourse: Evidence from ERPs and reading times. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 31(3), 443. <https://doi.org/10.1037/0278-7393.31.3.443>
- Van Berkum, J. J., Hagoort, P., & Brown, C. M. (1999). Semantic integration in sentences and discourse: Evidence from the N400. *Journal of Cognitive Neuroscience*, 11(6), 657–671. <https://doi.org/10.1162/089892999563724>
- Whiteside, D. M., Kealey, T., Semla, M., Luu, H., Rice, L., Basso, M. R., et al. (2016). Verbal fluency: Language or executive function measure? *Applied Neuropsychology: Adult*, 23(1), 29–34. <https://doi.org/10.1080/23279095.2015.1004574>
- Wicha, N. Y., Bates, E. A., Moreno, E. M., & Kutas, M. (2003). Potato not pope: Human brain potentials to gender expectation and

- agreement in Spanish spoken sentences. *Neuroscience Letters*, 346(3), 165–168. [https://doi.org/10.1016/S0304-3940\(03\)00599-8](https://doi.org/10.1016/S0304-3940(03)00599-8)
- Willems, R. M., Frank, S. L., Nijhof, A. D., Hagoort, P., & Van den Bosch, A. (2016). Prediction during natural language comprehension. *Cerebral Cortex*, 26(6), 2506–2516. <https://doi.org/10.1093/cercor/bhv075>
- Wlotko, E. W., & Federmeier, K. D. (2012). Age-related changes in the impact of contextual strength on multiple aspects of sentence comprehension. *Psychophysiology*, 49(6), 770–785. <https://doi.org/10.1111/j.1469-8986.2012.01366.x>