

Rationalizing the Decavanadate(V) and Oxidovanadium(IV) Binding to G-actin and the Competition with Decaniobate(V) and ATP

Giuseppe Sciortino,^{,§,#} Manuel Aureliano,^{*,†} and Eugenio Garribba^{*,§}*

[§] Dipartimento di Chimica e Farmacia, Università di Sassari, Via Vienna 2, I-07100 Sassari, Italy

[#] Institute of Chemical Research of Catalonia (ICIQ), Avda. Països Catalans, 16, 43007 Tarragona,
Spain

[†] CCMar, FCT, Faculdade de Ciências e Tecnologia, Universidade do Algarve, 8000-139 Faro,
Portugal

Corresponding authors. E-mail: gsciortino@iciq.es (G.S.); maalves@ualg.pt (M.A.);
garribba@uniss.it (E.G.).

Reviewer 1 Reviewer 2 Editor My/Giuseppe changes

Abstract

In this study the experimental data collected over the last fifteen years on the interaction of decavanadate(V) ($V_{10}O_{28}^{6-}$; V_{10}), a polyoxometalate (POM) with promising anticancer and antibacterial action, with G-actin, were rationalized using several computational approaches (docking, density functional theory (DFT) and molecular dynamics (MD)). Moreover, a comparison with the isostructural and more stable decaniobate(V) ($Nb_{10}O_{28}^{6-}$; Nb_{10}) was carried out. Four binding sites were identified, named α , β , γ and δ , being located the site α in the catalytic cleft of the enzyme at the interface of the subdomains II and IV. It was observed that the site α is preferred by V_{10} , whereas Nb_{10} is more stable at the site β ; this indicates that, differently from other proteins, G-actin contemporaneously binds the two POMs, whose action is synergistic. Both decavanadate and decaniobate induce conformational rearrangements in G-actin, larger for V_{10} than Nb_{10} . Moreover, the binding mode of oxido vanadium(IV) ion, $V^{IV}O^{2+}$, formed upon the reduction of decavanadate(V) by the $-SH$ groups of accessible cysteine residues, is also found in the catalytic site α with (His161, Asp154) coordination; this adduct overlaps significantly with the region where ATP is bound, accounting for the competition between V_{10} and its reduction product $V^{IV}O^{2+}$ with ATP, as previously observed by EPR spectroscopy. Finally, the competition with ATP was rationalized: since V_{10} most stable binding site is α , Ca^{2+} -ATP prevents the V_{10} interaction with G-actin, while the competition is less important for Nb_{10} once for this POM β is the primary site. A relevant consequence of this paper is that other metallodrug–protein systems, in the absence or presence of eventual inhibitors and/or competition with molecules of the organism, could be studied with the same approach, suggesting important elements for an explanation of the biological data and a rational drug design.

Introduction

After the discovery of cisplatin, inorganic medicinal chemistry is a field of extensive research. Potential antitumor drugs based on Pt, Ru, and Cu, antimicrobial and antiparasitic agents based on Sb, Ag, As, and Fe, antiarthritis on Au, antiulcers on Bi, and antivirals based on Co and Hg were proposed. In addition, diagnostic metallodrugs of Gd, Tc and In were also developed.^{1, 2} Similarly, vanadium compounds have long been studied as potential antidiabetic drugs and, more recently, as anticancer, antiparasitic, antiviral, antiHIV and antituberculosis agents.³

Interaction with proteins plays a crucial role in the transport and activity of metallodrugs in the organism, both for their high affinity toward the metals or high concentration in the bloodstream and cellular environment. Unfortunately, in most of cases, the experimental data cannot be rationalized through a description at molecular level of this interaction.⁴ Instrumental methods often give only information on the biological effects of a metallodrug after the binding with a target protein, but present severe limitations in order to unveil the type of interaction and the binding modes and sites, or to ascertain the competition between the metallospecies and other biomolecules of the organism.

Decavanadate ($V_{10}O_{28}^{6-}$, V_{10}) and decaniobate ($Nb_{10}O_{28}^{6-}$, Nb_{10}) are members of a large family of polyanionic clusters formed by group V and VI metals that are known as polyoxometalates (POMs). POMs have been found to have a wide range of biological activities such as antidiabetic, antibacterial, antiprotozoal, antiviral and anticancer, which have sparked interest in their use as bio-inorganic drugs.⁵ Recently, it has been demonstrated that POMs function as indirect activators of a G protein-coupled receptor.⁶ Moreover, POMs have also seen applications in other fields of research such as aerobic oxidation of starch, smart glasses, solar-driven photocatalysis, protein crystallography and synthesis of metal-organic frameworks (MOFs);⁷ for example, Nb_{10} can be used as building block to prepare 1D, 2D, and 3D frameworks.⁸

Based on an analysis of the antibacterial activity and POM structures, putative mode of actions toward potential targets were suggested and a structure-activity relationship was determined for a

series of POMs against *Helicobacter pylori* and *Streptococcus pneumoniae*.^{5e} The activity of POMs are found to be bacteria-dependent: for *Streptococcus pneumoniae* the most active POMs were polyoxovanadates (POVs), especially V₁₀, while for *Helicobacter pylori* the most of POMs are effective.^{5e}

The group of Aureliano extensively studied the influence of POMs, especially POVs and polyoxotungstates (POTs), on diverse biochemical mechanisms.⁹ In many cases, these POM-associated biological activities are the result of POM–protein/enzyme interactions, which are mainly of electrostatic nature as evidenced by crystallographic studies showing that the negatively charged metal clusters are generally found within or at positively charged regions of proteins.¹⁰ However, also covalent interactions between biomacromolecules and POMs are possible.¹⁰⁻¹¹ In this scenario, the inhibition of certain enzymes by POMs can trigger the impairment of vital cell functions.¹⁰⁻¹¹

Among the possible cellular targets for POMs, actins are worth of being considered. Actins are found in all life kingdoms and are responsible for many biologically essential processes, such as muscle contraction, cell adhesion, division and migration, thus assigning them important roles in health and diseases.¹² Globular actin (G-actin; 42 kDa), the actin monomer, is one of the most abundant proteins in cells, being involved in numerous cellular processes. Previously, it was shown that V₁₀–G-actin interaction affects the protein structure and dynamics, and also influences its transformation to filamentous F-actin (which is the polymerized form of actin).¹¹ Combining our experience regarding V₁₀–G-actin interactions,^{11, 13, 14, 15} and our expertizes in computational characterization of metallospecies–protein interactions,¹⁶ we further pursue the interaction between decavanadate and G-actin to deduce potential targets and provide possible modes of the biological activity of POMs.

Despite V₁₀ and Nb₁₀ are isoelectronic and isostructural isopolyoxoanions, V₁₀ is redox active and labile in aqueous solution, while Nb₁₀ is kinetically inert and redox stable.¹⁷ Although there has been growing interest in niobium chemistry in the last decade, research on decaniobate and its

interplay with biomolecules and/or biological effects is still scarce.^{9h, 18, 19} In the systems with Ca^{2+} -ATPase, Nb_{10} prevents the binding of V_{10} , suggesting the same target(s) in the protein structure;¹⁸ moreover, both of them cause comparable conformational changes.^{9h, 19} In addition, Nb_{10} constitutes an excellent model to explore the interplay between its V_{10} analogue and biologically vital proteins such as actin.

Even though some studies were carried out on the systems containing POMs and actin, up to now the detailed knowledge of the molecular basis of V_{10} -G-actin interactions and the possible involvement in protein polymerization are lacking. In fact, the exact mechanism of action of POMs is still elusive and thus also the POM's target enzymes, which are responsible for the observed biological effects. Molecular modelling could represent a valid tool to fill this gap and help to understand cellular processes providing valuable clues for the rational design of new metal-substituted POMs with specific interaction properties, pushing forward the comprehension of the molecular events governing POMs-protein binding. Even though the computational techniques have been largely applied to study the chemistry of POMs,²⁰ only few examples of molecular modelling application to POM-protein interaction have been published so far.²¹

Based on these premises described above, in this paper we tried to rationalize – using several computational approaches – the experimental data on the V_{10} -G-actin systems, collected over the last years in the literature, and compare the findings with the isostructural Nb_{10} . In particular, the aims of the present study are: i) to attain an understanding, at molecular level, of the interaction of V_{10} and Nb_{10} with G-actin starting from the available results already obtained by several spectroscopic techniques;^{11, 13-15} ii) to fully characterize the binding sites of G-actin for V_{10} and Nb_{10} , determine the number and type of neighbor atoms around POMs and evaluate their stabilization through secondary interactions; iii) to compare the behavior of V_{10} and Nb_{10} ; iv) to predict the binding site(s) of the reduction product of V_{10} , the oxidovanadium(IV) ion, $\text{V}^{\text{IV}}\text{O}^{2+}$; v) to study the $\text{V}_{10}/\text{Nb}_{10}$ -G-actin system in the presence of ATP to obtain a more accurate description of the mechanisms involved and possible competition/inhibition effects.

Computational Section

DFT calculations. $V_{10}O_{28}^{6-}$ and $Nb_{10}O_{28}^{6-}$ geometry and harmonic frequencies were computed through Gaussian 09 (revision D.01)²² using the B3LYP functional including Grimme's empirical correction for dispersion.²³ V and Nb atoms were described using the scalar-relativistic Stuttgart–Dresden SDD double- ζ basis set complemented with f polarization functions and its associated pseudopotential.²⁴ The 6-31g(d,p) basis set was used for the oxygen atoms. The structures of POMs were optimized in water described by the SMD continuum model.²⁵ Point charges of the complete POMs were derived using the RESP²⁶ (Restrained ElectroStatic Potential) model. DFT level of theory and RESP model to assign the atomic charges were selected following the protocol proposed to derive AMBER and TIP3P force fields for Molecular Dynamics (MD).²⁷

The refinement of the oxidovanadium(IV) adducts at the nucleotide site (site α), found by dockings, was performed cutting out the region with the $V^{IV}O^{2+}$ ion, modeled as $[V^{IV}O(H_2O)_4]^{2+}$, and neighbor interacting amino acid side chains and freezing the backbone atoms as reported by Siegbahn and Himo.²⁸ The geometry relaxation and ΔE calculations were performed using the doublet state in unrestricted simulations at B3P86/6-311++g(d,p) level of theory and describing water within the framework of SMD model²⁵ that gave excellent results for other V^{IV} –protein systems.^{16f, 16g, 29} For the geometry optimization of V^{IV} complexes, it was shown in the literature that B3P86 works better than other functionals like B3LYP.³⁰ Frequency calculations were computed to ensure that the structures were a minimum in the potential energy surface.

The prediction of the ^{51}V hyperfine coupling tensor $\mathbf{A}$ was performed on the optimized structures with Gaussian 09²² using BHandHLYP functional and 6-311+g(d) basis set.³¹ The spin contamination was negligible and the eigenvalue of the spin operator $\langle S^2 \rangle$ was 0.76 for

all the calculations, very close to the exact value of 0.75. This is in agreement with results obtained for hybrid functionals.³² The theory background can be found in refs. [32b](#), [33](#).

Molecular Dynamics. The model of apo monomeric actin, G-actin, were built removing Ca^{2+} -ATP ligand and adding non-characterized regions 39-51 and 1-5 with Modeller³⁴ on the unique native X-ray structure of mononuclear G-actin (PDB: 3hbt [35](#)). In the case of ATP bound simulations and competition studies, the Ca^{2+} -ATP present in the X-ray structure has been retained. Molecular Dynamics simulations were set up with Xleap,³⁶ which was instructed to solvate the protein with a cubic box of pre-equilibrated TIP3P water molecules and balance the total charge with Na^+ ions (ions94.lib library). The AMBER14SB force field³⁷ was used for the protein, while V- and Nb-bonding force constants and equilibrium parameters were obtained through the Seminario method, using the MCPB.py.³⁸ Ca^{2+} -ATP parameters from Meagher *et al.*³⁹ and Bradbrook *et al.*⁴⁰ have been used.

During MDs, solvent and the whole system were sequentially submitted to 3000 energy minimization steps. Then, thermalization of water molecules and side chains was achieved by increasing the temperature from 100 K up to 300 K. MDs under periodic boundary conditions were carried out during 100 ns with OpenMM engine⁴¹ through OMMProtocol.⁴²

Analysis of the trajectories was carried out by means of cpptraj implemented in ambertools16.³⁶ The MD trajectory was considered converged when a stable exploration of the conformational space was achieved; in particular, a stable conformation or a pool of relative stable conformations visited for a statistically consistent number of times were considered as convergence indicators.⁴³ Considering the alpha carbons of the G-actin backbone, root mean square deviation (RMSD) from the minimized structure, all-to-all frames RMSD⁴⁴ and cluster counting⁴⁵ analysis were performed. Moreover, to ensure that dynamic transitions occur between different conformations, a principal component analysis (PCA) was carried out plotting the two principal modes relative to each other.⁴⁴ The quality threshold (QT) clustering algorithm was used to generate representative clusters taking

the RMSD between frames as distance metric.⁴⁶ The representative structure of the most populated cluster was used in the further structural analysis.

Docking calculations. Docking calculations to the G-actin were carried out through GOLD 5.8 software⁴⁷ on the most sampled structure coming from MD trajectories. The proteins were docked with V₁₀ and Nb₁₀ optimized structures considering the whole rigid protein.

Dockings of V^{IV}O²⁺ moiety were carried out through GOLD 5.8⁴⁷ on the regions selected in the preliminary analysis by BioMetAll,⁴⁸ instructed to find latent binding motifs with a minimum of two coordinating residues considering His, Asp and Glu as potential donors. G-actin was docked with V^{IV}O²⁺, V^{IV}O(H₂O)²⁺, and V^{IV}O(H₂O)₂²⁺ moieties, obtained from the optimization of the oxido vanadium(IV) aquaion, [V^{IV}O(H₂O)₄]²⁺. The equatorial positions were activated replacing the equatorial water(s) with dummy hydrogen atoms according to what was recently established.^{16a-c, 16e-h} All dockings were computed considering both the protonation states at δ and ϵ nitrogens of His imidazole ring. Genetic algorithm (GA) parameters have been set to perform a minimum of 100000 operations during 50 GA runs for coordination dockings, and 100 GA runs for classic dockings. The other parameters of GA were set to default. The scoring (*Fitness* of GoldScore, F) was evaluated applying the modified versions of GoldScore scoring function, which was validated in previous published papers.^{16a, 16c-e} The best solutions (binding poses) were evaluated through three main criteria: i) the mean (F_{mean}) and the highest value (F_{max}) of the scoring associated with each pose; ii) the population of the cluster containing the best pose; iii) the position in the *Fitness* ranking of the computed cluster.

The POMs docking solutions were refined sending for each best proposal a 200 ns of Molecular Dynamics and ΔG_{bind} quantified by molecular mechanics-generalized Born surface area (MM-GBSA) including quasi-harmonic correction for entropy.⁴⁹ Average values have been obtained from 100 homogeneously distributed frames of the whole MD trajectories. The following parameters were used: $\text{igb} = 5$, $\text{saltcon} = 0.100$.

Results and discussion

(1) Determination of the binding sites. Experimental data reported so far pointed out that decavanadate anion (V_{10}) interacts strongly with G-actin.^{11, 13-14, 50} However, up to now, even if the biological effects are clear, the binding site(s) were not determined and this datum is lacking in the literature. Nevertheless, this information could be important and besides related to the subsequent reduction process to oxido vanadium(IV) cation, $V^{IV}O^{2+}$,¹³ very useful to gain insight for the rational design of new metal-substituted POMs (i.e. with two or more metals in their structure) with specific interaction with proteins.

To throw light on the unknown aspects, in the first step a blind docking screening was carried out on the equilibrated apo structure of G-actin, searching for putative binding sites for V_{10} and Nb_{10} . The results of this rigid docking essay suggest the presence of four potential binding sites, each of them involving at least two positive residues of Lys or Arg: site α is located in the catalytic cleft of the enzyme at the interface of subdomains II and IV; site β at the interface of subdomains I and II; site γ at the surface of subdomain III; and site δ at the C-terminus in the small cleft between subdomains I and III (Figure 1a). After this preliminary stage, further docking calculations on the identified regions were performed considering the flexibility of the side-chains: similar results for both POMs were obtained with scoring (F_{max}) ranging from 24.8 up to 36.4 Gold Score *Fitness* units. For the site α , the pose (solution) with the best scoring is stabilized by hydrogen bonds (Hbonds) with Arg207, Arg211, Arg62, Thr204, Gln59 and Tyr69 residues. The site β , with similar scoring, presents Hbond with Lys68, Arg37, Lys50 and Lys84. The sites γ and δ show lower scorings and Hbonds with Arg147, Lys329, Thr148 and Asn297 or Arg113, Lys374 and Tyr170, respectively (Figure 1a). Comparison of the DFT optimized structure of V_{10} and Nb_{10} is shown in Figure 1b; Nb_{10} displays slightly wider volume and surface area than its vanadium homologue V_{10} and, moreover, accumulates more surface negative charge (Figure S1 of the Supporting Information). However, since only slight differences in the behavior of V_{10} and Nb_{10} are observed,

because of the similarity between the two structures, dockings can be useful at most for defining the possible binding regions but do not allow for ascertaining their relative affinities toward the four sites. To move beyond docking and obtain reliable values for ΔG_{bind} for the two POMs, MDs were performed and residence time analysis and MM-GBSA calculations carried out.




Figure 1. a) The four binding sites found by docking for V_{10} and Nb_{10} in the structure of G-actin. The positively charged side-chains of Arg and Lys, which interact with POMs in each site, are depicted as blue balls. Subdomains (I to IV) of G-actin are shown with different colours (orange, blue, purple and green). b) Comparison of the DFT optimized structure of V_{10} and Nb_{10} . Volume (V in $\text{\AA}^3$) and surface area (A in $\text{\AA}^2$) were computed considering van der Waals (vdW) radii. V is shown in grey, Nb in mint and O in red.

(2) Conformational changes induced by V_{10} and Nb_{10} and relative stability of the binding sites. It was previously described in 2009 that the experimental exchange rate and apparent

dissociation constant for ATP, measured from fluorescence decrease (Figure S2 of the Supporting Information), indicated that V_{10} binding induce an opening of the G-actin cleft.¹³ Subsequently, in 2011, the ^1H NMR spectra of G-actin treated with V_{10} suggested that this latter causes major alterations in the protein structure (Figure S3 of the Supporting Information);^{50, 51} whereas in contrast, no significant changes in the ^{51}V NMR signals were observed for other vanadates, such as monomeric ($\text{H}_2\text{V}^{\text{V}}\text{O}_4^-/\text{HV}^{\text{V}}\text{O}_4^{2-}$; V_1) and tetrameric species ($\text{V}_4\text{O}_{12}^{4-}$; V_4).¹³ More recently, EXAFS and XANES experiments established that the interaction of V_{10} -G-actin triggers a protein conformational reorientation that causes an oxidation of Cys residues and formation of $\text{V}^{\text{IV}}\text{O}^{2+}$ cation (Figure 2).¹⁵




Figure 2. EXAFS profile of the systems containing V_{10} (in black), V_{10} -G-actine (in brown) and V_{10} -G-actin-ATP (in purple). Adapted from ref. ¹⁵.

The change of actin conformation was further confirmed by circular dichroism (CD) spectroscopy.⁵² Upon addition of V_{10} to G-actin, the melting points determined from the CD

melting curves were observed at lower values compared with the system with ATP, respectively T_{m1} 31.7 vs. 48.9 °C and T_{m2} 51.7 vs. 64.3 °C. The decrease (in the range 12.6-17.2 °C) in the protein melting temperature could probably be due to actin conformational changes and/or partial protein unfolding induced by V_{10} .⁵²

To assess the stability of the oxidovanadium(IV) adducts and the effect of the two POMs on the protein folding, subsequent MD simulations were carried out starting from V_{10} and Nb_{10} bound to each of the four possible sites and compared with the unbound state of G-actin. Concerning the site α , adducts are highly stable both for V_{10} and Nb_{10} during the whole explored trajectory (200 ns) and the folding of the protein is generally retained. It must be noted that a slight opening of the nucleotide binding site is observed after the interaction with V_{10} and Nb_{10} . Subdomain IV moves from subdomain II to accommodate the POMs. This movement can be described with the angle ϕ , defined as $C(\alpha)_{Lys67}-C(\alpha)_{Val339}-C(\alpha)_{Thr203}$, that varies from 28.6° (apo form) to 30.5° (Nb_{10} -G-actin) and 35.6° (V_{10} -G-actin) (Figure 3a). These evidences highly match with what is observed in this study by MDs simulations upon V_{10} binding to the site α . The Hbond interactions observed by docking are almost retained along the whole simulation time. The differences are as follows: for V_{10} the interaction with Thr203 is replaced by Arg184, while for Nb_{10} that with Tyr69 is lost and the moiety slightly moves outside the cleft toward the subdomain II where Nb_{10} is rotated $\sim 90^\circ$ compared to V_{10} on its principal axes (Figure 3b and 3c).




Figure 3. Molecular Dynamics (MD) most sampled structures showing: a) the cleft opening upon the POMs binding (apo G-actin, V₁₀-G-actin, Nb₁₀-G-actin are shown in green, orange and blue, respectively); b) binding mode of V₁₀ at the site α ; c) binding mode of Nb₁₀ at the site α . Hbond donors are explicitly depicted.

At the site β , both POMs form stable adducts during 200 ns of simulation. The Hbond interactions found by docking are generally retained with the addition in several frames of Lys50 coming from the flexible 39-51 loop (Figure 4a and 4b). The general G-actin folding is also maintained. The site γ shows differences between the two moieties: while the V₁₀ leaves the surface of subdomain III at 75 ns and after 170 ns moves to the site β where it is stabilized, Nb₁₀ forms a stable adduct during the whole trajectory. The Hbond network highlighted by docking is retained with the replacement of Thr148 with the positive side-chain of Lys327 (Figure 4c). Finally, the site δ at the C-terminus is unstable for both the POMs. V₁₀ at 40 ns moves away and, after 123 ns, it forms again a stable adduct at the site α . Nb₁₀, due to its higher volume, shows a faster unbinding process and, in the first steps of the simulation, leaves C-terminus to accommodate at the site γ , where it is stable along the entire trajectory.

From the energetic point of view the stable adducts formed by the two POMs at the binding regions of G-actin, behave differently. While V₁₀ shows the best affinity for the site α (ΔG_{bind} for

the site α is lower than *ca.* 23 kcal mol⁻¹ than the site β), the Nb₁₀ affinity order is different: the site β is more stable than the site α of 2.0 kcal·mol⁻¹ and than the site γ of 17.2 kcal·mol⁻¹ (Table 1). Even if the value of 2 kcal mol⁻¹ is close to the uncertainty for a DFT prediction, from these data it emerges clearly the different affinity order of V₁₀ and Nb₁₀ for the two sites. Moreover, these results are in good agreement with the previous studies which suggested that V₁₀ binds to proteins whose natural ligands are nucleotides, such as ribonucleases and myosin.⁵³

Table 1. MD residence time of V₁₀ and Nb₁₀ at the identified sites (α , β , γ and δ) and binding effects on the G-actin folding.

Site	Subdomain(s)	V ₁₀ (t, ns)	$\Delta(\Delta G_{\text{bind}})^a$	Nb ₁₀ (t, ns)	$\Delta(\Delta G_{\text{bind}})^a$	Fold
α	I/IV	Stable, 200	0.0	Stable, 200	2.0	ϕ variation ^b
β	II	Stable, 200	22.9	Stable, 200	0.0	Retained
γ	III	Unstable, < 75	–	Stable, 200	17.2	Retained
δ	I/III	Unstable, < 40	–	Unstable, < 5	–	–

^a MM-GBSA relative values including quasi-harmonic entropic correction computed on 100 homogeneously distributed frames of the whole MD trajectories (values in kcal·mol⁻¹). ^b ϕ = 28.6° (apo form), ϕ = 30.5° (adduct with Nb₁₀) and ϕ = 35.6° (adduct with V₁₀).

These results can rationalize the kinetic measurements of V₁₀ decomposition as a function of G-actin concentration, monitored by UV/Vis spectroscopy.¹¹ In fact, G-actin causes marked effects in stability of V₁₀, increasing its half-life from 5 h in solution up to 27 h (Figure S4 of the Supporting Information); this demonstrates a clear protection effect exerted by G-actin toward V₁₀. The best binding site proposed by simulations, site α , shows characteristics compatible with a protective environment being buried into the catalytic cleft of the proteins, and able to form a Hbond network,

beside electrostatic interactions, strong enough to stabilize V_{10} into the pocket preventing its exposition to the solvent.




Figure 4. Molecular Dynamics (MD) most sampled structures showing: a) the binding mode of V_{10} at the site β ; b) the binding mode of Nb_{10} at the site β ; c) the binding mode of Nb_{10} at the site γ . H bond donors are explicitly depicted.

(3) Reduction of V_{10} to V^{IV} and binding sites of $V^{IV}O^{2+}$ ion. After the incubation of V_{10} with G-actin, a rigid-limit EPR spectrum is revealed with eight resonances in the parallel region and eighth in the perpendicular one (trace b of Figure 5).¹³⁻¹⁴ This indicates a reduction process of V^V ($3d^0$, EPR-silent) to $V^{IV}O$ ($3d^1$, EPR-active), probably through the reaction of decavanadate(V) with the accessible $-SH$ groups of cysteine residues. The most plausible candidates involved in the redox reaction are Cys272 and Cys374.⁵⁴ The spin Hamiltonian parameters for the adduct formed upon the binding of $V^{IV}O^{2+}$ to G-actin are $g_z = 1.940$ and $|A_z| = 177.1 \times 10^{-4} \text{ cm}^{-1}$;¹⁴ these values would indicate weak donors or – in agreement to the reinterpretation of the "additivity relationship", the empirical rule verified for a lot of $V^{IV}O$ complexes which allows to determine the ^{51}V hyperfine coupling constant along the z axis, $|A_z|^{estmtd}$, from the sum of the equatorial donor contributions within $ca. 3 \times 10^{-4} \text{ cm}^{-1}$ from the experimental value^{55, 56} – coordination of His residues with the

aromatic plane perpendicular to the $V^{IV}=O$ bond.⁵⁵ At the moment, the binding site and the residues interacting with $V^{IV}O^{2+}$ ion are not known; however, the effect of the addition of an excess of ATP (Figure 5c) suggests that the site could be the same as that of the nucleotide.




Figure 5. X-band EPR spectra recorded on frozen aqueous solution containing: a) 2.5 mM V_{10} in 2 mM Tris, 0.2 mM $CaCl_2$, 0.2 mM ATP (pH 7.5); b) 2.5 mM V_{10} in 2 mM Tris, 0.2 mM $CaCl_2$, plus 100 μ M G-actin (pH 7.5); c) 2.5 mM V_{10} in 2 mM Tris, 0.2 mM $CaCl_2$, plus 50 μ M G-actin and 0.2 mM of ATP to prevent oxidovanadium(IV) formation (pH 7.5). Adapted from refs. [13](#) and [14](#).

In this study, computational methods have been applied to determine the latent $V^{IV}O^{2+}$ binding site into the nucleotide binding region. The catalytic pocket was first probed for zones displaying at least two donors-containing residues with a sufficient backbone preorganization to bind $V^{IV}O^{2+}$.^{48, 29a} A docking assay was carried out and two couples emerged as potential candidate sites, (His161; His73) and (His161; Asp154), both of them displaying the ability to coordinate $V^{IV}O^{2+}$ in a square pyramidal arrangement (Table 2).

Table 2. Full DFT refined structures (from the docking proposals), estimated (estmtd) and DFT calculated (calcd) $|A_z|$ values and electronic binding energies for the binding of $V^{IV}O^{2+}$ ion to the nucleotide site α of G-actin.

Donors	Distances ^a	$F_{\max}$ ^b	F_{mean} ^c	Pop. % ^d	ΔE_{bind} ^e	$ A_z _{\text{estmtd}}$ ^e	$ A_z _{\text{calcd}}$ ^f
(N _{H161} , COO ⁻ _{D154}); O _W ; O _W ^g	2.100, 1.964 ^h	34.1	31.8	60	-28.4	178.4 ⁱ	178.6 ⁱ
(N _{H73} , N _{H161}); O _W ; O _W ^j	2.128, 2.101 ^k	45.4	39.5	64	-13.2	173.4 ⁱ	169.4 ⁱ

^a Distance in Å. ^b GoldScore *Fitness* value obtained for the more stable pose of each cluster. ^c Average value of GoldScore *Fitness* for each cluster. ^d Percentage computed considering the total of the solutions reported (numbers of solutions per cluster). ^e ΔE_{bind} in kcal·mol⁻¹. ^f $|A_z|$ estimated (estmtd.) with the “additivity relationship” in 10⁻⁴ cm⁻¹ units. ^g H₂O molecules *cis* to each other. ^h V–N_{H161} and V–O_{D154} distances, respectively. ⁱ The experimental value of $|A_z|$ is 177.1×10^{-4} cm⁻¹. ^j H₂O molecules *trans* to each other. ^k V–N_{H73} and V–N_{H161} distances, respectively.

The binding mode with two His, (His 73, His 161); O_W; O_W, is characterized by the *trans* arrangement of the two H₂O molecules and by an additional Hbond stabilization between the oxido ligand and Arg177 (Figure 6a). The mode, in which His161 and Asp154 are bound to $V^{IV}O^{2+}$, displays a *cis* arrangement of the two aqua ligands with the aromatic ring of the His perpendicular to the V=O (θ , the calculated dihedral angle O=V^{IV}–N–C – where C is the carbon atom bridging the two nitrogens – is 72.1°; Figure 6b). It results located deeper inside the nucleotide binding site α and overlaps significantly with the region with ATP (Figure 6c), making it a potential candidate to rationalize the V₁₀/ATP competition observed by EPR. In fact, as reported by Ramos *et al.*, and described in Figure 5, the addition of ATP to a solution containing V₁₀ and G-actin prevents the stabilization of paramagnetic $V^{IV}O^{2+}$, the product of the reduction of V₁₀.^{13, 14}




Figure 6. Full DFT optimized structures of the two docking proposals for the binding of $V^{IV}O^{2+}$ moiety to the nucleotide binding site α of G-actin: a) the binding mode with (His73, His161); b) the binding mode with (His161, Asp154) and c) surface volume of ATP in its native site superimposed with the favored $V^{IV}O^{2+}$ binding.

The predicted structures were refined at full DFT level following the cluster method proposed by Siegbahn and Himo,²⁸ and QM binding energy were computed; finally, EPR spin Hamiltonian parameters were calculated for the predicted sites. From an energetic point of view, both sites show negative $\Delta E_{\text{binding}}$ values, but that with (His161, Asp154) is favored by more than 15 kcal·mol⁻¹. Its $|A_z|_{\text{calcd}}$ ($178.6 \times 10^{-4} \text{ cm}^{-1}$) matches well the experimental value of $177.1 \times 10^{-4} \text{ cm}^{-1}$ measured by Ramos *et al.*¹⁴ Moreover, $|A_z|$, estimated with the "additivity relationship"^{55, 56}, is $178.4 \times 10^{-4} \text{ cm}^{-1}$, closer to the experimental one than that expected for (His 73, His161) mode; the contributions of $45.1 \times 10^{-4} \text{ cm}^{-1}$ for His-N (derived from the equation $42.72 + 2.96 \times \sin[2\theta - 90]$,⁵⁵ θ being the calculated dihedral angle O= V^{IV} -N-C where C is the carbon atom bridging the two nitrogens), $42.1 \times 10^{-4} \text{ cm}^{-1}$ for Asp-COO⁻ and $45.6 \times 10^{-4} \text{ cm}^{-1}$ for water-O were used to determine $|A_z|_{\text{estmtd}}$.^{55, 56, 57}

These findings fortify the hypothesis that the binding mode (His161, Asp73), in the internal region of the site α , is the best candidate to explain the experimental EPR data collected so far. In

fact, the simultaneous binding of $V^{IV}O^{2+}$ and Ca^{2+} -ATP results hindered for the steric clashes and electrostatic repulsion into the enzymatic cleft (Figure 6c).

(4) Competition between V_{10} and ATP or Nb_{10} for the binding sites. Some years ago Aureliano *et al.* showed that, while G-actin increases the stability of V_{10} in solution, the addition of ATP to the medium has the opposite effect decreasing its half-life from 27 to 10 h (Figure S4 of the Supporting Information).¹¹ This insight suggested, for the first time, that V_{10} and ATP could compete for the same binding site, being ATP more affine to its natural site and able to partially displace V_{10} to a less protective and exposed binding region.

The possible competition between V_{10} with ATP for the nucleotide site was further evidenced in 2009, by EPR and ^{51}V NMR spectroscopy.¹³ The EPR spectra recorded on frozen aqueous solution containing increasing ATP/ V_{10} ratios in the presence and absence of G-actin indicated that V_{10} induces actin cysteine oxidation and formation of paramagnetic oxidovanadium(IV) ion and both the processes are prevented by the addition of ATP in solution (trace c of Figure 5). ^{51}V NMR spectra recorded at room temperature on the same systems confirmed this finding and showed that it is necessary to increase the ATP concentration up to 5 mM to broaden the V_{10} signal.¹³ Moreover, the major alterations in protein structure suggested by ^1H NMR spectra of G-actin treated with V_{10} (Figure S3 of the Supporting Information), which are not or hardly visible in the system with ATP, confirmed the competition between nucleotide and decavanadate in the critical binding site.⁵⁰

CD spectra show that G-actin is less stable in the absence (melting point, T_m , of 44.8 °C) than in the presence of ATP (T_{m1} 48.9 and T_{m2} 64.3 °C, *vide supra*), whose interaction results in two melting points which were extracted from recorded melting curves. Finally, recent EXAFS and XANES experiments established that the interaction of V_{10} with G-actin triggers a protein conformational reorientation that causes an oxidation of Cys residues and formation of $V^{IV}O^{2+}$. The

variations induced in V_{10} 's EXAFS profile in the presence of ATP suggest that the nucleotide and V_{10} compete for the same binding site (see the EXAFS profile in Figure 2).¹⁵

In order to deduce if the native actin substrate (ATP) interferes with V_{10} binding and to rationalize the experimental evidences (*vide supra*), additional MD simulations were carried out. ATP is found at the enzymatic cleft of G-actin in correspondence of the site α and, in principle, could compete with V_{10} for this region. From a first MD of 200 ns, the binding of Ca^{2+} -ATP to the site α is highly stable along the whole trajectory. The ΔG_{bind} of Ca^{2+} -ATP to its biological site α is the lowest of the series and is more favorable by 14.1 kcal·mol⁻¹ than V_{10} .

It must be highlighted that the catalytic cleft is more closed than the apo or POMs loaded forms with an angle ϕ of 22.0° (cfr. with the data in Table 1). This observation is coherent with ¹H NMR spectra of G-actin recorded in the presence and absence of V_{10} . Major alteration patterns attributable to a protein conformational change are negligible upon addition of ATP (Figure S3 of the Supporting Information).⁵¹ Therefore, ATP could displace V_{10} from the site α causing the closure of the cleft to its natural conformation and giving the normal pattern observed in ¹H NMR experiments.

Aimed to prove this hypothesis, further MDs were carried out. On the most sampled structure of the trajectory with Ca^{2+} -ATP bound to the site α , a subsequent docking of V_{10} was carried out to evaluate if the contemporaneous binding of V_{10} and ATP at the site α is possible. Additional 200 ns of MD unveil that the presence of ATP hinders the V_{10} binding; indeed, at 9 ns, V_{10} leaves the site α and moves to the site β after 53 ns, where it is stabilized (Figure 7).




Figure 7. Molecular representation of: a) the contemporaneous binding of V₁₀ at the site **α** and Nb₁₀ at the site **β**; b) the binding mode of Nb₁₀ at the site **β** in the presence of Ca²⁺-ATP.

These data can be rationalized assuming that, in the case of V₁₀, the effect of ATP is of clear inhibition. The catalytic site becomes populated preferentially by ATP with concomitant closure of the cleft which, even for the increased negative charge in the region, induces the displacement of V₁₀ from the site **α** to the site **β** less affine of *ca.* 23 kcal·mol⁻¹ (see Table 1).

As discussed above, the site **β** is found at the interface of subdomains I and II in the surface region of G-actin (Figure 1a). Therefore, the reason behind the increasing of the half-life of V₁₀ bound to G-actin from 5 to 27 h and its decreasing up to 10 h upon addition of ATP,¹¹ should be found on the ATP induced displacement of V₁₀ from the protective environment of the site **α** to the more solvent exposed site **β**. This event may favor the Cys oxidation and V₁₀ reduction to oxidovanadium(IV) cation, V^{IV}O²⁺, and the aggregation of G-actin to form F-actin observed in recent EXAFS and XANES study.¹⁵

Concerning Nb₁₀, being site **α** its secondary site, the competition with ATP is less important, suggesting no competition between the two POMs (Figure 7). This finding allows to establish a different behavior with respect to what is observed for the binding with Ca²⁺-ATPase. In that case,

Nb₁₀ prevents the binding of V₁₀, indicating that both isostructural POMs share the same primary binding site in the protein structure.¹⁸ Moreover, both of them cause comparable conformational changes.¹⁹

Conclusions

This study represents an unusual attempt to rationalize the experimental data – collected in the last 15 years – between the interaction of a POM, decavanadate(V) or V₁₀, and a very important cellular protein, G-actin, through computational methods. In fact, despite various instrumental techniques can ascertain the binding between a potential metallodrug and a protein, nevertheless they often cannot provide the description of this interaction at molecular level. For the system V₁₀–G-actin, the binding sites and interacting residues are not known nor the reason of the competition with other molecules of the organism, for example the physiological ligand ATP. In this context, molecular modelling represents a valid tool to understand these processes, providing new information for the comprehension of the molecular events governing POMs–protein and, in general, metallodrugs–protein binding).

For a careful application of POMs in biology and medicine there is a great need for understanding which cellular proteins represent the potential targets and often POMs–proteins interaction did not get the attention it deserves. Specifically, regarding POMs–actin systems described in the present study, the computational approach has proven to be an excellent tool to determine the binding sites and understand the different V₁₀ and Nb₁₀ mode of action. Summarizing the obtained results, it was elucidated and compared at the molecular level the behaviour of Nb₁₀ and V₁₀ in the G-actin conformational rearrangements. Two putative V₁₀/Nb₁₀–actin binding sites were fully characterised: V₁₀ prefers the catalytic nucleotide site (site α), whereas Nb₁₀ is more stable at the site β . Therefore, in contrast with other proteins such as Ca²⁺-ATPase, they could contemporaneously bind to G-actin and their biological and/or pharmacological action could

be synergistic. These findings should be considered for other studies in inorganic medicinal chemistry, administering POMs or other potential metallodrugs compatible with the same target protein, i.e. with affinity for different sites, to amplify their action. Concerning the presence of the native actin ligand ATP, it prevents the V_{10} binding to the site α , whereas it does not affect Nb_{10} –actin interaction, since β is the primary site for this species.

We believe that the present paper will facilitate the deeper understanding of the role POMs in various applications, giving valuable insight for a rational design of new metal-substituted POMs, i.e. with different metals in their structure, with specific interaction properties toward the putative proteins targets. In addition, it could be used as a "guide" for the study of the interactions (metal compounds)–(cellular proteins) and of the phenomena of inhibition and competition with the biomolecules at physiological conditions.

Finally, an important outcome of this paper is that other metallodrug–protein systems, in the absence or presence of eventual inhibitors and/or competition with molecules of the organism, could be studied with the same approach, suggesting important elements to explain the biological data and design more active and specific potential drugs.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.inorgchem.xxxxxxx. Comparison of the DFT optimized structure of V_{10} and Nb_{10} (Figure S1), exchange rate and apparent dissociation constant for the exchange of bound ϵ -ATP of G-actin with ATP (Figure S2), ^1H NMR spectra of G-actin and V_{10} –G-actin (Figure S3), V_{10} half-life time in presence of G-actin and ATP (Figure S4), Cartesian coordinates of V_{10} and Nb_{10} .

Author Information

Corresponding Authors

* G.S.: phone, +34 977 920229; E-mail, gsciortino@iciq.es

* M.A.: phone, +351 289 800977 ; E-mail, maalves@ualg.pt

* E.G.: phone, +39 079 229487; E-mail, garribba@uniss.it

ORCID

Giuseppe Sciortino: 0000-0001-9657-1788

Manuel Aureliano: 0000-0003-4858-3201

Eugenio Garribba: 0000-0002-7229-5966

Notes

The authors declare no competing financial interest.

Acknowledgments

E.G. and G.S. thank Regione Autonoma della Sardegna (grant RASSR79857) and Università di Sassari (fondo di Ateneo per la ricerca 2020). M.A. is grateful to Fundação para a Ciência e a Tecnologia (FCT) project UIDB/04326/2020.

References

- (1) (a) Barry, N. P. E.; Sadler, P. J. Exploration of the medical periodic table: towards new targets. *Chem. Commun.* **2013**, 49, 5106-5131. (b) Mjos, K. D.; Orvig, C. Metallodrugs in Medicinal Inorganic Chemistry. *Chem. Rev.* **2014**, 114, 4540-4563. (c) Medici, S.; Peana, M.; Nurchi, V. M.; Lachowicz, J. I.; Crisponi, G.; Zoroddu, M. A. Noble metals in medicine: Latest advances. *Coord. Chem. Rev.* **2015**, 284, 329-350. (d) Zhang, P.; Sadler, P. J. Advances in the design of organometallic anticancer complexes. *J. Organomet. Chem.* **2017**, 839, 5-14.
- (2) (a) Jones, C.; Thornback, J. *Medicinal Applications of Coordination Chemistry*. The Royal Society of Chemistry: Cambridge, 2007. (b) Dabrowiak, J. C. *Metals in Medicine*. John Wiley & Sons, Ltd: Chichester, 2009. (c) *Bioinorganic Medicinal Chemistry*. Alessio, E., Ed. Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, 2011. (d) *Interrelations between Essential Metal Ions and Human Diseases*. Springer Science+Business Media: Dordrecht, 2013. (e) *Metal-based Anticancer Agents*. Casini, A.; Vessières, A.; Meier-Menches, S. M., Eds. The Royal Society of Chemistry: Croydon, 2019.
- (3) (a) Gambino, D. Potentiality of vanadium compounds as anti-parasitic agents. *Coord. Chem. Rev.* **2011**, 255, 2193-2203. (b) Kioseoglou, E.; Petanidis, S.; Gabriel, C.; Salifoglou, A. The chemistry and biology of vanadium compounds in cancer therapeutics. *Coord. Chem. Rev.* **2015**, 301-302, 87-105. (c) Costa Pessoa, J.; Etcheverry, S.; Gambino, D. Vanadium compounds in medicine. *Coord. Chem. Rev.* **2015**, 301-302, 24-48. (d) Rehder, D. Perspectives for vanadium in health issues. *Future Med. Chem.* **2016**, 8, 325-338. (e) Rehder, D. Vanadium in health issues. *ChemTexts* **2018**, 4, 20. (f) Crans, D., C.; Yang, L.; Haase, A.; Yang, X. Health Benefits of Vanadium and Its Potential as an Anticancer Agent. In *Metallo-Drugs: Development and Action of Anticancer Agents*; Sigel, A., Sigel, H., Freisinger, E., Sigel, R. K. O., Ed.; De Gruyter GmbH: Berlin, 2018; Vol. 18, pp 251-280. (g) Crans, D. C.; Henry, L.; Cardiff, G.; Posner, B. I. Developing Vanadium as Antidiabetic and Anticancer Drugs: A Clinical and Historical Perspective In *Essential Metals in Medicine: Therapeutic Use and Toxicity of Metal Ions in the Clinic*; Carver, P. L., Ed.; De Gruyter GmbH: Berlin, 2019; pp 203-230.
- (4) (a) Levina, A.; Crans, D. C.; Lay, P. A. Speciation of metal drugs, supplements and toxins in media and bodily fluids controls in vitro activities. *Coord. Chem. Rev.* **2017**, 352, 473-498. (b) Doucette, K. A.; Hassell, K. N.; Crans, D. C. Selective speciation improves efficacy and lowers toxicity of platinum anticancer and vanadium antidiabetic drugs. *J. Inorg. Biochem.* **2016**, 165, 56-70.
- (5) (a) Rhule, J. T.; Hill, C. L.; Judd, D. A.; Schinazi, R. F. Polyoxometalates in Medicine. *Chem. Rev.* **1998**, 98, 327-358. (b) Yamase, T. Anti-tumor, -viral, and -bacterial activities of polyoxometalates for realizing an inorganic drug. *J. Mater. Chem.* **2005**, 15, 4773-4782. (c) Turner, T. L.; Nguyen, V. H.; McLauchlan, C. C.; Dymon, Z.; Dorsey, B. M.; Hooker, J. D.; Jones, M. A. Inhibitory effects of decavanadate on several enzymes and *Leishmania tarentolae* In Vitro. *J. Inorg.*

- Biochem.* **2012**, *108*, 96-104. (d) Treviño, S.; Velázquez-Vázquez, D.; Sánchez-Lara, E.; Diaz-Fonseca, A.; Flores-Hernandez, J. Á.; Pérez-Benítez, A.; Brambila-Colombres, E.; González-Vergara, E. Metforminium Decavanadate as a Potential Metallopharmaceutical Drug for the Treatment of Diabetes Mellitus. *Oxidative Med. Cell. Longev.* **2016**, *2016*, 6058705. (e) Bijelic, A.; Aureliano, M.; Rompel, A. The antibacterial activity of polyoxometalates: structures, antibiotic effects and future perspectives. *Chem. Commun.* **2018**, *54*, 1153-1169. (f) Qi, Y.; Han, L.; Qi, Y.; Jin, X.; Zhang, B.; Niu, J.; Zhong, J.; Xu, Y. Anti-flavivirus activity of polyoxometalate. *Antiviral Res.* **2020**, *179*, 104813. (g) Gu, Y.; Li, Q.; Huang, Y.; Zhu, Y.; Wei, Y.; Ruhlmann, L. Polyoxovanadate-iodobodipy supramolecular assemblies: new agents for high efficiency cancer photochemotherapy. *Chem. Commun.* **2020**, *56*, 2869-2872.
- (6) Althumairy, D.; Postal, K.; Barisas, B. G.; Nunes, G. G.; Roess, D. A.; Crans, D. C. Polyoxometalates function as indirect activators of a G protein-coupled receptor. *Metallomics* **2020**, *12*, 1044-1061.
- (7) (a) Bijelic, A.; Rompel, A. The use of polyoxometalates in protein crystallography – An attempt to widen a well-known bottleneck. *Coord. Chem. Rev.* **2015**, *299*, 22-38. (b) Chen, X.; Yan, S.; Wang, H.; Hu, Z.; Wang, X.; Huo, M. Aerobic oxidation of starch catalyzed by isopolyoxovanadate $\text{Na}_4\text{Co}(\text{H}_2\text{O})_6\text{V}_{10}\text{O}_{28}$. *Carbohydr. Polym.* **2015**, *117*, 673-680. (c) Zhou, H.; Li, J.; Bao, S.; Li, J.; Liu, X.; Jin, P. Use of ZnO as antireflective, protective, antibacterial, and biocompatible multifunction nanolayer of thermochromic VO_2 nanofilm for intelligent windows. *Appl. Surf. Sci.* **2016**, *363*, 532-542.
- (8) Nyman, M.; Martin, N. P. Directional bonding in $\{\text{Nb}_{10}\}$ inorganic frameworks. *Angew. Chem., Int. Ed.* **2020**, DOI: 10.1002/anie.202010902.
- (9) (a) Gândara, R. M. C.; Soares, S. S.; Martins, H.; Gutiérrez-Merino, C.; Aureliano, M. Vanadate oligomers: In vivo effects in hepatic vanadium accumulation and stress markers. *J. Inorg. Biochem.* **2005**, *99*, 1238-1244. (b) Soares, S. S.; Martins, H.; Duarte, R. O.; Moura, J. J. G.; Coucelo, J.; Gutiérrez-Merino, C.; Aureliano, M. Vanadium distribution, lipid peroxidation and oxidative stress markers upon decavanadate in vivo administration. *J. Inorg. Biochem.* **2007**, *101*, 80-88. (c) Tiago, D. M.; Laizé, V.; Cancela, M. L.; Aureliano, M. Impairment of mineralization by metavanadate and decavanadate solutions in a fish bone-derived cell line. *Cell Biol. Toxicol.* **2008**, *24*, 253-263. (d) Soares, S. S.; Martins, H.; Gutiérrez-Merino, C.; Aureliano, M. Vanadium and cadmium in vivo effects in teleost cardiac muscle: Metal accumulation and oxidative stress markers. *Comp. Biochem. Physiol., Part C: Toxicol. Pharmacol.* **2008**, *147*, 168-178. (e) Aureliano, M.; Fraqueza, G.; Ohlin, C. A. Ion pumps as biological targets for decavanadate. *Dalton Trans.* **2013**, *42*, 11770-11777. (f) Aureliano, M.; Ohlin, C. A. Decavanadate in vitro and in vivo effects: facts and opinions. *J. Inorg. Biochem.* **2014**, *137*, 123-130. (g) Gumerova, N.; Krivosudský, L.; Fraqueza, G.; Breibeck, J.; Al-Sayed, E.; Tanuhadi, E.; Bijelic, A.; Fuentes, J.; Aureliano, M.; Rompel, A. The P-type ATPase inhibiting potential of polyoxotungstates. *Metallomics* **2018**, *10*, 287-295. (h) Marques-da-Silva, D.; Fraqueza, G.; Lagoa, R.; Vannathan, A. A.; Mal, S. S.; Aureliano, M. Polyoxovanadate inhibition of Escherichia coli growth shows a reverse correlation

with Ca^{2+} -ATPase inhibition. *New J. Chem.* **2019**, 43, 17577-17587. (i) Fraqueza, G.; Fuentes, J.; Krivosudský, L.; Dutta, S.; Mal, S. S.; Roller, A.; Giester, G.; Rompel, A.; Aureliano, M. Inhibition of Na^+/K^+ - and Ca^{2+} -ATPase activities by phosphotetradecavanadate. *J. Inorg. Biochem.* **2019**, 197, 110700. (j) Pimpão, C.; da Silva, I. V.; Mósca, A. F.; Pinho, J. O.; Gaspar, M. M.; Gumerova, N. I.; Rompel, A.; Aureliano, M.; Soveral, G. The Aquaporin-3-Inhibiting Potential of Polyoxotungstates. *Int. J. Mol. Sci.* **2020**, 21, 2467.

(10) (a) Molitor, C.; Bijelic, A.; Rompel, A. In situ formation of the first proteinogenically functionalized $[\text{TeW}_6\text{O}_{24}\text{O}_2(\text{Glu})]^{7-}$ structure reveals unprecedented chemical and geometrical features of the Anderson-type cluster. *Chem. Commun.* **2016**, 52, 12286-12289. (b) Molitor, C.; Bijelic, A.; Rompel, A. The potential of hexatungstotellurate(VI) to induce a significant entropic gain during protein crystallization. *IUCrJ* **2017**, 4, 734-740.

(11) Ramos, S.; Manuel, M.; Tiago, T.; Duarte, R.; Martins, J.; Gutiérrez-Merino, C.; Moura, J. J. G.; Aureliano, M. Decavanadate interactions with actin: Inhibition of G-actin polymerization and stabilization of decameric vanadate. *J. Inorg. Biochem.* **2006**, 100, 1734-1743.

(12) Stevenson, R. P.; Veltman, D.; Machesky, L. M. Actin-bundling proteins in cancer progression at a glance. *J. Cell Sci.* **2012**, 125, 1073-1079.

(13) Ramos, S.; Duarte, R. O.; Moura, J. J. G.; Aureliano, M. Decavanadate interactions with actin: cysteine oxidation and vanadyl formation. *Dalton Trans.* **2009**, 7985-7994.

(14) Ramos, S.; Moura, J. J. G.; Aureliano, M. Actin as a potential target for decavanadate. *J. Inorg. Biochem.* **2010**, 104, 1234-1239.

(15) Marques, M. P. M.; Gianolio, D.; Ramos, S.; Batista de Carvalho, L. A. E.; Aureliano, M. An EXAFS Approach to the Study of Polyoxometalate-Protein Interactions: The Case of Decavanadate-Actin. *Inorg. Chem.* **2017**, 56, 10893-10903.

(16) (a) Sciortino, G.; Sanna, D.; Ugone, V.; Micera, G.; Lledós, A.; Maréchal, J.-D.; Garribba, E. Elucidation of Binding Site and Chiral Specificity of Oxidovanadium Drugs with Lysozyme through Theoretical Calculations. *Inorg. Chem.* **2017**, 56, 12938-12951. (b) Sciortino, G.; Rodríguez-Guerra Pedregal, J.; Lledós, A.; Garribba, E.; Maréchal, J.-D. Prediction of the interaction of metallic moieties with proteins: an update for protein-ligand docking techniques. *J. Comput. Chem.* **2018**, 39, 42-51. (c) Sciortino, G.; Sanna, D.; Ugone, V.; Lledós, A.; Maréchal, J.-D.; Garribba, E. Decoding Surface Interaction of $\text{V}^{\text{IV}}\text{O}$ Metallodrug Candidates with Lysozyme. *Inorg. Chem.* **2018**, 57, 4456-4469. (d) Sanna, D.; Ugone, V.; Sciortino, G.; Buglyo, P.; Bihari, Z.; Parajdi-Losoncz, P. L.; Garribba, E. $\text{V}^{\text{IV}}\text{O}$ complexes with antibacterial quinolone ligands and their interaction with serum proteins. *Dalton Trans.* **2018**, 47, 2164-2182. (e) Sciortino, G.; Garribba, E.; Maréchal, J.-D. Validation and Applications of Protein-Ligand Docking Approaches Improved for Metalloligands with Multiple Vacant Sites. *Inorg. Chem.* **2019**, 58, 294-306. (f) Sciortino, G.; Sanna, D.; Ugone, V.; Marechal, J. D.; Garribba, E. Integrated ESI-MS/EPR/computational characterization of the binding of metal species to proteins: vanadium drug-myoglobin application. *Inorg. Chem. Front.* **2019**, 6, 1561-1578. (g) Ugone, V.; Sanna, D.; Sciortino, G.; Marechal, J. D.; Garribba, E. Interaction of Vanadium(IV) Species with Ubiquitin: A Combined Instrumental and

- Computational Approach. *Inorg. Chem.* **2019**, *58*, 8064-8078. (h) Sciortino, G.; Sanna, D.; Ugone, V.; Maréchal, J.-D.; Alemany-Chavarria, M.; Garribba, E. Effect of secondary interactions, steric hindrance and electric charge on the interaction of V^{IV}O species with proteins. *New J. Chem.* **2019**, *43*, 17647-17660.
- (17) Aureliano, M.; Ohlin, C. A.; Vieira, M. O.; Marques, M. P. M.; Casey, W. H.; Batista de Carvalho, L. A. E. Characterization of decavanadate and decaniobate solutions by Raman spectroscopy. *Dalton Trans.* **2016**, *45*, 7391-7399.
- (18) Fraqueza, G.; Ohlin, C. A.; Casey, W. H.; Aureliano, M. Sarcoplasmic reticulum calcium ATPase interactions with decaniobate, decavanadate, vanadate, tungstate and molybdate. *J. Inorg. Biochem.* **2012**, *107*, 82-89.
- (19) Fraqueza, G.; Batista de Carvalho, L. A. E.; Marques, M. P. M.; Maia, L.; Ohlin, C. A.; Casey, W. H.; Aureliano, M. Decavanadate, decaniobate, tungstate and molybdate interactions with sarcoplasmic reticulum Ca²⁺-ATPase: quercetin prevents cysteine oxidation by vanadate but does not reverse ATPase inhibition. *Dalton Trans.* **2012**, *41*, 12749-12758.
- (20) López, X.; Carbó, J. J.; Bo, C.; Poblet, J. M. Structure, properties and reactivity of polyoxometalates: a theoretical perspective. *Chem. Soc. Rev.* **2012**, *41*, 7537-7571.
- (21) (a) Solé-Daura, A.; Poblet, J. M.; Carbó, J. J. Structure–Activity Relationships for the Affinity of Chaotropic Polyoxometalate Anions towards Proteins. *Chem.–Eur. J.* **2020**, *26*, 5799-5809. (b) Solé-Daura, A.; Goovaerts, V.; Stroobants, K.; Absillis, G.; Jiménez-Lozano, P.; Poblet, J. M.; Hirst, J. D.; Parac-Vogt, T. N.; Carbó, J. J. Probing Polyoxometalate–Protein Interactions Using Molecular Dynamics Simulations. *Chem.–Eur. J.* **2016**, *22*, 15280-15289.
- (22) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M. L., X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, J. A., Jr.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Keith, T.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C. J., J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. *Gaussian 09, revision D.01*. Gaussian, Inc.: Wallingford, CT, 2010.
- (23) Grimme, S. Semiempirical GGA-type density functional constructed with a long-range dispersion correction. *J. Comput. Chem.* **2006**, *27*, 1787-1799.
- (24) Ehlers, A. W.; Böhme, M.; Dapprich, S.; Gobbi, A.; Höllwarth, A.; Jonas, V.; Köhler, K. F.; Stegmann, R.; Veldkamp, A.; Frenking, G. A set of f-polarization functions for pseudo-potential basis sets of the transition metals Sc-Cu, Y-Ag and La-Au. *Chem. Phys. Lett.* **1993**, *208*, 111-114.
- (25) Marenich, A. V.; Cramer, C. J.; Truhlar, D. G. Universal Solvation Model Based on Solute

Electron Density and on a Continuum Model of the Solvent Defined by the Bulk Dielectric Constant and Atomic Surface Tensions. *J. Phys. Chem. B* **2009**, *113*, 6378-6396.

(26) Bayly, C. I.; Cieplak, P.; Cornell, W.; Kollman, P. A. A well-behaved electrostatic potential based method using charge restraints for deriving atomic charges: the RESP model. *J. Phys. Chem.* **1993**, *97*, 10269-10280.

(27) (a) Cabré, A.; Sciortino, G.; Ujaque, G.; Verdaguer, X.; Lledós, A.; Riera, A. Iridium-Catalyzed Isomerization of N-Sulfonyl Aziridines to Allyl Amines. *Org. Lett.* **2018**, *20*, 5747-5751.

(b) Salomó, E.; Gallen, A.; Sciortino, G.; Ujaque, G.; Grabulosa, A.; Lledós, A.; Riera, A.; Verdaguer, X. Direct Asymmetric Hydrogenation of N-Methyl and N-Alkyl Imines with an Ir(III)H Catalyst. *J. Am. Chem. Soc.* **2018**, *140*, 16967-16970. (c) Alonso-Cotchico, L.; Sciortino, G.; Vidossich, P.; Rodríguez-Guerra Pedregal, J.; Drienovská, I.; Roelfes, G.; Lledós, A.; Maréchal, J.-D. Integrated Computational Study of the Cu-Catalyzed Hydration of Alkenes in Water Solvent and into the Context of an Artificial Metallohydratase. *ACS Catalysis* **2019**, *9*, 4616-4626.

(28) Siegbahn, P. E. M.; Himo, F. The quantum chemical cluster approach for modeling enzyme reactions. *Wiley Interdiscip. Rev. Comput. Mol. Sci.* **2011**, *1*, 323-336.

(29) (a) Sciortino, G.; Garribba, E. The binding modes of $V^{IV}O^{2+}$ ions in blood proteins and enzymes. *Chem. Commun.* **2020**, *56*, 12218-12221. (b) Sciortino, G.; Sanna, D.; Lubinu, G.; Maréchal, J.-D.; Garribba, E. Unveiling $V^{IV}O^{2+}$ Binding Modes to Human Serum Albumins by an Integrated Spectroscopic–Computational Approach. *Chem.–Eur. J.* **2020**, *26*, 11316-11326.

(30) Micera, G.; Garribba, E. The effect of the functional, basis set, and solvent in the simulation of the geometry and spectroscopic properties of $V^{IV}O^{2+}$ complexes. Chemical and biological applications. *Int. J. Quantum Chem.* **2012**, *112*, 2486-2498.

(31) (a) Gorelsky, S.; Micera, G.; Garribba, E. Equilibrium Between the Octahedral and Square Pyramidal Form and the Influence of an Axial Ligand on the Molecular Properties of $V^{IV}O$ Complexes: A Spectroscopic and DFT Study. *Chem.–Eur. J.* **2010**, *16*, 8167-8180. (b) Sanna, D.; Pecoraro, V. L.; Micera, G.; Garribba, E. Application of DFT methods to the study of the coordination environment of the VO^{2+} ion in V proteins. *J. Biol. Inorg. Chem.* **2012**, *17*, 773-790.

(32) (a) Remenyi, C.; Reviakine, R.; Arbuznikov, A. V.; Vaara, J.; Kaupp, M. Spin–Orbit Effects on Hyperfine Coupling Tensors in Transition Metal Complexes Using Hybrid Density Functionals and Accurate Spin–Orbit Operators. *J. Phys. Chem. A* **2004**, *108*, 5026-5033. (b) Sanna, D.; Sciortino, G.; Ugone, V.; Micera, G.; Garribba, E. Nonoxido V^{IV} Complexes: Prediction of the EPR Spectrum and Electronic Structure of Simple Coordination Compounds and Amavadin. *Inorg. Chem.* **2016**, *55*, 7373-7387.

(33) Micera, G.; Garribba, E. Is the Spin-Orbit Coupling Important in the Prediction of the ^{51}V Hyperfine Coupling Constants of $V^{IV}O^{2+}$ Species? ORCA Versus Gaussian Performance and Biological Applications. *J. Comput. Chem.* **2011**, *32*, 2822-2835.

(34) Šali, A.; Blundell, T. L. Comparative Protein Modelling by Satisfaction of Spatial Restraints. *J. Mol. Biol.* **1993**, *234*, 779-815.

(35) Wang, H.; Robinson, R. C.; Burtnick, L. D. The structure of native G-actin. *Cytoskeleton*

2010, 67, 456-465.

- (36) Case, D. A.; Betz, R. M.; Cerutti, D. S.; Cheatham III, T. E.; Darden, T. A.; Duke, R. E.; Giese, T. J.; Gohlke, H.; Goetz, A. W.; Homeyer, N.; Izadi, S.; Janowski, P.; Kaus, J.; Kovalenko, A.; Lee, T. S.; LeGrand, S.; Li, P.; C. Lin; Luchko, T.; Luo, R.; Madej, B.; Mermelstein, D.; Merz, K. M.; Monard, G.; Nguyen, H.; Nguyen, H. T.; Omelyan, I.; Onufriev, A.; Roe, D. R.; Roitberg, A.; Sagui, C.; Simmerling, C. L.; Botello-Smith, W. M.; Swails, J.; Walker, R. C.; Wang, J.; Wolf, R. M.; Wu, X.; Xiao, L.; Kollman, P. A. *AMBER 2016*. University of California, San Francisco, CA, 2016.
- (37) Hornak, V.; Abel, R.; Okur, A.; Strockbine, B.; Roitberg, A.; Simmerling, C. Comparison of multiple Amber force fields and development of improved protein backbone parameters. *Proteins: Struct., Funct., Bioinf.* **2006**, 65, 712-725.
- (38) Li, P.; Merz, K. M. MCPB.py: A Python Based Metal Center Parameter Builder. *J. Chem. Inf. Model.* **2016**, 56, 599-604.
- (39) Meagher, K. L.; Redman, L. T.; Carlson, H. A. Development of polyphosphate parameters for use with the AMBER force field. *J. Comput. Chem.* **2003**, 24, 1016-1025.
- (40) Bradbrook, G. M.; Gleichmann, T.; Harrop, S. J.; Habash, J.; Raftery, J.; Kalb, J.; Yariv, J.; Hillier, I. H.; Helliwell, J. R. X-Ray and molecular dynamics studies of concanavalin-A glucoside and mannoside complexes Relating structure to thermodynamics of binding. *J. Chem. Soc., Faraday Trans.* **1998**, 94, 1603-1611.
- (41) Eastman, P.; Swails, J.; Chodera, J. D.; McGibbon, R. T.; Zhao, Y.; Beauchamp, K. A.; Wang, L.-P.; Simmonett, A. C.; Harrigan, M. P.; Stern, C. D.; Wiewiora, R. P.; Brooks, B. R.; Pande, V. S. OpenMM 7: Rapid development of high performance algorithms for molecular dynamics. *PLoS Comput. Biol.* **2017**, 13, e1005659.
- (42) Rodríguez-Guerra Pedregal, J.; Alonso-Cotchico, L.; Velasco-Carneros, L.; Maréchal, J.-D. *OMMProtocol: A Command Line Application to Launch Molecular Dynamics Simulations with OpenMM*. 2018.
- (43) Sciortino, G.; Sánchez-Aparicio, J.-E.; Rodríguez-Guerra Pedregal, J.; Garribba, E.; Maréchal, J.-D. Computational insight into the interaction of oxaliplatin with insulin. *Metallomics* **2019**, 11, 765-773.
- (44) Grossfield, A.; Zuckerman, D. M. Quantifying uncertainty and sampling quality in biomolecular simulations. *Annu. Rep. Comput. Chem.* **2009**, 5, 23-48.
- (45) Smith, L. J.; Daura, X.; van Gunsteren, W. F. Assessing equilibration and convergence in biomolecular simulations. *Proteins: Struct., Funct., Bioinf.* **2002**, 48, 487-496.
- (46) González-Alemán, R.; Hernández-Castillo, D.; Caballero, J.; Montero-Cabrera, L. A. Quality Threshold Clustering of Molecular Dynamics: A Word of Caution. *J. Chem. Inf. Model.* **2020**, 60, 467-472.
- (47) Jones, G.; Willett, P.; Glen, R. C.; Leach, A. R.; Taylor, R. Development and validation of a genetic algorithm for flexible docking1. *J. Mol. Biol.* **1997**, 267, 727-748.
- (48) Sánchez-Aparicio, J.-E.; Tiessler-Sala, L.; Velasco-Carnero, L.; Roldán-Martín, L.;

- Sciortino, G.; Maréchal, J.-D. BioMetAll: Identifying Metal-Binding Sites in Proteins from Backbone Preorganization. *ChemRxiv* **2020**, DOI: 10.26434/chemrxiv.12668651.v1.
- (49) Genheden, S.; Ryde, U. The MM/PBSA and MM/GBSA methods to estimate ligand-binding affinities. *Expert Opin. Drug Deliv.* **2015**, *10*, 449-461.
- (50) Ramos, S.; Moura, J. J. G.; Aureliano, M. Recent advances into vanadyl, vanadate and decavanadate interactions with actin. *Metallomics* **2012**, *4*, 16-22.
- (51) Ramos, S.; Almeida, R. M.; Moura, J. J. G.; Aureliano, M. Implications of oxidovanadium(IV) binding to actin. *J. Inorg. Biochem.* **2011**, *105*, 777-783.
- (52) Aureliano, M.; Breibeck, J.; Puchinger, M.; Djinovic-Carugo, K.; Rompel, A. New insights into actin interactions of decavanadate. *2nd International Caparica Christmas Congress on Translational Chemistry, Lisbon, 2017*.
- (53) (a) Pezza, R. J.; Villarreal, M. A.; Montich, G. G.; Argaraña, C. E. Vanadate inhibits the ATPase activity and DNA binding capability of bacterial MutS. A structural model for the vanadate–MutS interaction at the Walker A motif. *Nucleic Acids Res.* **2002**, *30*, 4700-4708. (b) Tiago, T.; Martel, P.; Gutiérrez-Merino, C.; Aureliano, M. Binding modes of decavanadate to myosin and inhibition of the actomyosin ATPase activity. *Biochim. Biophys. Acta, Proteins Proteom.* **2007**, *1774*, 474-480.
- (54) Lassing, I.; Schmitzberger, F.; Björnstedt, M.; Holmgren, A.; Nordlund, P.; Schutt, C. E.; Lindberg, U. Molecular and Structural Basis for Redox Regulation of β -Actin. *J. Mol. Biol.* **2007**, *370*, 331-348.
- (55) (a) Smith II, T. S.; LoBrutto, R.; Pecoraro, V. L. Paramagnetic spectroscopy of vanadyl complexes and its applications to biological systems. *Coord. Chem. Rev.* **2002**, *228*, 1-18. (b) Smith II, T. S.; Root, C. A.; Kampf, J. W.; Rasmussen, P. G.; Pecoraro, V. L. Reevaluation of the Additivity Relationship for Vanadyl–Imidazole Complexes: Correlation of the EPR Hyperfine Constant with Ring Orientation. *J. Am. Chem. Soc.* **2000**, *122*, 767-775.
- (56) Chasteen, D. N. Vanadyl(IV) EPR spin probe. Inorganic and Biochemical Aspects. In *Biological Magnetic Resonance*; Berliner, L. J. J., Reuben, J., Ed.; Plenum Press: New York, 1981; Vol. 3, pp 53-119.
- (57) Garribba, E.; Lodyga-Chruscinska, E.; Micera, G.; Panzanelli, A.; Sanna, D. Binding of oxovanadium(IV) to dipeptides containing histidine and cysteine residues. *Eur. J. Inorg. Chem.* **2005**, 1369-1382.

TEXT FOR GRAPHICAL ABSTRACT.

Four putative binding sites in globular G-actin (named α - δ) were identified for two biologically active polyoxometalates (POMs), decavanadate(V) (V_{10}) and decaniobate(V) (Nb_{10}). Among the binding sites V_{10} prefers the catalytic nucleotide site (site α), while Nb_{10} the site β . ATP inhibits mostly the binding of V_{10} than Nb_{10} , competing with decavanadate for the site α .

GRAPHICAL ABSTRACT

