




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Nasal high flow oxygen therapy in COVID-19 patients does not cause environmental surface contamination

Oriol Roca, MD, PhD, Andrés Pacheco, MD, Jordi Rodon, BSc, Andrés Antón, PhD, Júlia Vergara-Alert, PhD, Lluís Armadans, MD PhD, Joaquim Segalés, PhD, Tomás Pumarola, MD PhD, Magda Campins, MD PhD, Ricard Ferrer, MD PhD, Virginia Rodríguez-Garrido, PhD



PII: S0195-6701(21)00216-4

DOI: <https://doi.org/10.1016/j.jhin.2021.04.034>

Reference: YJHIN 6414

To appear in: *Journal of Hospital Infection*

Received Date: 14 February 2021

Revised Date: 29 April 2021

Accepted Date: 30 April 2021

Please cite this article as: Roca O, Pacheco A, Rodon J, Antón A, Vergara-Alert J, Armadans L, Segalés J, Pumarola T, Campins M, Ferrer R, Rodríguez-Garrido V, Nasal high flow oxygen therapy in COVID-19 patients does not cause environmental surface contamination, *Journal of Hospital Infection*, <https://doi.org/10.1016/j.jhin.2021.04.034>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd on behalf of The Healthcare Infection Society.

Nasal high flow oxygen therapy in COVID-19 patients does not cause environmental surface contamination

Oriol Roca MD, PhD^{1,2}, Andrés Pacheco MD¹, Jordi Rodon BSc³, Andrés Antón PhD⁴, Júlia Vergara-Alert PhD³, Lluís Armadans MD PhD⁴, Joaquim Segalés PhD^{5,6}, Tomás Pumarola MD PhD⁴, Magda Campins MD PhD⁷, Ricard Ferrer MD PhD^{1,2}, Virginia Rodríguez-Garrido PhD⁴

¹Servei de Medicina Intensiva, Hospital Universitari Vall d'Hebron, Institut de Recerca Vall d'Hebron, Barcelona, Spain.

²Centro de Investigación Biomédica en Red de Enfermedades Respiratorias (CIBER), Madrid, Spain

³IRTA, Centre de Recerca en Sanitat Animal (CReSA, IRTA-UAB), Campus de la Universitat Autònoma de Barcelona, Barcelona, Spain.

⁴Servei de Microbiologia, Hospital Universitari Vall d'Hebron, Institut de Recerca Vall d'Hebron, Barcelona, Spain

⁵UAB, Centre de Recerca en Sanitat Animal (CReSA, IRTA-UAB), Campus de la Universitat Autònoma de Barcelona, Barcelona, Spain.

⁶Departament de Sanitat i Anatomia Animals, Facultat de Veterinària, UAB, 08193 Bellaterra (Cerdanyola del Vallès), Spain

⁷Servei de Medicina Preventiva, Hospital Universitari Vall d'Hebron, Barcelona, Spain

Corresponding author:

Oriol Roca, MD, PhD
Servei de Medicina Intensiva
Hospital Universitari Vall d'Hebron
Passeig de la Vall d'Hebron, 119-129
08035 Barcelona, Spain.
E-mail: oroca@vhebron.net

Declaration of Interests: None

To the Editor:

Nasal high flow oxygen therapy (NHF) may spread the severe acute respiratory syndrome coronavirus (SARS-CoV-2) virus through respiratory droplets and contact with fomites. This concern leads to significant controversies regarding recommendations of its use in different evidence-based guidelines of therapy of coronavirus infectious disease 2019 (COVID-19) patients¹. We, therefore, aimed to determine if NHF use for COVID-19 patients was associated with higher environmental surface SARS-CoV-2 contamination compared with those who were intubated and mechanically ventilated (MV).

We performed an observational, single-centre study initially including 10 consecutive critically ill COVID-19 patients who required ICU admission (5 NHF and 5 MV). All patients were admitted to ICU negative pressure single rooms with 12 air changes per hour. Closed-aspiration systems were used in mechanically ventilated patients. Patients with NHF only wore surgical masks above the NHF cannula when healthcare personnel were inside the room. The Ethics Committee of Vall d'Hebron University Hospital approved this study (PR(AG)225/2020), and the need for informed consent was waived due to the observational nature of the study.

Confirmation of SARS-CoV-2 persistence on environmental surfaces was based on the detection of viral genome by reverse-transcription quantitative polymerase chain reaction (RT-qPCR). Sample collection was performed within the first 24 hours of admission and after 3 days, before the daily room cleaning, from the following sites: bedside monitor, mechanical ventilator screen, computer keyboard, nurse medication trolley, infusion pumps, patient sheet, bed handle, personnel protective equipment (gloves, coat, protection glasses, hat), stethoscope, room floor at 1.5 metres distance from the patient, and a filter paper placed immediately before the room high efficiency particulate air (HEPA) filter. Environmental and equipment surfaces (approximately 5x5 cm) were sampled using moistened sterile swabs.

If all samples were negative, inclusion of two additional patients (one of each group [NHF and MV]) was planned. Two more samples were collected for these two additional patients. First, air sampling of room air was performed using a Sartorius MD8 microbiological sampler, containing a gelatine membrane filter for a fixed given sample volume of 1000l by placing the sampler near the patient (50 cm). After sampling, the filter was immediately sent to the lab where 3 ml of Dulbecco's modified Eagle medium (DMEM; Lonza, Allendale, NJ) was added and frozen at -80 °C until the analysis was performed. A second additional sample from the NHF cannula of the non-intubated patient was also obtained. The swabs were transferred immediately into 3 ml of DMEM, sent out to the microbiology laboratory and frozen at -80°C until RT-qPCR analyses were done. Total nucleic acids were extracted using NucliSENS EasyMAG (BioMérieux, Marcy l'Etoile, France) according to the manufacturer's instructions. Detection of SARS-CoV-2 genome was carried out by a commercial multiplex RT-qPCR assay (TaqMan™ 2019-nCoV Assay Kit v2, ThermoFisher, USA).

SARS-CoV-2 samples containing viral RNA, as determined by RT-qPCR, were assessed for the presence of infectious virus by titration in Vero E6 cells (ATCC® CRL384 1586™), as previously reported ^{2, 3}. Vero E6 cells were cultured in DMEM (Lonza) supplemented with 2% foetal calf serum (FCS; EuroClone), 100 U/mL penicillin (ThermoFisher Scientific, Life Technologies), 100 µg/ml streptomycin (ThermoFisher Scientific, Life Technologies), and 2 mM glutamine (ThermoFisher Scientific, Life Technologies).

Direct samples and serial ten-fold dilutions in supplemented DMEM were transferred to Vero E6 monolayers. Plates were monitored daily under a light microscope and wells were evaluated for the presence of cytopathic effect (CPE) at 6 days after infection. Results are expressed as median (25th percentile–75th percentile).

The main characteristics of the patients included in this study are summarized in Table 1. Two of the patients were initially included as MV patients (only have samples at day 1 of MV) and

subsequent samples were collected when they were supported with NHF after extubation (cases 1 and 2 among MV patients that correspond to cases 6 and 8 among NHF patients, respectively). A total of 252 environmental samples were collected from the first 10 patients. No differences between the time of positive nasopharyngeal swab collection and time of environmental sample collection were observed (median of 3 days [1-5] in NHF and 4 [3-9] in MV patients; $p = 0.222$). At the time of first collection, patients with NHF were supported with a median flow of 55 Lpm (50-60) and had a median respiratory rate of 23 (22-24) bpm and a SpO_2/FiO_2 ratio of 154 (145-192). The RT-qPCR reactions were negative for all the tested surfaces.

An additional 29 samples were obtained from two more patients. Only the sample collected from the NHF cannula was positive for SARS-CoV-2 RNA. This NHF cannula positive sample was inoculated to susceptible Vero E6 cells, but no CPE was observed after a 6-day observation period. Therefore, no infectious SARS-CoV-2 was found in any of the environmental samples collected in this study. However, the degree of surface and air contamination from SARS-CoV-2 in a previous study was highly heterogeneous⁴. The singularity of the present study is that it includes the largest data on the environmental persistence of SARS-CoV-2 in the rooms of COVID-19 patients being treated with NHF. Moreover, we compared the environmental contamination associated with NHF use to MV. Lack of environmental contamination in an ICU environment with negative pressure rooms was observed in both groups. Indeed, it has recently been shown that NHF by itself may not increase aerosol generation in healthy volunteers in a negative-pressure room⁵. Interestingly, despite it being impossible to place living SARS-CoV-2 on a surface to obtain a positive control for obvious safety reasons, we could obtain one positive RT-qPCR from the nasal cannula of one of the non-intubated patients suggesting that the sample collection and PCR technique was correctly performed. These

results were similar to the previous reported by Colaneri et al⁶ that obtained a unique positive PCR sample from inside the helmet of one non-intubated patient. However, The limitations of this study are that, first, air sampling was performed in only two patients and we cannot exclude that with increasing the litres of air sampling, we could likely obtain a positive sample. Secondly, the study was performed in ICU rooms with negative pressure and use of closed-aspiration systems in intubated patients and therefore its results may not be generalizable to other hospital environments. Despite these limitations, our results suggest that, with adequate infrastructure, NHF does not increase environmental surface contamination in critically ill patients with COVID-19.

Authors' Contributions:

OR, VRG, JVA contributed to the conception and design of the study, the acquisition, analysis and interpretation of the data and drafting the manuscript. AP, JR, contributed to the acquisition, analysis and interpretation of the data and drafting the manuscript. AA, LA, MC, JS, TP, RF, contributed to the conception and design of the study and interpretation of the data. All authors critically revised and approved the final version of the manuscript.

Conflicts of interest: OR reports speaker fees from Hamilton Medical, Ambu and Aerogen Ltd and research grant by Hamilton Medical. He also received non-financial research support from Timpel and Masimo Corporation in the last five years all outside the submitted work.

Funding:

This study has been funded by a grant of Instituto de Salud Carlos III of the Spanish Health Ministry (COV20-01054).

Journal Pre-proof

References

1. Raoof S, Nava S, Carpati C, Hill NS High-Flow, Noninvasive Ventilation and Awake (Nonintubation) Proning in Patients With Coronavirus Disease 2019 With Respiratory Failure. *Chest* 2020; **158**: 1992-2002.
2. Rodon J, Muñoz-Basagoiti J, Perez-Zsolt D *et al.* Identification of Plitidepsin as Potent Inhibitor of SARS-CoV-2-Induced Cytopathic Effect after a Drug Repurposing Screen. 2021; doi <https://doi.org/10.1101/2020.04.23.055756>.
3. Brustolin M, Rodon J, Rodríguez de la Concepción M *et al.* Protection against reinfection with D614- or G614-SARS-CoV-2 isolates in hamsters. 2021; doi <https://doi.org/10.1101/2021.01.07.425729>.
4. Birgand G, Peiffer-Smadja N, Fournier S, Kerneis S, Lescure FX, Lucet JC Assessment of Air Contamination by SARS-CoV-2 in Hospital Settings. *JAMA Netw Open* 2020; **3**: e2033232.
5. Gaeckle NT, Lee J, Park Y, Kreykes G, Evans MD, Hogan CJ, Jr. Aerosol Generation from the Respiratory Tract with Various Modes of Oxygen Delivery. *Am J Respir Crit Care Med* 2020; **202**: 1115-24.
6. Colaneri M, Seminari E, Novati S *et al.* Severe acute respiratory syndrome coronavirus 2 RNA contamination of inanimate surfaces and virus viability in a health care emergency unit. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases* 2020; **26**: 1094.e1-.e5.

Table 1. General characteristics of the population included

Case	Age	SAPS III	APACHE II	N° of quadrants in Chest-X- Ray	D-dimer (ng/ml)	IL-6 (mg/dL)	Type of room	Type of respiratory support	Days from positive RT-qPCR to first sample collection
1	37	23	19	4	171	100	ICU single-room with negative pressure	MV	3
2	56	49	18	2	6329	2359	ICU single-room with negative pressure	MV	3
3	49	13	22	4	221	195	ICU single-room with negative pressure	MV	9
4	55	30	22	4	3177	53	ICU single-room with negative pressure	MV	3
5	71	67	31	4	953	103	ICU single-room with negative pressure	MV	10
6	37	23	20	4	171	100	ICU single-room with negative pressure	NHF	1
7	44	32	21	2	225	108	ICU single-room with negative pressure	NHF	6
8	56	49	18	2	6329	2359	ICU single-room with negative pressure	NHF	5
9	54	25	22	4	1321	4541	ICU single-room with negative pressure	NHF	4
10	69	30	31	4	456	365	ICU single-room with negative pressure	NHF	1
11	72	23	26	2	170	22	Intermediate care, double-room	NHF	1
12	39	38	19	4	3319	2656	ICU single-room with negative pressure	MV	5

SAPS III: Simplified Acute Physiology Score; APACHE II: Acute Physiology and Chronic Health disease Classification System; IL: interleukin; MV: mechanical ventilation; NHF: nasal high flow; RT-qPCR: real-time reverse-transcription polymerase chain reaction.