
This is the **accepted version** of the journal article:

Villanueva, Yovani; Llibre, Jaume; Euzébio, Rodrigo. «Limit cycles of generic piecewise center-type vector fields in R^3 separated by either one plane or by two parallel planes». Bulletin des Sciences Mathématiques, Vol. 179 (October 2022), art. 103173. DOI 10.1016/j.bulsci.2022.103173

This version is available at <https://ddd.uab.cat/record/267140>

under the terms of the  license

Limit cycles of generic piecewise center-type vector fields in \mathbb{R}^3 separated by either one plane or by two parallel planes

Yovani Villanueva^a, Jaume Llibre^b, Rodrigo Euzébio^{a,*}

^a*Instituto de Matemática e Estatística, Universidade Federal de Goiás, R. Jacarandá, Campus Samambaia, 74690-900, Goiânia, GO, Brazil*

^b*Departament de Matemàtiques. Universitat Autònoma de Barcelona, Edifici C, 08193 Barcelona, Catalonia, Spain*

Abstract

While the limit cycles of the piecewise differential systems in the plane \mathbb{R}^2 have been studied intensively during these last twenty years, this is not the case for the limit cycles of the piecewise differential systems in the space \mathbb{R}^3 .

The goal of this article is to study the continuous and discontinuous piecewise differential systems in \mathbb{R}^3 , formed by linear vector fields similar to planar centers separated by one or two parallel planes. We call those “center-type” differential systems, which have two pure imaginary numbers and zero as eigenvalues. When these kinds of piecewise differential systems are continuous or discontinuous separated by one plane, then they have no limit cycles. Also, if they are continuous separated by two planes, then generically they do not have limit cycles. But when the piecewise differential systems are discontinuous separated two parallel planes, we show that generically they can have at most four limit cycles, and that there exist such systems with four limit cycles. The genericity here means that the statements hold in a residual set of the space of parameters associated to the differential system.

We recall that the same problem but for discontinuous piecewise differential systems in \mathbb{R}^2 formed by linear differential centers separated by two parallel

*Corresponding author

Email addresses: yovaniing@discente.ufg.br (Yovani Villanueva), jlllibre@mat.uab.cat (Jaume Llibre), euzebio@ufg.br (Rodrigo Euzébio)

straight lines have at most one limit cycle.

Keywords: Filippov vector fields, limit cycles, continuous piecewise vector fields, discontinuous piecewise vector fields.

2020 MSC: 37G15, 34C05, 34A36, 37M05

1. Introduction and statement of the main results

The study of piecewise vector fields (PVF) goes back to Andronov, Vitt, and Khaikin [1] and still continues to receive strong attention from researchers. These last years a renewed interest has appeared in the mathematical community working in differential equations for understanding the dynamical richness of the piecewise vector fields, because these vector fields are widely used to model processes appearing in electronics, mechanics, economy, etc., see for instance the books [2] and [3] and the survey of [4] and the hundreds of references quoted in these last three works.

10 In this paper, we shall work with piecewise vector fields in \mathbb{R}^2 and \mathbb{R}^3 , and the definition of these vector fields on the separation line of their pieces in \mathbb{R}^2 , or on the discontinuity region of their pieces in \mathbb{R}^3 follow the rules of Filippov [5], see a summary of these rules in Section 2.

These last two decades the limit cycles of the piecewise differential systems in the plane \mathbb{R}^2 have been studied intensively, see for instance [6–30]. This is not the case for the limit cycles of the piecewise differential systems in the space \mathbb{R}^3 .

A *center* of a differential system in the plane \mathbb{R}^2 is an equilibrium point \mathbf{x} having a neighbourhood U such that $U \setminus \{\mathbf{x}\}$ is filled of periodic orbits. A *global* center is a center \mathbf{x} such that $\mathbb{R}^2 \setminus \{\mathbf{x}\}$ is filled of periodic orbits. The notion of a center appeared already in the works of Poincaré [31] in 1881 and Dulac [32] in 1908.

One of the main objects in the study of the vector fields is the limit cycles. A *limit cycle* of a vector field is a periodic orbit isolated in the set of all periodic orbits of the vector field.

In the paper [28] it is proved that continuous and discontinuous piecewise vector fields in \mathbb{R}^2 formed by two pieces separated by one straight line and formed by two arbitrary linear centers cannot have limit cycles, and that also the continuous piecewise vector fields in \mathbb{R}^2 formed by three pieces separated
30 by two parallel straight lines and formed by three arbitrary linear centers have no limit cycles. But the discontinuous piecewise differential systems separated by two parallel straight lines and formed by three arbitrary linear centers can have at most one limit cycle, and there are such systems which one limit cycle. The objective of this paper is to study a similar problem for continuous and
35 discontinuous piecewise vector fields in \mathbb{R}^3 .

In \mathbb{R}^3 there are no centers in the sense that there are no equilibrium points \mathbf{x} having a neighborhood U such that $U \setminus \{\mathbf{x}\}$ is filled with periodic orbits, see for instance [33].

The main goal of this paper is to study the limit cycles of the continuous
40 and discontinuous piecewise vector fields in \mathbb{R}^3 separated by one plane or by two parallel planes and formed by linear “center-type” vector fields of \mathbb{R}^3 .

More precisely, we consider the linear vector field

$$X(\mathbf{x}) = \begin{pmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} + \begin{pmatrix} B_1 \\ B_2 \\ B_3 \end{pmatrix} = A\mathbf{x} + B, \quad (1)$$

where

$$\begin{aligned} A_{11} &= a_1 a_3 c_2 + b_1 b_3 c_2 - a_1 a_2 c_3 - b_1 b_2 c_3, \\ A_{12} &= a_2 a_3 c_2 + b_2 b_3 c_2 - a_2^2 c_3 - b_2^2 c_3, \\ A_{13} &= a_3^2 c_2 + b_3^2 c_2 - a_2 a_3 c_3 - b_2 b_3 c_3, \\ A_{21} &= -a_1 a_3 c_1 - b_1 b_3 c_1 + a_1^2 c_3 + b_1^2 c_3, \\ A_{22} &= -a_2 a_3 c_1 - b_2 b_3 c_1 + a_1 a_2 c_3 + b_1 b_2 c_3, \\ A_{23} &= -a_3^2 c_1 - b_3^2 c_1 + a_1 a_3 c_3 + b_1 b_3 c_3, \\ A_{31} &= a_1 a_2 c_1 + b_1 b_2 c_1 - a_1^2 c_2 - b_1^2 c_2, \\ A_{32} &= a_2^2 c_1 + b_2^2 c_1 - a_1 a_2 c_2 - b_1 b_2 c_2, \\ A_{33} &= a_2 a_3 c_1 + b_2 b_3 c_1 - a_1 a_3 c_2 - b_1 b_3 c_2, \end{aligned}$$

$$\begin{aligned}
B_1 &= a_3a_4c_2 + b_3b_4c_2 - a_2a_4c_3 - b_2b_4c_3, \\
B_2 &= -a_3a_4c_1 - b_3b_4c_1 + a_1a_4c_3 + b_1b_4c_3, \\
B_3 &= a_2a_4c_1 + b_2b_4c_1 - a_1a_4c_2 - b_1b_4c_2.
\end{aligned}$$

The vector field (1) is obtained after applying the affine transformation

$$(x, y, z) \mapsto (a_1x + a_2y + a_3z + a_4, b_1x + b_2y + b_3z + b_4, c_1x + c_2y + c_3z + c_4)$$

with inverse

$$\begin{aligned}
(x, y, z) \mapsto & \frac{-1}{\det(A)} (a_4b_3c_2 - a_3b_4c_2 - a_4b_2c_3 + a_2b_4c_3 + a_3b_2c_4 - a_2b_3c_4 \\
& - b_3c_2x + b_2c_3x + a_3c_2y - a_2c_3y - a_3b_2z + a_2b_3z, -a_4b_3c_1 \\
& + a_3b_4c_1 + a_4b_1c_3 - a_1b_4c_3 - a_3b_1c_4 + a_1b_3c_4 + b_3c_1x - b_1c_3x \\
& - a_3c_1y + a_1c_3y + a_3b_1z - a_1b_3z, a_4b_2c_1 - a_2b_4c_1 - a_4b_1c_2 \\
& + a_1b_4c_2 + a_2b_1c_4 - a_1b_2c_4 - b_2c_1x + b_1c_2x + a_2c_1y - a_1c_2y \\
& - a_2b_1z + a_1b_2z).
\end{aligned}$$

to the linear differential system

$$\dot{x} = -y, \quad \dot{y} = x, \quad \dot{z} = 0, \quad (2)$$

which has the two independent first integrals $f_1(\mathbf{x}) = x^2 + y^2$ and $f_2(\mathbf{x}) = z$. Here we assume that $\det(A) = -a_3b_2c_1 + a_2b_3c_1 + a_3b_1c_2 - a_1b_3c_2 - a_2b_1c_3 + a_1b_2c_3 \neq 0$. We emphasize the genericity in this paper means that some result

45 holds in a residual set.

Note that the linear differential system (2) in \mathbb{R}^3 has the full z -axis filled with singular points, and on the invariant planes $z = z_0 = \text{constant}$ we have a global center at $(0, 0, z_0)$, i.e. all the periodic orbits surrounding the center $(0, 0, z_0)$ in the plane $z = z_0$ fill this plane with the exception of the singular
50 point $(0, 0, z_0)$.

In this paper the linear differential system in \mathbb{R}^3 defined by a vector fields X will be called a *linear center* or simply a *center* in \mathbb{R}^3 .

Changing the parameters (a_i, b_i, c_i) of the vector field $X(\mathbf{x})$ to $(\alpha_i, \beta_i, \gamma_i)$ for $i = 1, 2, 3$ we get another linear vector field $Y(\mathbf{x}) = C\mathbf{x} + D$.

55 For any piecewise vector field in \mathbb{R}^3 formed by two pieces and separated by one plane we can assume without loss of generality, after an affine change in \mathbb{R}^2 if necessary, that such a plane is the plane $x = 0$.

We define the *piecewise vector field* $N = (X, Y)$ in \mathbb{R}^3 of two pieces separated by the plane $x = 0$ and formed by the two centers X and Y of \mathbb{R}^3 as

$$N(\mathbf{x}) = \begin{cases} X(\mathbf{x}) = A\mathbf{x} + B, & \text{if } x \geq 0, \\ Y(\mathbf{x}) = C\mathbf{x} + D, & \text{if } x \leq 0. \end{cases}$$

If $X = Y$ on the plane $x = 0$ we say that the piecewise vector field N is *continuous*, and if $X \neq Y$ on the plane $x = 0$ we say that the piecewise vector
60 field N is *discontinuous*. For these two kinds of piecewise vector fields N we have the following two results.

Theorem 1.1. *A piecewise vector field in \mathbb{R}^3 separated by one plane and formed by two linear centers X and Y has no limit cycles.*

The limit cycles of Theorem 1.1 are crossing limit cycles, see for details the
65 last part of Section 2.

If we assume that the discontinuity region of a piecewise vector field in \mathbb{R}^3 is formed by two parallel planes, then without loss of generality we can assume, after an affine change in \mathbb{R}^3 if necessary, that these parallel planes are the planes $x = 1$ and $x = -1$.

70 We consider piecewise vector fields separated by the planes $x = \pm 1$ and formed by three centers X , Y and Z , where $Z(\mathbf{x}) = E\mathbf{x} + F$ is obtained from X changing the coefficients (a_i, b_i, c_i) by (A_i, B_i, C_i) for $i = 1, 2, 3$.

More precisely, we define the *piecewise vector field* $M = (X, Y, Z)$ in \mathbb{R}^3 of three pieces separated by the two planes $x = \pm 1$ and formed by the three centers X , Y and Z of \mathbb{R}^3 as

$$M(\mathbf{x}) = \begin{cases} X(\mathbf{x}) = A\mathbf{x} + B, & \text{if } x \geq 1, \\ Y(\mathbf{x}) = C\mathbf{x} + D, & \text{if } -1 \leq x \leq 1, \\ Z(\mathbf{x}) = E\mathbf{x} + F, & \text{if } x \leq -1. \end{cases}$$

As in the case when the discontinuity line was formed by a unique plane, now we can consider continuous and discontinuous piecewise vector fields. Our
75 main results for the piecewise vector fields M are the following two theorems.

Theorem 1.2. *A continuous piecewise vector field separated by two parallel planes and formed by three linear centers X , Y and Z generically has no limit cycles.*

We notice that our parameter space is \mathbb{R}^{36} . Thus, when we claim that the
80 continuous piecewise vector field has generically no limit cycle we mean that it has no limit cycles in a residual set of \mathbb{R}^{36} . In other words, the possible scenario where the result could not be verified has measure zero (here we can assume Lebesgue measure).

Theorem 1.3. *A discontinuous piecewise vector field separated by two parallel
85 planes and formed by three linear centers X , Y and Z generically has at most four limit cycles, and we provide one of these piecewise vector fields with exactly four limit cycles.*

The meaning of genericity in the statement of Theorem 1.3 is the same as in Theorem 1.2. Again the limit cycles of Theorems 1.2 and 1.3 are crossing limit
90 cycles, see for details the last part of Section 2. The proofs of Theorems 1.1, 1.2 and 1.3 are given in the Section 3.

2. Filippov rules for defining the piecewise vector fields

Following the Filippov rules introduced in [5] we first consider an open set $U \subset \mathbb{R}^3$ and the *discontinuity region* $\Sigma = f^{-1}(0)$ being 0 a regular value of a
95 C^r smooth function $f : U \subset \mathbb{R}^3 \rightarrow \mathbb{R}$, for $1 \leq r \leq \infty$. As usual a C^r vector field is a C^r function $X : U \rightarrow \mathbb{R}^3$. The set of C^r vector fields over \mathbb{R}^3 will be denoted by $\mathfrak{X}^r(\mathbb{R}^3)$.

Let $X_i \in \mathfrak{X}^r(\mathbb{R}^3)$ be arbitrary vector fields, with $i = 1, \dots, n$ where n is the number of connected components D_i of $\mathbb{R}^3 \setminus \Sigma$. We denote a (non-smooth)

piecewise vector field in \mathbb{R}^3 by the n -tuple $N = (X_1, \dots, X_n)$ where

$$N(\mathbf{x}) = X_i(\mathbf{x}), \quad \text{if } \mathbf{x} \text{ in } D_i,$$

here $\mathbf{x} = (x, y, z) \in \mathbb{R}^3$. We note that on the discontinuity region Σ the piecewise vector field N is bi-valuated.

100 Now we precisely define $N = (X, Y)$ over Σ . A point $\mathbf{x} \in \Sigma$ is of *crossing* type if vector fields $X(\mathbf{x})$ and $Y(\mathbf{x})$ points in the same direction respect to Σ . It is of *sliding* type if both $X(\mathbf{x})$ and $Y(\mathbf{x})$ points inward Σ and it is of *escaping* type if $X(\mathbf{x})$ and $Y(\mathbf{x})$ points outward Σ . In each situation we are assuming that the trajectories of X and Y are transversal to Σ . Otherwise we say that a
105 point $\mathbf{x} \in \Sigma$ is a *tangency point* of X or Y .

An effective criterion for classifying points on the discontinuity region Σ can be established in terms of the Lie derivatives as follows. We define the Lie derivative at $\mathbf{x} \in \Sigma$ as

$$Xf(\mathbf{x}) = \langle \nabla f(\mathbf{x}), X(\mathbf{x}), \rangle$$

and for $k \geq 2$ we define $X^k f(\mathbf{x}) = \langle \nabla X^{k-1} f(\mathbf{x}), X(\mathbf{x}) \rangle$. The transversal points on Σ with respect to the vector fields X and Y are classified as follows:

- *Crossing region:* $\Sigma^c = \{\mathbf{x} \in \Sigma, (Xf(\mathbf{x})).(Yf(\mathbf{x})) > 0\}$, formed by crossing points.
- 110 • *Sliding region:* $\Sigma^s = \{\mathbf{x} \in \Sigma, Xf(\mathbf{x}) < 0 \text{ and } Yf(\mathbf{x}) > 0\}$, formed by sliding points.
- *Escaping region:* $\Sigma^e = \{\mathbf{x} \in \Sigma, Xf(\mathbf{x}) > 0 \text{ and } Yf(\mathbf{x}) < 0\}$, formed by escaping points.

Filippov's convention [5] allows to define of two kinds of limit cycles for the
115 piecewise vector fields, the so-called *sliding limit cycles* and the *crossing limit cycles*. Sliding limit cycles contain *sliding points* on the line of discontinuity and *crossing limit cycles* contain only *crossing points*. Here we only work with crossing limit cycles, or simply *limit cycles*.

3. Proof of the results

120 *Proof of Theorem 1.1. The continuous case.* Consider a continuous piecewise linear differential system in \mathbb{R}^3 separated by one plane and formed by two centers X and Y . Without loss of generality, we can suppose that the plane is $x = 0$.

The continuous hypothesis, $X(\mathbf{x}) = Y(\mathbf{x})$ at $(x, y, z) = (0, y_0, z_0)$, provides 3 polynomial equations of degree 1 in the variables y_0 and z_0 , each equation corresponding to the coordinates of the involved vector fields. They write

$$(B_1 - D_1) + (A_{12} - C_{12})y_0 + (A_{13} - C_{13})z_0 = 0$$

$$(B_2 - D_2) + (A_{22} - C_{22})y_0 + (A_{23} - C_{23})z_0 = 0$$

$$(B_3 - D_3) + (A_{32} - C_{32})y_0 + (A_{33} - C_{33})z_0 = 0$$

Thus we get 9 cubic equations for the parameters, looking at each coefficient, $B_i - D_i = 0$, $A_{i2} - C_{i2} = 0$ and $A_{i3} - C_{i3} = 0$ for $i = 1, 2, 3$. From those
125 equations we get 7 parameters $b_4, b_1, a_1, b_2, a_2, \alpha_1, \beta_1$ in terms of the other ones, as solution of the system.

If the continuous piecewise differential system has a limit cycle, this must intersect the plane $x = 0$ in two points $(0, y_0, z_0)$ and $(0, y_1, z_1)$ with $(y_0, z_0) \neq (y_1, z_1)$. For the vector field X we have the first integrals

$$130 \quad F_1(\mathbf{x}) = (a_4 + a_1x + a_2y + a_3z)^2 + (b_4 + b_1x + b_2y + b_3z)^2,$$

$$F_2(\mathbf{x}) = c_4 + c_1x + c_2y + c_3z,$$

and for the vector field Y the first integrals

$$G_1(\mathbf{x}) = (\alpha_1x + \alpha_2y + \alpha_3z + \alpha_4)^2 + (\beta_1x + \beta_2y + \beta_3z + \beta_4)^2,$$

$$G_2(\mathbf{x}) = \gamma_1x + \gamma_2y + \gamma_3z + \gamma_4.$$

135 Clearly that the two points $(0, y_0, z_0)$ and $(0, y_1, z_1)$ with $(y_0, z_0) \neq (y_1, z_1)$ where the limit cycles intersects the plane $x = 0$ must satisfy the system of four equations

$$e_1 = F_1(0, y_0, z_0) - F_1(0, y_1, z_1) = 0,$$

$$e_2 = F_2(0, y_0, z_0) - F_2(0, y_1, z_1) = 0,$$

$$140 \quad e_3 = G_1(0, y_0, z_0) - G_1(0, y_1, z_1) = 0,$$

$$e_4 = G_2(0, y_0, z_0) - G_2(0, y_1, z_1) = 0.$$

Since the unique solution of this system is $y_0 = y_1$ and $z_0 = z_1$, it follows that the continuous piecewise differential system formed by the centers X and Y has no limit cycles.

145 **The discontinuous case.** Assume that we have a discontinuous piecewise linear differential system in \mathbb{R}^3 separated by one plane and formed by two centers X and Y . Without loss of generality we can suppose that the plane is $x = 0$.

Then a limit cycle intersects the plane $x = 0$ in the two points $(0, y_0, z_0)$ and $(0, y_1, z_1)$ with $(y_0, z_0) \neq (y_1, z_1)$. The vector fields X and Y have the
150 first integrals $F_1(\mathbf{x}), F_2(\mathbf{x})$ and $G_1(\mathbf{x}), G_2(\mathbf{x})$ respectively, given in the proof of Theorem 1.1. As in that proof the two points $(0, y_0, z_0)$ and $(0, y_1, z_1)$ must satisfy

$$e_1 = F_1(0, y_0, z_0) - F_1(0, y_1, z_1) = 0,$$

$$e_2 = F_2(0, y_0, z_0) - F_2(0, y_1, z_1) = 0,$$

$$155 \quad e_3 = G_1(0, y_0, z_0) - G_1(0, y_1, z_1) = 0,$$

$$e_4 = G_2(0, y_0, z_0) - G_2(0, y_1, z_1) = 0.$$

Computing the Gröebner basis of the polynomials e_1, e_2, e_3 and e_4 respect to the variables y_0, z_0, y_1 and z_1 , we obtain an equivalent polynomial system to $e_1 = e_2 = e_3 = e_4 = 0$ formed by 59 equations. One of those ones is
160 $(z_1 - z_0)(c_3\gamma_2 - c_2\gamma_3) = 0$.

Assume that $c_3\gamma_2 - c_2\gamma_3$ is not zero. Then $z_1 = z_0$. Therefore $e_4 = (y_0 - y_1)\gamma_2$. Since y_1 cannot be equal to y_0 , otherwise the point $(0, y_0, z_0)$ would be equal to $(0, y_1, z_1)$, we have that $\gamma_2 = 0$. Then $e_2 = (y_0 - y_1)c_2 = 0$, so $c_2 = 0$, in contradiction with the assumption that $c_3\gamma_2 - c_2\gamma_3$ is not zero. Therefore in

165 what follows we can assume that $c_3\gamma_2 - c_2\gamma_3 = 0$ and that $z_0 \neq z_1$. Now we consider two cases.

Case 1: c_2 is not zero. Then $\gamma_3 = c_3\gamma_2/c_2$. Doing again the Gröebner basis of the polynomials e_1, e_2, e_3 and e_4 respect to the variables y_0, z_0, y_1 and z_1 , we obtain an equivalent polynomial system with eight equations. After removing
170 the non-zero factor $z_0 - z_1$ from the equations having such a factor, all these are linear in the variables y_0, z_0, y_1 and z_1 except one equation. These equations can be solved and we obtain a continuum of solutions. Consequently, again we do not have limit cycles in this case.

Case 2: $c_2 = 0$. Then $e_2 = c_3(z_0 - z_1) = 0$. Hence $c_3 = 0$. It remains three
175 equations $e_1 = e_3 = e_4 = 0$ and four unknowns. we obtain a continuum of solutions. Consequently again we do not have limit cycles in this case. \square

Proof of Theorem 1.2. The continuous hypotheses that $X(\mathbf{x}) = Y(\mathbf{x})$ at $(x, y, z) = (1, y_0, z_0)$ and $Y(\mathbf{x}) = Z(\mathbf{x})$ at $(x, y, z) = (-1, y_1, z_1)$, provide 6 polynomial equations of degree 1 in the variables y_0, z_0, y_1 and z_1 , each equation corresponding to the coordinates of the involved vector fields. Thus, similarly to the
180 proof of Theorem 1.1, we get 18 equations for the parameters, looking at each coefficient. From that system we get 13 parameters, which satisfy the whole equations, $b_4, b_1, a_1, b_2, a_2, \alpha_1, \beta_1, B_4, B_1, A_1, B_2, A_2$ and γ_1 in terms of the other ones.

185 The vector fields X and Y have the first integrals $F_1(\mathbf{x})$, $F_2(\mathbf{x})$ and $G_1(\mathbf{x})$, $G_2(\mathbf{x})$, respectively. For the vector field Z we have the firsts integrals

$$H_1(\mathbf{x}) = (A_1x + A_2y + A_3z + A_4)^2 + (B_1x + B_2y + B_3z + B_4)^2,$$

$$H_2(\mathbf{x}) = C_1x + C_2y + C_3z + C_4.$$

A possible limit cycle intersects at the points $(1, y_0, z_0)$ and $(1, y_3, z_3)$ the
190 plane $x = 1$, and at the points $(-1, y_1, z_1)$ and $(-1, y_2, z_2)$ in the plane $x = -1$, with $(y_0, z_0) \neq (y_3, z_3)$ and $(y_1, z_1) \neq (y_2, z_2)$. These four points have to satisfy the following system of equations for $(y_0, y_1, y_2, y_3, z_0, z_1, z_2, z_3)$.

$$e_1 = F_1(1, y_3, z_3) - F_1(1, y_0, z_0) = 0,$$

$$\begin{aligned}
e_2 &= F_2(1, y_3, z_3) - F_2(1, y_0, z_0) = 0, \\
195 \quad e_3 &= G_1(1, y_0, z_0) - G_1(-1, y_1, z_1) = 0, \\
e_4 &= G_2(1, y_0, z_0) - G_2(-1, y_1, z_1) = 0, \\
e_5 &= H_1(-1, y_1, z_1) - H_1(-1, y_2, z_2) = 0, \\
e_6 &= H_2(-1, y_1, z_1) - H_2(-1, y_2, z_2) = 0, \\
e_7 &= G_1(-1, y_2, z_2) - G_1(1, y_3, z_3) = 0, \\
200 \quad e_8 &= G_2(-1, y_2, z_2) - G_2(1, y_3, z_3) = 0.
\end{aligned}$$

Using the even equations we get that

$$\begin{aligned}
z_1 &= \frac{2(c_1 C_2 \gamma_3 - c_1 \gamma_2 C_3 - C_1 c_2 \gamma_3 + C_1 \gamma_2 c_3)}{\gamma_3(C_2 c_3 - c_2 C_3)} + \frac{\gamma_2 y_0}{\gamma_3} - \frac{\gamma_2 y_1}{\gamma_3} + z_0, \\
z_2 &= \frac{2(c_1 C_2 \gamma_3 - c_1 \gamma_2 C_3 - C_1 c_2 \gamma_3 + C_1 \gamma_2 c_3)}{\gamma_3(C_2 c_3 - c_2 C_3)} - \frac{y_1(\gamma_2 C_3 - C_2 \gamma_3)}{\gamma_3 C_3} \\
&\quad - \frac{C_2 y_2}{C_3} + \frac{\gamma_2 y_0}{\gamma_3} + z_0, \\
y_3 &= -\frac{c_3 y_1(\gamma_2 C_3 - C_2 \gamma_3)}{C_3(\gamma_2 c_3 - c_2 \gamma_3)} + \frac{c_3 y_2(\gamma_2 C_3 - C_2 \gamma_3)}{C_3(\gamma_2 c_3 - c_2 \gamma_3)} + y_0, \\
z_3 &= \frac{c_2 y_1(\gamma_2 C_3 - C_2 \gamma_3)}{C_3(\gamma_2 c_3 - c_2 \gamma_3)} - \frac{c_2 y_2(\gamma_2 C_3 - C_2 \gamma_3)}{C_3(\gamma_2 c_3 - c_2 \gamma_3)} + z_0.
\end{aligned}$$

Note that we are assuming that all the denominators which appear in the previous expressions of z_1 , z_2 , y_3 and z_3 are non-zero. That is, we are solving the system $e_1 = 0, \dots, e_8 = 0$, in the more generic case, i.e. when the mentioned denominators do not vanish, and the denominators which will appear solving the remaining equations $e_1 = e_3 = e_5 = e_7 = 0$ do not vanish.

Then we have $e_1 = e_3 = e_5 = e_7 = 0$ for solving y_0, y_1, y_2, z_0 . From $e_1 = 0$ and $e_3 = 0$ we can find y_2 and z_0 , respectively. Substituting y_2 and z_0 in e_5 and e_7 , we find that $y_1 = y_0 + 2(C_1 c_3 - c_1 C_3)/(C_2 c_3 - c_2 C_3)$ vanishes both equations. So we have a continuum of periodic orbits and no limit cycles. \square

210 *Proof of Theorem 1.3.* We consider such a possible limit cycle which intersects at the points $(1, y_0, z_0)$ and $(1, y_3, z_3)$ the plane $x = 1$ and at the points

$(-1, y_1, z_1)$ and $(-1, y_2, z_2)$ the plane $x = -1$, with $(y_0, z_0) \neq (y_3, z_3)$ and $(y_1, z_1) \neq (y_2, z_2)$. The vector fields X , Y and Z have the first integrals $F_1(\mathbf{x}), F_2(\mathbf{x}), G_1(\mathbf{x}), G_2(\mathbf{x})$ and $H_1(\mathbf{x}), H_2(\mathbf{x})$, respectively.

215 These four points must satisfy the following systems of equations for the variables $(y_0, y_1, y_2, y_3, z_0, z_1, z_2, z_3)$.

$$e_1 = F_1(1, y_3, z_3) - F_1(1, y_0, z_0) = 0,$$

$$e_2 = F_2(1, y_3, z_3) - F_2(1, y_0, z_0) = 0,$$

$$e_3 = G_1(1, y_0, z_0) - G_1(-1, y_1, z_1) = 0,$$

$$220 \quad e_4 = G_2(1, y_0, z_0) - G_2(-1, y_1, z_1) = 0,$$

$$e_5 = H_1(-1, y_1, z_1) - H_1(-1, y_2, z_2) = 0,$$

$$e_6 = H_2(-1, y_1, z_1) - H_2(-1, y_2, z_2) = 0,$$

$$e_7 = G_1(-1, y_2, z_2) - G_1(1, y_3, z_3) = 0,$$

$$e_8 = G_2(-1, y_2, z_2) - G_2(1, y_3, z_3) = 0.$$

Using the even equations we get

$$\begin{aligned} z_1 &= z_0 + \frac{2\gamma_1}{\gamma_3} + \frac{y_0\gamma_2}{\gamma_3} - \frac{y_1\gamma_2}{\gamma_3}, \\ z_2 &= -\frac{C_2y_2}{C_3} + z_0 + \frac{2\gamma_1}{\gamma_3} + \frac{y_0\gamma_2}{\gamma_3} - \frac{y_1(C_3\gamma_2 - C_2\gamma_3)}{C_3\gamma_3}, \\ y_3 &= y_0 - \frac{c_3y_1(C_3\gamma_2 - C_2\gamma_3)}{C_3(c_3\gamma_2 - c_2\gamma_3)} + \frac{c_3y_2(C_3\gamma_2 - C_2\gamma_3)}{C_3(c_3\gamma_2 - c_2\gamma_3)}, \\ z_3 &= z_0 + \frac{c_2y_1(C_3\gamma_2 - C_2\gamma_3)}{C_3(c_3\gamma_2 - c_2\gamma_3)} - \frac{c_2y_2(C_3\gamma_2 - C_2\gamma_3)}{C_3(c_3\gamma_2 - c_2\gamma_3)}. \end{aligned}$$

225 Again note that we are assuming that all the denominators which appear in the previous expressions of z_1 , z_2 , y_3 and z_3 , are non-zero. That is, we are solving the system $e_1 = 0, \dots, e_8 = 0$, in the more generic case, i.e. when the mentioned denominators do not vanish, and the denominators which will appear solving the remaining equations $e_1 = e_3 = e_5 = e_7 = 0$ do not vanish.

EqVar	y_0	y_1	y_2	z_0
e_1	1	1	1	1
e_3	2	2	0	1
e_5	1	1	1	1
e_7	2	2	2	1

Table 1: Maximum exponent for each variable in the polynomials e_k for $k = 1, 3, 5, 7$.

230 We substitute the obtained values of y_3, z_1, z_2 and z_3 in the equations $e_k = 0$ for $k = 1, 3, 5, 7$.

It remains to solve the equations $e_1 = e_3 = e_5 = e_7 = 0$ with respect to the unknowns y_0, y_1, y_2 and z_0 . We do not provide the big explicit expressions of the equations $e_1 = e_3 = e_5 = e_7 = 0$, which are easy to obtain with some
235 algebraic manipulator as Mathematica or Mapple.

The equations $e_1 = 0$ and $e_5 = 0$ are both of degree one, see Table 1. Thus we can find the variables y_0 and y_1 as linear functions of the variables y_2 and of z_0 .

Substituting the expressions of y_0 and y_1 into the remaining equations $e_3 = 0$
240 and $e_7 = 0$ we obtain two polynomial equations of degree two in the variables z_0 and y_2 . Again we do not provide the huge explicit expressions of the equations $e_3 = 0$ and $e_7 = 0$ which will need several pages for writing them.

Applying the Bézout Theorem (see [34]) to this system we know that at most there are four real solutions for (y_2, z_0) , which can produce four limit
245 cycles substituting these solutions in the previously obtained expressions of $(y_0, y_1, y_3, z_1, z_2, z_3)$. Hence the first part of Theorem 1.3 is proved.

In order to complete the proof of Theorem 1.3 we provide a discontinuous piecewise differential system in \mathbb{R}^3 formed by three centers and separated by two parallel planes, obtained numerically following the ideas described in [35] with an absolute tolerance of 10^{-15} . Using the notation of the first part of the proof of Theorem 1.3, the center given by the vector field X is obtained for the

values of the parameters

$$\begin{aligned}
a_1 &= 13.708561862548212..., & b_1 &= -2.0668139685572826..., \\
a_2 &= -0.46436203202470083..., & b_2 &= -2.1162121013313744..., \\
a_3 &= -1.5737561143110694..., & b_3 &= -0.30359559321560803..., \\
a_4 &= -5.2597752179200175..., & b_4 &= -3.837509139027315..., \\
c_1 &= 1, & c_2 &= -0.938643702906351..., & c_3 &= 1, & c_4 &= 1.
\end{aligned}$$

For the vector field Y we have that

$$\begin{aligned}
\alpha_1 &= 2, & \alpha_2 &= -5, & \alpha_3 &= -7, & \alpha_4 &= -11, \\
\beta_1 &= -3, & \beta_2 &= 1, & \beta_3 &= 3, & \beta_4 &= 1, \\
\gamma_1 &= 0.4545454545454544, & \gamma_2 &= -0.5454545454545444, & \gamma_3 &= 1, & \gamma_4 &= 6.
\end{aligned}$$

And the vector field Z is given by

$$\begin{aligned}
A_1 &= -3.840000000006582, & A_2 &= 5, & A_3 &= 0, & A_4 &= 14, \\
B_1 &= 1, & B_2 &= 12, & B_3 &= 11, & B_4 &= 14, \\
C_1 &= 2, & C_2 &= -4, & C_3 &= 5, & C_4 &= -8.
\end{aligned}$$

When we have computed (y_3, z_1, z_2, z_3) from $e_2 = e_4 = e_6 = e_8 = 0$ in the last proof, it remains the equations $e_1 = e_3 = e_5 = e_7 = 0$ for computing (y_0, y_1, y_2, z_0) . From the equation $e_1 = 0$ the variable y_2 appears linearly and we obtain $y_2 = P_{y_2}(y_0, y_1, z_0)$. Now from the equation $e_5 = 0$ the variable y_1 appears linearly and we have $y_1 = P_{y_1}(y_0, z_0)$. From the polynomial equations $e_3 = e_7 = 0$ both of degree two, we can obtain the four solutions for (y_0, z_0) , and from these four solutions we obtain the intersection points $(y_0, z_0, y_1, z_1, y_2, z_2, y_3, z_3)$ of our limit cycles with the two planes of discontinuity, see Figure 1, which are

$$\begin{aligned}
&(0.8978733932366719..., -2.2444618685679187..., -0.8476262083331095..., \\
&-2.2874616512423445..., 0.885305277605572..., -0.901116462491399..., \\
&2.0197502924959307..., -1.1914191816421125...), \\
&(2.618593513041539..., -2.017041157299665..., 1.6304557768236465..., \\
&-1.6469344679639704..., -0.25095559475947454..., -3.152063565230467..., \\
&1.4005927017518383..., -3.1603099489515696...),
\end{aligned}$$




Figure 1: The four limit cycles, vector fields X (red), Y (blue) and Z (green).

$$\begin{aligned}
& (0.5348395678273713..., -1.7034390938102253..., -1.2941046025617466..., \\
& -1.7919540958406528..., 0.47916792990915597..., -0.3733360698639303..., \\
& 1.6828327264792675..., -0.6258825444620513...), \\
& (1.7558413358089768..., -3.9459570275133773..., 0.15273809964405125..., \\
& -3.911286065421518..., 2.3088449504708346..., -2.1864005847600914..., \\
& 3.1516763632780256..., -2.635765268683441...).
\end{aligned}$$

□

4. Conclusions

We have shown that from the four classes of piecewise differential systems in \mathbb{R}^3 here studied, generically the only ones having limit cycles are the discontinuous piecewise differential systems in \mathbb{R}^3 separated by two parallel planes and formed by three centers having at most four limit cycles, see Theorem 1.3.

We remark that the discontinuous piecewise differential systems in \mathbb{R}^2 separated by two parallel straight lines and formed by three linear differential centers can have at most one limit cycle, see [28].

Acknowledgments

The first author is partially supported by PDSE-CAPES grant 88881. 624523/2021-01 and DS-CAPES 88882.386238/2019-01.

The second author is partially supported by the Agencia Estatal de Investigación grant PID2019-104658GB-I00, and the H2020 European Research Council grant MSCA-RISE-2017-777911.

The third author is partially supported by Pronex/FAPEG/CNPq grant 2012 10 26 7000 803 and grant 2017 10 26 7000 508, Capes grant 88881.068462/2014-01 and Universal/CNPq grant 420858/2016-4.

- [1] A. A. Andronov, A. A. Vitt, S. E. Khaikin, Theory of oscillators, Pergamon Press, Oxford-New York-Toronto, Ont., 1966, translated from the Russian by F. Immirzi; translation edited and abridged by W. Fishwick.

- [2] M. di Bernardo, C. J. Budd, A. R. Champneys, P. Kowalczyk, Piecewise-smooth dynamical systems, Vol. 163 of Applied Mathematical Sciences, Springer-Verlag London, Ltd., London, 2008, theory and applications.
- [3] D. J. W. Simpson, Bifurcations in piecewise-smooth continuous systems, Vol. 70 of World Scientific Series on Nonlinear Science. Series A: Monographs and Treatises, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2010. doi:10.1142/7612.
- [4] O. Makarenkov, J. S. W. Lamb, Dynamics and bifurcations of nonsmooth systems: a survey, Phys. D 241 (22) (2012) 1826–1844. doi:10.1016/j.physd.2012.08.002.
- [5] A. F. Filippov, Differential equations with discontinuous righthand sides, in: Mathematics and Its Applications, 1988.
- [6] J. C. Artés, J. Llibre, J. C. Medrado, M. A. Teixeira, Piecewise linear differential systems with two real saddles, Math. Comput. Simulation 95 (2014) 13–22. doi:10.1016/j.matcom.2013.02.007.
- [7] A. Bakhshalizadeh, J. Llibre, Limit cycles of piecewise differential equations on the cylinder, Bull. Sci. Math. 170 (2021) Paper No. 103013, 13. doi:10.1016/j.bulsci.2021.103013.
- [8] D. de Carvalho Braga, L. F. Mello, Limit cycles in a family of discontinuous piecewise linear differential systems with two zones in the plane, Nonlinear Dynam. 73 (3) (2013) 1283–1288. doi:10.1007/s11071-013-0862-3.
- [9] C. Buzzi, C. Pessoa, J. Torregrosa, Piecewise linear perturbations of a linear center, Discrete Contin. Dyn. Syst. 33 (9) (2013) 3915–3936. doi:10.3934/dcds.2013.33.3915.
- [10] J. Castillo, J. Llibre, F. Verduzco, The pseudo-Hopf bifurcation for planar discontinuous piecewise linear differential systems, Nonlinear Dynam. 90 (3) (2017) 1829–1840. doi:10.1007/s11071-017-3766-9.

- 295 [11] M. Esteban, J. Llibre, C. Valls, The 16th Hilbert problem for discontinuous
piecewise isochronous centers of degree one or two separated by a straight
line, *Chaos* 31 (4) (2021) Paper No. 043112, 18. doi:10.1063/5.0023055.
- [12] R. D. Euzébio, J. Llibre, On the number of limit cycles in discontinuous
piecewise linear differential systems with two pieces separated by a straight
300 line, *J. Math. Anal. Appl.* 424 (1) (2015) 475–486. doi:10.1016/j.jmaa.
2014.10.077.
- [13] E. Freire, E. Ponce, F. Rodrigo, F. Torres, Bifurcation sets of continuous
piecewise linear systems with two zones, *Internat. J. Bifur. Chaos Appl.
Sci. Engrg.* 8 (11) (1998) 2073–2097. doi:10.1142/S0218127498001728.
- 305 [14] E. Freire, E. Ponce, F. Torres, Canonical discontinuous planar piecewise
linear systems, *SIAM J. Appl. Dyn. Syst.* 11 (1) (2012) 181–211. doi:
10.1137/11083928X.
- [15] E. Freire, E. Ponce, F. Torres, A general mechanism to generate three limit
cycles in planar Filippov systems with two zones, *Nonlinear Dynam.* 78 (1)
310 (2014) 251–263. doi:10.1007/s11071-014-1437-7.
- [16] F. Giannakopoulos, K. Pliete, Planar systems of piecewise linear differential
equations with a line of discontinuity, *Nonlinearity* 14 (6) (2001) 1611–1632.
doi:10.1088/0951-7715/14/6/311.
- [17] M. R. A. Gouveia, J. Llibre, D. D. Novaes, On limit cycles bifurcating from
315 the infinity in discontinuous piecewise linear differential systems, *Appl.
Math. Comput.* 271 (2015) 365–374. doi:10.1016/j.amc.2015.09.022.
- [18] S.-M. Huan, X.-S. Yang, On the number of limit cycles in general planar
piecewise linear systems, *Discrete Contin. Dyn. Syst.* 32 (6) (2012) 2147–
2164. doi:10.3934/dcds.2012.32.2147.
- 320 [19] S.-M. Huan, X.-S. Yang, Existence of limit cycles in general planar piece-
wise linear systems of saddle-saddle dynamics, *Nonlinear Anal.* 92 (2013)
82–95. doi:10.1016/j.na.2013.06.017.

- [20] S.-M. Huan, X.-S. Yang, On the number of limit cycles in general planar piecewise linear systems of node-node types, *J. Math. Anal. Appl.* 411 (1) (2014) 340–353. doi:10.1016/j.jmaa.2013.08.064.
- [21] L. Li, Three crossing limit cycles in planar piecewise linear systems with saddle-focus type, *Electron. J. Qual. Theory Differ. Equ.* (2014) No. 70, 14doi:10.14232/ejqtde.2014.1.70.
- [22] J. Llibre, D. D. Novaes, M. A. Teixeira, Limit cycles bifurcating from the periodic orbits of a discontinuous piecewise linear differentiable center with two zones, *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* 25 (11) (2015) 1550144, 11. doi:10.1142/S0218127415501448.
- [23] J. Llibre, D. D. Novaes, M. A. Teixeira, Maximum number of limit cycles for certain piecewise linear dynamical systems, *Nonlinear Dynam.* 82 (3) (2015) 1159–1175. doi:10.1007/s11071-015-2223-x.
- [24] J. Llibre, D. D. Novaes, M. A. Teixeira, On the birth of limit cycles for non-smooth dynamical systems, *Bull. Sci. Math.* 139 (3) (2015) 229–244. doi:10.1016/j.bulsci.2014.08.011.
- [25] J. Llibre, M. Ordóñez, E. Ponce, On the existence and uniqueness of limit cycles in planar continuous piecewise linear systems without symmetry, *Nonlinear Anal. Real World Appl.* 14 (5) (2013) 2002–2012. doi:10.1016/j.nonrwa.2013.02.004.
- [26] J. Llibre, E. Ponce, Three nested limit cycles in discontinuous piecewise linear differential systems with two zones, *Dyn. Contin. Discrete Impuls. Syst. Ser. B Appl. Algorithms* 19 (3) (2012) 325–335.
- [27] J. Llibre, M. A. Teixeira, Piecewise linear differential systems without equilibria produce limit cycles?, *Nonlinear Dynam.* 88 (1) (2017) 157–164. doi:10.1007/s11071-016-3236-9.

- [28] J. Llibre, M. A. Teixeira, Piecewise linear differential systems with only
 350 centers can create limit cycles?, *Nonlinear Dynam.* 91 (1) (2018) 249–255.
 doi:10.1007/s11071-017-3866-6.
- [29] J. Llibre, M. A. Teixeira, J. Torregrosa, Lower bounds for the maximum
 number of limit cycles of discontinuous piecewise linear differential systems
 with a straight line of separation, *Internat. J. Bifur. Chaos Appl. Sci. Engrg.*
 355 23 (4) (2013) 1350066, 10. doi:10.1142/S0218127413500661.
- [30] J. Llibre, X. Zhang, Limit cycles for discontinuous planar piecewise linear
 differential systems separated by one straight line and having a center, *J.*
Math. Anal. Appl. 467 (1) (2018) 537–549. doi:10.1016/j.jmaa.2018.
 07.024.
- [31] H. Poincaré, Mémoire sur les courbes définies par les équations
 360 différentielles (i), *Journal de Mathématiques Pures et Appliquées* 7 (1881)
 375–422.
- [32] H. Dulac, Détermination et intégration d’une certaine classe d’équations
 différentielles ayant pour point singulier un centre, Gauthier-Villars, 1908.
- [33] M. Brunella, Instability of equilibria in dimension three, *Ann. Inst. Fourier*
 365 (Grenoble) 48 (5) (1998) 1345–1357.
- [34] I. R. Shafarevich, Basic algebraic geometry, Springer Study Edition,
 Springer-Verlag, Berlin-New York, 1977, translated from the Russian by
 K. A. Hirsch, Revised printing of *Grundlehren der mathematischen Wis-*
 370 *senschaften*, Vol. 213, 1974.
- [35] L. M. Echeverry, Y. Villanueva, Minimizing fuel consumption in orbit
 transfers using universal variable and particle swarm optimization, *Re-*
vista mexicana de astronomía y astrofísica 55 (2019) 177 – 184. doi:
 10.22201/ia.01851101p.2019.55.02.05.