



ICTV Virus Taxonomy Profile: *Potyviridae* 2022

Alice K. Inoue-Nagata^{1,*}, Ramon Jordan², Jan Kreuze³, Fan Li⁴, Juan José López-Moya⁵, Kristiina Mäkinen⁶, Kazusato Ohshima⁷, Stephen J. Wylie⁸ and ICTV Report Consortium

Abstract

The family *Potyviridae* includes plant viruses with single-stranded, positive-sense RNA genomes of 8–11 kb and flexuous filamentous particles 650–950 nm long and 11–20 nm wide. Genera in the family are distinguished by the host range, genomic features and phylogeny of the member viruses. Most genomes are monopartite, but those of members of the genus *Bymovirus* are bipartite. Some members cause serious disease epidemics in cultivated plants. This is a summary of the International Committee on Taxonomy of Viruses (ICTV) Report on the family *Potyviridae*, which is available at [ictv.global/report/potyviridae](https://www.ictv.global/report/potyviridae).

Table 1. Characteristics of members of the family *Potyviridae*

Example	potato virus Y-O (U09509), species <i>Potato virus Y</i> , genus <i>Potyvirus</i>
Virion	Non-enveloped, flexuous and filamentous particles, 650–950 nm long and 11–20 nm in diameter with a core capsid protein
Genome	8.2–11.5 kb of positive-sense, single-stranded, usually monopartite RNA (bipartite in the genus <i>Bymovirus</i>)
Replication	Cytoplasmic, initiated in virus replication complexes on 6K2-induced membranous vesicles at endoplasmic reticulum exit sites
Translation	Directly from genomic RNA
Host range	Plants; transmission by arthropods, plasmodiophorids (<i>Bymovirus</i>), seeds or pollen
Taxonomy	Realm <i>Riboviria</i> , kingdom <i>Orthornavirae</i> , phylum <i>Pisuviricota</i> , class <i>Stelpaviricetes</i> , order <i>Patatavirales</i> : >10 genera including >230 species

VIRION

The flexuous, filamentous particles are typically 650–950 nm long and 11–20 nm wide, with helical symmetry and a pitch of about 3.4–3.5 nm (Fig. 1, Table 1), but members of the genus *Bymovirus* are bipartite with particles of two modal lengths of 250–300 nm and 500–600 nm. Virions contain a core capsid protein (CP) of 28.5–47 kDa [1]; the tip of the particle may contain the virus-encoded proteins genome-linked protein (VPg), helper-component protease (HC-Pro), cylindrical inclusion (CI) and the host factor eIF4E. There are serological relationships among many members.

GENOME

The positive-sense ssRNA genome ranges from 8.2 kb (members of the species *Artichoke latent virus*, genus *Macluravirus*, and species *Bellflower veinal mottle virus*, genus *Bevemovirus*) to 11.5 kb (species *Celery latent virus*, genus *Celavirus*), with a VPg covalently linked to the 5'-end, and a polyadenylated 3'-terminus (Fig. 2). The genome of the celavirus celery latent virus is not polyadenylated. Most genomes are monopartite, but those of members of the genus *Bymovirus* are bipartite. The major large ORF of monopartite genomes encodes a polyprotein that is cleaved into functional proteins at semi-conserved sites by two or three viral proteases [2]. Bipartite *Bymovirus* genomes encode

Received 03 February 2022; Accepted 06 February 2022; Published 05 May 2022

Author affiliations: ¹Embrapa Vegetables, Brasília, Brazil; ²USDA, Beltsville, MD, USA; ³CIP, Lima, Peru; ⁴Yunnan Agricultural University, Yunnan, PR China; ⁵CRAG-CSIC, Barcelona, Spain; ⁶University of Helsinki, Helsinki, Finland; ⁷Saga University, Saga, Japan; ⁸Murdoch University, Perth, Australia.

*Correspondence: Alice K. Inoue-Nagata, alice.nagata@embrapa.br

Keywords: ICTV Profile; taxonomy; *Potyviridae*.

Abbreviations: CI, cylindrical inclusion; CP, capsid protein; HC-Pro, helper-component protease; VPg, virus protein genome-linked; VRC, viral replication complex.

001738 © 2022



This is an open-access article distributed under the terms of the Creative Commons Attribution License.




Fig. 1. (Left) Cryo-electron microscopy reconstruction of a section of a virion of turnip mosaic virus (TuMV). Depicted are the ssRNA (red), and the central globular domain (pale yellow), N-terminal arm (blue) and C-terminal region (grey) of the capsid protein. In the upper section some of the densities are cut-away. (Right) Cryo-electron micrograph of TuMV virions where the periodic organization of the helical filaments is apparent. Bar, ~500 nm. Images are courtesy of R. Cuesta and M. Valle, CIC bioGUNE, Spain, based on [7].

two polyproteins that are cleaved by two proteases. A second small ORF, PIPO, is generated by a polymerase slippage mechanism and is expressed as the *trans*-frame protein P3N-PIPO [3]. Another additional small ORF, PISPO, is generated through a transcriptional slippage mechanism among sweet potato-subgroup




Fig. 2. Schematic diagram of the genome organization of potato virus Y. VPg, virus protein genome-linked (indicated by an ellipse at the 5'-terminus); P1, protein 1 protease; HC-Pro, helper component protease; P3, protein 3; PIPO, pretty interesting *Potyviridae* ORF; 6K1, 6 kDa peptide 1; CI, cylindrical inclusion; 6K2, 6 kDa peptide 2; Nla-Pro, nuclear inclusion a-protease; Nlb, nuclear inclusion b, RNA-directed RNA polymerase; CP, capsid protein. Arrowheads indicate cleavage sites of P1 (blue), HC-Pro (red) and Nla-Pro (black).

References

- Martínez-Turiño S, García JA. Potyviral coat protein and genomic RNA: A striking partnership leading virion assembly and more. *Adv Virus Res* 2020;108:165–211.
- Adams MJ, Antoniw JF, Beaudoin F. Overview and analysis of the polyprotein cleavage sites in the family *Potyviridae*. *Mol Plant Pathol* 2005;6:471–487.
- Chung BY-W, Miller WA, Atkins JF, Firth AE. An overlapping essential gene in the *Potyviridae*. *Proc Natl Acad Sci U S A* 2008;105:5897–5902.

potyviruses and leads to the production of a *trans*-frame protein P1N-PISPO. Some viruses lack one or both of the P1 and HC-Pro N-terminal coding regions, and may be replaced by genus-specific or species-specific regions.

REPLICATION

Genomic RNA serves as a template for both translation and replication (reviewed by [4]). The genomic RNA replicates via the production of a full-length negative-sense RNA, which takes place in the cytoplasmic viral replication complex (VRC). The VRC is formed by virus proteins, such as NlB, 6K2, VPg, Nla-Pro, HC-Pro, CI, and possibly P3 and 6K1, and host factors. The viral genome is transported cell-to-cell through the plasmodesmata, probably in the form of particles, involving the coordinated action of CP, CI, P3, 6K2 and P3N-PIPO [5].

TAXONOMY

Current taxonomy: ictv.global/taxonomy. Members of different genera in the family are distinguished by host range, vector, genomic features and phylogeny, with species demarcation typically based upon sequence identity of the large ORF (or, if necessary, the CP-coding region) being <76% (nt) and <82% (aa) [6]. Thresholds for other coding regions are usually 58% (P1 coding region) or 74–78% (other regions). Members of the family *Potyviridae* are transmitted by aphids, eriophyid mites, whiteflies and plasmodiophorids, but the vector is not known for members of five genera. New species or genera in the family will be considered after analyses based on complete genome sequences and additional data about biological characteristics.

RESOURCES

Full ICTV Report on the family *Potyviridae*: ictv.global/report/potyviridae.

Funding information

Production of this summary, the online chapter and associated resources was funded by a grant from the Wellcome Trust (WT108418AIA).

Acknowledgements

Members of the ICTV Report Consortium are Stuart G. Siddell, Elliot J. Lefkowitz, Sead Sabanadzovic, Peter Simmonds, F. Murilo Zerbini, Donald B. Smith, and Luisa Rubino.

Conflicts of interest

The authors declare that there are no conflicts of interest.

- Mäkinen K, Hafrén A. Intracellular coordination of potyviral RNA functions in infection. *Front Plant Sci* 2014;5:110.
- Wang A. Cell-to-cell movement of plant viruses via plasmodesmata: a current perspective on potyviruses. *Curr Opin Virol* 2021;48:10–16.
- Adams MJ, Antoniw JF, Fauquet CM. Molecular criteria for genus and species discrimination within the family *Potyviridae*. *Arch Virol* 2005;150:459–479.
- Cuesta R, Yuste-Calvo C, Gil-Cartón D, Sánchez F, Ponz F, et al. Structure of Turnip mosaic virus and its viral-like particles. *Sci Rep* 2019;9:15396.