
This is the **accepted version** of the journal article:

Porti, Joan. «Dimension of representation and character varieties for two- and three- orbifolds». Algebraic and Geometric Topology, Vol. 22, Issue 4 (2022), p. 1905-1967. DOI 10.2140/agt.2022.22.1905

This version is available at <https://ddd.uab.cat/record/275429>

under the terms of the  license

Dimension of representation and character varieties for two and three-orbifolds

Joan Porti*

February 9, 2023

Abstract

We consider varieties of representations and characters of 2 and 3-dimensional orbifolds in semisimple Lie groups, and we focus on computing their dimension. For hyperbolic 3-orbifolds, we consider the component of the variety of characters that contains the holonomy composed with the principal representation, we show that its dimension equals half the dimension of the variety of characters of the boundary. We also show that this is a lower bound for the dimension of generic components. We furthermore provide tools for computing dimensions of varieties of characters of 2-orbifolds, including the Hitchin component. We apply this computation to the dimension growth of varieties of characters of some 3-dimensional manifolds in $\mathrm{SL}(n, \mathbb{C})$.

1 Introduction

In this paper we are interested in varieties of representations and characters of the fundamental group of two and three-dimensional orbifolds \mathcal{O} in complex semi-simple algebraic Lie groups G , denoted respectively by

$$R(\mathcal{O}, G) = \mathrm{hom}(\pi_1(\mathcal{O}), G) \quad \text{and} \quad X(\mathcal{O}, G) = R(\mathcal{O}, G) // G.$$

Since those are not irreducible, we use $\dim_\rho R(\mathcal{O}, G)$ and $\dim_{[\rho]} X(\mathcal{O}, G)$ to denote the dimension of a component that contains ρ .

We discuss first a particular representation for oriented hyperbolic 3-orbifolds, with holonomy in $\mathrm{Isom}^+(\mathbb{H}^3) \cong \mathrm{PSL}(2, \mathbb{C})$. Let G be an adjoint complex simple Lie group, eg. $\mathrm{PSL}(n, \mathbb{C})$, $\mathrm{PO}(n, \mathbb{C})$ or $\mathrm{PSp}(2m, \mathbb{C})$. Let

$$\tau: \mathrm{PSL}(2, \mathbb{C}) \rightarrow G$$

be the principal representation (see §2.2 for the definition), eg. $\mathrm{Sym}^{n-1}: \mathrm{PSL}(2, \mathbb{C}) \rightarrow \mathrm{PSL}(n, \mathbb{C})$, whose image is contained in $\mathrm{PSp}(2m, \mathbb{C})$ for $n = 2m$ or in $\mathrm{PO}(2m+1, \mathbb{C})$ for $n = 2m+1$. When $G_{\mathbb{R}}$ is a split real form of G , it restricts to $\tau: \mathrm{PGL}(2, \mathbb{R}) \rightarrow G_{\mathbb{R}}$, so that the Hitchin component of a surface or a 2-orbifold is the component of the variety of representations that contains the composition of τ with a Fuchsian representation. Here we consider orientable hyperbolic 3-orbifolds and we compose the holonomy in $\mathrm{PSL}(2, \mathbb{C})$ with the principal representation τ . We prove the following generalization of [25, 26]:

Theorem 1.1. *Let \mathcal{O}^3 be a compact orientable 3-orbifold whose interior is hyperbolic with holonomy $\mathrm{hol}: \pi_1(\mathcal{O}^3) \rightarrow \mathrm{PSL}(2, \mathbb{C})$. Then the character of $\tau \circ \mathrm{hol}$ is a smooth point of $X(\mathcal{O}^3, G)$ of dimension*

$$\dim_{[\tau \circ \mathrm{hol}]} X(\mathcal{O}^3, G) = \frac{1}{2} \dim X(\partial \mathcal{O}^3, G).$$

*Partially supported by the Micinn/FEDER grant PGC2018-095998-B-I00

Furthermore, the map induced by restriction $X(\mathcal{O}^3, G) \rightarrow X(\partial\mathcal{O}^3, G)$ is locally injective at the character of $\tau \circ \text{hol}$.

The proof of Theorem 1.1 is based on a vanishing theorem of L^2 -cohomology, as for manifolds in [25, 26]. Notice that for a Euclidean component \mathcal{O}^2 of $\partial\mathcal{O}^3$, the restriction of $\tau \circ \text{hol}$ can be a singular point of $X(\mathcal{O}^2, G)$, and to prove that $X(\mathcal{O}^3, G) \rightarrow X(\partial\mathcal{O}^3, G)$ is locally injective we use evaluation of characters.

A representation $\rho: \Gamma \rightarrow G$ is called *good* if it is irreducible and the centralizer of $\rho(\Gamma)$ in G is the center $\mathcal{Z}(G)$ of G , in particular $\mathfrak{g}^{\rho(\Gamma)} = 0$. We are interested in representations of Euclidean 2-orbifolds (eg. $\chi(\mathcal{O}^2) = 0$) that may be not good. For instance, nontrivial representations of a 2-torus are never good by Kolchin's theorem. A representation of a Euclidean 2-orbifold $\rho: \pi_1(\mathcal{O}^2) \rightarrow G$ is called *strongly regular* if for a maximal torsion free subgroup $\Gamma_0 < \pi_1(\mathcal{O}^2)$, it holds:

- $\dim \mathfrak{g}^{\rho(\Gamma_0)} = \text{rank } G$, and
- the projection of $\rho(\Gamma_0)$ is contained in a connected abelian subgroup of $G/\mathcal{Z}(G)$.

The following is a generalization of a theorem of Falbel and Guilloux for manifolds in [8]:

Theorem 1.2. *Let \mathcal{O}^3 be a compact orientable good 3-orbifold, with boundary $\partial\mathcal{O}^3 = \partial_1\mathcal{O}^3 \sqcup \dots \sqcup \partial_k\mathcal{O}^3$, and $\chi(\partial_i\mathcal{O}^3) \leq 0$ for $i = 1, \dots, k$. Let $\rho: \pi_1(\mathcal{O}^3) \rightarrow G$ be a good representation such that:*

- *If $\chi(\partial_i\mathcal{O}^3) < 0$, then $\rho|_{\pi_1(\partial_i\mathcal{O}^2)}$ is good.*
- *If $\chi(\partial_i\mathcal{O}^3) = 0$, then $\rho|_{\pi_1(\partial_i\mathcal{O}^2)}$ is strongly regular.*

Then

$$\dim_{[\rho]} X(\mathcal{O}^3, G) \geq \frac{1}{2} \dim X(\partial\mathcal{O}^3, G).$$

If $\partial\mathcal{O}^3 = \partial_1\mathcal{O}^3 \sqcup \dots \sqcup \partial_k\mathcal{O}^3$ denotes the decomposition in connected components, then $X(\partial\mathcal{O}^3, G) = X(\partial_1\mathcal{O}^3, G) \times \dots \times X(\partial_k\mathcal{O}^3, G)$ and

$$\dim X(\partial\mathcal{O}^3, G) = \dim X(\partial_1\mathcal{O}^3, G) + \dots + \dim X(\partial_k\mathcal{O}^3, G).$$

When the component $\partial_i\mathcal{O}^3$ is a (closed orientable) surface S :

$$\dim X(S, G) = \begin{cases} -\chi(S) \dim G & \text{if } \chi(S) < 0, \\ 2 \text{rank } G & \text{if } \chi(S) = 0, \end{cases}$$

which can be rewritten as

$$\dim X(S, G) = -\chi(S) \dim G + 2 \dim \mathfrak{g}^{\rho(\pi_1(S))},$$

where \mathfrak{g}^H denotes the centralizer in the Lie algebra of a subset $H \subset G$, via the adjoint action. When the $\partial_i\mathcal{O}^3$ is a (closed orientable) 2-orbifold \mathcal{O}^2 with branching locus $\Sigma \subset \mathcal{O}^2$, we prove in Theorems 1.3 and 1.4:

$$\dim X(\mathcal{O}^2, G) = -\chi(\mathcal{O}^2 \setminus \Sigma) \dim G + \sum_{x \in \Sigma} \dim \mathfrak{g}^{\rho(\text{Stab}(x))} + 2 \dim \mathfrak{g}^{\rho(\pi_1(\mathcal{O}^2))}. \quad (1)$$

Formula (1) suggests that, instead of the usual Euler characteristic of an orbifold, we need another quantity to compute dimensions of varieties of characters and representations. For

that purpose, consider K a CW -structure on a compact orbifold \mathcal{O}^n and $\rho: \pi_1(\mathcal{O}^n) \rightarrow G$ a representation. The *twisted Euler characteristic* is defined (Definition 3.1) as

$$\tilde{\chi}(\mathcal{O}^n, \text{Ad}\rho) = \sum_{e \text{ cell of } K} (-1)^{\dim e} \dim \mathfrak{g}^{\text{Ad}(\rho(\text{Stab}(\bar{e})))} \in \mathbb{Z}.$$

This is always an integer and should not be confused with the orbifold Euler characteristic, that is a rational number. For the trivial representation, this is just the Euler characteristic of the underlying space of the orbifold times $\dim G$. It is the alternated sum of dimensions of cohomology groups of \mathcal{O}^n twisted by $\text{Ad}\rho$ (Proposition 3.2):

$$\tilde{\chi}(\mathcal{O}^n, \text{Ad}\rho) = \sum_i (-1)^i \dim H^i(\mathcal{O}^n, \text{Ad}\rho).$$

In Section 4 we prove the following results, based on Goldman's work on surfaces [10] (see also [32]):

Theorem 1.3. *Let \mathcal{O}^2 be a compact connected 2-orbifold, with $\chi(\mathcal{O}^2) \leq 0$. Let $\rho \in R(\mathcal{O}^2, G)$ be a good representation. Then $[\rho]$ is a smooth point of $X(\mathcal{O}^2, G)$ of dimension $-\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho)$.*

Theorem 1.4. *Let \mathcal{O}^2 be a closed Euclidean 2-orbifold and $\rho: \pi_1(\mathcal{O}^2) \rightarrow G$ a strongly regular representation. Then it belongs to a single component of $X(\mathcal{O}^2, G)$ that has dimension*

- (a) $-\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) + 2 \dim \mathfrak{g}^{\rho(\pi_1(\mathcal{O}^2))}$ if \mathcal{O}^2 is orientable,
- (b) $-\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) + \dim \mathfrak{g}^{\rho(\pi_1(\widetilde{\mathcal{O}^2}))}$ if \mathcal{O}^2 is non-orientable, where $\widetilde{\mathcal{O}^2} \rightarrow \mathcal{O}^2$ denotes the orientation covering of \mathcal{O}^2 .

Notice that Theorem 1.4 does not conclude smoothness. For instance, a parabolic representation of \mathbb{Z}^2 in $\text{SL}(2, \mathbb{C})$ is strongly regular but its character (the trivial character) is a singular point of $X(\mathbb{Z}^2, \text{SL}(2, \mathbb{C}))$ and $X(\mathbb{Z}^2, \text{PSL}(2, \mathbb{C}))$.

Under the hypothesis of Theorem 1.3 or Theorem 1.4, when \mathcal{O}^2 is *closed* and *orientable* then $\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho)$ is even (Corollary 3.5) and therefore $\dim_{[\rho]} X(\mathcal{O}^2, G)$ is even.

Most of the computations apply to real Lie groups, in particular we spend a section discussing applications to the Hitchin component, that we denote by $\text{Hit}(\mathcal{O}^2, G_{\mathbb{R}})$, where $G_{\mathbb{R}}$ is the split real form of the adjoint group G . As we consider *non orientable* 2-orbifolds, we deal with *non connected* real forms $G_{\mathbb{R}}$: we require that $G_{\mathbb{R}}$ contains the image by the principal representation of $\text{PGL}(2, \mathbb{R})$, not only of its identity component $\text{PSL}(2, \mathbb{R})$. For instance $G_{\mathbb{R}} = \text{PSL}(n, \mathbb{R})$, $\text{PSp}^{\pm}(2m)$ or $\text{PO}(n, n+1)$.

The dimension of $\text{Hit}(\mathcal{O}^2, G_{\mathbb{R}})$ has already been computed in [1] by Alessandrini, Lee and Schaffhauser, but we give a different approach, closer to the one of Long and Thistlethwaite in [23] for turnovers. For instance, we show that

$$|\dim(\text{Hit}(\mathcal{O}^2, \text{PGL}(n, \mathbb{R}))) + \chi(\mathcal{O}^2) \dim(\text{PGL}(n, \mathbb{R}))| \leq C(\mathcal{O}^2),$$

where $C(\mathcal{O}^2)$ is a constant that depends only on \mathcal{O}^2 , Proposition 5.10. For $\text{PSp}^{\pm}(2m, \mathbb{R})$ the bound is not uniform but linear on m and we only have a weaker conclusion (Corollary 5.15), cf. [1, Proposition 5.15]:

$$\lim_{m \rightarrow \infty} \frac{\dim(\text{Hit}(\mathcal{O}^2, \text{PSp}^{\pm}(2m, \mathbb{R})))}{\dim(\text{PSp}^{\pm}(2m, \mathbb{R}))} = -\chi(\mathcal{O}^2).$$

We also show that $\text{Hit}(\mathcal{O}^2, G_{\mathbb{R}})$ maximizes the dimension among all components of the variety of (good) representations of \mathcal{O}^2 in $G_{\mathbb{R}}$, for $G_{\mathbb{R}} = \text{PGL}(n, \mathbb{R})$, Proposition 5.8, or $\text{PSp}^{\pm}(2m, \mathbb{R})$, Proposition 5.16.

For \mathcal{O}^2 closed and orientable, the differential form of Atiyah-Bott-Goldman gives naturally a symplectic structure on $\text{Hit}(\mathcal{O}^2, G_{\mathbb{R}})$. Furthermore (see Proposition 5.4):

Proposition 1.5. *When \mathcal{O}^2 is closed and non-orientable, with orientation covering $\widetilde{\mathcal{O}^2}$, then $\text{Hit}(\mathcal{O}^2, G_{\mathbb{R}})$ embeds in $\text{Hit}(\widetilde{\mathcal{O}^2}, G_{\mathbb{R}})$ as a Lagrangian submanifold.*

We apply these computations on 2-orbifolds to estimate the growth of $X(M^3, \text{SL}(n, \mathbb{C}))$ with respect to n for some 3-manifolds, in particular for some knot exteriors. For instance we show:

Proposition 1.6. *Let Γ be the fundamental group of the exterior of the figure eight knot. Besides the canonical component (that has dimension $n - 1$), for large n $X(\Gamma, \text{SL}(n, \mathbb{C}))$ has at least 3 components that contain irreducible representations, whose dimension grow respectively as $n^2/12$, $n^2/20$ and $n^2/42$.*

The paper is organized as follows. In Section 2 we recall some basic notions and tools on varieties of representations, and in Section 3 we introduce tools of orbifold cohomology (some well known, some other new) that we use later to compute Zariski tangent spaces to varieties of representations and characters. In Section 4 we discuss varieties of representations of two dimensional orbifolds, and we apply some of the ideas to compute dimensions of Hitchin components of orbifolds in Section 5. In Section 6 we prove the results on three-dimensional orbifolds. Finally, some explicit examples are computed in Section 7.

2 Varieties of representations and characters

Let Γ be a finitely generated group, we are mainly interested in the fundamental group of a compact orbifold. Let G be a complex semi-simple algebraic Lie group. The variety of representations

$$R(\Gamma, G) = \text{hom}(\Gamma, G)$$

is an affine algebraic set and its quotient by conjugation in the algebraic category is the variety of characters

$$X(\Gamma, G) = R(\Gamma, G) // G.$$

Namely, the algebra of invariant functions $\mathbb{C}[R(\Gamma, G)]^G$ is of finite type and $X(\Gamma, G)$ can be defined as the affine scheme with this function ring: $\mathbb{C}[X(\Gamma, G)] \cong \mathbb{C}[R(\Gamma, G)]^G$. When $\Gamma = \pi_1(\mathcal{O})$, they are denoted respectively by $R(\mathcal{O}, G)$ and $X(\mathcal{O}, G)$.

Following for instance [19], we define:

Definition 2.1. *A representation $\rho: \Gamma \rightarrow G$ is:*

- (a) *irreducible if $\rho(\Gamma)$ is not contained in a proper parabolic subgroup of G , and*
- (b) *good if it is irreducible and the centralizer of the image equals the center of G , $Z(G)$.*

When $G = \text{SL}(n, \mathbb{C})$, irreducible and good are equivalent.

Let $R^{\text{good}}(\Gamma, G)$ denote the subset of good representations in $R(\Gamma, G)$. The set of conjugacy classes

$$\mathcal{R}^{\text{good}}(\Gamma, G) = R^{\text{good}}(\Gamma, G) / G$$

is a Zariski open subset of $X(\Gamma, G)$, cf. [19].

We discuss also representations in real Lie groups. For $G_{\mathbb{R}}$, the variety of characters is more subtle, cf. [5], but we just consider $\mathcal{R}^{\text{good}}(\Gamma, G_{\mathbb{R}}) = R^{\text{good}}(\Gamma, G_{\mathbb{R}}) / G_{\mathbb{R}}$.

For a compact 2 orbifold \mathcal{O}^2 , we may also consider the *relative character variety*, defined as

$$\mathcal{R}^{\text{good}}(\mathcal{O}^2, \partial\mathcal{O}^2, G) = \{[\rho] \in \mathcal{R}^{\text{good}}(\mathcal{O}^2, G) \mid [\rho|_{\partial_1\mathcal{O}^2}], \dots, [\rho|_{\partial_k\mathcal{O}^2}] \text{ are constant}\},$$

where $\partial\mathcal{O}^2 = \partial_1\mathcal{O}^2 \sqcup \dots \sqcup \partial_k\mathcal{O}^2$ is the decomposition in connected components and $[\rho|_{\partial_i\mathcal{O}^2}]$ denotes the conjugacy class of the restriction to the i -th boundary component.

2.1 Zariski tangent space and cohomology

For a representation $\rho: \Gamma \rightarrow G$, its composition with the adjoint representation is denoted by

$$\text{Ad}\rho: \Gamma \rightarrow \text{Aut}(\mathfrak{g}).$$

The adjoint representation preserves the Killing form

$$\mathcal{B}: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C},$$

that is non-degenerate, as we assume G semi-simple.

A *derivation* or *crossed morphism* is a \mathbb{C} -linear map $d: \Gamma \rightarrow \mathfrak{g}$ that satisfies

$$d(\gamma_1 \gamma_2) = d(\gamma_1) + \text{Ad}_{\rho(\gamma_1)}(d(\gamma_2)), \quad \forall \gamma_1, \gamma_2 \in \Gamma.$$

The space of crossed morphisms is denoted by $Z^1(\Gamma, \text{Ad}\rho)$. A crossed morphism is called *inner* if there exists $a \in \mathfrak{g}$ such that

$$d(\gamma) = (\text{Ad}_{\rho(\gamma)} - 1)(a), \quad \forall \gamma \in \Gamma,$$

and the subspace of inner morphism is denoted by $B^1(\Gamma, \text{Ad}\rho)$. The quotient is the first cohomology group:

$$H^1(\Gamma, \mathfrak{g}_{\text{Ad}\rho}) = H^1(\Gamma, \text{Ad}\rho) \cong Z^1(\Gamma, \text{Ad}\rho) / B^1(\Gamma, \text{Ad}\rho).$$

Theorem 2.2 (Weil). *Let $T_\rho^{\text{Zar}} R(\Gamma, G)$ be the Zariski tangent space of $R(\Gamma, G)$ as a scheme at ρ . There is a natural isomorphism*

$$Z^1(\Gamma, \text{Ad}\rho) \cong T_\rho^{\text{Zar}} R(\Gamma, G)$$

that maps $B^1(\Gamma, \text{Ad}\rho)$ to the tangent space to the orbit by conjugation. Furthermore, when ρ is good,

$$H^1(\Gamma, \text{Ad}\rho) \cong T_\rho^{\text{Zar}} X(\Gamma, G).$$

We do not need the precise definition of scheme, just mention that the polynomial ideal that defines $R(\Gamma, G)$ may be nonreduced, and this is taken into account in the Zariski tangent space. In particular, when the dimension of the Zariski tangent space at some representation or character equals the dimension of the component, the variety is smooth (and the scheme is reduced and smooth) at this representation or character. For a discussion of Theorem 2.2, see for instance [17] and the references therein. There is a real version of this theorem: when the image of a good representations is contained in a real form $G_{\mathbb{R}}$, then

$$H^1(\Gamma, \text{Ad}\rho_{\mathbb{R}}) \cong T_\rho^{\text{Zar}} \mathcal{R}^{\text{good}}(\Gamma, G_{\mathbb{R}}).$$

When \mathcal{O}^2 is orientable, a representation $\rho: \pi_1(\mathcal{O}^2) \rightarrow G$ is called ∂ -regular if for each γ generator of a peripheral subgroup, $\rho(\gamma)$ is a regular element (where regular means $\dim \mathfrak{g}^{\rho(\gamma)} = \text{rank } G$). For non-orientable \mathcal{O}^2 , we call a representation ∂ -regular if its restriction to the orientation covering is ∂ -regular.

Proposition 2.3 ([17]). *Assume that \mathcal{O}^2 is orientable and that $\rho: \pi_1(\mathcal{O}^2) \rightarrow G$ is good and ∂ -regular. Then*

$$T_\rho^{\text{Zar}} \mathcal{R}^{\text{good}}(\mathcal{O}^2, \partial\mathcal{O}^2, G) \cong \ker \left(H^1(\pi_1(\mathcal{O}^2), \text{Ad}\rho) \rightarrow \bigoplus_{i=1}^k H^1(\pi_1(\partial_i \mathcal{O}^2), \text{Ad}\rho) \right).$$

2.2 The principal representation

Let G be a simple complex Lie group, with Lie algebra \mathfrak{g} . An element in \mathfrak{g} is called *regular* if its centralizer has minimal dimension, that is the rank of \mathfrak{g} . Given a regular nilpotent element, Jacobson-Morozov theorem provides a representation of Lie algebras $\tau: \mathfrak{sl}(2, \mathbb{C}) \rightarrow \mathfrak{g}$ whose image contains the given regular nilpotent element. Such a representation is unique up to conjugacy. Assume that G is an adjoint group, namely that has trivial center. Then the induced representation of Lie groups

$$\tau: \mathrm{PSL}(2, \mathbb{C}) \rightarrow G$$

is called the *principal representation*.

For $G = \mathrm{PSL}(n, \mathbb{C})$ the principal representation is Sym^{n-1} . When defined from $\mathrm{SL}(2, \mathbb{C})$ to $\mathrm{SL}(n, \mathbb{C})$, Sym^{n-1} preserves a bilinear form of \mathbb{C}^{n-1} , that is symmetric for n odd and skew-symmetric for n even, and its restriction yields the principal representations in $\mathrm{PSp}(2m, \mathbb{C})$ and $\mathrm{PO}(2m+1, \mathbb{C})$.

The principal representation restricts to $\tau: \mathrm{PGL}(2, \mathbb{R}) \rightarrow G_{\mathbb{R}}$, for $G_{\mathbb{R}}$ the split real form of G , perhaps not connected. The Hitchin component is the connected component of $\mathcal{R}^{\mathrm{good}}(\pi_1(\mathcal{O}^2), G_{\mathbb{R}})$ that contains the composition of τ with the holonomy representation of any Fuchsian structure on \mathcal{O}^2 .

We require that the real form contains the image of $\mathrm{PGL}(2, \mathbb{R})$ and therefore may be not connected. For instance $\mathrm{PGL}(n, \mathbb{R})$ contains two components, according to the sign of the determinant. We also use the notation

$$\mathrm{Sp}^{\pm}(2m) = \{A \in \mathrm{SL}(2m, \mathbb{R}) \mid A^t J A = \pm J\}$$

for J an antisymmetric, non-degenerate, bilinear form; the sign $+$ or $-$ corresponds to one or the other component of the group.

Remark 2.4. *The principal representation is irreducible, and its image consists of regular elements (and the trivial element).*

For the principal representation $\tau: \mathrm{PSL}(2, \mathbb{C}) \rightarrow G$ we have a decomposition

$$\mathrm{Ad} \circ \tau = \bigoplus_{i=1}^r \mathrm{Sym}^{2d_i}, \quad (2)$$

where $r = \mathrm{rank} G$. The $d_1, \dots, d_r \in \mathbb{N}$ are denoted the *exponents* of G . In particular,

$$\dim G = \sum_{i=1}^r (2d_i + 1). \quad (3)$$

For instance, the exponents of $\mathrm{PSL}(n, \mathbb{C})$ are $1, \dots, n-1$, because

$$\mathrm{Ad} \circ \mathrm{Sym}^{n-1} = \bigoplus_{i=1}^{n-1} \mathrm{Sym}^{2i}.$$

The exponents of simple Lie algebras are detailed in Tables 1 and 2, cf. [1].

The algebra of polynomial functions on the Cartan subalgebra invariant by the Weyl group $\mathbb{C}[\mathfrak{h}]^W$ is freely generated by r homogeneous polynomials. By a theorem of Kostant, the degree of these polynomials is $d_1 + 1, \dots, d_r + 1$.

Lie algebra	Exponents	Rank	Dimension
$\mathfrak{sl}(n, \mathbb{C})$	$1, 2, \dots, n-1$	$n-1$	n^2-1
$\mathfrak{sp}(2m, \mathbb{C})$	$1, 3, 5, \dots, 2m-1$	m	$2m^2+m$
$\mathfrak{so}(2m+1, \mathbb{C})$	$1, 3, 5, \dots, 2m-1$	m	$2m^2+m$
$\mathfrak{so}(2m, \mathbb{C})$	$1, 3, 5, \dots, 2m-3, m-1$	m	$2m^2-m$

Table 1: Exponents of simple classical Lie algebras

Lie algebra	Exponents	Rank	Dimension
\mathfrak{g}_2	$1, 5$	2	14
\mathfrak{f}_4	$1, 5, 7, 11$	4	52
\mathfrak{e}_6	$1, 4, 5, 7, 8, 11$	6	78
\mathfrak{e}_7	$1, 5, 7, 9, 11, 13, 17$	7	133
\mathfrak{e}_8	$1, 7, 11, 13, 17, 19, 23, 29$	8	248

Table 2: Exponents of exceptional Lie algebras

3 Orbifold cohomology

Group cohomology is useful to study dimensions, by Weil's theorem (Theorem 2.2). In this section we discuss a combinatorial approach to cohomology. We are interested in representations in semi-simple complex algebraic groups, but this section applies also to their real forms, by replacing complex dimension by real dimension. We focus in homology and cohomology with coefficients twisted by the adjoint of a representation, but the constructions and results can be easily adapted to coefficients twisted by other representations.

We recall some basic definitions of orbifolds [2, 31]. An orbifold \mathcal{O} is called:

- *good* if it has an orbifold cover that is a manifold.
- *very good* if it has an orbifold cover of finite index that is a manifold.
- *aspherical* if its universal covering is a contractible manifold.
- An orbifold is called *hyperbolic/Euclidean/spherical* if it is the quotient of hyperbolic space/Euclidean space/the round sphere by a discrete group of isometries.

In a cell decomposition K of an orbifold, we require that isotropy groups or stabilizers are unchanged along cells. For a cell e in K this isotropy group is denoted by $\text{Stab}(e)$. The *orbifold Euler characteristic* for a compact orbifold is defined as

$$\chi(\mathcal{O}) = \sum_{e \text{ cell of } K} \frac{1}{|\text{Stab}(e)|} (-1)^{\dim e} \in \mathbb{Q}. \quad (4)$$

This is well behaved under coverings: if $\mathcal{O}' \rightarrow \mathcal{O}$ is an orbifold covering of finite index k , then $\chi(\mathcal{O}') = k\chi(\mathcal{O})$. See [31, 2] for instance.

3.1 Combinatorial homology and cohomology for orbifolds

Let \mathcal{O}^n be a compact n -dimensional good orbifold, possibly not orientable. Fix a CW-complex structure K on \mathcal{O}^n . This means that K is a CW-complex with the same underlying space as \mathcal{O}^n ,

so that the branching locus of \mathcal{O}^n is a subcomplex of K . In particular, it lifts to a CW-complex structure \tilde{K} of the universal covering $\tilde{\mathcal{O}}^n$ of \mathcal{O}^n , so that each cell of K lifts to a disjoint union of homeomorphic cells of \tilde{K} (perhaps with nontrivial stabilizer).

The complex of chains on the universal covering is the free \mathbb{Z} -module on the cells of \tilde{K} , equipped with the usual boundary operator, and it is denoted by $C_*(\tilde{K}, \mathbb{Z})$. It has a (non-free) action of $\Gamma = \pi_1(\mathcal{O}^n)$ induced by deck transformations. The *twisted* chain and cochain complexes are defined as:

$$C_*(K, \text{Ad}\rho) = \mathfrak{g} \otimes_{\Gamma} C(\tilde{K}, \mathbb{Z}), \quad (5)$$

$$C^*(K, \text{Ad}\rho) = \text{hom}_{\Gamma}(C(\tilde{K}, \mathbb{Z}), \mathfrak{g}). \quad (6)$$

The group Γ acts on \mathfrak{g} via $\text{Ad}\rho$ on the left, and for the tensor product Γ acts on \mathfrak{g} on the right using inverses. Those are complexes and cocomplexes of finite-dimensional vector spaces, and the corresponding homology and cohomology groups are denoted respectively by

$$H_*(K, \text{Ad}\rho) \quad \text{and} \quad H^*(K, \text{Ad}\rho).$$

3.2 Twisted orbifold Euler characteristic

For a compact manifold M and a representation $\rho: \pi_1(M) \rightarrow G$, as $\pi_1(M)$ acts freely on the universal covering \tilde{M} , we have

$$\chi(M) \dim(G) = \sum_i (-1)^i \dim H_i(M, \text{Ad}\rho) = \sum_i (-1)^i \dim H^i(M, \text{Ad}\rho).$$

To have a similar formula for an orbifold \mathcal{O} we need to take into account that $\Gamma = \pi_1(\mathcal{O})$ acts non freely on $\tilde{\mathcal{O}}$; this motivates the definition of twisted orbifold Euler characteristic below, Definition 3.1.

Let \mathcal{O} be compact oriented orbifold, very good, with a CW-structure K . Given a subset $\Gamma_0 \subset \Gamma = \pi_1(\mathcal{O})$, the space of *invariants* is

$$\mathfrak{g}^{\rho(\Gamma_0)} = \{v \in \mathfrak{g} \mid \text{Ad}_{\rho(g)}(v) = v, \forall g \in \Gamma_0\}, \quad (7)$$

and the quotient of *coinvariants*,

$$\mathfrak{g}_{\rho(\Gamma_0)} = \mathfrak{g} / \langle \text{Ad}_{\rho(g)}(v) - v \mid v \in \mathfrak{g}, g \in \Gamma_0 \rangle. \quad (8)$$

As the Killing form on \mathfrak{g} is nondegenerate (G is semisimple) and Ad -invariant,

$$\langle \text{Ad}_{\rho(g)}(v) - v \mid v \in \mathfrak{g}, g \in \Gamma_0 \rangle = (\mathfrak{g}^{\rho(\Gamma_0)})^{\perp}. \quad (9)$$

Therefore

$$\dim(\mathfrak{g}^{\rho(\Gamma_0)}) = \dim(\mathfrak{g}_{\rho(\Gamma_0)}). \quad (10)$$

Definition 3.1. *The orbifold Euler characteristic of \mathcal{O} twisted by $\text{Ad}\rho$ is*

$$\tilde{\chi}(\mathcal{O}, \text{Ad}\rho) = \sum_{e \text{ cell of } K} (-1)^{\dim e} \dim \mathfrak{g}^{\rho(\text{Stab}(\tilde{e}))} \in \mathbb{Z}.$$

Here \tilde{e} denotes any lift of e to the universal covering of \mathcal{O} and $\text{Stab}(\tilde{e}) \subset \Gamma$ its stabilizer, whose conjugacy class in Γ is independent of the choice of the lift.

It should not be confused with the usual orbifold Euler characteristic $\chi(\mathcal{O})$, recalled in (4), that is a rational number, whilst $\tilde{\chi}(\mathcal{O}, \text{Ad}\rho)$ is always an integer. Notice also that $\tilde{\chi}$ is not multiplicative under coverings either; its intended to be a tool to compute dimensions of cohomology groups:

Proposition 3.2. *For \mathcal{O} and ρ as above:*

$$\tilde{\chi}(\mathcal{O}, \text{Ad}\rho) = \sum_i (-1)^i \dim H_i(\mathcal{O}, \text{Ad}\rho) = \sum_i (-1)^i \dim H^i(\mathcal{O}, \text{Ad}\rho).$$

Proof. We compute the dimension of $C_*(K, \text{Ad}\rho)$ and $C^*(K, \text{Ad}\rho)$ as \mathbb{C} -vector spaces. We aim to show that, for $0 \leq i \leq \dim \mathcal{O}$,

$$\dim C_i(K, \text{Ad}\rho) = \dim C^i(K, \text{Ad}\rho) = \sum_{j=1}^{k_i} \dim \mathfrak{g}^{\rho(\text{Stab}(\tilde{e}_j^i))} \quad (11)$$

where $\{e_1^i, \dots, e_{k_i}^i\}$ are the i -cells of K and $\{\tilde{e}_1^i, \dots, \tilde{e}_{k_i}^i\}$ a choice of lifts in the universal covering. Equation (11) will imply that

$$\tilde{\chi}(\mathcal{O}, \text{Ad}\rho) = \sum_i (-1)^i \dim C_i(K, \text{Ad}\rho) = \sum_i (-1)^i \dim C^i(K, \text{Ad}\rho) \quad (12)$$

and then the proposition will follow from standard arguments in homological algebra.

To prove (11), use the decomposition as $\mathbb{Z}[\Gamma]$ -module of the chain complex on \tilde{K} :

$$C_i(\tilde{K}, \mathbb{Z}) = \bigoplus_{j=1}^{k_i} \mathbb{Z}[\Gamma] \tilde{e}_j^i, \quad (13)$$

because the Γ -orbits of the lifts $\{\tilde{e}_1^i, \dots, \tilde{e}_{k_i}^i\}$ give a partition of the i -cells of \tilde{K} . Then we apply the natural isomorphisms of \mathbb{C} -vector spaces, for each cell \tilde{e} of \tilde{K} :

$$\begin{array}{ccc} \text{hom}_{\Gamma}(\mathbb{Z}[\Gamma] \tilde{e}, \mathfrak{g}) & \cong & \mathfrak{g}^{\rho(\text{Stab}(\tilde{e}))} \\ \theta & \mapsto & \theta(\tilde{e}) \end{array} \quad \begin{array}{ccc} \mathfrak{g} \otimes_{\Gamma} \mathbb{Z}[\Gamma] \tilde{e} & \cong & \mathfrak{g}_{\rho(\text{Stab}(\tilde{e}))} \\ v \otimes \tilde{e} & \mapsto & v \end{array} \quad (14)$$

whose proof is straightforward and use Equality (10). \square

For 2-dimensional orientable orbifolds, Proposition 3.2 is essentially a formula computed by André Weil in [32, Sections 6 and 7].

3.3 Regular Coverings

Let $\mathcal{O}_0 \rightarrow \mathcal{O}$ be a finite regular covering. Namely, $\Gamma_0 = \pi_1(\mathcal{O}_0)$ is a normal subgroup of $\Gamma = \pi_1(\mathcal{O})$ of finite index. Though \mathcal{O} is very good, we do not require \mathcal{O}_0 to be a manifold in this subsection (for instance, \mathcal{O}_0 can be the orientation covering).

The *Galois group*, or *group of deck transformations*, of the covering is Γ/Γ_0 . It acts naturally on the chain and cochain complexes, $C_*(K_0, \text{Ad}\rho)$ and $C^*(K_0, \text{Ad}\rho)$. Namely, any $\gamma \in \Gamma$ maps a chain $m \otimes c \in C_*(K_0, \text{Ad}\rho)$ to $m \cdot \gamma^{-1} \otimes \gamma c = \text{Ad}\rho(\gamma)(m) \otimes \gamma c$, cf. [27]. The action of Γ_0 is trivial by construction and therefore we have an action of Γ/Γ_0 . Similarly, $\gamma \in \Gamma$ maps a cochain $\theta \in C^*(K_0, \text{Ad}\rho)$ to $\text{Ad}(\rho(\gamma)) \circ \theta \circ \gamma^{-1}$, which again induces an action of the Galois group Γ/Γ_0 . The subspace of elements *fixed* by this action in homology and cohomology are denoted respectively by

$$H_*(\mathcal{O}_0, \text{Ad}\rho)^{\Gamma/\Gamma_0} \quad \text{and} \quad H^*(\mathcal{O}_0, \text{Ad}\rho)^{\Gamma/\Gamma_0}.$$

Proposition 3.3. *The map $\pi: \mathcal{O}_0 \rightarrow \mathcal{O}$ induces an epimorphism $\pi_*: H_*(\mathcal{O}_0, \text{Ad}\rho) \rightarrow H_*(\mathcal{O}, \text{Ad}\rho)$ that restricts to an isomorphism*

$$H_*(\mathcal{O}_0, \text{Ad}\rho)^{\Gamma/\Gamma_0} \cong H_*(\mathcal{O}, \text{Ad}\rho).$$

Similarly, it induces a monomorphism $\pi^: H^*(\mathcal{O}, \text{Ad}\rho) \hookrightarrow H^*(\mathcal{O}_0, \text{Ad}\rho)$ that yields an isomorphism*

$$H^*(\mathcal{O}, \text{Ad}\rho) \cong H^*(\mathcal{O}_0, \text{Ad}\rho)^{\Gamma/\Gamma_0}.$$

Proof. The argument is standard and we just outline it. The proof is based in constructing a section to the maps induced by the projection π . The section at the chain and cochain level consists in taking an element of the inverse image of the map induced by π and averaging by the action of Γ/Γ_0 . One can check that the section for chains is a well defined chain morphism, induces sections in homology and has the required property, as well as for sections of cochains and cohomology. \square

The following proposition summarizes the main properties we need about orbifold cohomology:

Proposition 3.4. *Let \mathcal{O} be a compact very good orbifold and ρ a representation of $\Gamma = \pi_1(\mathcal{O})$ in G . The following hold:*

- (a) *$H^i(\mathcal{O}, \text{Ad}\rho)$ and $H_i(\mathcal{O}, \text{Ad}\rho)$ are dual, via the Kronecker pairing.*
- (b) *If \mathcal{O} is orientable, then the cup product induces a perfect pairing*

$$H^i(\mathcal{O}, \text{Ad}\rho) \times H^{\dim \mathcal{O} - i}(\mathcal{O}, \partial \mathcal{O}, \text{Ad}\rho) \rightarrow \mathbb{C}.$$

- (c) *There is a natural isomorphism $H^1(\Gamma, \text{Ad}\rho) \cong H^1(\mathcal{O}, \text{Ad}\rho)$.*
- (d) *If \mathcal{O} is aspherical (its universal covering is a contractible manifold), then there is a natural isomorphism $H^*(\Gamma, \text{Ad}\rho) \cong H^*(\mathcal{O}, \text{Ad}\rho)$.*

Proof. For assertion (a), we use the Kronecker pairing between chains and cochains via the Killing form \mathcal{B} :

$$\begin{array}{ccc} C_i(K, \text{Ad}\rho) \times C^i(K, \text{Ad}\rho) & \rightarrow & \mathbb{C} \\ (m \otimes c, \theta) & \mapsto & \mathcal{B}(m, \theta(c)) \end{array}$$

One checks that it is well defined and it is not degenerate at the level of chains, using (13), (14), and (9). Hence it induces a perfect pairing between homology and cohomology.

For (b), using Proposition 3.3, the strategy is to use the invariance by Γ/Γ_0 of the corresponding properties for a finite regular covering $\mathcal{O}_0 \rightarrow \mathcal{O}$ that is a manifold. More precisely, using that \mathcal{O}_0 is a manifold, the cup product defines a perfect pairing

$$H^i(\mathcal{O}_0, \text{Ad}\rho) \times H^{\dim \mathcal{O}_0 - i}(\mathcal{O}_0, \partial \mathcal{O}_0, \text{Ad}\rho) \rightarrow H^{\dim \mathcal{O}_0}(\mathcal{O}_0, \partial \mathcal{O}_0, \mathbb{C}) \cong \mathbb{C}.$$

This is compatible with the action of Γ/Γ_0 , therefore the assertion follows from Proposition 3.3. Notice that orientability of \mathcal{O} is relevant for saying that the action of the Galois group Γ/Γ_0 on $H^{\dim \mathcal{O}_0}(\mathcal{O}_0, \partial \mathcal{O}_0, \mathbb{C}) \cong \mathbb{C}$ is trivial.

Next we prove (c). Chose X a $K(\Gamma, 1)$, namely a CW-complex with $\pi_1(X) \cong \Gamma$ and trivial higher homotopy groups. Thus there is a natural isomorphism $H^*(X, \text{Ad}\rho) \cong H^*(\Gamma, \text{Ad}\rho)$. Furthermore, the covering corresponding to $\Gamma_0 < \Gamma$, $X_0 \rightarrow X$, is a $K(\Gamma_0, 1)$; in particular

$H^*(X_0, \text{Ad}\rho) \cong H^*(\Gamma_0, \text{Ad}\rho)$. As \mathcal{O}_0 is a manifold, a $K(\Gamma_0, 1)$ can be constructed from the 2-skeleton of \mathcal{O}_0 by adding cells of dimension ≥ 3 , therefore:

$$H^1(\mathcal{O}_0, \text{Ad}\rho) \cong H^1(X_0, \text{Ad}\rho).$$

Thus, by Proposition 3.3 we have natural isomorphisms

$$H^1(\mathcal{O}, \text{Ad}\rho) \cong H^1(\mathcal{O}_0, \text{Ad}\rho)^{\Gamma/\Gamma_0} \cong H^1(X_0, \text{Ad}\rho)^{\Gamma/\Gamma_0} \cong H^1(X, \text{Ad}\rho),$$

and $H^1(X, \text{Ad}\rho) \cong H^1(\Gamma, \text{Ad}\rho)$ because X is a $K(\Gamma, 1)$.

Finally, we prove (d). In the proof of (c), as \mathcal{O}_0 is already a $K(\Gamma_0, 1)$, and therefore homotopy equivalent to X_0 , hence

$$H^*(\mathcal{O}_0, \text{Ad}\rho) \cong H^*(X_0, \text{Ad}\rho).$$

and the conclusion holds for any degree in cohomology. \square

From the duality in Proposition 3.4 (b) we get:

Corollary 3.5. *When \mathcal{O}^n is closed and orientable, of dimension n , for n odd $\tilde{\chi}(\mathcal{O}^n, \text{Ad}\rho) = 0$, and for n even $\tilde{\chi}(\mathcal{O}^n, \text{Ad}\rho)$ is also even.*

3.4 Non orientable orbifolds

For a connected non-orientable orbifold \mathcal{O}^n , there exists a unique orientable covering $\widetilde{\mathcal{O}}^n \rightarrow \mathcal{O}^n$ of index 2, called the *orientation covering*.

Lemma 3.6. *Let \mathcal{O}^n be a compact, non-orientable, very good orbifold of dimension n with orientation covering $\widetilde{\mathcal{O}}^n \rightarrow \mathcal{O}^n$. Then*

$$\dim H^i(\mathcal{O}^n, \text{Ad}\rho) + \dim H^{n-i}(\mathcal{O}^n, \partial\mathcal{O}^n, \text{Ad}\rho) = \dim H^i(\widetilde{\mathcal{O}}^n, \text{Ad}\rho).$$

Proof. The nontrivial deck transformation of the orientation covering is an involution denoted by $\sigma: \widetilde{\mathcal{O}}^n \rightarrow \widetilde{\mathcal{O}}^n$. Let σ^* be the induced morphism in the cohomology group. By Proposition 3.3:

$$H^i(\mathcal{O}^n, \text{Ad}\rho) \cong H^i(\widetilde{\mathcal{O}}^n, \text{Ad}\rho)^{\sigma^*} \quad \text{and} \quad H^i(\mathcal{O}^n, \partial\mathcal{O}^n, \text{Ad}\rho) \cong H^i(\widetilde{\mathcal{O}}^n, \partial\widetilde{\mathcal{O}}^n, \text{Ad}\rho)^{\sigma^*}. \quad (15)$$

The cup product induces a non-degenerate pairing, Proposition 3.4:

$$H^i(\widetilde{\mathcal{O}}^n, \text{Ad}\rho) \times H^{n-i}(\widetilde{\mathcal{O}}^n, \partial\widetilde{\mathcal{O}}^n, \text{Ad}\rho) \rightarrow H^n(\widetilde{\mathcal{O}}^n, \partial\widetilde{\mathcal{O}}^n, \mathbb{C}) \cong \mathbb{C}. \quad (16)$$

Chose \mathbb{C} -basis for $H^i(\widetilde{\mathcal{O}}^n, \text{Ad}\rho)$ and for $H^{n-i}(\widetilde{\mathcal{O}}^n, \partial\widetilde{\mathcal{O}}^n, \text{Ad}\rho)$, and use them to get matrices; let

- J denote the matrix of the pairing (16),
- S^i denote the matrix of σ^* on $H^i(\widetilde{\mathcal{O}}^n, \text{Ad}\rho)$, and
- T^j denote the matrix of σ^* on $H^j(\widetilde{\mathcal{O}}^n, \partial\widetilde{\mathcal{O}}^n, \text{Ad}\rho)$.

Since σ reverses the orientation, σ^* acts as minus the identity on $H^n(\widetilde{\mathcal{O}}^n, \partial\widetilde{\mathcal{O}}^n, \mathbb{C})$:

$$(S^i)^t J T^{n-i} = -J.$$

As the pairing is non-degenerate, J is invertible, and since σ is an involution:

$$(S^i)^t = -J T^{n-i} J^{-1}.$$

Thus

$$\dim \ker(T^{n-i} - \text{Id}) = \dim \ker((S^i)^t + \text{Id}) = \dim \ker(S^i + \text{Id}). \quad (17)$$

Since $(S^i)^2 = \text{Id}$:

$$H^i(\widetilde{\mathcal{O}}^n, \text{Ad}\rho) = \ker(S^i + \text{Id}) \oplus \ker(S^i - \text{Id}). \quad (18)$$

Combining (17) and (18):

$$\dim \ker(T^{n-i} - \text{Id}) + \dim \ker(S^i - \text{Id}) = \dim H^i(\widetilde{\mathcal{O}}^n, \text{Ad}\rho).$$

With (15) this concludes the proof. \square

Corollary 3.7. *If \mathcal{O}^n is non-orientable, very good, closed and of even dimension n , then*

$$\dim H^{\frac{n}{2}}(\mathcal{O}^n, \text{Ad}\rho) = \frac{1}{2} \dim H^{\frac{n}{2}}(\widetilde{\mathcal{O}}^n, \text{Ad}\rho).$$

Remark 3.8. *The pairing in the proof of Lemma 3.6 induces a nondegenerate pairing between the kernels $\ker(H^i(\mathcal{O}^n, \text{Ad}\rho) \rightarrow H^i(\partial\mathcal{O}^n, \text{Ad}\rho))$ and $\ker(H^{n-i}(\mathcal{O}^n, \text{Ad}\rho) \rightarrow H^{n-i}(\partial\mathcal{O}^n, \text{Ad}\rho))$ [22, 17]. Thus the same argument in the proof of Lemma 3.6 yields*

$$\begin{aligned} & \dim \ker(H^i(\mathcal{O}^n, \text{Ad}\rho) \rightarrow H^i(\partial\mathcal{O}^n, \text{Ad}\rho)) \\ & \quad + \dim \ker(H^{n-i}(\mathcal{O}^n, \text{Ad}\rho) \rightarrow H^{n-i}(\partial\mathcal{O}^n, \text{Ad}\rho)) \\ & \quad = \dim \ker(H^i(\widetilde{\mathcal{O}}^n, \text{Ad}\rho) \rightarrow H^i(\partial\widetilde{\mathcal{O}}^n, \text{Ad}\rho)). \end{aligned}$$

4 Varieties of representations and characters of 2-orbifolds

We recall the possible stabilizers (or isotropy groups) of a point in a 2-orbifold, so that we fix notation. Besides the trivial one, the possible stabilizers of a point $x \in \mathcal{O}^2$ are:

- $\text{Stab}(x) \cong C_k$ is a cyclic group of rotations of the plane \mathbb{R}^2 of order k . Then the singular point is isolated and its is called a *cone point*. Those are the unique possible non-trivial stabilizers for an orientable 2-orbifold (Figure 1 left).
- $\text{Stab}(x) \cong C_2$ is a group of order two generated by a reflection of the plane or the semiplane, ie. modelled in the interior or in the boundary. This is called a *mirror point*. The singular locus is locally an open segment, or a proper half segment in the boundary (Figure 1, both pictures in the center).
- $\text{Stab}(x) \cong D_{2k}$ is a dihedral group of $2k$ elements, generated by two reflections at the plane of angle π/k . The point is called a *corner reflector* (Figure 1 right).



Figure 1: From left to right, a cone point, an interior mirror point, a mirror point in the boundary, and a corner reflector. Mirror points are represnetes by a double line.

The boundary of a compact 2-orbifold is a union of closed 1-orbifolds. There are two possibilities for a closed 1-orbifold, up to homeomorphism:

- A circle S^1 (Figure 2 left). Its fundamental group is \mathbb{Z} .
- An interval with mirror end-points $\llbracket 0, 1 \rrbracket$ (Figure 2 right). It is the quotient of S^1 by an orientation reversing involution. Its fundamental group is dihedral infinite $D_\infty \cong C_2 * C_2$, the free product of the stabilizers of the mirror points. It is called *full* 1-orbifolds in [1].




Figure 2: A circle S^1 and an interval with mirror end-points $\llbracket 0, 1 \rrbracket$ as boundaries.

In dimension two, good and very good are equivalent. Furthermore, a closed 2-orbifold is good if and only if it is hyperbolic, Euclidean or spherical. When $\chi(\mathcal{O}^2) \leq 0$, \mathcal{O}^2 is very good and either hyperbolic or Euclidean. In particular a closed 2-orbifold \mathcal{O}^2 is aspherical iff $\chi(\mathcal{O}^2) \leq 0$.

4.1 Varieties of representations of 2-orbifolds

In this subsection we give a couple of results on the tangent space of the variety of representations and characters of a 2-orbifold. Those are orbifold versions of theorems of Goldman for surfaces [10]. In the orientable case, those results go back to André Weil [32, Sections 6 and 7]. We start with the following formulas:

Proposition 4.1. *Let \mathcal{O}^2 be a closed aspherical 2-orbifold. Set $\Gamma = \pi_1(\mathcal{O}^2)$ and let $\rho \in R(\Gamma, G)$.*

a) If \mathcal{O}^2 is orientable, then

$$\dim T_\rho^{Zar} R(\Gamma, G) = -\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) + \dim G + \dim \mathfrak{g}^{\rho(\Gamma)}.$$

b) If \mathcal{O}^2 is non-orientable, with orientation covering $\tilde{\mathcal{O}}^2$, and $\tilde{\Gamma} = \pi_1(\tilde{\mathcal{O}}^2)$, then

$$\dim T_\rho^{Zar} R(\Gamma, G) = -\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) + \dim G + \dim \mathfrak{g}^{\rho(\tilde{\Gamma})} - \dim \mathfrak{g}^{\rho(\Gamma)}.$$

Proof. Using

$$T_\rho^{Zar} R(\Gamma, G) \cong Z^1(\Gamma, \text{Ad}\rho) = \dim H^1(\Gamma, \text{Ad}\rho) + \dim B^1(\Gamma, \text{Ad}\rho),$$

the isomorphism $H^*(\mathcal{O}^2, \text{Ad}\rho) \cong H^*(\Gamma, \text{Ad}\rho)$ (as \mathcal{O}^2 is aspherical we apply Proposition 3.4 (d)), and:

$$\dim H^1(\mathcal{O}^2, \text{Ad}\rho) = -\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) + \dim H^0(\mathcal{O}^2, \text{Ad}\rho) + \dim H^2(\mathcal{O}^2, \text{Ad}\rho),$$

$$\dim B^1(\Gamma, \text{Ad}\rho) = \dim \mathfrak{g} - \dim \mathfrak{g}^{\rho(\Gamma)} = \dim G - \dim H^0(\Gamma, \text{Ad}\rho).$$

we get

$$\dim T_\rho^{Zar} R(\Gamma, G) = -\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) + \dim G + \dim H^2(\mathcal{O}^2, \text{Ad}\rho).$$

In the orientable case, by duality (Proposition 3.4):

$$\dim H^2(\mathcal{O}^2, \text{Ad}\rho) = \dim H^0(\mathcal{O}^2, \text{Ad}\rho) = \dim \mathfrak{g}^{\rho(\Gamma)}.$$

In the non-orientable case, Lemma 3.6 yields

$$\dim H^2(\mathcal{O}^2, \text{Ad}\rho) = \dim H^2(\tilde{\mathcal{O}}^2, \text{Ad}\rho) - \dim H^0(\mathcal{O}^2, \text{Ad}\rho) = \dim \mathfrak{g}^{\rho(\tilde{\Gamma})} - \dim \mathfrak{g}^{\rho(\Gamma)},$$

and the proposition follows. \square

Proposition 4.2. *Let \mathcal{O}^2 be a compact aspherical 2-orbifold and $\rho: \Gamma \rightarrow G$ a good representation. When \mathcal{O}^2 is non-orientable, assume furthermore that the restriction of ρ to the orientation covering is also good.*

Then the conjugacy class $[\rho]$ is a smooth point of $X(\mathcal{O}^2, G)$ of dimension $-\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho)$.

Proof. As ρ is good, its centralizer is finite, hence

$$\dim H^0(\mathcal{O}^2, \text{Ad}\rho) = \dim H^0(\Gamma, \text{Ad}\rho) = \dim \mathfrak{g}^{\rho(\Gamma)} = 0.$$

When \mathcal{O}^2 is closed then $H^2(\mathcal{O}^2, \text{Ad}\rho) = 0$ (by duality in the orientable case, Proposition 3.4, or by Lemma 3.6 in the non-orientable case). When \mathcal{O}^2 is not closed, then it has virtually the homotopy type of a 1-complex and also $H^2(\mathcal{O}^2, \text{Ad}\rho) = 0$.

Moreover, as \mathcal{O}^2 is aspherical, by Proposition 3.4 $H^2(\Gamma, \text{Ad}\rho) = 0$. By Goldman's obstruction theory [10], this implies that $[\rho]$ is a smooth point of local dimension $\dim H^1(\Gamma, \text{Ad}\rho) = \dim H^1(\mathcal{O}, \text{Ad}\rho) = -\tilde{\chi}(\mathcal{O}, \text{Ad}\rho)$. \square

From Proposition 4.2 and Corollary 3.7:

Corollary 4.3. *Let \mathcal{O}^2 be a closed non-orientable 2-orbifold with $\chi(\mathcal{O}^2) \leq 0$ and $\rho: \Gamma \rightarrow G$. Assume that the restriction of ρ to the orientation covering $\tilde{\mathcal{O}}^2$ is good. Then*

$$\dim_{[\rho]} X(\mathcal{O}^2, G) = \frac{1}{2} \dim_{[\rho]} X(\tilde{\mathcal{O}}^2, G).$$

Proposition 4.4. *Let \mathcal{O}^2 be a compact aspherical 2-orbifold and $\rho: \Gamma \rightarrow G$ a good representation that is ∂ -regular. Assume that \mathcal{O}^2 is orientable. Then the cup product defines a \mathbb{C} -valued symplectic structure on $\mathcal{R}^{\text{good}}(\mathcal{O}^2, \partial\mathcal{O}^2, G)$. Furthermore, it is real valued for $G_{\mathbb{R}}$, a real form of G .*

Proof. From the construction, non-degeneracy and skew symmetry is clear. The non-trivial issue is to check that this differential form is closed. For manifolds with boundary, closedness is due to [13], as explained in [17]. For an orientable orbifold with boundary, we reduce to the manifold case. Let $\Sigma_{\mathcal{O}^2}$ denote its branching locus, it is a finite union of cone points. The restriction map

$$\mathcal{R}^{\text{good}}(\mathcal{O}^2, \partial\mathcal{O}^2, G) \rightarrow \mathcal{R}^{\text{good}}(\mathcal{O}^2 \setminus \mathcal{N}(\Sigma_{\mathcal{O}^2}), \partial(\mathcal{O}^2 \setminus \mathcal{N}(\Sigma_{\mathcal{O}^2})), G)$$

is a (local) isomorphism, as the conjugacy classes of finite order elements cannot be deformed. To check the isomorphism at the level of tangent spaces, notice that the map induced by inclusion

$$H^1(\mathcal{O}^2, \text{Ad}\rho) \rightarrow H^1(\mathcal{O}^2 \setminus \mathcal{N}(\Sigma_{\mathcal{O}^2}), \text{Ad}\rho)$$

yields an isomorphism

$$\begin{aligned} \ker (H^1(\mathcal{O}^2, \text{Ad}\rho) \rightarrow H^1(\partial\mathcal{O}^2, \text{Ad}\rho)) \\ \cong \ker (H^1(\mathcal{O}^2 \setminus \mathcal{N}(\Sigma_{\mathcal{O}^2}), \text{Ad}\rho) \rightarrow H^1(\partial(\mathcal{O}^2 \setminus \mathcal{N}(\Sigma_{\mathcal{O}^2})), \text{Ad}\rho)), \end{aligned}$$

that can be proved for instance by using Mayer-Vietoris exact sequence. Furthermore, by naturality, it can be shown that the isomorphism maps the cup product to the cup product. \square

Proposition 4.5. *Let \mathcal{O}^2 be a closed non-orientable 2-orbifold with $\chi(\mathcal{O}^2) \leq 0$. Let $\rho: \Gamma \rightarrow G$ be a representation whose restriction to the orientation covering $\tilde{\mathcal{O}}^2$ is good and ∂ -regular. Then the restriction map induces an immersion around $[\rho]$ of $\mathcal{R}^{\text{good}}(\mathcal{O}^2, \partial\mathcal{O}^2, G)$ in $\mathcal{R}^{\text{good}}(\mathcal{O}^2, \partial\tilde{\mathcal{O}}^2, G)$ as a Lagrangian submanifold.*

Proof. Let $\sigma: \widetilde{\mathcal{O}^2} \rightarrow \widetilde{\mathcal{O}^2}$ be the orienting reversing involution so that $\widetilde{\mathcal{O}^2}/\sigma = \mathcal{O}^2$. By Propositions 3.3 and 4.2

$$T_{[\rho]}^{Zar} \mathcal{R}(\mathcal{O}^2, G) \cong (T_{[\rho]}^{Zar} \mathcal{R}(\widetilde{\mathcal{O}^2}, G))^\sigma$$

As σ acts as minus the identity on $H^2(\widetilde{\mathcal{O}^2}, \mathbb{C})$ and by construction of the bilinear form (see the proof of Lemma 3.6), the bilinear form must be trivial on the space fixed by σ , ie $T_{[\rho]}^{Zar} \mathcal{R}(\mathcal{O}^2, G)$ is an isotropic subspace. By Corollary 4.3, it has the dimension to be Lagrangian. \square

4.2 Orbifolds with boundary

For an orientable 2-orbifold \mathcal{O}^2 a representation ρ is called ∂ -regular if for each γ generator of a peripheral subgroup, $\rho(\gamma)$ is a regular element (i.e. $\dim \mathfrak{g}^{\rho(\gamma)} = \text{rank } G$). For a non-orientable 2-orbifold, we will consider ∂ -regularity on the orientation covering.

We start with a result on the dimension of the relative variety of (conjugacy classes of) representations $\mathcal{R}^{\text{good}}(\mathcal{O}^2, \partial\mathcal{O}^2, G)$:

Proposition 4.6. *Let \mathcal{O}^2 be a compact 2-orbifold with boundary and $\chi(\mathcal{O}^2) \leq 0$. Assume $\rho: \Gamma \rightarrow G$ is good and ∂ -regular. When \mathcal{O}^2 is non-orientable, assume that the restriction to the orientation covering is good and ∂ -regular. Then $[\rho]$ is a smooth point of $\mathcal{R}^{\text{good}}(\mathcal{O}^2, \partial\mathcal{O}^2, G)$ of dimension*

$$-\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) - (\mathfrak{c} + \frac{\mathfrak{b}}{2}) \text{rank } G + \frac{1}{2} \tilde{\chi}(\partial\mathcal{O}^2, \text{Ad}\rho),$$

where \mathfrak{c} is the number of components of $\partial\mathcal{O}^2$ that are circles and \mathfrak{b} the number of components of $\partial\mathcal{O}^2$ that are intervals with mirror points $\llbracket 0, 1 \rrbracket$.

In this proposition, the components of the boundary that are circles do not contribute to $\tilde{\chi}(\partial\mathcal{O}^2, \text{Ad}\rho)$.

Proof. The same proof as in [17, Proposition 2.10] applies here, in particular we have smoothness and the following equalities

$$\begin{aligned} \dim(\mathcal{R}^{\text{good}}(\mathcal{O}^2, \partial\mathcal{O}^2, G)) &= \dim \ker (H^1(\mathcal{O}^2, \text{Ad}\rho) \rightarrow H^1(\partial\mathcal{O}^2, \text{Ad}\rho)) \\ &= \dim H^1(\mathcal{O}^2, \text{Ad}\rho) - \dim H^1(\partial\mathcal{O}^2, \text{Ad}\rho). \end{aligned}$$

As $H^0(\mathcal{O}^2, \text{Ad}\rho) = H^2(\mathcal{O}^2, \text{Ad}\rho) = 0$,

$$\dim H^1(\mathcal{O}^2, \text{Ad}\rho) = -\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho).$$

We count the contribution of each component $\partial_i \mathcal{O}^2$ of $\partial\mathcal{O}^2$ to $\dim H^1(\partial\mathcal{O}^2, \text{Ad}\rho)$.

- When $\partial_i \mathcal{O}^2 \cong S^1$, then, by duality and ∂ -regularity:

$$\dim H^1(\partial_i \mathcal{O}^2, \text{Ad}\rho) = \dim H^0(\partial_i \mathcal{O}^2, \text{Ad}\rho) = \dim \mathfrak{g}^{\rho(\partial_i \mathcal{O}^2)} = \text{rank } G. \quad (19)$$

- When $\partial_i \mathcal{O}^2 \cong \llbracket 0, 1 \rrbracket$, then

$$\begin{aligned} \tilde{\chi}(\partial_i \mathcal{O}^2, \text{Ad}\rho) &= \dim H^0(\partial_i \mathcal{O}^2, \text{Ad}\rho) - \dim H^1(\partial_i \mathcal{O}^2, \text{Ad}\rho) \\ \text{rank } G &= \dim H^0(\partial_i \mathcal{O}^2, \text{Ad}\rho) + \dim H^1(\partial_i \mathcal{O}^2, \text{Ad}\rho), \end{aligned}$$

where the last line follows from Lemma 3.6 and the assumption that ρ is ∂ -regular on the orientation covering. From this we deduce

$$\dim H^1(\partial_i \mathcal{O}^2, \text{Ad}\rho) = \frac{1}{2}(\text{rank } G - \tilde{\chi}(\partial_i \mathcal{O}^2, \text{Ad}\rho)). \quad (20)$$

From (19) and (20):

$$\dim H^1(\partial\mathcal{O}^2, \text{Ad}\rho) = (c + \frac{b}{2}) \text{rank} G - \frac{1}{2} \tilde{\chi}(\partial\mathcal{O}^2, \text{Ad}\rho),$$

which concludes the proof of the proposition. \square

Proposition 4.7. *Let \mathcal{O}^2 be a non-orientable compact 2-orbifold with orientation covering $\widetilde{\mathcal{O}^2}$. Let $\rho: \pi_1(\mathcal{O}^2) \rightarrow G$ be a representation whose restriction to $\widetilde{\mathcal{O}^2}$ is good and ∂ -regular. Then*

$$\dim_{[\rho]} X(\mathcal{O}^2, G) = \frac{1}{2} \dim_{[\rho]} X(\widetilde{\mathcal{O}^2}, G) + \frac{1}{2} \sum_{[0,1] \subset \partial\mathcal{O}^2} \tilde{\chi}([0,1], \text{Ad}\rho)$$

where the sum runs on the components of $\partial\mathcal{O}^2$ that are intervals with mirror boundary, $[0,1]$.

Proof. In the proof of Proposition 4.6 we use (based on [17]):

$$\begin{aligned} \dim_{[\rho]} X(\mathcal{O}^2, G) &= \dim_{[\rho]} \mathcal{R}^{\text{good}}(\mathcal{O}^2, \partial\mathcal{O}^2, G) + \dim H^1(\partial\mathcal{O}^2, \text{Ad}\rho), \\ \dim_{[\rho]} X(\widetilde{\mathcal{O}^2}, G) &= \dim_{[\rho]} \mathcal{R}^{\text{good}}(\widetilde{\mathcal{O}^2}, \partial\mathcal{O}^2, G) + \dim H^1(\partial\widetilde{\mathcal{O}^2}, \text{Ad}\rho). \end{aligned}$$

Furthermore, by Proposition 4.5:

$$\dim_{[\rho]} \mathcal{R}^{\text{good}}(\mathcal{O}^2, \partial\mathcal{O}^2, G) = \frac{1}{2} \dim_{[\rho]} \mathcal{R}^{\text{good}}(\widetilde{\mathcal{O}^2}, \partial\widetilde{\mathcal{O}^2}, G).$$

And from (19) and (20) in the proof of Proposition 4.6:

$$\begin{aligned} \dim H^1(\partial\mathcal{O}^2, \text{Ad}\rho) &= (c + \frac{b}{2}) \text{rank} G - \frac{1}{2} \tilde{\chi}(\partial\mathcal{O}^2, \text{Ad}\rho), \\ \dim H^1(\partial\widetilde{\mathcal{O}^2}, \text{Ad}\rho) &= (2c + b) \text{rank} G. \end{aligned}$$

Using these formulas, we just need to know that $\tilde{\chi}(\partial\mathcal{O}^2, \text{Ad}\rho)$ is the sum of the twisted Euler characteristics of each component, and that $\tilde{\chi}(S^1, \text{Ad}\rho) = 0$. \square




Figure 3: The orbifold $[0,1] = \mathbb{R}/D_\infty$.

The fundamental group of $[0,1]$ is the infinite dihedral group D_∞ , so that $\mathbb{R}/D_\infty \cong [0,1]$. In a group presentation

$$D_\infty = \langle \sigma_1, \sigma_2 \mid \sigma_1^2 = \sigma_2^2 = 1 \rangle$$

the stabilizers of the mirror points are the cyclic groups of order 2 generated by σ_1 and σ_2 respectively, see Figure 3. For further applications we need the following lemma:

Lemma 4.8. *For any representation $\rho: D_\infty = \langle \sigma_1, \sigma_2 \mid \sigma_1^2 = \sigma_2^2 = 1 \rangle \rightarrow G$:*

$$\begin{aligned} \tilde{\chi}(\mathbb{R}/D_\infty, \text{Ad}\rho) &= \dim \mathfrak{g}^{\rho(\sigma_1)} + \dim \mathfrak{g}^{\rho(\sigma_2)} - \dim \mathfrak{g}, \\ \dim H^0(\mathbb{R}/D_\infty, \text{Ad}\rho) &= (\dim \mathfrak{g}^{\rho(\sigma_1\sigma_2)} + \dim \mathfrak{g}^{\rho(\sigma_1)} + \dim \mathfrak{g}^{\rho(\sigma_2)} - \dim \mathfrak{g})/2, \\ \dim H^1(\mathbb{R}/D_\infty, \text{Ad}\rho) &= (\dim \mathfrak{g}^{\rho(\sigma_1\sigma_2)} - \dim \mathfrak{g}^{\rho(\sigma_1)} - \dim \mathfrak{g}^{\rho(\sigma_2)} + \dim \mathfrak{g})/2. \end{aligned}$$

Proof. We compute the twisted Euler characteristic by using the simplicial structure of \mathbb{R}/D_∞ with one 1-cell and two 0-cells (with stabilizers generated by σ_1 and σ_2 respectively):

$$\begin{aligned}\tilde{\chi}(\mathbb{R}/D_\infty, \text{Ad}\rho) &= \dim \mathfrak{g}^{\rho(\sigma_1)} + \dim \mathfrak{g}^{\rho(\sigma_2)} - \dim \mathfrak{g} \\ &= \dim H^0(\mathbb{R}/D_\infty, \text{Ad}\rho) - \dim H^1(\mathbb{R}/D_\infty, \text{Ad}\rho).\end{aligned}\tag{21}$$

The orientation covering is denoted by $S^1 \rightarrow \mathbb{R}/D_\infty$, and $\pi_1(S^1)$ corresponds to the infinite cyclic subgroup of D_∞ generated by $\sigma_1\sigma_2$. As S^1 is a manifold and $\chi(S^1) = 0$:

$$\dim H^1(S^1, \text{Ad}\rho) \cong \dim H^0(S^1, \text{Ad}\rho) \cong \mathfrak{g}^{\rho(\sigma_1\sigma_2)}.\tag{22}$$

By Lemma 3.6:

$$\dim H^0(\mathbb{R}/D_\infty, \text{Ad}\rho) + \dim H^1(\mathbb{R}/D_\infty, \text{Ad}\rho) = \dim \mathfrak{g}^{\rho(\sigma_1\sigma_2)}.\tag{23}$$

Then the lemma follows from (21) and (23). \square

Notice that in the previous lemma we do not require anything for the representation ρ , that could be trivial, but G is a semi-simple group. The first direct application goes to the stabilizer of a corner reflector, the dihedral group D_{2k} , by pre-composing any representation $D_{2k} \rightarrow G$ with the natural surjection from the infinite dihedral group $D_\infty \rightarrow D_{2k}$:




Figure 4: Corner reflector modeled on \mathbb{R}^2/D_{2k} , with the notation in Corollary 4.9.

Corollary 4.9. *Let $x_0 \in \mathcal{O}^2$ be a corner reflector and x_1 and x_2 mirror points in a neighborhood of x_0 separated by x_0 , see Figure 4. If $C_k \subset \text{Stab}(x_0)$ is the orientation preserving cyclic subgroup of index 2, then:*

$$\dim \mathfrak{g}^{\rho(\text{Stab}(x_0))} = (\dim \mathfrak{g}^{\rho(C_k)} + \dim \mathfrak{g}^{\rho(\text{Stab}(x_1))} + \dim \mathfrak{g}^{\rho(\text{Stab}(x_2))} - \dim \mathfrak{g})/2.$$

4.3 Euclidean 2-orbifolds

Along this subsection, let \mathcal{O}^2 be a Euclidean two orbifold without boundary. Set $\Gamma = \pi_1(\mathcal{O}^2)$. As $\text{Isom}^+(\mathbb{R}^2)$ is the semidirect product $\mathbb{R}^2 \rtimes \text{O}(2)$, we have an exact sequence (Bieberbach theorem):

$$1 \rightarrow \Gamma_0 \rightarrow \Gamma \rightarrow \Lambda \rightarrow 1,\tag{24}$$

with $\Gamma_0 \cong \mathbb{Z}^2$ and $\Lambda \subset \text{O}(2)$ the linear part. When \mathcal{O}^2 is orientable, Λ is cyclic and when \mathcal{O}^2 is non-orientable, Λ is dihedral (or cyclic of order 2). Furthermore, when \mathcal{O}^2 is orientable or a Coxeter group, Γ_0 is the maximal torsion free subgroup and the sequence (24) splits, as Λ corresponds to the stabilizer of a point in \mathbb{R}^2 . A Euclidean Coxeter group is a group generated by reflections along a square or a Euclidean triangle with angles an integer divisor of π .

There are precisely five closed orientable Euclidean 2-orbifolds (Figure 5):

- the 2-torus T^2 ,




Figure 5: The five closed, orientable, and Euclidean 2-orbifolds.

- a 2-sphere with 4 cone points of order 2, $S^2(2, 2, 2, 2)$, and
- three 2-spheres with cone points of order p , q and r satisfying $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 1$, $S^2(p, q, r)$ for $(p, q, r) = (3, 3, 3)$, $(2, 4, 4)$ and $(2, 3, 6)$.

The orbifold $S^2(2, 2, 2, 2)$ is sometimes called a *pillowcase*, geometrically is the double of a rectangle, and $S^2(p, q, r)$ is called a *turnover*, as it is the double of a triangle. For different values of the ramifications, turnovers can also be spherical or Euclidean. Table 3 gives the cardinality of $\Lambda \cong \Gamma/\Gamma_0$ for these five Euclidean orbifolds.

\mathcal{O}^2	$ \Gamma/\Gamma_0 $
T^2	1
$S^2(2, 2, 2, 2)$	2
$S^2(3, 3, 3)$	3
$S^2(2, 4, 4)$	4
$S^2(2, 3, 6)$	6

Table 3: Values of $k(\mathcal{O}^2) = |\Gamma/\Gamma_0|$ when \mathcal{O}^2 is orientable, for Γ_0 maximal torsion free.

As $\Gamma = \pi_1(\mathcal{O}^2)$ is virtually abelian, most of the representations we consider are not irreducible. Instead, we deal with strong regularity:

Definition 4.10. A representation of a Euclidean 2-orbifold $\rho: \pi_1(\mathcal{O}^2) \rightarrow G$ is called strongly regular if for a normal subgroup $\Gamma_0 < \pi_1(\mathcal{O}^2)$, $\Gamma_0 \cong \mathbb{Z}^2$,

(a) $\dim \mathfrak{g}^{\rho(\Gamma_0)} = \text{rank } G$, and

(b) the projection of $\rho(\Gamma_0)$ is contained in a connected abelian subgroup of $G/\mathcal{Z}(G)$.

Theorem 4.11. Assume that \mathcal{O}^2 is compact Euclidean and that $\rho: \pi_1(\mathcal{O}^2) \rightarrow G$ is strongly regular. Then:

(i) ρ is a smooth point of $R(\Gamma, G)$ of dimension

$$-\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) + \dim G + \dim \mathfrak{g}^{\rho(\Gamma)}.$$

(ii) The component $X_0(\Gamma, G)$ of $X(\Gamma, G)$ that contains the character of ρ has dimension:

$$\dim X_0(\Gamma, G) = -\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) + 2 \dim \mathfrak{g}^{\rho(\Gamma)} = \dim H^1(\mathcal{O}^2, \text{Ad}\rho).$$

Proof. (i) First consider the case $\mathcal{O}^2 = T^2$, here we adapt an argument of [16]. In this case $\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) = 0$, and by a theorem of Richardson [29, Theorem B], ρ is in the closure of $\mathbb{T} \times \mathbb{T}$

for $\mathbb{T} \subset G$ a maximal torus, $\mathbb{T} \cong (\mathbb{C}^*)^{\text{rank}(G)}$ (meaning torus of an algebraic group). Therefore ρ is contained in a component of dimension at least $\dim G + \text{rank } G$. As $\dim \mathfrak{g}^{\rho(\Gamma)} = \text{rank } G$, by Proposition 4.1, $\dim T_\rho^{\text{Zar}} R(\Gamma, G) = \dim G + \text{rank } G$. Therefore

$$\dim_\rho R(\Gamma, G) \geq \dim G + \text{rank } G = \dim T_\rho^{\text{Zar}} R(\Gamma, G).$$

Using that the dimension of the Zariski tangent space is always larger than or equal to the dimension of a variety, with equality only at smooth points, it follows that $\dim_\rho R(\Gamma, G) = \dim T_\rho^{\text{Zar}} R(\Gamma, G)$ and ρ is a smooth point.

For general \mathcal{O}^2 , notice that $-\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) + \dim G + \dim \mathfrak{g}^{\rho(\Gamma)} = \dim T_\rho^{\text{Zar}} R(\Gamma, G)$, by Proposition 4.1 and all we need to show is that ρ is a smooth point. For that purpose, we use that the variety of representations of Γ_0 at $\rho|_{\Gamma_0}$ is smooth. As Goldman obstructions are natural [18, §3], they are also equivariant. In particular if we start with a equivariant infinitesimal deformation of Γ_0 , then the sequence of obstructions to integrability is also equivariant. Thus, by Proposition 3.3 and Proposition 3.4, the sequence of obstructions to integrability of Γ vanishes. This proves (i).

For (ii), notice that $\dim \mathfrak{g}^{\rho(\Gamma)} = \dim H^0(\Gamma, \text{Ad}\rho)$ is upper semicontinuous on $R(\Gamma, G)$ [14, Ch. III, Theorem 12.8]. By smoothness, the dimension of the Zariski tangent space reaches its minimum (along the irreducible component) at ρ . Hence, as

$$\dim G + \dim \mathfrak{g}^{\rho(\Gamma)} - \tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) = \dim T_\rho^{\text{Zar}} R(\Gamma, G)$$

by Proposition 4.1, $\dim \mathfrak{g}^{\rho(\Gamma)}$ reaches its minimum along the irreducible component at ρ . (Here we use that $\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho)$ is constant along components, because the elements of finite order cannot be deformed.) Equivalently the dimension of the orbit by conjugation

$$\dim G - \dim \mathfrak{g}^{\rho(\Gamma)}$$

reaches its maximum at ρ . It follows (for instance from [6, Section 6.3]) that

$$\dim X_0(\Gamma, G) = \dim R_0(\Gamma, G) - (\dim G - \dim \mathfrak{g}^{\rho(\Gamma)}),$$

which proves (ii). □

Remark 4.12. *It does not follow from Theorem 4.11 that the variety of characters $X(\Gamma, G)$ is smooth at the character of ρ .*

To explain Remark 4.12, we point out that even if ρ is strongly regular, its orbit may be not closed, so ρ could be smooth but its character not. For instance, the parabolic representation of \mathbb{Z}^2 in $\text{SL}(2, \mathbb{C})$ is a smooth point of $R(\mathbb{Z}^2, \text{SL}(2, \mathbb{C}))$, but its character is a singular point of $X(\mathbb{Z}^2, \text{SL}(2, \mathbb{C}))$. The orbit of the parabolic representation is not closed because it accumulates to the trivial representation (that it is not regular).

Orientable Euclidean 2-orbifolds appear as peripheral subgroups of hyperbolic three-orbifolds of finite volume. Thus a natural representation of $\Gamma = \pi_1(\mathcal{O}^2)$ in $\text{PSL}(2, \mathbb{C})$ occurs as the holonomy of horospherical cusps, homeomorphic to $\mathcal{O}^2 \times [0, +\infty)$. We are interested in the composition of this holonomy with the principal representation $\tau: \text{PSL}(2, \mathbb{C}) \rightarrow G$, in view of further computations for hyperbolic three-orbifolds.

Proposition 4.13. *Let \mathcal{O}^2 be a closed orientable Euclidean 2-orbifold. Consider the holonomy $\text{hol}: \pi_1(\mathcal{O}^2) \rightarrow \text{PSL}(2, \mathbb{C})$ as the horospherical section of a cusp, composed with the principal representation $\tau: \text{PSL}(2, \mathbb{C}) \rightarrow G$. Then $\dim_{[\tau \circ \text{hol}]} X(\mathcal{O}^2, G)$ is given by Table 4. Furthermore, the dimension of $\mathfrak{g}^{\tau \circ \text{hol}(\Gamma)}$ is given by Table 5.*

$G \backslash \mathcal{O}^2$	T^2	$S^2(2, 2, 2, 2)$	$S^2(3, 3, 3)$	$S^2(2, 4, 4)$	$S^2(2, 3, 6)$
$\mathrm{PSL}(n, \mathbb{C})$	$2(n-1)$	$2\lfloor \frac{n}{2} \rfloor$	$2\lfloor \frac{n}{3} \rfloor$	$2\lfloor \frac{n}{4} \rfloor$	$2\lfloor \frac{n}{6} \rfloor$
$\mathrm{PSp}(2m, \mathbb{C})$	$2m$	$2m$	$2\lfloor \frac{m}{3} \rfloor$	$2\lfloor \frac{m}{2} \rfloor$	$2\lfloor \frac{m}{3} \rfloor$
$\mathrm{PO}(2m+1, \mathbb{C})$	$2m$	$2m$	$2\lfloor \frac{m}{3} \rfloor$	$2\lfloor \frac{m}{2} \rfloor$	$2\lfloor \frac{m}{3} \rfloor$
$\mathrm{PO}(2m, \mathbb{C})$	$2m+2$	$4\lfloor \frac{m}{2} \rfloor$	$2\lfloor \frac{m}{3} \rfloor$	$2(\lfloor \frac{m}{4} \rfloor + \lfloor \frac{m+1}{4} \rfloor)$	$2(\lfloor \frac{m}{6} \rfloor + \lfloor \frac{m+2}{6} \rfloor)$
G_2	4	4	2	0	2
F_4	8	8	4	4	4
E_6	12	8	4	4	6
E_7	14	14	6	4	6
E_8	16	16	8	8	8

Table 4: Dimension of $X(\mathcal{O}^2, G)$ at $[\tau \circ \mathrm{hol}]$, for $\mathrm{hol}: \Gamma \rightarrow \mathrm{PGL}(2, \mathbb{C})$ the holonomy of a horospherical cusp and $\tau: \mathrm{PGL}(2, \mathbb{C}) \rightarrow G$ the principal representation.

$\mathfrak{g} \backslash \mathcal{O}^2$	T^2	$S^2(2, 2, 2, 2)$	$S^2(3, 3, 3)$	$S^2(2, 4, 4)$	$S^2(2, 3, 6)$
$\mathfrak{sl}(n, \mathbb{C})$	$n-1$	$\lfloor \frac{n-1}{2} \rfloor$	$\lfloor \frac{n-1}{3} \rfloor$	$\lfloor \frac{n-1}{4} \rfloor$	$\lfloor \frac{n-1}{6} \rfloor$
$\mathfrak{sp}(2m, \mathbb{C})$	m	0	$\lfloor \frac{m+1}{3} \rfloor$	0	0
$\mathfrak{so}(2m+1, \mathbb{C})$	m	0	$\lfloor \frac{m+1}{3} \rfloor$	0	0
$\mathfrak{so}(2m, \mathbb{C})$	m	$\delta_{2\mathbb{Z}}^{m-1}$	$\lfloor \frac{m}{3} \rfloor + \delta_{3\mathbb{Z}}^{m-1}$	$\delta_{4\mathbb{Z}}^{m-1}$	$\delta_{6\mathbb{Z}}^{m-1}$
\mathfrak{g}_2	2	0	0	0	0
\mathfrak{f}_4	4	0	0	0	0
\mathfrak{e}_6	6	2	0	2	0
\mathfrak{e}_7	7	0	1	0	0
\mathfrak{e}_8	8	0	0	0	0

Table 5: Dimension of $\mathfrak{g}^{\tau \circ \mathrm{hol}(\Gamma)}$ for $\tau \circ \mathrm{hol}$ as in Table 4, where $\delta_{j\mathbb{Z}}^i = 1$ if $i \in j\mathbb{Z}$ and $\delta_{j\mathbb{Z}}^i = 0$ otherwise.

Proof. The principal representation τ maps regular elements to regular elements. Furthermore the parabolic holonomy of the maximal torsion-free subgroup of $\pi_1(\mathcal{O}^2)$ is contained in a connected abelian subgroup of $\mathrm{PGL}(2, \mathbb{C})$, hence $\tau \circ \mathrm{hol}|_{\pi_1(\mathcal{O}^2)}$ is strongly regular and Theorem 4.11 applies. The computation of dimension is based on the decomposition of $\mathrm{Ad} \circ \tau$ as a sum of Sym^{2d_α} , according to the exponents d_1, \dots, d_r of G , where r is the rank of G . Let V_{2d_α} is the space of the representation Sym^{2d_α} . We use the following three facts:

1. $\dim V_{2d_\alpha} = 2d_\alpha + 1$.
2. For a cyclic group of rotations $C_k < \mathrm{PSL}(2, \mathbb{C})$ of order k , with hyperbolic rotation angles in $\frac{2\pi}{k}\mathbb{Z}$, $\dim V_{2d_\alpha}^{C_k} = 2\lfloor \frac{d_\alpha}{k} \rfloor + 1$.
3. $\dim V_{2d_\alpha}^\Gamma = \delta_{k\mathbb{Z}}^i$, $\Gamma = \pi_1(\mathcal{O}^2)$, $\Gamma_0 \cong \mathbb{Z}^2$ a maximal torsion-free subgroup and $k = |\Gamma/\Gamma_0|$, where $\delta_{k\mathbb{Z}}^i = 1$ if $i \in k\mathbb{Z}$ and 0 if $i \notin k\mathbb{Z}$.

From this, the computation is elementary. □

Lemma 4.14. *Let $\mathcal{O}^2 = S^2(2, 2, 2, 2)$, $\mathcal{O}^2 = S^2(3, 3, 3)$, $\mathcal{O}^2 = S^2(2, 4, 4)$, or $\mathcal{O}^2 = S^2(2, 3, 6)$. There is an irreducible representation of $\Gamma = \pi_1(\mathcal{O}^2)$ in $\mathrm{PSL}(k, \mathbb{C})$ iff and only if $k = |\Gamma/\Gamma_0|$.*

Proof. We first construct the irreducible representation of Γ in $\mathrm{PSL}(k, \mathbb{C})$, for $k = |\Gamma/\Gamma_0|$. From the exact sequence (24), start with a diagonal representation of Γ_0 , then represent the cyclic group Γ/Γ_0 as a cyclic permutation the k coordinates of \mathbb{C}^k . (In each case, Proposition 4.2 applies and the dimension of the variety of characters is 2.)

Let $\rho: \Gamma \rightarrow \mathrm{PSL}(k, \mathbb{C})$ be an irreducible representation. Let $\mathrm{Fix}(\rho(\Gamma_0)) \subset \mathbb{CP}^{k-1}$ denote the fixed point set of $\rho(\Gamma_0)$, that is nonempty by Kolchin's theorem. We consider the action of the cyclic group Γ/Γ_0 on $\mathrm{Fix}(\rho(\Gamma_0))$. It can be checked that any possibility other than $k = |\Gamma/\Gamma_0|$ and that $\mathrm{Fix}(\rho(\Gamma_0))$ has k points in generic position that are permuted cyclically by Γ/Γ_0 yields a contradiction with irreducibility. \square

Example 4.15. We consider planar Euclidean Coxeter groups, namely generated by reflections along a square or a Euclidean triangle with angles an integer divisor of π , denoted by:

$$Q(2, 2, 2, 2), \quad T(3, 3, 3), \quad T(2, 4, 4), \quad T(2, 3, 6).$$

The respective orientation coverings are $S^2(2, 2, 2, 2)$, $S^2(3, 3, 3)$, $S^2(2, 4, 4)$ and $S^2(2, 3, 6)$. The group $\Gamma = \pi_1(\mathcal{O})$ acts naturally in the Euclidean plane, and one can construct representations in G by realizing this action on a flat of the symmetric space associated to G . On this flat, one allows translations, but the rotational part is restricted to the action of the Weyl group, which in its turn is a Coxeter group that acts faithfully on a maximal flat (eg the Cartan subalgebra). Hence one can construct representations by analyzing the Weyl group.

We discuss the case of rank 2. Here the Weyl group is a dihedral group of order $2m$ generated by reflections along two lines at angle π/m , the walls of the Weyl chamber. For type A_2 the angle is $\pi/3$, for B_2 , $\pi/4$, and for G_2 , $\pi/6$. This yields discrete faithful representations of the triangle group $T(3, 3, 3)$ in $\mathrm{SL}(3, \mathbb{C})$ and $G_2(\mathbb{C})$, of $T(2, 4, 4)$ in $\mathrm{Sp}(4, \mathbb{C})$, of $T(2, 3, 6)$ in $G_2(\mathbb{C})$ and of the quadrilateral group $Q(2, 2, 2, 2)$ in $\mathrm{Sp}(4, \mathbb{C})$ or $G_2(\mathbb{C})$ (or the corresponding split real forms).

The triangle groups constructed here have a one-parameter space of deformations (homoteties in the plane) and the quadrilateral group a 2-parameter space. Those are half the dimension of the deformation space of their orientable cover.

4.4 Spherical 2-orbifolds

Finally, we also need to consider spherical two-orbifolds, namely finitely covered by the 2-sphere. They are rigid, but we require a cohomological computation later in the paper.

Lemma 4.16. Let \mathcal{O}^2 be a spherical 2-orbifold and $\rho: \pi_1(\mathcal{O}^2) \rightarrow G$ a representation. Then

$$\dim H^i(\mathcal{O}^2, \mathrm{Ad}\rho) = \begin{cases} \dim \mathfrak{g}^{\rho(\pi_1(\mathcal{O}^2))}, & \text{for } i = 0, 2 \\ 0, & \text{for } i = 1 \end{cases},$$

and thus

$$\tilde{\chi}(\mathcal{O}^2, \mathrm{Ad}\rho) = 2 \dim \mathfrak{g}^{\rho(\pi_1(\mathcal{O}^2))}. \quad (25)$$

Proof. Let \mathcal{O}^2 be a spherical 2-orbifold. As it is finitely covered by S^2 , by Proposition 3.3 we have

$$H^i(\mathcal{O}^2, \mathrm{Ad}\rho) \cong H^i(S^2, \mathfrak{g})^{\pi_1(\mathcal{O}^2)} \cong H^i(S^2, \mathbb{Z}) \otimes \mathfrak{g}^{\rho(\pi_1(\mathcal{O}^2))}.$$

Hence the lemma follows. \square

Remark 4.17. For a spherical orbifold, the component of $R(\mathcal{O}^2, G)$ that contains ρ is an orbit by conjugation of dimension

$$\dim G - \dim \mathfrak{g}^{\rho(\pi_1(\mathcal{O}^2))}.$$

In particular, by (25) it is a smooth variety of dimension

$$-\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) + \dim G + \dim \mathfrak{g}^{\rho(\pi_1(\mathcal{O}^2))}.$$

5 Computing dimensions of the Hitchin component

Assume that \mathcal{O}^2 is hyperbolic, namely that the standard orbifold-Euler characteristic is negative, $\chi(\mathcal{O}^2) < 0$. For G a *simple* complex adjoint group, the principal representation $\tau: \text{PSL}(2, \mathbb{C}) \rightarrow G$ is constructed from Jacobson-Mozorov theorem, and it restricts to $\tau: \text{PGL}(2, \mathbb{R}) \rightarrow G_{\mathbb{R}}$, for $G_{\mathbb{R}}$ a (non connected) split real form of G . The Hitchin component

$$\text{Hit}(\mathcal{O}^2, G_{\mathbb{R}})$$

is the connected component of $\mathcal{R}^{\text{good}}(\mathcal{O}^2, G_{\mathbb{R}})$ that contains the composition of τ with the holonomy representation of any Fuchsian structure on \mathcal{O}^2 . Hitchin components for surface have been intensively studied. here just mention that Alessandrini, Lee, and Schaffhauser [1] have introduced them for (perhaps non-orientable) 2-orbifolds; furthermore they have shown that they are homeomorphic to balls, as for surfaces.

The purpose of this section is to provide formulas for the dimension of the Hitchin component of orbifolds, using the tools of Section 4. Many of the formulas are equivalent to the ones already computed in [1], but the approach and presentation of results is different. In particular we give some applications in Proposition 5.10 and Corollary 5.15 to the dimension growth of some families of Hitchin representations. Long and Thistlethwaite in [23] have also computed the dimension of Hitchin components in $\text{PGL}(n, \mathbb{R})$ for turnovers (2-spheres with three cone points).

Remark 5.1. The representation $\tau \circ \text{hol}$ is good, Remark 2.4, so the real dimension of the Hitchin component of \mathcal{O}^2 in $G_{\mathbb{R}}$ is precisely the complex dimension of $X(\mathcal{O}^2, G)$ at the character of $\tau \circ \text{hol}$, namely $-\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho)$; we often use:

$$\dim_{\mathbb{R}} \text{Hit}(\mathcal{O}^2, G_{\mathbb{R}}) = \dim_{\mathbb{C}, [\tau \circ \text{hol}]} X(\mathcal{O}^2, G) = -\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho).$$

Remark 5.2. By Corollary 3.5, for \mathcal{O}^2 orientable $\dim_{\mathbb{R}} \text{Hit}(\mathcal{O}^2, G_{\mathbb{R}})$ is even.

Remark 5.3. For \mathcal{O}^2 closed and orientable, Atiyah-Bott-Goldmann defines a symplectic structure on $\text{Hit}(\mathcal{O}^2, G_{\mathbb{R}})$, Proposition 4.4.

The following is Proposition 1.5 from the introduction:

Proposition 5.4. For \mathcal{O}^2 closed and non-orientable, with orientation covering $\widetilde{\mathcal{O}^2}$, via the restriction map $\text{Hit}(\mathcal{O}^2, G_{\mathbb{R}})$ embeds in $\text{Hit}(\widetilde{\mathcal{O}^2}, G_{\mathbb{R}})$ as a Lagrangian submanifold.

Proof. By Proposition 4.5 the restriction map yields a Lagrangian *immersion* from $\text{Hit}(\mathcal{O}^2, G_{\mathbb{R}})$ to $\text{Hit}(\widetilde{\mathcal{O}^2}, G_{\mathbb{R}})$. Furthermore, using the irreducibility of Hitchin representations, one can prove that this restriction map is injective (see for instance Lemma 2.9 and Proposition 2.10 in ??). The restriction map to a finite index subgroup is proper, therefore it is an embedding (or one can quote [1, Corollary 2.13]) \square

5.1 Hitchin component for $\mathrm{PGL}(n, \mathbb{R})$

Following Long and Thistlethwaite [23], for n, k positive integers we set

$$\sigma(n, k) = qn + (q + 1)r,$$

where q and r are nonnegative integers such that $n = qk + r$ (the quotient and remainder of integer division).

Proposition 5.5. *Let $x \in \mathcal{O}^2$ and $\tau = \mathrm{Sym}^{n-1}$. For $\rho = \tau \circ \mathrm{hol}: \pi_1(\mathcal{O}^2) \rightarrow \mathrm{PGL}(n, \mathbb{R})$,*

$$\dim \mathfrak{g}^{\rho(\mathrm{Stab}(x))} = \begin{cases} \sigma(n, k) - 1 & \text{if } x \text{ is a cone point with } \mathrm{Stab}(x) \cong C_k, \\ (\sigma(n, k) - 1)/2 & \text{if } x \text{ is a corner with } \mathrm{Stab}(x) \cong D_{2k} \text{ and } n \text{ is odd,} \\ (\sigma(n, k) - 2)/2 & \text{if } x \text{ is a corner with } \mathrm{Stab}(x) \cong D_{2k} \text{ and } n \text{ is even,} \\ (n^2 - 1)/2 & \text{if } x \text{ is a mirror point and } n \text{ is odd,} \\ (n^2 - 2)/2 & \text{if } x \text{ is a mirror point and } n \text{ is even.} \end{cases}$$

Proof. For an elliptic element $\gamma \in \Gamma$ of order k , we may compute $\dim \mathfrak{g}^{\rho(\gamma)}$ using that, up to conjugacy:

$$\rho(\gamma) = \pm \mathrm{Sym}^{n-1} \begin{pmatrix} e^{\frac{\pi i}{k}} & 0 \\ 0 & e^{-\frac{\pi i}{k}} \end{pmatrix} = \pm \mathrm{diag}(e^{\frac{\pi i}{k}(n-1)}, e^{\frac{\pi i}{k}(n-3)}, \dots, e^{\frac{\pi i}{k}(n-1)}).$$

As $n = qk + r$, among the k eigenvalues of this matrix, $(k - r)$ eigenvalues have multiplicity q and r eigenvalues have multiplicity $q + 1$. Thus

$$\dim \mathfrak{g}^{\rho(\gamma)} = (k - r)q^2 + r(q + 1)^2 - 1 = qn + (q + 1)r - 1 = \sigma(n, k) - 1.$$

This computation yields the formula for cone points and mirror points. For corner points, apply Corollary 4.9 and the previous computations. \square

We then have:

Proposition 5.6. *Let \mathcal{O}^2 be a connected hyperbolic 2-orbifold, with underlying surface $|\mathcal{O}^2|$. Then*

$$\dim \mathrm{Hit}(\mathcal{O}^2, \mathrm{PGL}(n, \mathbb{R})) = -(n^2 - 1)\chi(|\mathcal{O}^2|) + \sum_{i=1}^{\mathrm{cp}} (n^2 - \sigma(n, k_i)) + \sum_{j=1}^{\mathrm{cr}} \frac{n^2 - \sigma(n, l_j)}{2} + \mathbf{b} \lfloor \frac{n^2}{2} \rfloor,$$

where the cone points have order $k_1, k_2, \dots, k_{\mathrm{cp}}$, the corner reflectors have order $2l_1, 2l_2, \dots, 2l_{\mathrm{cr}}$ and \mathbf{b} is the number of components of $\partial\mathcal{O}^2$ homeomorphic to \mathbb{R}/D_∞ , an interval with mirror boundary.

Proof. We apply Proposition 4.2 to get that the dimension of the Hitchin component is $-\chi(\mathcal{O}^2, \mathrm{Ad}\rho)$, which, by Proposition 5.5, equals:

$$-(n^2 - 1)\chi(\mathcal{O}^2 \setminus \Sigma) - \sum_{i=1}^{\mathrm{cp}} (\sigma(n, k_i) - 1) + \begin{cases} -\sum_{j=1}^{\mathrm{cr}} \frac{\sigma(n, l_j) - 2}{2} + (\mathbf{me} - \mathbf{mp}) \frac{n^2 - 2}{2} & \text{if } n \text{ even} \\ -\sum_{j=1}^{\mathrm{cr}} \frac{\sigma(n, l_j) - 1}{2} + (\mathbf{me} - \mathbf{mp}) \frac{n^2 - 1}{2} & \text{if } n \text{ odd} \end{cases} \quad (26)$$

where \mathbf{me} is the number of mirror edges (joining corner reflectors or mirror points in the boundary), and \mathbf{mp} the number of boundary points that are also mirrors. Next we use the formulas (from additivity of the Euler characteristic):

$$\begin{aligned}\chi(|\mathcal{O}^2|) &= \chi(\mathcal{O}^2 \setminus \Sigma) + \mathbf{cp} + \mathbf{cr} + \mathbf{mp} - \mathbf{me}, \\ 0 &= \chi(\partial|\mathcal{O}^2|) = -\mathbf{me} - \mathbf{b} + \mathbf{cr} + \mathbf{mp},\end{aligned}$$

that are equivalent to

$$-\chi(\mathcal{O}^2 \setminus \Sigma) = -\chi(|\mathcal{O}^2|) + \mathbf{cp} + \mathbf{b}, \quad (27)$$

$$\mathbf{me} - \mathbf{mp} = \mathbf{cr} - \mathbf{b}. \quad (28)$$

Using (27) and (28), (26) becomes:

$$-(n^2 - 1)\chi(|\mathcal{O}^2|) + \sum_{i=1}^{\mathbf{cp}} (n^2 - \sigma(n, k_i)) + \sum_{j=1}^{\mathbf{cr}} \frac{n^2 - \sigma(n, l_j)}{2} + \begin{cases} \mathbf{b} \frac{n^2}{2} & \text{if } n \text{ even} \\ \mathbf{b} \frac{n^2-1}{2} & \text{if } n \text{ odd} \end{cases}$$

which proves the proposition. \square

Example 5.7 (Proved in [23] when $\mathbf{cp} = 3$). *Let $S^2(k_1, \dots, k_{\mathbf{cp}})$ be a sphere with $\mathbf{cp} \geq 3$ cone points of order $k_1, \dots, k_{\mathbf{cp}}$ respectively (and satisfying $\frac{1}{k_1} + \frac{1}{k_2} + \frac{1}{k_3} < 1$ for $\mathbf{cp} = 3$). Then, the dimension of the Hitchin component of $S^2(k_1, \dots, k_{\mathbf{cp}})$ in $\mathrm{PGL}(n, \mathbb{R})$ is*

$$n^2(\mathbf{cp} - 2) + 2 - \sum_{i=1}^{\mathbf{cp}} \sigma(n, k_i).$$

Consider now $P(k_1, \dots, k_{\mathbf{cr}})$ the orbifold generated by reflections on a hyperbolic polygon with angles $\frac{\pi}{k_1}, \dots, \frac{\pi}{k_{\mathbf{cr}}}$. In particular $S^2(k_1, \dots, k_{\mathbf{cr}})$ is its orientation covering and therefore the dimension of the Hitchin component of $P(k_1, \dots, k_{\mathbf{cr}})$ in $\mathrm{PGL}(n, \mathbb{R})$ is

$$\frac{n^2(\mathbf{cr} - 2) + 2 - \sum_{i=1}^{\mathbf{cr}} \sigma(n, k_i)}{2}.$$

Proposition 5.8. *Every component of $\mathcal{R}^{\mathrm{good}}(\mathcal{O}^2, \mathrm{PGL}(n, \mathbb{R}))$ that contains representations that are boundary regular has dimension at most the dimension of the Hitchin component.*

Proof. Assume first that $\partial\mathcal{O}^2 = \emptyset$. By Corollary 4.3, we may assume that \mathcal{O}^2 is orientable, hence the singular locus is a finite union of cone points, that have cyclic stabilizers. We claim that for an elliptic element γ , $\dim(\mathfrak{g}^{\rho(\gamma)})$ is minimized for ρ in the Hitchin component. Assuming the claim, $\tilde{\chi}(\mathcal{O}^2, \rho)$ is minimized for ρ in the Hitchin component, and apply Proposition 4.2. To prove the claim, since $\rho(\gamma)$ has order k , its eigenvalues have multiplicities $\{n_1, \dots, n_k\}$, with $n_1 + \dots + n_k = n$. Therefore $\dim(\mathfrak{g}^{\rho(\gamma)}) = n_1^2 + \dots + n_k^2 - 1$. This function is minimized when $|n_i - n_j| \leq 1$, because if we replace n_1 by $n_1 + 1$ and n_2 by $n_2 - 1$, then $n_1^2 + n_2^2$ increases by $2(n_1 - n_2 + 1)$. As, for a rotation of angle $2\pi/k$, $\mathrm{Sym}^{n-1}(\rho(\gamma))$ satisfies $|n_i - n_j| \leq 1$, it minimizes the required dimension.

When $\partial\mathcal{O}^2 \neq \emptyset$, in the orientable case we apply Proposition 4.6, knowing that by orientability $\partial\mathcal{O}^2$ is a union of circles and $\tilde{\chi}(\partial\mathcal{O}^2, \mathrm{Ad}\rho) = 0$. In the non-orientable case we apply Proposition 4.7 and use the claim we have proved, that for an elliptic element γ , $\dim(\mathfrak{g}^{\rho(\gamma)})$ is minimized for ρ in the Hitchin component. \square

Example 5.9. *In Table 6 we compute the dimension of $\mathrm{Hit}(S^2(3, 3, 4)\mathrm{PSL}(n, \mathbb{R}))$. Notice that the difference with $n^2/12$ is bounded, this is a particular case of the next proposition.*

$n \bmod 12$	$\dim \text{Hit}(S^2(3, 3, 4), \text{PGL}(n, \mathbb{R}))$
0	$n^2/12 + 2$
$\pm 1, \pm 5$	$(n^2 - 1)/12$
± 2	$(n^2 - 4)/12$
± 3	$(n^2 + 15)/12$
± 4	$(n^2 + 8)/12$
6	$n^2/12 + 1$

Table 6: Dimension of the Hitchin component of $S^2(3, 3, 4)$. The dimension for the group generated by reflections on a triangle of angles $(\frac{\pi}{3}, \frac{\pi}{3}, \frac{\pi}{4})$ is half of it.

Proposition 5.10. *There exists a uniform constant $C(\mathcal{O}^2)$ depending only on \mathcal{O}^2 such that*

$$|\dim \text{Hit}(\mathcal{O}^2, \text{PGL}(n, \mathbb{R})) + \chi(\mathcal{O}^2)(n^2 - 1)| \leq C(\mathcal{O}^2).$$

Here $0 > \chi(\mathcal{O}^2) \in \mathbb{Q}$ denotes the (untwisted) orbifold-Euler characteristic.

Proof. Let $\rho = \tau \circ \text{hol}$. Assume first that \mathcal{O}^2 is orientable. If γ is an elliptic element of order k , we shall prove that $\left| \frac{n^2 - 1}{k} - \dim \mathfrak{g}^{\rho(\gamma)} \right|$ is bounded uniformly on n . Namely, if $n = kq + r$, then $\dim \mathfrak{g}^{\rho(\gamma)} = kq^2 + 2rq + r - 1$. On the other hand $n^2 = k^2q^2 + 2rkq + r^2$, thus:

$$\left| \frac{n^2 - 1}{k} - \dim \mathfrak{g}^{\rho(\gamma)} \right| = \left| \frac{r^2 - 1}{k} - r + 1 \right| = (r - 1) \left(1 - \frac{r + 1}{k} \right) \leq k.$$

In the orientable case, as all stabilizers are cyclic this estimate tells that the difference $\tilde{\chi}(\mathcal{O}^2, \text{Ad}\rho) - \chi(\mathcal{O}^2)(n^2 - 1)$ is uniformly bounded, for $\rho = \text{Sym}^{n-1} \circ \text{hol}$. With Proposition 4.2, this yields the proposition for \mathcal{O}^2 orientable.

In the non-orientable case, consider $\widetilde{\mathcal{O}^2} \rightarrow \mathcal{O}^2$ the orientation cover. By Proposition 4.7

$$\left| \dim \text{Hit}(\mathcal{O}^2, \text{PGL}(n, \mathbb{R})) - \frac{1}{2} \dim \text{Hit}(\widetilde{\mathcal{O}^2}, \text{PGL}(n, \mathbb{R})) \right| \leq \frac{1}{2} (\dim G - 2 \dim \mathfrak{g}^{\rho(C_2)}) \mathbf{b}(\mathcal{O}^2)$$

where $\rho(C_2)$ denotes the image of any cyclic stabilizer of order 2 and $\mathbf{b}(\mathcal{O}^2)$ is the number of components of $\partial \mathcal{O}^2$ homeomorphic to $\mathbb{R}/D_\infty = \llbracket 0, 1 \rrbracket$ (an interval with mirror end-points). As

$$\frac{1}{2} (\dim G - 2 \dim \mathfrak{g}^{\rho(C_2)}) = \frac{n^2 - 1}{2} - (\sigma(n, 2) - 1) = \begin{cases} 1/2 & \text{for even } n \\ 0 & \text{for odd } n \end{cases}$$

the proposition follows. \square

5.2 Exponents of a simple Lie algebra

Recall from Proposition 5.5 that the exponents $d_1, \dots, d_r \in \mathbb{N}$ of the Lie algebra \mathfrak{g} are defined by the equation

$$\text{Ad} \circ \tau = \bigoplus_{\alpha=1}^r \text{Sym}^{2d_\alpha},$$

Equation (2), where τ is the principal representation. Here $r = \text{rank } \mathfrak{g}$ and $\sum_{\alpha=1}^r (2d_\alpha + 1) = \dim \mathfrak{g}$.

Lemma 5.11. For $\rho \in \text{Hit}(\mathcal{O}^2, G)$ and $x \in \Sigma_{\mathcal{O}^2}$ a point in the branching locus with nontrivial stabilizer Γ_x :

$$\dim \mathfrak{g}^{\rho(\Gamma_x)} = \begin{cases} \sum_{\alpha=1}^r (2\lfloor \frac{d_\alpha}{k} \rfloor + 1) & \text{if } \Gamma_x \cong C_k \\ (\sum_{\alpha=1}^r \lfloor \frac{d_\alpha}{k} \rfloor) + \#\{d_\alpha \mid d_\alpha \text{ is even}\} & \text{if } \Gamma_x \cong D_{2k} \end{cases}$$

where C_k is cyclic of order k , and D_{2k} dihedral of order $2k$, $r = \text{rank}(\mathfrak{g})$, and $\{d_1, \dots, d_r\}$ are the exponents of the Lie algebra.

Proof. For a Fuchsian representation, the image of the cyclic group C_k is generated by a matrix conjugate to

$$\pm \begin{pmatrix} e^{\frac{\pi i}{k}} & 0 \\ 0 & e^{-\frac{\pi i}{k}} \end{pmatrix}. \quad (29)$$

Using (2), the contribution of the α -th exponent to $\dim \mathfrak{g}^{\rho(\Gamma_x)}$ is the multiplicity of eigenvalue 1 in the $2d_\alpha$ -th symmetric power of (29). Namely, the number of appearances of 1 in $\{e^{\frac{2\pi i}{k}d_\alpha}, e^{\frac{2\pi i}{k}(d_\alpha-1)}, \dots, e^{\frac{2\pi i}{k}(-d_\alpha)}\}$ which is precisely $2\lfloor \frac{d_\alpha}{k} \rfloor + 1$.

For the dihedral group, Corollary 4.9, the cyclic case, and (3) yield that the dimension is

$$\dim \mathfrak{g}^{\rho(D_k)} = (2 \dim \mathfrak{g}^{\rho(C_2)} + \dim \mathfrak{g}^{\rho(C_k)} - \dim \mathfrak{g})/2 = \sum_{\alpha=1}^r (2\lfloor \frac{d_\alpha}{2} \rfloor + 1 + \lfloor \frac{d_\alpha}{k} \rfloor - d_\alpha)$$

$$\text{and } 2\lfloor \frac{d_\alpha}{2} \rfloor + 1 - d_\alpha = \begin{cases} 1 & \text{for } d_\alpha \text{ even,} \\ 0 & \text{for } d_\alpha \text{ odd.} \end{cases} \quad \square$$

Lemma 5.11 allows to give a new proof of Proposition 5.5.

As the exponents for $\mathfrak{sp}(2m)$ and $\mathfrak{so}(m, m+1)$ are the same, $(1, 3, 5, \dots, 2m-1)$, by Lemma 5.11 we have:

Remark 5.12. Let $\rho_1 \in \text{Hit}(\mathcal{O}^2, \text{PSp}^\pm(2m))$ and $\rho_2 \in \text{Hit}(\mathcal{O}^2, \text{PO}(m+1, m))$. For any $x \in \mathcal{O}^2$,

$$\dim \mathfrak{g}^{\rho_1(\Gamma_x)} = \dim \mathfrak{g}^{\rho_2(\Gamma_x)}.$$

Another consequence of Lemma 5.11 is the following.

Corollary 5.13 ([1]). Let \mathcal{O}^2 be a connected hyperbolic 2-orbifold. If $\{d_1, \dots, d_{\text{rank} G}\}$ are the exponents of G . Then the dimension of $\text{Hit}(\mathcal{O}^2, G)$ is

$$-\chi(|\mathcal{O}^2|) \dim G + \sum_{\alpha=1}^{\text{rank} G} \left(\sum_{i=1}^{\text{cp}} 2(d_\alpha - \lfloor \frac{d_\alpha}{k_i} \rfloor) + \sum_{j=1}^{\text{cr}} (d_\alpha - \lfloor \frac{d_\alpha}{l_j} \rfloor) + 2\mathbf{b} \lfloor \frac{d_\alpha + 1}{2} \rfloor \right)$$

where the cone points have order $k_1, k_2, \dots, k_{\text{cp}}$, the corner reflectors have order $2l_1, 2l_2, \dots, 2l_{\text{cr}}$, and \mathbf{b} is the number of components of $\partial \mathcal{O}^2$ homeomorphic to \mathbb{R}/D_∞ , an interval with mirror boundary.

Proof. We follow the same scheme as the proof of Proposition 5.6. Set $\delta_{j\mathbb{Z}}^i = 1$ if $i \in j\mathbb{Z}$ and $\delta_{j\mathbb{Z}}^i = 0$ otherwise. Firstly we apply Proposition 4.2 and Lemma 5.11 to get that the dimension of the Hitchin component is $-\chi(\mathcal{O}^2, \text{Ad}\rho)$, which equals:

$$-\chi(\mathcal{O} \setminus \Sigma) \dim G + \sum_{\alpha=1}^{\text{rank} G} \left(-\sum_{i=1}^{\text{cp}} (2\lfloor \frac{d_\alpha}{k_i} \rfloor + 1) - \sum_{j=1}^{\text{cr}} (\lfloor \frac{d_\alpha}{l_j} \rfloor + \delta_{2\mathbb{Z}}^{d_\alpha}) + (\mathbf{me} - \mathbf{mp})(2\lfloor \frac{d_\alpha}{2} \rfloor + 1) \right)$$

where \mathbf{me} is the number of mirror edges (joining corner reflectors or mirror points in the boundary), and \mathbf{mp} the number of boundary points that are also mirrors, and $\delta_{2\mathbb{Z}}^{d_\alpha} = 1$ if d_α is even, and 0 if d_α is odd. Then the corollary follows from (27) and (28). \square

This computation of dimensions is contained in [1, Theorem 1.2], where it is proved that the Hitchin component is a cell.

For exceptional groups the dimension can be directly computed from this corollary and the exponents in Table 2. For instance, for (the real split form of the adjoint group) G_2 , as the indices are 1 and 5 the dimension of the Hitchin component is

$$-14\chi(|\mathcal{O}^2|) + 8\text{cp}_2 + 10\text{cp}_{3,4,5} + 12\text{cp}_{\geq 6} + 4\text{cr}_4 + 5\text{cr}_{6,8,20} + 6\text{cr}_{\geq 12} + 8b,$$

where cp_i is the number of cone points with stabilizer C_i and cr_{2i} of corner reflectors with stabilizer D_{2i} .

It can also be checked that for G_2 and F_4 that dimension of the variety of characters is maximized by the Hitchin component, but this is a rather tedious computation.

5.3 Hitchin component for $\text{PSp}^\pm(2m)$ and $\text{PO}(m, m+1)$

For both $G = \text{PSp}^\pm(2m)$ and $G = \text{PO}(m, m+1)$, the principal representation $\tau: \text{PGL}(2, \mathbb{R}) \rightarrow G$ is the restriction of $\text{Sym}^{n-1}: \text{PGL}(2, \mathbb{R}) \rightarrow \text{PGL}(n, \mathbb{R})$, with $n = 2m$ and $n = 2m+1$, respectively. Since $\text{SL}(2, \mathbb{R})$ preserves the skew bilinear form

$$J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$

$\text{Sym}^{n-1}(\text{SL}(2, \mathbb{R}))$ preserves the symmetric power of J , which is symmetric or skew, according to the parity. Similarly one easily argues that $\text{Sym}^{n-1}(\text{SL}(2, \mathbb{R}))$ is contained in either $\text{PSp}^\pm(2m)$ or $\text{PO}(m, m+1)$, according to whether $n = 2m$ or $n = 2m+1$.

Write

$$2m = kq + r$$

with $q, m \in \mathbb{Z}$, $q, r \geq 0$ and $r < k$. Recall that:

$$\sigma(2m, k) = 2mq + r(q+1).$$

Proposition 5.14. *Let $\rho \in \text{Hit}(\mathcal{O}^2, G)$, for $G = \text{PSp}^\pm(2m)$ or $\text{PO}(m, m+1)$ and let $x \in \mathcal{O}^2$.*

1. *If x is a cone point with $\text{Stab}(x) \cong C_k$ then*

$$\dim \mathfrak{g}^{\rho(C_k)} = \begin{cases} \frac{\sigma(2m, k)}{2} & \text{for } k \text{ even,} \\ \frac{\sigma(2m, k)}{2} + \lfloor \frac{q+1}{2} \rfloor & \text{for } k \text{ odd.} \end{cases}$$

2. *If x is a mirror point with $\text{Stab}(x) \cong C_2$ then $\dim \mathfrak{g}^{\rho(C_2)} = m^2$.*

3. *If x is a corner reflector with $\text{Stab}(x) \cong D_{2k}$ then*

$$\dim \mathfrak{g}^{\rho(D_k)} = \begin{cases} \frac{\sigma(2m, k)}{4} - \frac{m}{2} & \text{for } k \text{ even,} \\ \frac{\sigma(2m, k)}{4} - \frac{m}{2} + \frac{1}{2} \lfloor \frac{q+1}{2} \rfloor & \text{for } k \text{ odd.} \end{cases}$$

Proof. By Remark 5.12, it is sufficient to make the computations for $G = \text{PSp}(2m, \mathbb{C})$. The bilinear form $\text{Sym}^{2m-1}(J)$ has matrix

$$\begin{pmatrix} 0 & \cdots & 0 & 0 & 1 \\ 0 & \cdots & 0 & -1 & 0 \\ 0 & \cdots & 1 & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ -1 & \cdots & 0 & 0 & 0 \end{pmatrix}.$$

thus

$$\mathfrak{sp}(2m) = \{(a_{i,j} \in M_{2m,2m}(\mathbb{C}) \mid a_{i,j} = (-1)^{i+j+1} a_{2m+1-j, 2m+1-i}\}.$$

From this expression, as a diagonal matrix $D \in \mathrm{PSp}(2m, \mathbb{C})$ is

$$D = \mathrm{diag}(\lambda_1, \dots, \lambda_m, \lambda_m^{-1}, \dots, \lambda_1^{-1}),$$

we have that

$$\dim(\mathfrak{sp}(2m)^D) = \frac{1}{2} \dim(\mathfrak{gl}(2m)^D) + \dim(\mathfrak{ad}^D)$$

where $\mathfrak{ad} = \{(a_{i,j} \in M_{2m,2m}(\mathbb{C}) \mid a_{ij} = 0 \text{ if } i + j \neq 2m + 1\} \subset \mathfrak{sp}(2m)$ is the anti-diagonal, namely the subspace of matrices

$$\begin{pmatrix} 0 & \cdots & 0 & 0 & * \\ 0 & \cdots & 0 & * & 0 \\ 0 & \cdots & * & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ * & \cdots & 0 & 0 & 0 \end{pmatrix}.$$

A rotation of angle $\frac{2\pi}{k}$ in $\mathrm{PGL}(2, \mathbb{C})$ is conjugate to $\pm \begin{pmatrix} e^{\frac{\pi i}{k}} & 0 \\ 0 & e^{-\frac{\pi i}{k}} \end{pmatrix}$ and

$$\begin{aligned} D &= \mathrm{Sym}^{2m-1} \left(\pm \begin{pmatrix} e^{\frac{\pi i}{k}} & 0 \\ 0 & e^{-\frac{\pi i}{k}} \end{pmatrix} \right) \\ &= \pm \mathrm{diag}(e^{\frac{\pi i}{k}(2m-1)}, e^{\frac{\pi i}{k}(2m-3)}, \dots, e^{\frac{\pi i}{k}}, e^{-\frac{\pi i}{k}}, \dots, e^{-\frac{\pi i}{k}(2m-1)}). \end{aligned}$$

Therefore, the (adjoint) action of D , on the antidiagonal \mathfrak{ad} has eigenvalues

$$\{e^{\frac{2\pi i}{k}(2m-1)}, e^{\frac{2\pi i}{k}(2m-3)}, \dots, e^{\frac{2\pi i}{k}}, e^{-\frac{2\pi i}{k}}, \dots, e^{-\frac{2\pi i}{k}(2m-1)}\}.$$

Thus, $\dim \mathfrak{ad}^D$ is the number appearances of 1 in this list of eigenvalues:

$$\dim \mathfrak{ad}^D = \begin{cases} 0 & \text{for } k \text{ even,} \\ 2 \lfloor \frac{2m-1+k}{2k} \rfloor = 2 \lfloor \frac{q+1}{2} \rfloor & \text{for } k \text{ odd.} \end{cases}$$

On the other hand, $\mathfrak{gl}(2m)^D = 2mq + r(q+1)$ (because D has r eigenvalues with multiplicity $q+1$ and $k-r$ eigenvalues with multiplicity q), what concludes item 1.

Item 2 is a particular case of item 1, and item 3 follows from Corollary 4.9 and the previous cases. \square

If we look at the growth of $\dim \mathrm{Hit}(\mathcal{O}^2, \mathrm{PSp}^\pm(2m))$, the difference

$$|\dim \mathrm{Hit}(\mathcal{O}^2, \mathrm{PSp}^\pm(2m)) + \chi(\mathcal{O}^2) \dim(\mathrm{PSp}^\pm(2m))|$$

may be unbounded because of cyclic stabilizers of odd order (and dihedral stabilizers of order a multiple of 4). But instead we have (already proved in [1]):

Corollary 5.15 ([1]). *For a compact hyperbolic two orbifold \mathcal{O}^2 :*

$$\lim_{m \rightarrow \infty} \frac{\dim \mathrm{Hit}(\mathcal{O}^2, \mathrm{PSp}^\pm(2m))}{\dim(\mathrm{PSp}^\pm(2m))} = \lim_{m \rightarrow \infty} \frac{\dim \mathrm{Hit}(\mathcal{O}^2, \mathrm{PO}(m+1, m))}{\dim(\mathrm{PO}(m+1, m))} = -\chi(\mathcal{O}^2).$$

Proof. For every stabilizer $\text{Stab}(x)$ we check from Proposition 5.14 that

$$\lim_{m \rightarrow \infty} \frac{\dim \mathfrak{g}^{\rho(\text{Stab}(x))}}{2m^2} = \frac{1}{|\text{Stab}(x)|},$$

because $2m/q \rightarrow k$. The corollary follows using Proposition 4.2. \square

Proposition 5.16. *Every component of $\mathcal{R}^{\text{good}}(\mathcal{O}^2, \text{PSp}^\pm(2m))$ that contains representations that are boundary regular has dimension at most the dimension of the Hitchin component.*

Proof. We use the same argument as in the proof of Proposition 5.8. The same discussion applies here, the difference being in the proof that the Hitchin component minimizes the dimension of the centralizer of elliptic elements of given order k . We follow the description in the proof of Proposition 5.14, in particular a diagonal element in $\text{PSp}(2m, \mathbb{C})$ is

$$D = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_m, \lambda_m^{-1}, \dots, \lambda_2^{-1}, \lambda_1^{-1}).$$

Hence $\{\pm \text{Id}\}$ is the center of $\text{Sp}(2m, \mathbb{C})$. To compute $\dim \mathfrak{Sp}(2m, \mathbb{C})^D$, we distinguish the case k even from the case k odd.

For k even we assume first that $\lambda_i^k = -1$, then there are no elements in the antidiagonal \mathfrak{ad} fixed by D . Therefore $\dim \mathfrak{Sp}(2m, \mathbb{C})^D = \frac{1}{2} \dim \mathfrak{gl}(2m, \mathbb{C})^D$. Hence if n_i is the multiplicity of λ_i plus the multiplicity of λ_i^{-1} (in this notation we chose half of the eigenvalues $\{\lambda_1, \dots, \lambda_{\frac{k}{2}}\}$ disjoint from their inverses):

$$\dim \mathfrak{Sp}(2m, \mathbb{C})^D = n_1^2 + n_2^2 + \dots + n_k^2.$$

As seen in the proof of Proposition 5.8 this is minimized when $|n_i - n_j| \leq 1$. Next assume that k is even and that $\lambda_i^k = 1$. Then ± 1 are possible eigenvalues of D , and they yield fixed elements in the antidiagonal. Let n_0 be the multiplicity of 1, n_1 the multiplicity of -1 , and let for $i = 2, \dots, \frac{k}{2}$ the addition of the multiplicities of each eigenvalue and its inverse. Then

$$\dim \mathfrak{Sp}(2m, \mathbb{C})^D = 2n_0^2 + n_0 + 2n_1^2 + n_1 + n_2^2 + \dots + n_k^2$$

An elementary computation shows that $2n_0^2 + n_0 + 2n_1^2 + n_1 \geq (n_0 + n_1)^2$ (with equality iff $n_0 = n_1 = 0$), therefore the dimension of the invariant subspace is higher than when $\lambda_i^k = -1$.

For k odd, up to changing D by $-D$ we may assume that $\lambda_i^k = 1$ (and -1 is not an eigenvalue). We count again multiplicities: we take n_0 the multiplicity of 1. We group the remaining eigenvalues in pairs, each eigenvalue with its inverse, and we add the multiplicities of each pair to get $n_1, \dots, n_{(k-1)/2}$. Then a computation yields

$$\dim \mathfrak{Sp}(2m, \mathbb{C})^D = 2n_0^2 + n_0 + n_1^2 + n_2^2 + \dots + n_{(k-1)/2}^2.$$

To minimize it, we notice that when we replace n_0 by $n_0 + 1$ and n_1 by $n_1 - 1$ then this quantity increases by $2(2n_0 - n_1 + 2)$. Furthermore, if we replace n_0 by $n_0 - 1$ and n_1 by $n_1 + 1$ then it increases by $2(-2n_0 + n_1 + 1)$. Combined with the previous computations, this quantity is minimized when

$$2n_0 - 1 \leq n_i \leq 2n_0 + 2 \text{ and } |n_i - n_j| \leq 1, \quad \text{for } 1 \leq i, j \leq (k-1)/2.$$

This is realized by the symmetric power of a fuchsian element of order k . \square

Remark 5.17. *Proposition 5.16 also holds true for $\text{PO}(m+1, m)$ with a similar proof. However the argument does not work for $\text{PO}(m+1, m+1)$.*

6 Representations of 3-orbifolds

We restrict to orientable 3-orbifolds. In particular their singular locus is a union of circles, proper segments, and trivalent graphs, where the isotropy groups or stabilizers of edges are cyclic, and the isotropy groups or stabilizers of trivalent vertices are non-cyclic finite subgroups of $\mathrm{SO}(3)$ (hence dihedral, tetrahedral, octahedral or icosahedral). See Figure 6.




Figure 6: Local models for the branching locus, with the corresponding isotropy groups: the cyclic group C_n , the dihedral group D_{2n} , the tetrahedral group T_{12} , the octahedral group O_{24} , and the icosahedral group I_{60} . The subindex denotes the order of the group. The label on an edge denotes the order of the cyclic isotropy group of points in the interior of the edge.

6.1 Geometric representations

Let \mathcal{O}^3 denote a hyperbolic 3-orbifold of finite type, possibly of infinite volume. In particular it has a compactification $\overline{\mathcal{O}^3}$ that is an orbifold with boundary. Let

$$\partial\overline{\mathcal{O}^3} = \partial_1\overline{\mathcal{O}^3} \sqcup \dots \sqcup \partial_k\overline{\mathcal{O}^3}$$

denote its decomposition in connected components.

Consider the composition of the holonomy $\mathrm{hol}: \pi_1(\mathcal{O}^3) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ with the principal representation $\tau: \mathrm{PSL}(2, \mathbb{C}) \rightarrow G$. The following is part of Theorem 1.1.

Theorem 6.1. *The character of $\rho = \tau \circ \mathrm{hol}$ is a smooth point of $X(\mathcal{O}^3, G)$. Furthermore*

$$\begin{aligned} \dim_{[\rho]} X(\mathcal{O}^3, G) &= -\frac{1}{2} \sum_{i=1}^k \tilde{\chi}(\partial_i\overline{\mathcal{O}^3}, \mathrm{Ad}\rho) + \sum_{i=1}^k \dim \mathfrak{g}^{\rho(\pi_1(\partial_i\overline{\mathcal{O}^3}))} \\ &= \frac{1}{2} \sum_{i=1}^k \dim_{[\rho|_{\partial_i}]} X(\partial_i\overline{\mathcal{O}^3}, G) = \frac{1}{2} \dim_{[\rho|_{\partial}]} X(\partial\overline{\mathcal{O}^3}, G). \end{aligned}$$

Proof. We follow the proof for the manifold case in [26] for representations in $\mathrm{PSL}(n, \mathbb{C})$, or [20] in $\mathrm{PSL}(2, \mathbb{C})$. In particular we use Selberg's lemma: there exists $\mathcal{O}_0^3 \rightarrow \mathcal{O}^3$ a finite regular covering which is a manifold.

The first step is to prove that the inclusion of $\partial\overline{\mathcal{O}^3}$ in $\overline{\mathcal{O}^3}$ induces an injection in cohomology

$$H^1(\overline{\mathcal{O}^3}, \mathrm{Ad}\rho) \hookrightarrow H^1(\partial\overline{\mathcal{O}^3}, \mathrm{Ad}\rho).$$

When \mathcal{O}^3 is a manifold, then this is proved in [26] using a theorem of vanishing in L^2 -cohomology (as, using de Rham cohomology and a metric on the bundle and differential forms, every element in the kernel is represented by an L^2 -form), by using the decomposition (2) of $\mathrm{Ad} \circ \tau$. For orbifolds in general, using Selberg's lemma, the inclusion follows from the manifold case and Proposition 3.3.

The next step is to prove that

$$\dim H^1(\overline{\mathcal{O}^3}, \text{Ad}\rho) = \frac{1}{2} \dim H^1(\partial\overline{\mathcal{O}^3}, \text{Ad}\rho).$$

As in the manifold case, this follows from Poincaré duality, Proposition 3.4, and the long exact sequence of the pair $(\overline{\mathcal{O}^3}, \partial\overline{\mathcal{O}^3})$. Namely, if the morphisms in the long exact sequence in cohomology are

$$H^1(\overline{\mathcal{O}^3}, \text{Ad}\rho) \xrightarrow{i^*} H^1(\partial\overline{\mathcal{O}^3}, \text{Ad}\rho) \xrightarrow{\beta} H^2(\overline{\mathcal{O}^3}, \partial\overline{\mathcal{O}^3}; \text{Ad}\rho)$$

then

$$\langle i^*(x), y \rangle_{\partial\overline{\mathcal{O}^3}} = \langle x, \beta(y) \rangle_{\overline{\mathcal{O}^3}} \quad \forall x \in H^1(\overline{\mathcal{O}^3}, \text{Ad}\rho), \forall y \in H^1(\partial\overline{\mathcal{O}^3}, \text{Ad}\rho),$$

where \langle, \rangle denotes the perfect pairing in Proposition 3.4, and the pairing on $\partial\overline{\mathcal{O}^3}$ is the sum of pairings on each connected component. As the pairings are nondegenerate and $\ker \beta = \text{im } i^*$, the claim on the dimension follows from a linear algebra argument.

Finally, as $H^1(\overline{\mathcal{O}^3}, \text{Ad}\rho) \cong H^1(\Gamma, \text{Ad}\rho)$ by Proposition 3.4, with $\Gamma = \pi_1(\mathcal{O}^3)$, we use Goldman's obstruction theory to integrability to prove that $[\rho]$ is a smooth point of the character variety of local dimension $H^1(\overline{\mathcal{O}^3}, \text{Ad}\rho)$. For $\Gamma_0 < \Gamma$ a torsion-free subgroup, any infinitesimal deformation in $H^1(\Gamma, \text{Ad}\rho)$ yields a Γ/Γ_0 -equivariant infinitesimal deformation of Γ_0 , and we may apply the same argument as in Theorem 4.11. Alternatively as $X(\Gamma_0, G)$ is analytically smooth at the character of the restriction of ρ , by Cartan's linearization [4, Lemma 1] there exist local analytic coordinates that linearize the action of Γ/Γ_0 in a neighborhood of the character of $\rho|_{\Gamma_0}$ in $X(\Gamma_0, G)$. Hence the fixed point set of Γ/Γ_0 is a smooth subvariety and, as $\rho|_{\Gamma_0}$ is good this fixed point set can be locally identified with $X(\Gamma, G)$. \square

Definition 6.2. *The component of Theorem 6.1 is called the canonical or distinguished component.*

To complete the proof of Theorem 1.1 we need to show that $X(\mathcal{O}^3, G) \rightarrow X(\partial\mathcal{O}^3, G)$ is locally an injection. This does not follow directly from the injection $H^1(\mathcal{O}^3, \text{Ad}\rho) \rightarrow H^1(\partial\mathcal{O}^3, \text{Ad}\rho)$, as $X(\partial\mathcal{O}^3, G)$ can be non smooth because of rank 2 cusps. (Rank one cusps are part of components with negative Euler characteristic and we discuss only rank 2 cusps.) Consider a nonsingular (manifold) cusp, $T^2 \times [0, +\infty)$ equipped with the warped product metric $e^{-2t}g_{T^2} + dt^2$, where g_{T^2} is a flat metric on the torus and $t \in [0, +\infty)$ is the Busemann function coordinate. Set $\rho = \tau \circ \text{hol}: \pi_1(T^2) \rightarrow G$. Using de Rham cohomology, we may talk about L^2 -forms on $T^2 \times [0, +\infty)$.

Lemma 6.3. *For any non-contractible loop $l \subset T^2$, the restriction induces a surjection*

$$H^1(T^2 \times [0, +\infty), \text{Ad}\rho) \rightarrow H^1(l, \text{Ad}\rho)$$

whose kernel consists of the cohomology classes represented by L^2 -forms.

Proof. Since $\text{Ad} \circ \tau = \oplus_{\alpha} \text{Sym}^{2d_{\alpha}}$ by (2) and $\text{Ad} \circ \text{Sym}^{n-1} = \oplus_{i=1}^{n-1} \text{Sym}^{2i}$, it suffices to prove the lemma for $\tau = \text{Sym}^{n-1}$ and $\mathfrak{g} = \mathfrak{sl}(n, \mathbb{C})$. In [25] precisely the situation when $\tau = \text{Sym}^{n-1}$ is discussed. In particular by [25, Lemma 3.3] the image of the map induced by the inclusion $\mathfrak{sl}(n, \mathbb{C})^{\pi_1(T^2)} \subset \mathfrak{sl}(n, \mathbb{C})$,

$$H^1(T^2, \mathfrak{sl}(n, \mathbb{C})^{\pi_1(T^2)}) \rightarrow H^1(T^2, \mathfrak{sl}(n, \mathbb{C})_{\text{Ad}\rho}) \quad (30)$$

is represented by L^2 -forms. (Here $\mathfrak{sl}(n, \mathbb{C})_{\text{Ad}\rho}$ denotes the coefficients on the Lie algebra, usually just denoted by $\text{Ad}\rho$, whilst on the invariant subspace $\mathfrak{sl}(n, \mathbb{C})^{\pi_1(T^2)}$ the action is trivial by construction.) Furthermore, either by [25, Lemma 3.4] or by a straightforward computation,

$$H^1(T^2, \mathfrak{sl}(n, \mathbb{C})_{\text{Ad}\rho}) \rightarrow H^1(l, \mathfrak{sl}(n, \mathbb{C})_{\text{Ad}\rho})$$

is a surjection with kernel precisely the image of (30). \square

Remark 6.4. Let $\gamma \in \pi_1(T^2)$ be an element represented by the loop l as in Lemma 6.3. We have a natural isomorphism:

$$H^1(l, \text{Ad}\rho) \cong H_0(l, \text{Ad}\rho) \cong H_0(\langle \gamma \rangle, \text{Ad}\rho) \cong \mathfrak{g}_{\rho(\gamma)} = \mathfrak{g}_{\rho(\pi_1(T^2))}$$

Let \tilde{G} be the universal covering of G and let χ_1, \dots, χ_r denote its fundamental characters (the characters of the fundamental representations). E.g. for $\tilde{G} = \text{SL}(n, \mathbb{C})$ the fundamental characters are the symmetric functions on the eigenvalues. By a theorem of Steinberg, they define an isomorphism:

$$(\chi_1, \dots, \chi_r): \tilde{G}^{\text{reg}}/G \cong \mathbb{C}^r, \quad (31)$$

where \tilde{G}^{reg} denotes the set of regular elements in \tilde{G} . When the group G is not simply connected, the characters (χ_1, \dots, χ_r) still define local functions in a neighborhood of $\rho(\gamma)$ in G , where $\gamma \in \pi_1(T^2)$ is represented by the loop l . We identify the variety of representations of the cyclic group $\langle \gamma \rangle \cong \mathbb{Z}$ with G via the image of γ . In particular the differential form $d\chi_i: \mathfrak{g} \rightarrow \mathbb{C}$ may be viewed as a linear map

$$d\chi_i: H^1(\langle \gamma \rangle, \text{Ad}\rho) \cong \mathfrak{g}_{\rho(\gamma)} \rightarrow \mathbb{C},$$

because the characters χ_i are constant on orbits by conjugation (hence the characters are trivial on coboundaries). This uses the natural identification between $H^1(\langle \gamma \rangle, \text{Ad}\rho)$ and the space of coinvariants $\mathfrak{g}_{\rho(\gamma)}$ in Remark 6.4.

Lemma 6.5. $(d\chi_1, \dots, d\chi_r): H^1(\langle \gamma \rangle, \text{Ad}\rho) \rightarrow \mathbb{C}^r$ is an isomorphism.

Proof. We follow the proof of [17, Corollary 19]. Steinberg has shown that the map (31) has a smooth section, $\mathbb{C}^r \rightarrow \tilde{G}^{\text{reg}}$ [30]. In particular $(d\chi_1, \dots, d\chi_r): \mathfrak{g} \rightarrow \mathbb{C}^r$ is surjective and so is the induced map $(d\chi_1, \dots, d\chi_r): H^1(\langle \gamma \rangle, \text{Ad}\rho) \cong \mathfrak{g}_{\rho(\gamma)} \rightarrow \mathbb{C}^r$. Hence the lemma follows from equality of dimensions. \square

Consider $\partial_{\text{hyp}} \overline{\mathcal{O}^3}$ the union of components of $\partial \overline{\mathcal{O}^3}$ that have negative Euler characteristic. In particular, for a manifold $\mathcal{O}^3 = M^3$,

$$\partial \overline{M^3} = \partial_{\text{hyp}} \overline{M^3} \sqcup T_1^2 \sqcup \dots \sqcup T_k^2$$

where k is the number of rank-2 cusps of M^3 .

Proposition 6.6. When $M^3 = \mathcal{O}^3$ is a manifold, then the restriction to the boundary and the characters of peripheral curves of cusps yield a local embedding

$$X(M^3, G) \rightarrow X(\partial_{\text{hyp}} \overline{M^3}, G) \times \mathbb{C}^{rk},$$

where k is the number of rank-2 cusps and the coordinates in \mathbb{C}^{rk} are (locally defined) fundamental characters of chosen peripheral elements, one for each cusp.

Proof. The proof of Theorem 6.1 is based in the vanishing of L^2 -cohomology, here we use the fact that a form is L^2 on M^3 if it is so in the restriction to each end. Hence using Lemmas 6.3 and 6.5 the kernel of the differential of the map of the proposition is represented by L^2 -forms, hence trivial. \square

The following concludes the proof of Theorem 1.1.

Corollary 6.7. The restriction to the boundary yields a local embedding $X(\mathcal{O}^3, G) \rightarrow X(\partial \overline{\mathcal{O}^3}, G)$.

Proof. In the manifold case, we use that the injection in Proposition 6.6 factors through the restriction to $X(\partial\overline{M^3}, G)$:

$$X(M^3, G) \rightarrow X(\partial\overline{M^3}, G) \rightarrow X(\partial_{\text{hyp}}\overline{M^3}, G) \times \mathbb{C}^{rk},$$

Thus the map induced by restriction $X(M^3, G) \rightarrow X(\partial\overline{M^3}, G)$ must be locally an injection. For an orbifold \mathcal{O}^3 , consider a finite manifold covering $\mathcal{O}_0^3 \rightarrow \mathcal{O}^3$, we have the commutative diagram:

$$\begin{array}{ccc} X(\mathcal{O}_0^3, G) & \longrightarrow & X(\partial\overline{\mathcal{O}_0^3}, G) \\ \uparrow & & \uparrow \\ X(\mathcal{O}^3, G) & \longrightarrow & X(\partial\overline{\mathcal{O}^3}, G) \end{array}$$

As \mathcal{O}_0^3 is a manifold, we have already shown that $X(\mathcal{O}_0^3, G) \rightarrow X(\partial\overline{\mathcal{O}_0^3}, G)$ is locally an inclusion. Furthermore, as the restriction of ρ to \mathcal{O}_0^3 is good, the restriction map $X(\mathcal{O}^3, G) \rightarrow X(\mathcal{O}_0^3, G)$ is also locally injective, and we are done from the commutativity of the diagram. \square

A particular case of Proposition 6.6 is the following corollary (proved for representations of manifold groups in $\text{SL}(2, \mathbb{C})$ in [3, 20], and for representations of manifold groups in $\text{SL}(n, \mathbb{C})$ in [25]).

Corollary 6.8. *When M^3 is a manifold whose interior has finite volume, chose $\gamma_1, \dots, \gamma_k$ a nontrivial peripheral element for each cusp. Let χ_1, \dots, χ_r denote the fundamental characters of \widehat{G} . Then*

$$(\chi_1, \gamma_1, \dots, \chi_r, \gamma_k): U \subset X(M^3, G) \rightarrow \mathbb{C}^{kr}.$$

define a local homeomorphism for a neighborhood $U \subset X(M^3, G)$ of $[\tau \circ \text{hol}]$.

Recall that the adjoint or Sym^2 gives an isomorphism $\text{PSL}(2, \mathbb{C}) \cong \text{SO}(3, \mathbb{C})$. By an argument on dimensions (see Table 4):

Corollary 6.9. *Let \mathcal{O}^3 be a compact orientable three-orbifold, with hyperbolic interior of finite volume. If all components of $\partial\mathcal{O}^3$ are homeomorphic to $S^2(2, 2, 2, 2)$, $S^2(2, 4, 4)$ or $S^2(2, 3, 6)$, then*

$$X_0(\mathcal{O}^3, \text{SL}(3, \mathbb{C})) \cong X_0(\mathcal{O}^3, \text{SO}(3, \mathbb{C})) \cong X_0(\mathcal{O}^3, \text{PSL}(2, \mathbb{C})),$$

where X_0 denotes the distinguished component.

6.2 Lower bound of the dimension

In this subsection we provide a lower bound à la Thurston (for $\text{PSL}(2, \mathbb{C})$) or à la Falbel-Guilloux (for general G), in both cases for manifolds. For a compact three orbifold \mathcal{O}^3 , let

$$\partial\mathcal{O}^3 = \partial_1\mathcal{O}^3 \sqcup \dots \sqcup \partial_k\mathcal{O}^3$$

denote the decomposition in connected components of its boundary.

Theorem 6.10. *Let G be semi-simple \mathbb{C} -algebraic Lie group, \mathcal{O}^3 a compact, orientable very good orbifold. Let $\rho: \pi_1(\mathcal{O}^3) \rightarrow G$ be a good representation. Assume that:*

- (a) *if $\partial_i\mathcal{O}^3$ is a hyperbolic boundary component, then the centralizer of its image is zero-dimensional;*
- (b) *if $\partial_i\mathcal{O}^3$ is a Euclidean boundary component, then the restriction $\rho|_{\pi_1(\mathcal{O}^3)}$ is strongly regular.*

Then

$$\begin{aligned}\dim_{[\rho]} X_0(\Gamma, G) &\geq \frac{1}{2} \sum_{i=1}^k \dim_{[\rho|\partial_i]} X_0(\partial_i \mathcal{O}^3, G) \\ &= -\frac{1}{2} \sum_{i=1}^k \tilde{\chi}(\partial_i \mathcal{O}^3, \text{Ad}\rho) + \sum_{i=1}^k \dim \mathfrak{g}^{\rho(\pi_1(\partial_i \mathcal{O}^3))}.\end{aligned}$$

Proof. The branching locus Σ of the compact orbifold \mathcal{O}^3 is a trivalent graph. Chose a finite subset $\Sigma_0 \subset \Sigma$ as follows: for each component of Σ that is a circle chose precisely one point, and chose also the trivalent vertices of Σ . So Σ_0 has the smallest cardinality so that $\Sigma \setminus \Sigma_0$ is a disjoint union of edges (open or not, as they can meet $\partial \mathcal{O}^3$). Consider the orbifold

$$\mathcal{O}_0 = \mathcal{O}^3 \setminus \mathcal{N}(\Sigma_0),$$

ie. remove a tubular neighborhood for each point in Σ_0 . The branching locus of \mathcal{O}_0 is a union of (proper) edges (it contains no vertices nor circles). The boundary components of \mathcal{O}_0 are either the boundary components of \mathcal{O} or the boundary of a neighborhood of a point in Σ_0 (spherical). For each Euclidean or spherical boundary component $\partial_i \mathcal{O}_0$, chose k_i disjoint embedded loops $\gamma_{i,1}, \dots, \gamma_{i,k_i}$ in \mathcal{O}_0 based at $\partial_i \mathcal{O}_0$ so that

$$\mathfrak{g}^{\rho(\pi_1(\partial \mathcal{O}_0))} \cap \mathfrak{g}^{\rho(\gamma_{i,1})} \cap \dots \cap \mathfrak{g}^{\rho(\gamma_{i,k_i})} = 0.$$

Namely, the centralizer in \mathfrak{g} of the image of the group generated by $\pi_1(\partial_i \mathcal{O}_0)$ and $\gamma_{i,1}, \dots, \gamma_{i,k_i}$ is trivial. The next step in the construction is to remove a tubular neighborhood of the $\gamma_{i,j}$:

$$\mathcal{O}_1 = \mathcal{O}_0 \setminus \bigcup_{i,j} \mathcal{N}(\gamma_{i,j}).$$

Finally remove a tubular neighborhood of the branching locus of \mathcal{O}_1 (which is a union of edges):

$$\mathcal{O}_2 = \mathcal{O}_1 \setminus \mathcal{N}(\Sigma_{\mathcal{O}_1}).$$

In particular \mathcal{O}_2 is a manifold.

The connected 3-manifold with non-empty boundary \mathcal{O}_2 retracts to a 2-dimensional CW-complex with a single 0-cell. This gives a presentation of $\pi_1(\mathcal{O}_2)$ from this CW-complex (1-cells are generators and 2-cells relations). Each generator contributes with a copy of G in the variety of representations, and each relation decreases at most $\dim G$ the dimension; this gives the standard bound:

$$\begin{aligned}\dim R(\mathcal{O}_2, G) &\geq (1 - \chi(\mathcal{O}_2)) \dim G = \left(1 - \frac{\chi(\partial \mathcal{O}_2)}{2}\right) \dim G \\ &= -\frac{1}{2} \tilde{\chi}(\partial \mathcal{O}_2, \text{Ad}\rho) + \dim G. \quad (32)\end{aligned}$$

Next we compute lower bounds for the orbifolds \mathcal{O}_1 , \mathcal{O}_0 , and finally for \mathcal{O} . We shall use the following key lemma of Falbel and Guilloux, in fact it is a local version of Proposition 1 in [8]:

Lemma 6.11 ([8]). *Let W be a smooth complex (analytic) variety and $W' \subset W$ a smooth subvariety. Let X be a complex variety with an analytic map $f: X \rightarrow W$. For $p \in X$ there exists a neighborhood $U \subset X$ of p such that*

$$\text{codim}_p(f^{-1}(W') \cap U, X) \leq \text{codim}_{f(p)}(W', W).$$

Proof. By the implicit function theorem, there exists a neighborhood $V \subset W$ of $f(p)$ and an analytic map $F: V \rightarrow \mathbb{C}^k$ such that $V \cap W' = F^{-1}(0)$, where $k = \text{codim}_{f(p)}(W', W)$. Then, for a neighborhood of p , $U \subset f^{-1}(V) \subset X$,

$$f^{-1}(W') \cap U = (F \circ f)^{-1}(0) \cap U,$$

and the estimate follows, because $f^{-1}(W') \cap U$ is a fiber of a map to \mathbb{C}^k . \square

To get \mathcal{O}_1 from \mathcal{O}_2 , we add the edges of $\Sigma_{\mathcal{O}_1}$. Topologically, we add 2-handles whose co-core is an edge of $\Sigma_{\mathcal{O}_1}$. For each singular edge e of $\Sigma_{\mathcal{O}_1}$ consider a meridian $\mu \in \pi_1(\mathcal{O}_2)$, which is represented by the attaching circle of the 2-handle, and has a finite order in $\pi_1(\mathcal{O}_1)$. By homogeneity, the conjugation orbit $G \cdot \rho(\mu) = O(\rho(\mu)) \subset G$ is a smooth analytic submanifold of G , of codimension equal the dimension of the centralizer $\dim(\mathfrak{g}^{\rho(\mu)})$. Thus, by Lemma 6.11, when we add the edge e to \mathcal{O}_2 , the dimension of the variety of representations decreases at most by $\dim(\mathfrak{g}^{\rho(\mu)})$:

$$\dim R(\mathcal{O}_2, G) - \dim R(\mathcal{O}_1, G) = \text{codim}(R(\mathcal{O}_1, G), R(\mathcal{O}_2, G)) \leq \sum_{\mu} \dim(\mathfrak{g}^{\rho(\mu)}). \quad (33)$$

Furthermore, adding the 2-handle to \mathcal{O}_2 corresponding to μ means replacing an annulus by two cone points with cyclic stabilizer generated by μ . Hence, the twisted Euler characteristic increases by $2 \dim(\mathfrak{g}^{\rho(\mu)})$. Counting the contribution of all edges of $\Sigma_{\mathcal{O}_1}$:

$$\tilde{\chi}(\partial \mathcal{O}_1, \text{Ad} \rho) = \tilde{\chi}(\partial \mathcal{O}_2, \text{Ad} \rho) + 2 \sum_{\mu} \dim(\mathfrak{g}^{\rho(\mu)}). \quad (34)$$

Hence from (32), (33), and (34):

$$\dim R(\mathcal{O}_1, G) \geq -\frac{1}{2} \tilde{\chi}(\partial \mathcal{O}_1, \text{Ad} \rho) + \dim G. \quad (35)$$

To get \mathcal{O}_0 from \mathcal{O}_1 , we add the neighborhoods of the loops $\gamma_{i,j}$, i.e. we add 2-handles (without singular co-core). Consider the i -th boundary component $\partial_i \mathcal{O}_1$. The corresponding boundary component $\partial_i \mathcal{O}_0$ is obtained from $\partial_i \mathcal{O}_1$ by the surgery corresponding to the addition of k_i 2-handles. Namely k_i annuli in $\partial_i \mathcal{O}_1$ corresponds to $2k_i$ smooth disks in $\partial_i \mathcal{O}_0$. Thus

$$\tilde{\chi}(\partial_i \mathcal{O}_0, \text{Ad} \rho) = \tilde{\chi}(\partial_i \mathcal{O}_1, \text{Ad} \rho) + 2k_i \dim G. \quad (36)$$

On the other hand, following the idea of [8], we apply Lemma 6.11 to $\partial_i \mathcal{O}_1$ and the free product $\pi_1(\partial_i \mathcal{O}_0) * \langle \gamma_{i,1} \rangle * \cdots * \langle \gamma_{i,k_i} \rangle$, that is, the group obtained by filling the k_i meridians of the surface $\partial_i \mathcal{O}_1$:

$$W' = R(\pi_1(\partial_i \mathcal{O}_0) * \langle \gamma_{i,1} \rangle * \cdots * \langle \gamma_{i,k_i} \rangle, G) \subset R(\partial_i \mathcal{O}_1, G) = W. \quad (37)$$

By Proposition 4.1, and since we assume that the centralizer of $\rho(\pi_1(\partial_i \mathcal{O}_1))$ in \mathfrak{g} is trivial,

$$\dim W = \dim R(\partial_i \mathcal{O}_1, G) = -\tilde{\chi}(\partial_i \mathcal{O}_1, \text{Ad} \rho) + \dim G. \quad (38)$$

By Proposition 4.1 when $\partial_i \mathcal{O}_0$ is hyperbolic, Theorem 4.11 when Euclidean, and Remark 4.17 when spherical, $R(\pi_1(\partial_i \mathcal{O}_0), G)$ is smooth and

$$\dim R(\partial_i \mathcal{O}_0, G) = -\tilde{\chi}(\partial_i \mathcal{O}_0, \text{Ad} \rho) + \dim G + \dim \mathfrak{g}^{\rho(\pi_1(\partial_i \mathcal{O}_0))}. \quad (39)$$

As the variety of representations of a free product is the Cartesian product of varieties or representations of its factors, W' is smooth and:

$$\dim W' = \dim R(\pi_1(\partial_i \mathcal{O}_0) * \langle \gamma_{i,1} \rangle * \cdots * \langle \gamma_{i,k_i} \rangle, G) = \dim R(\partial_i \mathcal{O}_0, G) + k_i \dim G. \quad (40)$$

From (39) and (40):

$$\dim W' = -\tilde{\chi}(\partial_i \mathcal{O}_0, \text{Ad}\rho) + (k_i + 1) \dim G + \dim \mathfrak{g}^{\rho(\pi_1(\partial_i \mathcal{O}_0))}. \quad (41)$$

It follows from (38) and (41) that the codimension of the inclusion (37) is

$$\text{codim}(W', W) = \dim W - \dim W' = \tilde{\chi}(\partial_i \mathcal{O}_0, \text{Ad}\rho) - \tilde{\chi}(\partial_i \mathcal{O}_1, \text{Ad}\rho) - k_i \dim G - \dim \mathfrak{g}^{\rho(\pi_1(\partial_i \mathcal{O}_0))},$$

that combined with (36) yields

$$\text{codim}(W', W) = k_i \dim G - \dim \mathfrak{g}^{\rho(\pi_1(\partial_i \mathcal{O}_0))}. \quad (42)$$

So when we apply Lemma 6.11 we get from the contribution (42) of each boundary component:

$$\text{codim}(R(\mathcal{O}_0, G), R(\mathcal{O}_1, G)) \leq \sum_i \left(k_i \dim G - \dim \mathfrak{g}^{\rho(\pi_1(\partial_i \mathcal{O}_0))} \right). \quad (43)$$

Thus, with (35) and (36), (43) becomes:

$$\begin{aligned} \dim R(\mathcal{O}_0, G) &\geq \dim R(\mathcal{O}_1, G) - \sum_i \left(k_i \dim G - \dim \mathfrak{g}^{\rho(\pi_1(\partial_i \mathcal{O}_0))} \right) \\ &\geq \sum_i \left(-\frac{1}{2} \tilde{\chi}(\partial_i \mathcal{O}_1, \text{Ad}\rho) - k_i \dim G + \dim \mathfrak{g}^{\rho(\pi_1(\partial_i \mathcal{O}_0))} \right) + \dim G \\ &= \sum_i \left(-\frac{1}{2} \tilde{\chi}(\partial_i \mathcal{O}_0, \text{Ad}\rho) + \dim \mathfrak{g}^{\rho(\pi_1(\partial_i \mathcal{O}_0))} \right) + \dim G. \end{aligned}$$

Finally, for each spherical boundary component $\partial_i \mathcal{O}_0$, by (25)

$$-\frac{1}{2} \tilde{\chi}(\partial_i \mathcal{O}_0, \text{Ad}\rho) + \dim \mathfrak{g}^{\rho(\pi_1(\partial_i \mathcal{O}_0))} = 0.$$

Hence we can get rid of the contribution of the neighborhoods of vertices and get the initial orbifold \mathcal{O}^3 :

$$\begin{aligned} \dim R(\mathcal{O}^3, G) &\geq \sum_i \left(-\frac{1}{2} \tilde{\chi}(\partial_i \mathcal{O}^3, \text{Ad}\rho) + \dim \mathfrak{g}^{\rho(\pi_1(\partial_i \mathcal{O}^3))} \right) + \dim G \\ &= \dim X(\partial \mathcal{O}^3, G) + \dim G. \end{aligned}$$

As $\dim X(\mathcal{O}^3, G) = \dim R(\mathcal{O}^3, G) - \dim G$, because the representation is good, this concludes the proof of the theorem. \square

6.3 Dimension growth

Let M^3 be an orientable hyperbolic 3-manifold of finite volume with $k \geq 1$ cusps. As M^3 is a manifold, its holonomy lifts to $\text{SL}(2, \mathbb{C})$.

Let $X_0(M, \text{SL}(n, \mathbb{C}))$ be the *canonical* or *distinguished component* of $X(M, \text{SL}(n, \mathbb{C}))$ (Definition 6.2), i.e. the component that contains the composition of a lift of the holonomy with Sym^{n-1} . By Theorem 6.1,

$$\dim X_0(M, \text{SL}(n, \mathbb{C})) = (n-1)k,$$

where k is the number of cusps. This linear growth differs from the quadratic growth of 2-orbifolds in Section 5. Those 2-orbifolds may appear as basis of Seifert fibered Dehn filling and yield components in the variety of characters of higher dimension. Next we discuss two examples, the figure eight knot and the Whitehead link exteriors. The following is Proposition 1.6 from the introduction.

Proposition 6.12. *Let Γ be the fundamental group of the exterior of the figure eight knot. Besides the canonical component (that has dimension $n - 1$), for large n $X(\Gamma, \mathrm{SL}(n, \mathbb{C}))$ has at least 3 components that contain irreducible representations, whose dimension grow respectively as $n^2/12$, $n^2/20$ and $n^2/42$.*

Proof. Let $K_{p/q}$ denote the Dehn surgery on the figure eight knot with coefficients p/q . There are 6 Dehn fillings on the figure eight knot that yield small Seifert manifolds [11, 24]: $K_{\pm 3}$ are small Seifert manifolds fibered over the 2-orbifold $\mathcal{O}_3^2 = S^2(3, 3, 4)$, $K_{\pm 2}$ over $\mathcal{O}_2^2 = S^2(2, 4, 5)$, and $K_{\pm 1}$ over $\mathcal{O}_1^2 = S^2(2, 3, 7)$. They induce representations in $\mathrm{SL}(2, \mathbb{R})$ of the filled 3-manifolds K_p , that map the fiber to $-\mathrm{Id}$ and induce the holonomy representation of the (unique) hyperbolic structure on \mathcal{O}_p^2 . The composition with Sym^{n-1} is a representation that maps the fiber to $(-1)^{n-1} \mathrm{Id}$ and it induces a representation from $\pi_1(\mathcal{O}_p^2)$ to $\mathrm{PSL}(n, \mathbb{R})$ in the Hitchin component. Therefore the complexification of the Hitchin component of \mathcal{O}_p^2 yields a component of $X(K_p, \mathrm{SL}(n, \mathbb{C}))$ of the same dimension, and hence a subvariety of $X(\Gamma, \mathrm{SL}(n, \mathbb{C}))$. Let $X_p \subset X(M, \mathrm{SL}(n, \mathbb{C}))$ be the component of $X(\Gamma, \mathrm{SL}(n, \mathbb{C}))$ that contains the subvariety induced by $X(K_p, \mathrm{SL}(n, \mathbb{C}))$. Next we estimate the dimension of X_1 , X_2 and X_3 , which allow to distinguish them, but does not allow to distinguish the component induced by X_p from X_{-p} , as the dimension estimates are the same.

The lower bound on the dimension of X_p is given by Proposition 5.10:

$$\dim X_p \geq -\chi(\mathcal{O}_p^2)n^2 - c(\mathcal{O}_p^2).$$

Aiming to find an upper bound of $\dim X_p$, we bound above the dimension of its Zariski tangent space, which is isomorphic to $H^1(\Gamma, \mathrm{Ad}\rho) \cong H^1(M, \mathrm{Ad}\rho)$ for M the (compact) exterior of the figure eight knot, $M = S^3 \setminus \mathcal{N}(K)$. In particular $\partial M \cong T^2$. Here $\rho = \mathrm{Sym}^{n-1} \circ \rho_0$, for $\rho_0: \Gamma \rightarrow \mathrm{SL}(2, \mathbb{R})$ the representation that factors through K_p .

First we bound $\dim H^1(\partial M, \mathrm{Ad}\rho)$. Notice that since $\chi(T^2) = 0$,

$$\dim H^1(\partial M, \mathrm{Ad}\rho) = 2 \dim H^0(\partial M, \mathrm{Ad}\rho) = 2 \dim \mathfrak{g}^{\mathrm{Ad}\rho(\pi_1(\partial M))}.$$

To compute $\dim \mathfrak{g}^{\mathrm{Ad}\rho(\pi_1(\partial M))}$, one may check explicitly that the image of $\rho_0(\pi_1(\partial M))$ contains an element of $\mathrm{SL}(2, \mathbb{R})$ of infinite order (one may use for instance the A-polynomial). As the symmetric power of a hyperbolic matrix in $\mathrm{SL}(2, \mathbb{C})$ is regular, $\rho(\pi_1(\partial M))$ contains regular elements and $\dim \mathfrak{g}^{\mathrm{Ad}\rho(\pi_1(\partial M))} = \mathrm{rank} G = n - 1$. Thus $\dim H^1(\partial M, \mathrm{Ad}\rho) = 2(n - 1)$.

Let

$$i^*: H^1(M, \mathrm{Ad}\rho) \rightarrow H^1(\partial M, \mathrm{Ad}\rho)$$

be the morphism induced by inclusion. Using the long exact sequence of the pair $(M, \partial M)$ and Poincaré duality as in the proof of Theorem 6.1, its rank is

$$\mathrm{rank}(i^*) = n - 1. \tag{44}$$

Furthermore, by Mayer-Vietoris exact sequence applied to the decomposition $K_p = M \cup_{\partial M} (D^2 \times S^1)$, we get that

$$\dim \ker i^* \leq \dim H^1(K_p, \mathrm{Ad}\rho). \tag{45}$$

Next we need a lemma:

Lemma 6.13. *The projection $\pi_1(K_p) \rightarrow \pi_1(\mathcal{O}_p^2)$ induces an isomorphism*

$$H^1(\pi_1(K_p), \mathrm{Ad}\rho) \cong H^1(\pi_1(\mathcal{O}_p^2), \mathrm{Ad}\rho). \tag{46}$$

Assuming the lemma, we conclude the proof of Proposition 6.12. Putting together (44), (45) and (46):

$$\dim H^1(M, \text{Ad}\rho) \leq \dim H^1(\mathcal{O}_p^2, \text{Ad}\rho) + (n-1).$$

It follows that

$$-n^2\chi(\mathcal{O}_p^2) - c(\mathcal{O}_p^2) \leq \dim X_p \leq -n^2\chi(\mathcal{O}_p^2) + c(\mathcal{O}_p^2) + n - 1.$$

For large n this allows to distinguish three components, according to the different values for $\chi(\mathcal{O}_p^2)$. \square

Proof of Lemma 6.13. We use the central exact sequence

$$1 \rightarrow \mathcal{Z}(\pi_1(K_p)) \rightarrow \pi_1(K_p) \rightarrow \pi_1(\mathcal{O}_p^2) \rightarrow 1,$$

where the center $\mathcal{Z}(\pi_1(K_p)) \cong \mathbb{Z}$ is the group generated by the fiber. To prove the lemma, we use crossed morphisms or derivations, as in Subsection 2.1. Since $\pi_1(K_p) \rightarrow \pi_1(\mathcal{O}_p^2)$ is a surjection, we have an injection of spaces of cocycles

$$Z^1(\pi_1(\mathcal{O}_p^2), \mathfrak{g}) \hookrightarrow Z^1(\pi_1(K_p), \mathfrak{g}). \quad (47)$$

We prove surjectivity of the map in (47): let $d: \pi_1(K_p) \rightarrow \mathfrak{g}$ be a crossed morphism (namely, satisfying $d(\gamma_1\gamma_2) = d(\gamma_1) + \text{Ad}_{\rho(\gamma_1)}d(\gamma_2)$, $\forall \gamma_1, \gamma_2 \in \pi_1(K_p)$). Let $t \in \pi_1(K_p)$ be a generator of the center $\mathcal{Z}(\pi_1(K_p)) \cong \mathbb{Z}$; from the relation

$$t\gamma t^{-1} = \gamma \quad \forall \gamma \in \pi_1(K_p),$$

we deduce

$$d(t) + \text{Ad}_{\rho(t)}d(\gamma) - \text{Ad}_{\rho(t\gamma t^{-1})}d(t) = d(\gamma), \quad \forall \gamma \in \pi_1(K_p).$$

Since $\rho(t) = \pm \text{Id}$, $\text{Ad}_{\rho(t)}$ is the identity on \mathfrak{g} , hence

$$(1 - \text{Ad}_{\rho(\gamma)})d(t) = 0, \quad \forall \gamma \in \pi_1(K_p).$$

This equality implies that $d(t) = 0$, because $\mathfrak{g}^{\rho(\pi_1(K_p))} = 0$ by irreducibility of ρ . This proves that every crossed morphism of $\pi_1(K_p)$ factors through a crossed morphism of $\pi_1(\mathcal{O}_p^2)$, hence (47) is surjective. This isomorphism between cocycle spaces induces an isomorphism of coboundary space and this proves the lemma. \square

Remark 6.14. For small values of n , the arguments do not apply, but for $n = 3$ the canonical component has dimension 2, and according to Table 6, (± 3) -Dehn fillings also provide two subvarieties of dimension 2. It is proved in [9] and [15] that the canonical component and the subvarieties induced by the (± 3) -Dehn filling are precisely the three components of $X(\Gamma, \text{SL}(3, \mathbb{C}))$ that contain irreducible representations.

Remark 6.15. At the moment, we do not know whether for large n the component that contains representations that factor through K_p is the same as the component corresponding to K_{-p} , for $p = 1, 2, 3$. To distinguish them would require a better upper bound of the dimension of their Zariski tangent space. This would allow also to distinguish components corresponding to Galois conjugates (5-th or 7-th roots of unity).

Remark 6.16. An analog of Proposition 6.12 can be obtained for the variety of characters of the figure eight knot in $\text{Sp}(2m, \mathbb{C})$, for large m .

Proposition 6.17. *Let Γ be the fundamental group of the exterior of the Whitehead link. Besides the canonical component (that has dimension $2n - 2$), for large n , $X(\Gamma, \mathrm{SL}(n, \mathbb{C}))$ has at least 3 components that contain irreducible characters, whose dimension grow respectively as $n^2/3$, $n^2/4$ and $n^2/6$.*

The proof of Proposition 6.17 is the same as Proposition 6.12. Here there are partial Dehn fillings that yield Seifert fibered manifolds, with basis a disc with two cone points: the (-3) -filling fibers over $D^2(3, 3)$, the (-2) -filling over $D^2(2, 4)$, and the (-1) -filling is fibered over $D^2(2, 3)$ [24, Table A.1]. This yields the three rational orbifold Euler characteristics, $1/3$, $1/4$ and $1/6$.

As the basis orbifolds are rather simple, it is also easy to state the dimension of their Hitchin component. The dimension of the Hitchin component of $D^2(3, 3)$ is

$$\begin{cases} \frac{n^2}{3} + 1 & \text{for } n \equiv 0 \pmod{3} \\ \frac{n^2-1}{3} & \text{for } n \not\equiv 0 \pmod{3} \end{cases}$$

The dimension of the Hitchin component of $D^2(2, 4)$ is

$$\begin{cases} \frac{n^2}{4} + 1 & \text{for } n \equiv 0 \pmod{4} \\ \frac{n^2-1}{4} & \text{for } n \equiv 1 \pmod{4} \\ \frac{n^2}{4} & \text{for } n \equiv 2 \pmod{4} \end{cases}$$

The dimension of the Hitchin component of $D^2(2, 3)$ is

$$\begin{cases} \frac{n^2}{6} + 1 & \text{for } n \equiv 0 \pmod{6} \\ \frac{n^2-1}{6} & \text{for } n \equiv \pm 1 \pmod{6} \\ \frac{n^2+2}{6} & \text{for } n \equiv 2 \pmod{6} \\ \frac{n^2+3}{6} & \text{for } n \equiv 3 \pmod{6} \end{cases}$$

Remark 6.18. *When we apply these computation to $\mathrm{SL}(3, \mathbb{C})$, for $D^2(2, 4)$ and $D^2(2, 3)$ it yields dimension 2, but for $D^2(3, 3)$ it has a component of dimension 4, the same as the dimension of the canonical component. It has been proved by Guilloux and Will [12] that this is in fact a whole component of $X(\Gamma, \mathrm{SL}(3, \mathbb{C}))$.*

For Montesinos links, the same arguments yield the following:

Proposition 6.19. *Let $L \subset S^3$ be a Montesinos link. Assume that either it has at least 4 tangles, or that it has three tangles (eg. a pretzel link) with indices α_1 , α_2 , and α_3 satisfying $\frac{1}{\alpha_1} + \frac{1}{\alpha_2} + \frac{1}{\alpha_3} \leq 1$. Then $\dim X(S^3 \setminus L, \mathrm{SL}(n, \mathbb{C}))$ grows quadratically with n .*

7 Representations in $\mathrm{SL}(3, \mathbb{C})$

In this section we give explicit computations of some varieties of characters in $\mathrm{SL}(3, \mathbb{C})$. We compute varieties of characters of groups generated by two elements, using the description of $X(F_2, \mathrm{SL}(3, \mathbb{C}))$ due to Lawton [21], where $F_2 = \langle a, b \mid \rangle$ is the free group of rank two.

Setting coordinates

$$\begin{aligned} x &= t_a, & y &= t_b, & z &= t_{ab}, & r &= t_{ab^{-1}}, & \tau &= t_{[a,b]}, \\ u &= t_{a^{-1}}, & v &= t_{b^{-1}}, & w &= t_{(ab)^{-1}}, & s &= t_{a^{-1}b}, \end{aligned} \tag{48}$$

Lawton has proved:

Theorem 7.1 ([21]). $X(F_2, \text{SL}(3, \mathbb{C}))$ is the hypersurface of \mathbb{C}^9 defined by

$$\tau^2 - P\tau + Q = 0$$

for some polynomials $P, Q \in \mathbb{Z}[x, y, z, u, v, w, r, s]$. The solutions of $\tau^2 - P\tau + Q = 0$ are $t_{[a,b]}$ and $t_{[b,a]}$. Namely $t_{[a,b]} + t_{[b,a]} = P$ and $t_{[a,b]}t_{[b,a]} = Q$.

The polynomials P and Q in Theorem 7.1 can be found in [21]:

$$P = xuyv - uyr - xvs - uvz - xyw + rs + xu + yv + zw - 3$$

and

$$\begin{aligned} Q = & vu^2x^2y + uv^2y^2x - ryxu^2 - uxvrw - ry^2vu + rx^2v^2 - rzyxv + su^2y^2 - swvuy - svux^2 \\ & - uxyzs - sv^2yx - u^3vy + u^2v^2w - zu^2xv - uv^3x - zv^2yu - wx^2uy - uy^3x - wy^2vx \\ & - x^3vy + x^2y^2z + uwr^2 - 2r^2xv + r^2zy + surx + svry + wszr + rzu^2 + ruv^2 + rvz^2 \\ & + xrw^2 + rwy^2 + x^2yr - 2s^2uy + s^2wv + xzs^2 + u^2vs + usz^2 + szv^2 + syw^2 + swx^2 \\ & + sxy^2 + u^2wy - 2w^2vu + xuyv + wuzx + uy^2z + xv^2w + wvzy + x^2zv - 2z^2yx \\ & + r^3 + 3uyr - 3rvw - 3rzx + s^3 - 3swu + 3xvs - 3syz + u^3 + 3uvz + v^3 + w^3 \\ & + 3xyw + x^3 + y^3 + z^3 - 6rs - 6xu - 6yv - 6zw + 9 \end{aligned}$$

This theorem tells that an $\text{SL}(3, \mathbb{C})$ -character of F_2 is a polynomial on the coordinates $x, y, z, u, v, w, r, s, \tau$. Its expression can be computed by using basic identities on traces, including Cayley-Hamilton's formula

$$A^3 - \text{tr}(A)A^2 + \text{tr}(A^{-1})A - \text{Id} = 0, \quad \forall A \in \text{SL}(3, \mathbb{C}), \quad (49)$$

and elementary identities on traces.

7.1 Two dimensional examples

Let us compute the variety of $\text{SL}(3, \mathbb{C})$ -characters of some 2-orbifolds.

We start with the turnover $S^2(3, 3, 4)$, Figure 7. Its fundamental group has presentation

$$\pi_1(S^2(3, 3, 4)) \cong \langle a, b \mid a^3 = b^3 = (ab)^4 = 1 \rangle$$

We look at the possible eigenvalues of the elements of finite order:

- (a) A matrix in $\text{SL}(3, \mathbb{C})$ of order 3 is either central or has eigenvalues $\{1, \omega, \omega^2\}$, for $\omega \in \mathbb{C}$ a primitive third root of unity: $\omega^3 = 1$ and $\omega \neq 1$.
- (b) A matrix in $\text{SL}(3, \mathbb{C})$ of order 4 is either trivial or has eigenvalues

$$\{1, i, -i\}, \{-1, i, i\}, \{-1, -i, -i\}, \text{ or } \{1, -1, -1\}.$$

Remark 7.2. Let $\rho: \pi_1(S^2(3, 3, 4)) \rightarrow \text{SL}(3, \mathbb{C})$ be an irreducible representation. The eigenvalues of $\rho(a)$ and $\rho(b)$ are $\{1, \omega, \omega^2\}$ and the eigenvalues of $\rho(ab)$ are $\{1, i, -i\}$, $\{-1, i, i\}$, $\{-1, -i, -i\}$, or $\{1, -1, -1\}$. Furthermore, using that for an irreducible representation

$$\dim_{[\rho]} X(S^2(3, 3, 4), \text{SL}(3, \mathbb{C})) = -\tilde{\chi}(S^2(3, 3, 4), \text{Ad}\rho),$$

and $\dim \mathfrak{g}^{\rho(ab)}$ depends on whether $\rho(ab)$ has repeated eigenvalues or not, we have:

$$\dim_{[\rho]} X(S^2(3, 3, 4), \text{SL}(3, \mathbb{C})) = \begin{cases} 2 & \text{if the eigenvalues of } \rho(ab) \text{ are } \{1, i, -i\}, \\ 0 & \text{otherwise.} \end{cases}$$




Figure 7: The turnover $S^2(3, 3, 4)$ and two nonorientable quotients: a triangle with mirror boundary $T(3, 3, 4)$, which is a Coxeter group, and a disc with a cone point, mirror boundary, and a corner reflector, $D(3; 4)$.

Using the coordinates (48), the two dimensional components of the variety of characters are obtained by setting $x = y = u = v = 0$ and $z = w = 1$, because this fixes the eigenvalues of $\rho(a)$, $\rho(b)$ and $\rho(ab)$. By replacing those values in Theorem 7.1, we obtain:

Example 7.3. *The component of $X(S^2(3, 3, 4), \text{SL}(3, \mathbb{C}))$ that has positive dimension is isomorphic to*

$$\{(r, s, \tau) \in \mathbb{C}^3 \mid \tau^2 - (rs - 2)\tau + (r^3 + s^3 - 5rs + 5) = 0\}.$$

For the components that are isolated points, we set, again $x = y = u = v = 0$ and, according to the eigenvalues of $\rho(ab)$:

$$(z, w) = (-1 + 2i, -1 - 2i), \quad (z, w) = (-1 - 2i, -1 + 2i), \quad \text{or} \quad (z, w) = (-1, -1).$$

Notice that this does not fix the conjugacy class of $\rho(ab)$, because it has eigenvalues of multiplicity 2. We need to find further restrictions on the traces: by taking traces on the relation $(ab)^{-2}a = (ab)^2a$, using (49) and replacing $x = y = u = v = 0$ we get

$$s = zs \quad \text{and} \quad r = wr. \tag{50}$$

For the 2-dimensional component it holds $z = w = 1$, hence (50) do not give any further information. For the isolated components, this yields $r = s = 0$. For these values $P^2 - 4Q = 0$ and therefore there is a unique value for τ . Hence we get:

Example 7.4. *There are three components of $X(S^2(3, 3, 4), \text{SL}(3, \mathbb{C}))$ that contain irreducible representations and are zero-dimensional. They have coordinates $x = y = u = v = r = s = 0$ and*

$$(z, w, \tau) = (-1 + 2i, -1 - 2i, 1), \quad (-1 - 2i, -1 + 2i, 1), \quad \text{or} \quad (-1, -1, -1).$$

Next we describe the Hitchin component $\text{Hit}(S^2(3, 3, 4), \text{PSL}(3, \mathbb{R}))$. Notice that the representation $\text{Sym}^2: \text{SL}(2, \mathbb{R}) \rightarrow \text{SL}(3, \mathbb{R})$ factors through $\text{PSL}(2, \mathbb{R})$, hence $\text{Hit}(S^2(3, 3, 4), \text{PSL}(3, \mathbb{R}))$ lifts to a component of $X(S^2(3, 3, 4), \text{SL}(3, \mathbb{R}))$. We consider the 2-dimensional component of Example 7.3 and we require that its coordinates are real:

$$\{(r, s, \tau) \in \mathbb{R}^3 \mid \tau^2 - (rs - 2)\tau + (r^3 + s^3 - 5rs + 5) = 0\}.$$

By looking at the discriminant of this equation, this real set has three components. One of them is the isolated point $(r, s, \tau) = (2, 2, 1)$, which is in fact a singular point of the defining surface. The other two components are homeomorphic to \mathbb{R}^2 , one of them is the lift of the Hitchin component. By looking at the symmetric power of the holonomy, we deduce:

Example 7.5. *The Hitchin component $\text{Hit}(S^2(3, 3, 4), \text{PSL}(3, \mathbb{R}))$ is isomorphic to the component of*

$$\{(r, s, \tau) \in \mathbb{R}^3 \mid \tau^2 - (rs - 2)\tau + (r^3 + s^3 - 5rs + 5) = 0\}$$

that contains the point of coordinates $(r, s, \tau) = (2 + 2\sqrt{2}, 2 + 2\sqrt{2}, 5 + 4\sqrt{2})$.

Next we consider the turnover $T(3, 3, 4)$, Figure 7. Its fundamental group is an extension of $\pi_1(S^2(3, 3, 4))$ by an involution σ , that satisfies

$$\sigma^2 = 1, \quad \sigma a \sigma = a^{-1}, \quad \text{and} \quad \sigma b \sigma = b^{-1}. \quad (51)$$

Denote by σ_* the involution induced on the fundamental group:

$$\sigma_*(\gamma) = \sigma \gamma \sigma^{-1}, \quad \forall \gamma \in \pi_1(S^2(3, 3, 4)).$$

This involution σ_* permutes the coordinates r and s and preserves τ , the trace of the commutator. Therefore, the component of $X(S^2(3, 3, 4), \text{SL}(3, \mathbb{C}))^{\sigma_*}$ of positive dimension is

$$\{(r, \tau) \in \mathbb{C}^2 \mid \tau^2 - (r^2 - 2)\tau + (2r^3 - 5r^2 + 5) = 0\}.$$

This curve is singular at precisely at $(r, \tau) = (2, 1)$, and the singularity is an ordinary double point (a self intersection with transverse tangents).

Lemma 7.6. *The restriction map*

$$X(T(3, 3, 4), \text{SL}(3, \mathbb{C})) \rightarrow X(S^2(3, 3, 4), \text{SL}(3, \mathbb{C}))^{\sigma_*}$$

desingularizes the curve $\tau^2 - (r^2 - 2)\tau + (2r^3 - 5r^2 + 5) = 0$.

The component of $X(T(3, 3, 4), \text{SL}(3, \mathbb{C}))$ of positive dimension that contains irreducible representations is this desingularization.

Proof. We look at the fibre of the restriction map. We show that:

- (a) the fibre of an irreducible character consists precisely of one point,
- (b) $(r, \tau) = (2, 1)$ is the only reducible character in $X(S^2(3, 3, 4), \text{SL}(3, \mathbb{C}))^{\sigma_*}$, and
- (c) the fibre of $(r, \tau) = (2, 1)$ consists precisely of 2 points.

Those three items prove the lemma, because the singularity is an ordinary double point.

We prove (a): For an irreducible representation ρ of $\pi_1(S^2(3, 3, 4))$ in $\text{SL}(3, \mathbb{C})$ we show that there is a unique choice of $A \in \text{SL}(3, \mathbb{C})$ such that mapping σ to A defines an extension of ρ to $\pi_1(T(3, 3, 4))$. By irreducibility of ρ , as ρ and $\rho \circ \sigma_*$ have the same character, there exists a matrix $A \in \text{SL}(3, \mathbb{C})$ such that $A\rho A^{-1} = \rho \circ \sigma_*$. Furthermore, A is unique up multiplication by a matrix in the center (the center of $\text{SL}(3, \mathbb{C})$ is $\{\text{Id}, \omega \text{Id}, \omega^2 \text{Id}\}$). As σ is an involution, A^2 commutes with ρ , that is irreducible, and therefore $A^2 \in \{\text{Id}, \omega \text{Id}, \omega^2 \text{Id}\}$. Hence among $A, \omega A$ or $\omega^2 A$ there is a unique choice whose square is the identity. Thus there is a unique choice that satisfies (51).

To prove (b) we use that for a reducible representation the trace of a commutator and its inverse are the same, thus it must be a zero of the discriminant of the defining equation. This discriminant is

$$(r - 2)^2(r^2 - 4r - 4) \quad (52)$$

and its zeros are $r = 2$ and $r = 2 \pm 2\sqrt{2}$. The value $r = 2 \pm 2\sqrt{2}$ corresponds to the symmetrization of the fuchsian holonomy (and its Galois conjugate), hence it is irreducible. Thus $r = 2$ is the only reducible character.

To prove (c), we check that $r = 2$ corresponds to a representation $\rho = \rho_2 \oplus \text{Id}$, for $\rho_2: \pi_1(S^2(3, 3, 4)) \rightarrow \text{SL}(2, \mathbb{C})$ an irreducible representation (for a reducible representation in $\text{SL}(2, \mathbb{C})$ the trace of any commutator is 2 and, as $\tau = 1$, $\text{trace}(\rho_2([a, b])) = \tau - 1 \neq 2$). Then, reproducing the argument of the irreducible case (a), there exist a matrix $A \in \text{SL}(2, \mathbb{C})$ such that $A\rho_2 A^{-1} = \rho_2 \circ \sigma_*$ and, by irreducibility of ρ_2 , A is unique up to sign. Furthermore, again by irreducibility of ρ_2 , A^2 is central in $\text{SL}(2, \mathbb{C})$, namely $A^2 = \pm \text{Id}$. The case $A^2 = +\text{Id}$ does not occur, because this would imply that $A = \pm \text{Id}$ and with (51) this would yield that ρ_2 itself would be central, but it is irreducible. Therefore $A^2 = -\text{Id}$ and the choices for $\rho(\sigma)$ are

$$\rho(\sigma) = \begin{pmatrix} \pm i A & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

This concludes the proof of (c) and of the lemma. \square

Next we look for the Hitchin component of the turnover. We look therefore at the real points

$$\{(r, \tau) \in \mathbb{R}^2 \mid \tau^2 - (r^2 - 2)\tau + (2r^3 - 5r^2 + 5) = 0\}.$$

By looking at the discriminant (52), the set of real points has three components, the isolated point $(r, \tau) = (2, 1)$ and two lines, defined by $r \leq 2 - 2\sqrt{2}$ and $r \geq 2 + 2\sqrt{2}$. As $r = 2 + 2\sqrt{2}$ is the symmetric power of the holonomy:

Example 7.7. *The Hitchin component $\text{Hit}(T(3, 3, 4), \text{PSL}(3, \mathbb{R}))$ is isomorphic (via the restriction to $S^2(3, 4, 4)$) to the line*

$$\{(r, \tau) \in \mathbb{R}^2 \mid \tau^2 - (r^2 - 2)\tau + (2r^3 - 5r^2 + 5) = 0, r \geq 2 + 2\sqrt{2}\}.$$

Finally, we consider $D(3; 4)$, the disc with a cone point of order 3, mirror boundary, and a corner reflector of order 4 (with isotropy group the dihedral group of 8 elements). It is again the quotient of $S^2(3, 3, 4)$ by an involution. This involution maps a to b^{-1} and b to a^{-1} . Therefore it fixes the coordinates r and s but permutes the trace of $[a, b]$ with the trace of its inverse $[b, a]$. The fixed point set of this involution in $X(S^2(3, 3, 4), \text{SL}(3, \mathbb{C}))$ is obtained by looking at the zero locus of the discriminant of the variable τ :

$$\{(r, s) \in \mathbb{C}^2 \mid r^2 s^2 - 4r^3 - 4s^3 + 16rs - 16 = 0\}.$$

With the very same discussion as in Lemma 7.6:

Lemma 7.8. *The restriction map*

$$X(D(3; 4), \text{SL}(3, \mathbb{C})) \rightarrow X(S^2(3, 3, 4), \text{SL}(3, \mathbb{C}))^{\sigma'}$$

desingularizes the curve $r^2 s^2 - 4r^3 - 4s^3 + 16rs - 16 = 0$.

The component of $X(D(3; 4), \text{SL}(3, \mathbb{C}))$ of positive dimension that contains irreducible characters is this desingularization.

The set of real points of $r^2 s^2 - 4r^3 - 4s^3 + 16rs - 16 = 0$ is the discriminant locus of the set of real points for $S^2(3, 3, 4)$: it has three components, the isolated point $r = s = 2$ and two unbounded curves. Thus, similarly to $S^2(3, 3, 4)$ we have:

Example 7.9. *The Hitchin component $\text{Hit}(D(3; 4), \text{PSL}(3, \mathbb{R}))$ is isomorphic (via the restriction to $S^2(3, 4, 4)$) to the component of*

$$\{(r, s) \in \mathbb{R}^2 \mid r^2 s^2 - 4r^3 - 4s^3 + 16rs - 16 = 0\}$$

that contains $r = s = 2 + 2\sqrt{2}$.

7.2 A family of 3-dimensional orbifolds

Consider the family of examples with underlying space S^3 (perhaps punctured twice) and three branching arcs as in Figure 8, with branching labels n_0 , n_1 and $n_2 \geq 2$. Whether a singular vertex is included or not depends on the orders of the adjacent branching locus: if $1/n_0 + 2/n_1 > 1$, then the three edges meet at a vertex, otherwise the vertex is removed, and similarly for the other vertex. In other words, we consider the twice punctured 3-sphere, and we add the vertices needed for the orbifold to be irreducible.




Figure 8: The underlying space of the orbifold is a 3-sphere, possibly punctured once or twice, according to the branching indices, the branching locus consists of three arcs as in the picture.

Let \mathcal{O}^3 denote this orbifold. The geometry of \mathcal{O}^3 depends on the indices:

- (a) When $n_2 = 2$ and $1/n_0 + 2/n_1 \leq 1$, it is an interval bundle, over a disc with mirror boundary, with an interior cone point of order n_1 and a corner reflector of order $2n_0$. It is doubly covered by the product of a turnover $S^2(n_0, n_1, n_1)$ by an interval.
- (b) When $n_2 = 2$ and $1/n_0 + 2/n_1 > 1$, it is obtained by coning the boundary of the previous example, and it is spherical. It is doubly covered by the suspension over a turnover $S^2(n_0, n_1, n_1)$.
- (c) When $(n_0, n_1, n_2) = (2, 3, 3)$ the orbifold is spherical [7].
- (d) If $n_1, n_2 \neq 2$ and $(n_0, n_1, n_2) \neq (2, 3, 3)$, then it is hyperbolic.

The hyperbolicity is realized by a tetrahedron as in Figure 9. The vertices are possibly finite, ideal or exterior (hence truncated by geodesic triangles perpendicular to the other sides). Poincaré fundamental domain theorem yields the following presentation for the fundamental group

$$\pi_1(\mathcal{O}^3) \cong \langle a, b \mid a^{n_1} = b^{n_2} = [a, b]^{n_0} = 1 \rangle.$$

In case (a) the variety of characters of \mathcal{O}^3 in $\mathrm{SL}(3, \mathbb{C})$ is the variety of characters of a non-orientable 2-orbifold. It has dimension 1 or 0 according to the values of n_0 and n_1 . Cases (b) and (c) are spherical, hence the variety of characters is finite. In the hyperbolic case, (d), when $n_0 = 2$ then the canonical component is an isolated point, by Theorem 6.1. For $n_i \geq 3$, it has dimension 2.

Example 7.10. Consider the case $n_0 = n_1 = n_2 = 3$. The fundamental group is

$$\pi_1(\mathcal{O}^3) = \langle a, b \mid a^3 = b^3 = [a, b]^3 = 1 \rangle.$$

On the canonical component, the image of any of a , b and $[a, b]$ is not central, therefore by Theorem 7.1, using the notation in (48) we impose the following equalities:

$$x = y = u = v = P = Q = 0.$$

Thus the canonical component $X_0(\mathcal{O}^3, \mathrm{SL}(3, \mathbb{C}))$ is the following complex surface:

$$\begin{cases} rs + zw - 3 = 0 \\ r^3 + s^3 + z^3 + w^3 + rszw - 6rs - 6zw + 9 = 0 \end{cases}$$




Figure 9: The fundamental domain of the orbifold in this subsection. It is an ideal tetrahedron in hyperbolic 3-space, with vertices perhaps ideal or hyperideal. The side parings, a and b , are rotations of order n_1 and n_2 around two opposite edges. Those edges have dihedral angle $2\pi/n_1$ and $2\pi/n_2$ respectively, the remaining four edges have dihedral angle $\pi/(2n_0)$.

The symmetric power of the complete structure has coordinates

$$r = s = \frac{-3 + \sqrt{-3}}{2}, \quad z = w = \frac{-3 - \sqrt{-3}}{2}$$

(The complex conjugate corresponds to a change of orientation).

There are other components of $X(\mathcal{O}^3, \mathrm{SL}(3, \mathbb{C}))$ but it may be checked that they are isolated points. For instance, if $\rho(a)$ is central, then $\rho(b)$ has order three at this gives finitely many conjugacy classes for ρ (whether $\rho(b)$ is central or not). If $\rho(a)$ and $\rho(b)$ are noncentral but $\rho([a, b])$ is central, then it can be computed that $x = y = z = r = u = v = w = s = 0$. This is realized by the representation

$$\rho(a) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix} \quad \text{and} \quad \rho(b) = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

When $n_0 = n_1 = n_2 = 3$, a component of the variety of characters in $\mathrm{SL}(4, \mathbb{R})$ corresponding to projective structures has been computed in [28].

References

- [1] Daniele Alessandrini, Gye-Seon Lee, and Florent Schaffhauser. Hitchin components for orbifolds. arXiv:1811.05366, 2018.
- [2] Michel Boileau, Sylvain Maillot, and Joan Porti. *Three-dimensional orbifolds and their geometric structures*, volume 15 of *Panoramas et Synthèses [Panoramas and Syntheses]*. Société Mathématique de France, Paris, 2003.
- [3] K. Bromberg. Rigidity of geometrically finite hyperbolic cone-manifolds. *Geom. Dedicata*, 105:143–170, 2004.
- [4] Henri Cartan. Quotient d’une variété analytique par un groupe discret d’automorphismes. *Séminaire Henri Cartan*, 6, 1953-1954. talk:12.

- [5] Suhyoung Choi, Gye-Seon Lee, and Ludovic Marquis. Deformations of convex real projective manifolds and orbifolds. In *Handbook of group actions. Vol. III*, volume 40 of *Adv. Lect. Math. (ALM)*, pages 263–310. Int. Press, Somerville, MA, 2018.
- [6] Igor Dolgachev. *Lectures on invariant theory*, volume 296 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2003.
- [7] William D. Dunbar. Nonfibered spherical 3-orbifolds. *Trans. Amer. Math. Soc.*, 341(1):121–142, 1994.
- [8] E. Falbel and A. Guilloux. Dimension of character varieties for 3-manifolds. *Proc. Amer. Math. Soc.*, 145(6):2727–2737, 2017.
- [9] E. Falbel, A. Guilloux, P.-V. Koseleff, F. Rouillier, and M. Thistlethwaite. Character varieties for $\mathrm{SL}(3, \mathbb{C})$: the figure eight knot. *Exp. Math.*, 25(2):219–235, 2016.
- [10] William M. Goldman. The symplectic nature of fundamental groups of surfaces. *Adv. in Math.*, 54(2):200–225, 1984.
- [11] Cameron Gordon. Dehn surgery and 3-manifolds. In *Low dimensional topology*, volume 15 of *IAS/Park City Math. Ser.*, pages 21–71. Amer. Math. Soc., Providence, RI, 2009.
- [12] Antonin Guilloux and Pierre Will. On $\mathrm{SL}(3, \mathbb{C})$ -representations of the Whitehead link group. *Geom. Dedicata*, 202:81–101, 2019.
- [13] K. Guruprasad, J. Huebschmann, L. Jeffrey, and A. Weinstein. Group systems, groupoids, and moduli spaces of parabolic bundles. *Duke Math. J.*, 89(2):377–412, 1997.
- [14] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [15] Michael Heusener, Vicente Muñoz, and Joan Porti. The $\mathrm{SL}(3, \mathbb{C})$ -character variety of the figure eight knot. *Illinois J. Math.*, 60(1):55–98, 2016.
- [16] Michael Heusener and Joan Porti. Representations of knot groups into $\mathrm{SL}_n(\mathbb{C})$ and twisted Alexander polynomials. *Pacific J. Math.*, 277(2):313–354, 2015.
- [17] Michael Heusener and Joan Porti. Holomorphic volume forms on representation varieties of surfaces with boundary. *Annales Henri Lebesgue*, 3:341–380, 2020.
- [18] Michael Heusener, Joan Porti, and Eva Suárez Peiró. Deformations of reducible representations of 3-manifold groups into $\mathrm{SL}_2(\mathbb{C})$. *J. Reine Angew. Math.*, 530:191–227, 2001.
- [19] Dennis Johnson and John J. Millson. Deformation spaces associated to compact hyperbolic manifolds. In *Discrete groups in geometry and analysis (New Haven, Conn., 1984)*, volume 67 of *Progr. Math.*, pages 48–106. Birkhäuser Boston, Boston, MA, 1987.
- [20] Michael Kapovich. *Hyperbolic manifolds and discrete groups*, volume 183 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 2001.
- [21] Sean Lawton. Generators, relations and symmetries in pairs of 3×3 unimodular matrices. *J. Algebra*, 313(2):782–801, 2007.
- [22] Sean Lawton. Poisson geometry of $\mathrm{SL}(3, \mathbb{C})$ -character varieties relative to a surface with boundary. *Trans. Amer. Math. Soc.*, 361(5):2397–2429, 2009.

- [23] D. D. Long and M. B. Thistlethwaite. The dimension of the Hitchin component for triangle groups. *Geom. Dedicata*, 200:363–370, 2019.
- [24] Bruno Martelli and Carlo Petronio. Dehn filling of the “magic” 3-manifold. *Comm. Anal. Geom.*, 14(5):969–1026, 2006.
- [25] Pere Menal-Ferrer and Joan Porti. Local coordinates for $\mathrm{SL}(n, \mathbf{C})$ -character varieties of finite-volume hyperbolic 3-manifolds. *Ann. Math. Blaise Pascal*, 19(1):107–122, 2012.
- [26] Pere Menal-Ferrer and Joan Porti. Twisted cohomology for hyperbolic three manifolds. *Osaka J. Math.*, 49(3):741–769, 2012.
- [27] Joan Porti. Torsion de Reidemeister pour les variétés hyperboliques. *Mem. Amer. Math. Soc.*, 128(612):x+139, 1997.
- [28] Joan Porti and Stephan Tillmann. Projective structures on a hyperbolic 3-orbifold. arXiv 2004.06242, 2020.
- [29] R. W. Richardson. Commuting varieties of semisimple Lie algebras and algebraic groups. *Compositio Math.*, 38(3):311–327, 1979.
- [30] Robert Steinberg. Regular elements of semisimple algebraic groups. *Inst. Hautes Études Sci. Publ. Math.*, (25):49–80, 1965.
- [31] William P. Thurston. The Geometry and Topology of Three-Manifolds. (Electronic version from 2002 available at <http://www.msri.org/publications/books/gt3m/>), 1980.
- [32] André Weil. Remarks on the cohomology of groups. *Ann. of Math. (2)*, 80:149–157, 1964.

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 Cerdanyola del Vallès, AND
 BARCELONA GRADUATE SCHOOL OF MATHEMATICS (BGSMATH)
 porti@mat.uab.cat