



## Research article

## Prefrontal reactivity to TMS perturbation as a toy model of mental health outcomes during the COVID-19 pandemic



Ruben Perellón-Alfonso<sup>a,b</sup>, María Redondo-Camós<sup>c,d,e</sup>, Kilian Abellaneda-Pérez<sup>a,b,c</sup>, Gabriele Cattaneo<sup>c,d,e</sup>, Selma Delgado-Gallén<sup>c,d,e</sup>, Goretti España-Irla<sup>c,d,e</sup>, Javier Solana Sánchez<sup>c,d,e</sup>, José M. Tormos<sup>c,d,e</sup>, Alvaro Pascual-Leone<sup>c,f,g,\*\*,1</sup>, David Bartrés-Faz<sup>a,b,c,\*</sup>

<sup>a</sup> Faculty of Medicine and Health Sciences, and Institute of Neurosciences, University of Barcelona, Barcelona, Spain

<sup>b</sup> Institute of Biomedical Research August Pi i Sunyer (IDIBAPS), Barcelona, Spain

<sup>c</sup> Institut Guttmann, Institut Universitari de Neurorehabilitació adscrit a la UAB, Badalona, Barcelona, Spain

<sup>d</sup> Universitat Autònoma de Barcelona, Bellaterra (Cerdanyola del Vallès), Spain

<sup>e</sup> Fundació Institut d'Investigació en Ciències de la Salut Germans Trias i Pujol, Badalona, Barcelona, Spain

<sup>f</sup> Hinda and Arthur Marcus Institute for Aging Research and Deanna and Sidney Wolk Center for Memory Health, Hebrew Senior Life, Boston, MA, USA

<sup>g</sup> Department of Neurology, Harvard Medical School; Boston, MA, USA

## ARTICLE INFO

## Keywords:

TMS-EEG  
Mental health  
Resilience  
COVID-19

## ABSTRACT

Psychosocial hardships associated with the COVID-19 pandemic led many individuals to suffer adverse mental health consequences, however, others show no negative effects. We hypothesized that the electroencephalographic (EEG) response to transcranial magnetic stimulation (TMS) could serve as a toy-model of an individual's capacity to resist psychological stress, in this case linked to the COVID-19 pandemic. We analyzed data from 74 participants who underwent mental health monitoring and concurrent electroencephalography with transcranial magnetic stimulation of the left dorsolateral prefrontal cortex (L-DLPFC) and left inferior parietal lobule (L-IPL). Within the following 19 months, mental health was reassessed at three timepoints during lock-down confinement and different phases of de-escalation in Spain. Compared with participants who remained stable, those who experienced increased mental distress showed, months earlier, significantly larger late EEG responses locally after L-DLPFC stimulation (but not globally nor after L-IPL stimulation). This response, together with years of formal education, was significantly predictive of mental health status during the pandemic. These findings reveal that the effect of TMS perturbation offers a predictive toy model of psychosocial stress response, as exemplified by the COVID-19 pandemic.

## 1. Introduction

The stressors associated with the coronavirus disease (COVID-19) pandemic, as well as the restrictions imposed to contain the spread of the virus, are expected to increase the global burden on mental health (Pfefferbaum and North, 2020; Torales et al., 2020). The World Health Organization has acknowledged this fact (Giacalone et al., 2020) and highlighted the importance of integrating mental health into the preparedness and response plans to public health emergencies (WHO, 2021). Some studies estimate a 25% increase in the general prevalence of

depression and anxiety symptoms (Bueno-Notivol et al., 2021). However, whereas some individuals' mental wellbeing will be negatively impacted, others - on account of their 'resilience' - will not be affected, or even thrive in the face of adversity (Pascual-Leone and Bartres-Faz, 2021).

The concept of resilience is highly heterogeneous with various meanings across different fields of study (Pascual-Leone and Bartres-Faz, 2021). Here we use resilience to refer to the processes that enable an individual to resist the development of illness, mental health problems or distress when confronted with stressful events or trauma (Moore et al., 2020; Russo et al., 2012). Conversely, psychological vulnerability is

\* Corresponding author.

\*\* Corresponding author.

E-mail addresses: [apleone@hsl.harvard.edu](mailto:apleone@hsl.harvard.edu) (A. Pascual-Leone), [dbartres@ub.edu](mailto:dbartres@ub.edu) (D. Bartrés-Faz).

<sup>1</sup> Drs. Alvaro Pascual-Leone and David Bartrés-Faz contributed equally to this manuscript.

reduced ability to cope with stressors, which would constitute a risk factor for developing psychopathology (Wright et al., 2013). Rather than a dichotomy, resilience and vulnerability can be understood as opposite ends of a continuum, which likely reflects the dynamic product of a complex interplay of individual and environmental factors (Cathomas et al., 2019; Rutter, 2012; Tost et al., 2015), including genetic and demographic characteristics, socio-economic status, developmental circumstances, access to health care, living conditions, adherence to certain lifestyle factors (e.g. cognitive, physical, nutritional and sleep habits), engagement in emotion-regulation practices such as meditation, social relations and support (particularly early in life), and years of education (Campbell-Sills et al., 2009; Di Marco et al., 2014; Frankish and Horton, 2017; Gelfo et al., 2018; Livingston et al., 2017).

Evidence from animal models and human neuroimaging studies have identified several brain regions and networks that likely play a role in the continuum of resilience and vulnerability. Converging evidence points at the crucial roles of anterior cingulate and insular cortices and their connections within the salience network (Menon, 2015; Menon and Uddin, 2010), as well as limbic structures, such as the amygdala and the ventral striatum (Holz et al., 2020). Additionally, the prefrontal cortex has been identified as an important structure (Maier and Watkins, 2010). Specifically, prefrontal cortical volume, activation and connectivity with limbic structures, positively correlate with resilience to traumatic events (Bolsinger et al., 2018); and longitudinal studies using functional magnetic resonance imaging (fMRI) found that children who were better able to regulate amygdala's emotional response through recruitment of the frontoparietal network exhibited greater resilience to developing depressive symptoms following maltreatment (Rodman et al., 2019). Prefrontal function is also involved in the pathophysiology and treatment of psychiatric conditions, most prominently, major depressive disorder (Hare and Duman, 2020) and schizophrenia (Selemon and Zecevic, 2015). Finally, the prefrontal cortex appears to not only play a central role in psychological resilience and psychiatric pathophysiology, but has been also proposed as a hub region for cognitive resilience in normal aging (Franzmeier et al., 2017b) as well as to brain atrophy and pathology associated with Alzheimer's disease (Franzmeier et al., 2018; Neitzel et al., 2019). Therefore, in the present study we focused on the prefrontal cortex to investigate the neural substrate of resilience to mental health impact of the COVID-19 pandemic and, argued that single pulse transcranial magnetic stimulation (TMS) in combination with electroencephalography (EEG) could be used in human experimental designs akin to the intervention-based animal studies of the neural substrate of resilience.

Animal studies that employ stimuli that can be precisely quantified and controlled, for example using tail-shock stress paradigms (Seligman et al., 1975) in which a mild electric shock is given to the tail of rats or mice, illustrate the power of such interventional experimental approaches to gain mechanistic insights into the substrate of resilience. Similar approaches in human research combining non-invasive brain stimulation with neuroimaging are possible. For example, Shafi and colleagues (Shafi et al., 2015) have shown that brain responses to TMS allow identifying abnormal cortical activity patterns before the manifestation of clinical symptoms in some forms of epilepsy. More recently, Abellana-Pérez and colleagues (Abellana-Pérez et al., 2019) have shown that the default mode network's response profile to intermittent theta burst stimulation of the inferior parietal lobule can be used to predict cognitive decline or maintenance after a three-year follow-up in an aging population, well over and above of what baseline neuroimaging data alone could predict. Furthermore, recent methodological advances (Ozdemir et al., 2020, 2021b), have revealed that single pulse TMS can be used concurrently with EEG to produce highly specific and reliable cortical response profiles.

We propose that TMS-EEG can be a 'toy model' of the impact of a perturbation onto an individual brain and provide a quantitative observation of the effect of the controlled external perturbation on brain dynamics that can be used to test specific predictions about a complex

system. In theoretical physics 'toy-models' refer to simple models which nevertheless provide a quantitative explanation and reliable prediction of a given phenomenon (Georgescu, 2012; Marzuoli, 2008). Specifically, as illustrated in Figure 1, here we use the EEG response to TMS as a 'toy model' predictive of the eventual (months later) impact of the COVID pandemic and confinement on mental health. We hypothesized that individual differences in the electrophysiological cortical response to single pulse TMS brain perturbation of the left dorsolateral prefrontal cortex, compared to another control cortical target (i.e., inferior parietal lobule), and recorded using EEG, would be predictive of psychological distress outcomes during the COVID-19 pandemic and confinement. Our findings contribute to the understanding of biological brain mechanisms of resilience processes and identify a potential target and novel strategy to promote individual resilience.


## 2. Results

### 2.1. The dynamics of the EEG response to TMS perturbation differentiate individuals eventually found to have a 'negative' impact on mental health status, from those who remained 'stable', during the COVID-19 pandemic

Assessments of mental health using the four-item patient health questionnaire (PHQ-4), an ultra-brief depression and anxiety screening self-report questionnaire, were obtained prior to the COVID-19 pandemic and up to three additional times during the pandemic. If during all timepoints across the confinement, the PHQ-4 score was lower or equal than before the pandemic outbreak, subjects were classified as 'stable' ( $n = 32$ ). Conversely, if a given subject had a higher score at any timepoint during the pandemic, they were classified as having a 'negative' impact ( $n = 32$ ). Because not all participants completed stimulation at both target locations, the subgroups used in this analysis were actually smaller for each stimulation target (for L-DLPFC, 23 stable and 25 negative; for L-IPL, 22 stable and 23 negative). To make sure that the mental health impact of the pandemic was related to the levels of stress perceived during the outbreak, we tested for correlation between the average score of the three pandemic PHQ-4 timepoints, and the scores of the 14-item perceived stress scale (Cohen et al., 1983), which was also completed by participants during the pandemic, and found a strong positive correlation ( $R_s = .69$ ;  $p < .001$ ), indicating that subjects experiencing more stress during the pandemic also had a larger mental health impact. Overall, participants had a low to moderate level of perceived stress during the pandemic ( $Mdn = 14$ ; range from 2 to 32). Additionally, to rule out that the groups significantly differed demographically, in resting motor threshold or pre-COVID PHQ-4 score, a multivariate ANOVA was used to compare both groups regarding four independent variables (age, years of education, pre-COVID PHQ4 score and resting motor threshold; see supplementary table S1 for descriptive statistics), which showed that the groups did not differ in any of these variables ( $F(4, 63) = 3.869$ ,  $p = 0.130$ ; Wilk's lambda = 0.888, partial  $\eta^2 = 7.111$ ).

Point-by-point non-parametric permutation testing (1000 permutations) with cluster correction for multiple comparisons (Cohen, 2014) on the TMS-EEG evoked time-series, revealed a single broad cluster (i.e., 202–269 ms post-stimulus) surviving correction for multiple comparisons, only during stimulation of the left dorsolateral prefrontal cortex (L-DLPFC) (Figure 2, A). Inspection of the topographical distributions in source space for the surviving cluster, confirms that individuals whose mental health was negatively impacted had a qualitatively stronger frontal activation than those who remained stable (Figure 2, B). There were no significant clusters revealed after analysis of the responses to the left inferior parietal lobule (L-IPL) control target (Figure 2, C and D), nor for the distributed responses to stimulation on either target (Figure S2).

Finally, given that only a subsample (i.e., 50% of participants) completed stimulation of both targets, we additionally run permutation testing of the local TMS-EEG evoked time-series after DLPFC stimulation, but including only the participants that had undergone stimulation of both targets. The results on this subsample of participants are comparable



**Figure 1.** Schematic Illustration of the toy-model approach in our study design. MRI-guided single pulse TMS perturbation is used to experimentally model as the stressor, captured by evoked EEG reactivity, and measured at local and distributed (i.e., global) levels. We examined whether this ‘toy-model’ can predict the eventual impact of the COVID-19 on mental health assessed months later. We further hypothesized the stressor would be moderated by demographic and individual factors such as years of formal education. Modified from Pascual-Leone and Bartrés-Faz (Pascual-Leone and Bartrés-Faz, 2021).

to the full sample (Figure S3), since the broad cluster surviving multiple comparisons correction remains in the same location, albeit slightly shorter in duration (i.e., from 200 to 253 ms after the TMS pulse).

## 2.2. Local EEG response to TMS perturbation of the left dorsolateral prefrontal cortex predicts mental health during the pandemic's lockdown confinement

Without dividing the sample into two groups, a multiple linear regression model was fit to determine the potential of TMS evoked EEG perturbation of the L-DLPFC to predict mental health outcomes after the COVID-19 pandemic outbreak and the strict lock-down confinement imposed to curb community transmission of the virus. The model's response variable was the mean of the total scores for the three PHQ-4 questionnaires, which were completed by participants during the lockdown confinement. Candidate predictors were the local and global (i.e., distributed) brain EEG reactivity to the TMS pulse — recorded before the pandemic outbreak —, as well as their interaction with the stimulation target definition method (i.e., functional or anatomical). Additionally, we included age, gender, and years of formal education as predictors, because these are demographic and individual factors partially predictive of resilience to stress (Campbell-Sills et al., 2009). Finally, we included as a predictor the number of months before the pandemic since each subject underwent TMS-EEG. This was included to control for the possibility that the amount of time passed from stimulation to pandemic would have an impact in the prediction.


The full linear regression model for the L-DLPFC stimulation target significantly predicted mental health during the pandemic ( $F(8,47) = 3.1, p = .007, R^2_{adj} = .234$ ), and revealed as significant predictors the local brain reactivity to TMS ( $t = 3.27, p < .002$ ) and years of formal education ( $t = -2.86, p = .006$ ). See supplementary Table S2 (model ‘Full DLPFC’) for detailed results. Both predictors were independent from

each other, as revealed by the lack of correlation between them ( $R^2 = .187, p = .167$ ). Subsequently, we tested a reduced model ( $F(2,53) = 10.5, p < .001, R^2_{adj} = .257$ ) retaining only as predictors local brain reactivity ( $t = 3.66, p < .001$ ) and education ( $t = -3.40, p = .001$ ). See supplementary table S2 (model ‘Reduced DLPFC’) for detailed results. Likelihood ratio test comparing the two models showed that the full model did not provide a better fit than the reduced one ( $\chi^2(6) = 4.98, p = .546$ ). The lower Akaike and Bayesian information criteria (AIC and BIC, respectively) values for the reduced model further suggest a better and more parsimonious fit ( $AIC_{full} = 79.83, AIC_{reduced} = 72.82; BIC_{full} = 98.06, BIC_{reduced} = 78.89$ ). Figure 3 illustrates the linear relationship between the significant predictors and the response variable in the reduced model. Analysis of variance of the reduced model revealed that local L-DLPFC reactivity explained 12.79% of total variance in mental health during the pandemic, while education explained 15.64%.


To confirm that our findings were specifically associated with prefrontal reactivity, we fitted a model replacing the predictors for local and global EEG reactivity with those measured when stimulating the L-IPL. The regression model for this control stimulation target did not significantly predict mental health during the pandemic ( $F(8,46) = 0.4, p = .915, R^2_{adj} = -.097$ ). See supplementary table S2 (model ‘Full IPL’) for detailed results.

To demonstrate the specificity of the stimulation itself, a model was fitted where we added the local baseline pre-TMS activity as an additional predictor to the reduced L-DLPFC model. The resulting model, while still significant ( $F(3,47) = 7.09, p < .001, R^2_{adj} = .25$ ), revealed that baseline pre-TMS EEG activity did not significantly contribute to predict mental health during the pandemic ( $t = .67, p = .507$ ). See supplementary table S2 (model ‘Reduced + Baseline’) for detailed results.

Finally, to ensure that the results are consistent, even when only considering the subsample of participants who completed stimulation of both targets, a model was fitted with only this subsample and the local



**Figure 2.** Results of permutation testing of the difference between stable and negatively impacted individuals on the TMS evoked EEG time-series. A) shows the significant differences at the local EEG time-series during L-DLPFC stimulation in grey vertical bands, while the vertical green band highlights the cluster surviving correction for multiple comparisons. B) shows the topographical distribution in source space of the response to L-DLPFC stimulation during the green shaded time-window in A, for both groups and their difference. C) depicts the results of permutation testing for the control stimulation of the L-IPL. D) shows the topographical distribution for the same time window as B, for both groups and their difference. Red and blue contours along the plot lines in A and C depict the standard error of the mean. DLPFC, dorsolateral prefrontal cortex; IPL, inferior parietal lobule.



**Figure 3.** Scatter plots illustrating the linear relationship between the predictors and the response variable in the reduced model. Black line in each plot depicts the least squares regression line; shaded grey contours depict 95% confidence intervals. The response variable depicted here in the y-axis is Box-Cox transformed ( $\lambda = -0.078$ ). PHQ-4, four item patient health questionnaire; DLPFC, dorsolateral prefrontal cortex; AUC, area under the curve.

response to IPL stimulation added as a predictor to the reduced model. The resulting model was still significant ( $F(3,33) = 3.25; p = .034$ ) and revealed that, while local DLPFC reactivity ( $t = 2.44, p = .021$ ) and education ( $t = -2.46, p = .019$ ) were significant, local IPL reactivity was not ( $t = -1.03, p = .311$ ). See supplementary table S2 (model “Reduced Subsample + IPL”) for detailed results.

### 3. Discussion

We tested the EEG brain reactivity to TMS perturbation as a toy model of mental health outcomes in the face of the COVID-19 pandemic and lockdown confinement. The results show that the local response to TMS perturbation of the left DLPFC — measured months *before* the pandemic outbreak — offers a predictive marker of the future mental health impact of the pandemic and confinement. These results serve as a proof of concept that understanding the TMS pulse as an external transitory insult allows quantification of the brain responses and identifies critical and specific substrates of susceptibility to more complex stressors. At the core of this ‘toy-model’ is the assumption that single TMS stimuli can be understood as transitory perturbations or insults in themselves. This is supported by evidence showing that the stimuli interfere with ongoing brain activity by suddenly injecting an amount of current into the neural circuitry that disrupts ongoing brain activity, resulting in phase resetting and TMS-evoked perturbation of the ongoing EEG (Rocchi et al., 2018). Indeed, failure to suppress this perturbation can lead — in the presence of pathological conditions such as stroke or epilepsy — to a cascading synchronization of neuronal activity that in turn might lead to a seizure (Kimiskidis, 2019; Kimiskidis et al., 2017). Thus, we can interpret the present results as showing that in the presence of such a brain state disruption, a more resilient brain is better able to tolerate the perturbation. A link between the ability of the brain to withstand a targeted attack and resilience, has also been proposed by Santarnecchi and colleagues employing in-silico models (Santarnecchi et al., 2015), but the present study is the first to offer direct experimental support on a topic of substantial timely relevance.

Our findings are specific to TMS prefrontal perturbation, because neither the response to L-IPL stimulation, nor the pre-TMS baseline EEG activity held significant predictive value. Moreover, we show that the findings are restricted locally to the stimulated area, because the distributed measure of response could not differentiate stable from negatively impacted individuals and did not yield significant predictive value. Nevertheless, years of formal education was also found to be predictive of mental health during the pandemic, which is unsurprising given the well-known epidemiological-level notion that individuals with higher socioeconomic status (encompassing, among others, educational attainment and income) have lower odds of being depressed (Lorant et al., 2003), and that education might be a protective factor both against cognitive as well as emotional vulnerability, by boosting a higher efficiency on top-down emotional regulatory processes (Huang et al., 2019). In this context, and while education and TMS-EEG reactivity were independent from one another in this analysis, it is still plausible that both reflect prefrontal function, and is therefore, their joint inclusion that best predicted mental health status during the pandemic.

Dividing the sample into participants who remained stable and those who had a negative change in mental health during the pandemic, allowed us to directly compare the dynamics of the EEG response to the TMS perturbation, revealing that the most discriminative time segment after TMS perturbation of the local L-DLPFC is the late TMS evoked response between 202 ms and 269 ms post-TMS. Interestingly, this occurs in the vicinity of the commonly found P180 EEG evoked component in response to single TMS pulses of the primary motor cortex (Lioumis et al., 2009). This component has been found to significantly decrease after application of voltage-gated sodium channel blockers, such as lamotrigine and carbamazepine (Darmani and Ziemann, 2019; Premoli et al., 2017), indicating that this component reflects cortical excitability. Therefore, we propose that higher amplitudes found in vulnerable

individuals may be reflective of cortical hyperexcitability. Furthermore, the amplitude of late TMS-EEG responses ( $\geq 180$  ms) is related to GABA-B mediated inhibition, as it is significantly reduced after long interval intracortical inhibition (de Goede et al., 2020). Therefore, the increased amplitude in late EEG responses found in individuals that had a negative impact, when compared to the ones who remained stable, might reflect a relatively lower intracortical inhibitory capacity and point to differential levels of activation of parvo-albumin positive cells and integrity of peri-neural nets — main substrates of intracortical excitability-inhibition balance (Favuzzi et al., 2017; Xue et al., 2014). This would predict that conditions that alter and disrupt parvo-albumin positive cells and integrity of peri-neural nets, such as status post traumatic brain injury, early stages of Alzheimer’s disease, or schizophrenia, would be associated with a loss of resilience and increased vulnerability to stressors. Epidemiologic data appear to support such notions (Buckley et al., 2009; Ehrenberg et al., 2018; Hammond et al., 2019).

Our results are novel and relevant in advancing our understanding of the neural mechanisms of resilience. However, this study has some limitations. We had to conduct the regression analysis on each stimulation target separately due to missing data, which may have hindered statistical power. The regression analysis results would benefit from further validation on a separate independent sample to be able to make reliable predictions of mental health outcomes based on the response to TMS perturbation. Moreover, due to the limited number of participants relative to the number of mental health monitoring timepoints during the pandemic, we were unable to account for all the mental health trajectories that have been described in the literature, such as people who have a negative impact but eventually recover (e.g., Gambin et al., 2021). Instead, in the permutation analysis, we focused on dichotomizing the two main trajectories (i.e., resilient vs vulnerable). Future studies should investigate the neurophysiological signature in response to stimulation of subjects who recover in the medium or long term, and that here were all classified as vulnerable. Finally, the changes in mental health observed during the pandemic were small overall, with most participants not surpassing clinical screening thresholds for depression and anxiety, therefore, a sample with a broader range of mental health impact could provide a clearer picture of the neurophysiological determinants of such impact. However, despite the narrow range of mental health changes, we are still able to show that TMS-EEG can detect a neurophysiological signature underlying the future differential impact of the pandemic on mental health.

The presented results are not only relevant as a proof of concept for using intervention-based designs in neuroimaging investigations of the neural basis of resilience, but also add to the existing evidence of a primary role of the left prefrontal cortex in resilience processes (Bolsinger et al., 2018; Dedovic et al., 2009; Franzmeier et al., 2017a, 2017b, 2017c, 2018, 2017b; Holz et al., 2020; Neitzel et al., 2019; Rodman et al., 2019; Stern et al., 2018). This, in turn, singles out the prefrontal cortex as a promising target for interventions aiming to promote positive outcomes after disrupting events such as the pandemic and associated social restrictions, including the potentially transformative possibility of using non-invasive stimulation to promote brain resilience by modulating prefrontal brain activity. Several of our results and other lines of evidence support such potentially transformative therapeutic intervention: (1) the known protective role of higher prefrontal function to the deleterious mental health effects of stress and trauma (Bolsinger et al., 2018; Rodman et al., 2019); (2) the link we have demonstrated between the L-DLPFC response to a brain perturbation and the mental health outcomes when facing the stressors associated with the pandemic; (3) our finding of exaggerated response to TMS perturbation of the L-DLPFC in individuals that would be negatively impacted by pandemic related stress; and (4) the established ability of non-invasive stimulation techniques to induce long lasting brain plastic changes (Huang et al., 2017). Finally, the identified electrophysiological dynamics in the local DLPFC response to TMS perturbation is a potential neurophysiological marker that might be useful in a preventive precision medicine framework, when

assessing the potential risk of deleterious mental health impacts for a given individual, when exposed to future stressful events such as a new pandemic.

## 4. Methods

### 4.1. Study design

In the present study we analyzed existing data from participants of the longitudinal study ‘Barcelona Brain Health Initiative’, BBHI for short (Cattaneo et al., 2018). In mid-March 2020, during the COVID-19 epidemic, the BBHI launched a longitudinal substudy to investigate the mental and brain health impact of societal and personal restrictions imposed by the pandemic (Bartrés-Faz et al., 2021; Pascual-Leone et al., 2021). For the present report, we selected those BBHI participants who had undergone concurrent TMS-EEG between July 2018 and February 2020, before the COVID-19 pandemic outbreak, as well as mental health monitoring before and during the lockdown using the Patient Health Questionnaire for Depression and Anxiety (PHQ-4), a standardized ultra-brief tool for detecting both anxiety and depressive disorders (Kroenke et al., 2009). The scale was administered at four different timepoints; one between November 2018 and January 2020, hence before a mandatory lockdown that was issued by the Spanish Government on March 14th 2020, and another three timepoints during the pandemic, spanning a total of 3 months during the strictest home-confinement and initial phases of de-escalation (Figure 4).


The sample included 74 healthy adults (45 male) ranging from 42 to 66 years ( $M = 55.07$ ;  $SD = 7.1$ ), with a range of years of formal education from 8 to 28 years ( $M = 18.01$ ;  $SD = 3.85$ ). Consistent with the BBHI general inclusion criteria, none of these individuals reported a medical diagnosis of any major neuropsychiatric disorder (including mood and anxiety disorders) at study entrance and had normal cognitive function as assessed by comprehensive neuropsychological testing (Cattaneo et al., 2018). All participants gave written informed consent, and the local ethics committee (Comité d'Ètica i Investigació Clínica de la Unió Catalana d'Hospitals) approved the protocols here described and conformed to the Declaration of Helsinki for research involving human subjects.

The objective of this analysis was to evaluate the potential of using the brain response to TMS perturbation – quantified by EEG – as a toy model of the mental health impact of a complex stressor, namely, the COVID-19 pandemic and confinement, the impact of which was quantified with the PHQ-4 questionnaires. Given the known involvement of the prefrontal cortex in various forms of resilience, we hypothesized that the

EEG response to left dorsolateral prefrontal stimulation would be predictive of mental health during the pandemic. Stimulation on the left inferior parietal lobule was included in the analysis as a control stimulation condition.

### 4.2. Neuronavigated TMS-EEG

Transcranial magnetic stimulation was delivered over the left dorsolateral prefrontal cortex (L-DLPFC) and the left inferior parietal lobule (L-IPL). Stimulation was guided by a BrainSight neuronavigation system (RogueResearch, Inc., Canada). Targets were determined for each individual based on either anatomy or the cortical parcellation by Yeo and colleagues (Thomas Yeo et al., 2011). See supplementary materials for MRI acquisition parameters and target determination procedures. Stimulation intensity was 120% of resting motor threshold, determined as the minimum intensity required to elicit motor evoked potentials in the first dorsal interosseous muscle of the relaxed right hand, of at least 50 $\mu$ V peak-to-peak, in at least five out of ten trials (Rossini et al., 2015). For each target, 120 single biphasic pulses were delivered through an MCF-B65 butterfly coil, using a MagPro X100 stimulator (Magventure, Inc., Denmark), with a random inter-pulse interval between four and six seconds. The order of targets was randomized for each participant. In order to attenuate auditory evoked responses induced by the TMS coil click, participants listened to white-noise through earplug-earbuds at their maximum comfortable volume. Stimulation was performed concurrently with EEG using a TMS compatible ActiChamp 64-channel amplifier system, coupled with an ActiCap Slim with active electrodes (BrainProducts, GmbH., Germany). While the use of active electrodes is relatively novel in the context of TMS-EEG, recent research has successfully used them to evaluate TMS evoked brain reactivity (to cite some, Gamboa Arana et al., 2020; Ozdemir et al., 2021b, 2021a, 2020; Redondo-Camós et al., 2022; Rocchi et al., 2021). Moreover, it has been recently shown that the TMS evoked potential waveforms are reliable and comparable to those obtained with passive electrodes (Mancuso et al., 2021), provided that interelectrode impedance is kept low. Therefore, we monitored electrode impedance to make sure it was kept under 5k $\Omega$  for all electrodes and throughout the experiment. EEG data was recorded DC to 500Hz and digitized at a 1KHz sampling rate. While 76% of participants completed stimulation of the L-DLPFC and 74% completed stimulation of the L-IPL, only 50% of participants completed stimulation on both targets. For this reason, statistical analysis was conducted separately for each stimulation target.



**Figure 4.** Timeline of relevant events for the study cohort. According to the Spanish Government state of alarm dictation orders and subsequent de-escalation (<https://www.boe.es/eli/es/rd/2020/03/14/463/con>). 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> indicate the three PHQ-4 based mental health monitoring timepoints during the pandemic. PHQ-4, four item patient health questionnaire; COVID-19, coronavirus disease 2019; TMS-EEG, transcranial magnetic stimulation with concurrent electroencephalography.

#### 4.3. Mental health assessment

Mental health was measured using the PHQ-4, an ultra-brief four item depression and anxiety screening self-report questionnaire, that consists of a 2-item depression scale (PHQ-2) and a 2-item anxiety scale (GAD-2). Each subitem scores in the range of 0–6, with combined range from 0 to 12. On each subscale a score of 3 or greater is considered positive for screening purposes. The test was administered a total of four times in an online format, once before the pandemic and at three timepoints during the confinement and de-escalation. All participants in this analysis completed the pre-pandemic questionnaire and most completed the three additional ones during the pandemic (69%), however, few participants completed only two (23%) or one (8%) of them. For the purposes of quantifying mental health status during the pandemic in the regression models, we used the mean of total scores from the completed questionnaires during the pandemic.

#### 4.4. EEG data preprocessing and analysis

All EEG data was preprocessed using functions from the EEGLAB toolbox (Delorme and Makeig, 2004) and the TESA plugin (Mutanen et al., 2020; Rogasch et al., 2017). Source reconstruction and analysis was performed using Brainstorm (Tadel et al., 2019) and custom made Matlab (The MathWorks, Inc., USA) scripts.

First, the data was segmented around the TMS pulse (−1000 to 1000 ms from the pulse) and baseline corrected (−900 ms to −100 ms from the pulse). Then the direct electrical pulse artifact (between −2 ms and 14 ms from pulse) was zero-padded. Bad channels were then identified via visual inspection and removed (range from 0 to 3;  $M = 0.49$ ,  $SD = 0.76$ ). Bad epochs were first tagged based on threshold voltage ( $>100 \mu\text{V}$ ), probability and kurtosis using the inbuilt TESA plugin functions, visual inspection ensured that the epochs were correctly tagged and that no bad epochs were missed, then they were removed from further analysis (range from 0 to 53;  $M = 20.59$ ;  $SD = 9.66$ ). A first round of fast independent component analysis (ICA) was used to reject any remains of the immediate electrical pulse artifact (range from 0 to 3;  $M = 0.67$ ;  $SD = 0.65$ ). The zero-padded pulse artifact was then linearly interpolated, and the data was re-referenced to the average of all channels. Finally, a second round of ICA was used to reject any other remaining artifacts (e.g., muscle, eye-movements, heartbeat and others), as well as the somatosensory and auditory potentials evoked by transcutaneous scalp nerve excitation and coil firing sounds, respectively (range from 21 to 23;  $M = 28.37$ ;  $SD = 2.77$ ). These are commonplace preprocessing procedures for TMS-EEG data and have been described in greater detail elsewhere (Rogasch et al., 2017).

The cleaned preprocessed data was then used for source reconstruction in Brainstorm. For each subject a forward model was estimated via the openMEEG algorithm (Kybic et al., 2005) using the default settings (i.e., 3 layers with 1922 vertices each; skull and scalp conductivities of 1 and brain conductivity of 0.0125; adaptive integration), and based on each subject's T1 and T2 weighted MRI images and digitized real electrode locations, when available (the 29 subjects for which anatomical target determination was used, had no digitized electrode locations, therefore, the standard 10-10 electrode locations were used instead). The inverse solution was estimated using the minimum norm imaging method (Salmelin and Baillet, 2009). Sources were then computed as current density maps for constrained orientations only (i.e., normal to cortex). These are commonplace source reconstruction procedures for TMS-EEG data and have been described in greater detail elsewhere (Ozdemir et al., 2020, 2021a, 2021b).

#### 4.5. Local and distributed EEG measures of the response to TMS perturbation

To quantify EEG derived brain reactivity measures to the TMS pulse we computed the following global (i.e., distributed) and local reactivity measures:

**Global response:** the global mean field amplitude (GMFA; Lehmann and Skrandies, 1980) of the TMS evoked potentials (TEPs) in sensor space, was taken as the estimated time-series of the global brain response to the TMS pulse. This measure was computed according to the following formula:

$$GMFA(t) = \sqrt{\frac{\sum_i^k (V_i(t) - V_{\text{mean}(t)})^2}{K}}$$

Where  $V_i(t)$  is the voltage at electrode  $i$  at a certain point in time,  $V_{\text{mean}(t)}$  is the mean of instantaneous TEP across electrodes, and  $K$  is the number of electrodes.

**Local response:** To extract local measures, we first defined a region of interest (ROI) of 100 vertices around each subject's stimulation target coordinate, which corresponds to a cortex surface area of approximately  $10 \text{ cm}^2$ . Then the TEP response at the targeted location in source space was taken as the local reactivity time-series. To allow group level statistics, the TEP time-series in source space of each vertex within the target ROI were rectified, averaged together, and then normalized via z-score transformation:

$$z = (TEP - \mu) / \sigma$$

Where  $\mu$  is the average of the pre-stimulus baseline (−500 ms to −3 ms) and  $\sigma$  is the standard deviation of the baseline.

For both global and local TMS response measures, the trapezoidal integration from 15 ms to 400 ms post-TMS stimulus was used in the regression analyses as an estimate of the overall response to TMS perturbation.

#### 4.6. Statistical analysis

Statistical analysis was performed in RStudio (RStudio Team, 2020) and Matlab 2020b.

To compare the global and local TMS evoked time-series for each target we conducted four permutation-based tests. In each test, we computed the difference of means for each data point within the time-series time-window of interest (from 15 ms to 400 ms after the TMS pulse). In each of the 1000 permutations, the labels for each group (resilient or vulnerable) were scrambled. The resulting p-values were adjusted for multiple comparisons using cluster correction (Cohen, 2014), whereby the size and magnitude of a given cluster of significant timepoints is considered to survive correction if the size and magnitude of the cluster is above 95% of all cluster sizes and magnitudes discovered during permutation testing. To test the consistency of the significant results on the subsample of participants that completed stimulation on both targets, an additional permutation-based test was run on this subsample. To test the correlation between perceived stress and mental health during the pandemic a Spearman rank correlation was used.

To investigate the predictive value of TMS reactivity at a local and global levels we used two multiple linear regression models, one for each stimulation target (i.e., L-DLPFC and L-IPL). The full model included as predictors the local and global TMS brain reactivity measures, age, gender, and years of education. Additionally, and to control for the possibility that the target definition method influenced the candidate reactivity measures, we included the interaction between the targeting method (2 levels: anatomical or functional) and both local and global reactivity measures, for each model. In each regression model the response variable was each subject's mean of the completed pandemic PHQ-4 scores. In the presence of significant predictors, a reduced model including only those was defined and compared against the full model. To determine the better fitting model, we used the likelihood ratio test and further confirmed the result based on the AIC and BIC values. To assert the TMS induced specificity of the findings, we fitted an additional model where the possible contribution of the pre-TMS stimulus baseline to the prediction of mental health was tested. Finally, to confirm the

consistency of the results on the subsample of participants that completed stimulation on both targets, a model was fitted adding the local IPL reactivity as a predictor and including only the subsample participants.

Lilliefors test on each model's residuals revealed that they were not normally distributed, therefore, we transformed the response variable in each model using Box-Cox transformation, resulting in normally distributed residuals, therefore, the results reported in this work correspond to the regression models with the transformed response variable. Assumptions of multicollinearity, autocorrelation and heteroscedasticity were met in each model.

## Declarations

### Author contribution statement

Ruben Perellón-Alfonso: Conceived and designed the experiments; Analyzed and interpreted the data; Contributed analysis tools; Wrote the paper.

María Redondo-Camós: Conceived and designed the experiments; Performed the experiments; Analyzed and interpreted the data.

Kilian Abellana-Pérez: Conceived and designed the experiments; Interpreted the data; Wrote the paper.

Gabriele Cattaneo, Javier Solana Sánchez: Conceived and designed the experiments; Interpreted the data.

Selma Delgado-Gallén, Goretta España-Irla: Performed the experiments.

José M. Tormos: Conceived and designed the experiments; Contributed materials and data.

Alvaro Pascual-Leone, David Bartrés-Faz: Conceived and designed the experiments; Interpreted the data; Contributed materials and data; Wrote the paper.

### Funding statement

David Bartrés-Faz and Kilian Abellana-Pérez were supported by Ministerio de Ciencia, Innovación y Universidades de España [RTI2018-095181-B-C21].

Ruben Perellón-Alfonso was supported by a fellowship from "la Caixa" Foundation [ID 100010434. Fellowship code: LCF/BQ/DI19/11730050].

Alvaro Pascual-Leone was supported by the National Institutes of Health, United States [R24AG06142 and P01AG031720].

The research leading to these results has received funding from "la Caixa" Foundation [LCF/PR/PR16/11110004].

### Data availability statement

The authors encourage interested investigators to reach out and we will honor all reasonable and scientifically motivated requests for data access and make the raw data available when required.

### Declaration of interest's statement

The authors declare the following conflict of interests: Dr. A. Pascual-Leone is listed as an inventor on several issued and pending patents on the real-time integration of noninvasive brain stimulation with electroencephalography and magnetic resonance imaging. He is co-founder of Linus Health and TI Solutions AG; serves on the scientific advisory boards for Starlab Neuroscience, Magstim Inc., Radiant Hearts, and MedRhythms, and is an Associate Editor for *Annals of Neurology*. None of the other authors reports any conflict of interest.

### Additional information

Supplementary content related to this article has been published online at <https://doi.org/10.1016/j.heliyon.2022.e10208>.

## Acknowledgements

Authors thank the Barcelona Brain Health Initiative project participants, whose invaluable contribution made this research possible. We are thankful for the contribution of Indre Pileckyte to the statistical analysis in this work, as well as for her review of the manuscript.

## References

- Abellana-Pérez, K., Vaqué-Alcázar, L., Vidal-Piñero, D., Jannati, A., Solana, E., Bargalló, N., Santarnecchi, E., Pascual-Leone, A., Bartrés-Faz, D., 2019. Age-related differences in default-mode network connectivity in response to intermittent theta-burst stimulation and its relationships with maintained cognition and brain integrity in healthy aging. *Neuroimage* 188, 794–806.
- Bartrés-Faz, D., Macià, D., Cattaneo, G., Borrás, R., Tarrero, C., Solana, J., Tormos, J.M., Pascual-Leone, A., 2021. The paradoxical effect of COVID-19 outbreak on loneliness. *BJ Psych Open* 7, 1–4.
- Bolsinger, J., Seifritz, E., Kleim, B., Manoliu, A., 2018. Neuroimaging correlates of resilience to traumatic events—a comprehensive review. *Front. Psychiatr.* 9.
- Buckley, P.F., Miller, B.J., Lehrer, D.S., Castle, D.J., 2009. Psychiatric comorbidities and schizophrenia. *Schizophr. Bull.* 35, 383–402.
- Bueno-Notivol, J., Gracia-García, P., Olaya, B., Lasheras, I., López-Antón, R., Santabàrbara, J., 2021. Prevalence of depression during the COVID-19 outbreak: a meta-analysis of community-based studies. *Int. J. Clin. Health Psychol.* 21.
- Campbell-Sills, L., Forde, D.R., Stein, M.B., 2009. Demographic and childhood environmental predictors of resilience in a community sample. *J. Psychiatr. Res.* 43, 1007–1012.
- Cathomas, F., Murrugh, J.W., Nestler, E.J., Han, M.H., Russo, S.J., 2019. Neurobiology of resilience: interface between mind and body. *Biol. Psychiatr.* 86, 410–420.
- Cattaneo, G., Bartrés-Faz, D., Morris, T.P., Sánchez, J.S., Macià, D., Tarrero, C., Tormos, J.M., Pascual-Leone, A., 2018. The Barcelona brain health initiative: a cohort study to define and promote determinants of brain health. *Front. Aging Neurosci.* 10.
- Cohen, M.X., 2014. *Analyzing Neural Time Series Data: Theory and Practice*. MIT press.
- Cohen, S., Kamarck, T., Mermelstein, R., 1983. A global measure of perceived stress. *J. Health Soc. Behav.* 24, 385–396.
- Darmani, G., Ziemann, U., 2019. Pharmacophysiology of TMS-evoked EEG potentials: a mini-review. *Brain Stimul.* 12, 829–831.
- de Goede, A.A., Cumplido-Mayoral, I., van Putten, M.J.A.M., 2020. Spatiotemporal dynamics of single and paired pulse TMS-EEG responses. *Brain Topogr.* 33, 425–437.
- Dedovic, K., D'Aguiar, C., Pruessner, J.C., 2009. What stress does to your brain: a review of neuroimaging studies. *Can. J. Psychiatr.* 54, 6–15.
- Delorme, A., Makeig, S., 2004. *Eeglab Jnm03.Pdf*. Tech. Soc 134, 9–21.
- Di Marco, L.Y., Marzo, A., Muñoz-Ruiz, M., Ikram, M.A., Kivipelto, M., Ruefenacht, D., Venneri, A., Soininen, H., Wanke, I., Ventikos, Y.A., Frangi, A.F., 2014. Modifiable lifestyle factors in dementia: a systematic review of longitudinal observational cohort studies. *J. Alzheim. Dis.* 42, 119–135.
- Ehrenberg, A.J., Suemoto, C.K., França Resende, E. de P., Petersen, C., Leite, R.E.P., Rodriguez, R.D., Ferretti-Rebustini, R.E. de L., You, M., Oh, J., Nitrini, R., Pasqualucci, C.A., Jacob-Filho, W., Kramer, J.H., Gatchel, J.R., Grinberg, L.T., 2018. Neuropathologic correlates of psychiatric symptoms in Alzheimer's disease. *J. Alzheimers. Dis.* 66, 115–126.
- Favuzzi, E., Marques-Smith, A., Deogracias, R., Winterflood, C.M., Sánchez-Aguilera, A., Mantoan, L., Maeso, P., Fernandes, C., Ewers, H., Rico, B., 2017. Activity-Dependent gating of parvalbumin interneuron function by the perineuronal net protein brevican. *Neuron* 95, 639–655.e10.
- Frankish, H., Horton, R., 2017. Prevention and management of dementia: a priority for public health. *Lancet* 390, 2614–2615.
- Franzmeier, N., Göttler, J., Grimm, T., Drzega, A., Araque-Caballero, M.A., Simon-Vermot, L., Taylor, A.N.W., Bürger, K., Catak, C., Janowitz, D., Müller, C., Duering, M., Sorg, C., Ewers, M., 2017a. Resting-state connectivity of the left frontal cortex to the default mode and dorsal attention network supports reserve in mild cognitive impairment. *Front. Aging Neurosci.* 9, 1–11.
- Franzmeier, N., Hartmann, J., Taylor, A.N.W., Araque-Caballero, M., Simon-Vermot, L., Kambeitz-Ilanovic, L., Bürger, K., Catak, C., Janowitz, D., Müller, C., Ertl-Wagner, B., Stahl, R., Dichgans, M., Duering, M., Ewers, M., 2018. The left frontal cortex supports reserve in aging by enhancing functional network efficiency Rik Ossenkoppele. *Alzheimer's Res. Ther.* 10, 1–12.
- Franzmeier, N., Hartmann, J.C., Taylor, A.N.W., Caballero, M.Á.A., Simon-Vermot, L., Buerger, K., Kambeitz-Ilanovic, L.M., Ertl-Wagner, B., Mueller, C., Catak, C., Janowitz, D., Stahl, R., Dichgans, M., Duering, M., Ewers, M., 2017b. Left frontal hub connectivity during memory performance supports reserve in aging and mild cognitive impairment. *J. Alzheim. Dis.* 59, 1381–1392.
- Gambin, M., Oleksy, T., Sękowski, M., Wnuk, A., Woźniak-Prus, M., Kmita, G. 1/4yna, Holas, P., Pisula, E., Łojek, E., Hansen, K., Gorgol, J., Kubicka, K., Huflejt-Łukasik, M., Cudo, A., Łyś, A.E., Szczepaniak, A., Bonanno, G.A., 2021. Pandemic trajectories of depressive and anxiety symptoms and their predictors: five-wave study during the COVID-19 pandemic in Poland. *Psychol. Med.* 19, 1–3.
- Gamboa Arana, O.L., Palmer, H., Dannhauer, M., Hile, C., Liu, S., Hamdan, R., Brito, A., Cabeza, R., Davis, S.W., Peterchev, A.V., Sommer, M.A., Appelbaum, L.G., 2020. Intensity- and timing-dependent modulation of motion perception with transcranial magnetic stimulation of visual cortex. *Neuropsychologia* 147, 107581.



- Gelfo, F., Mandolesi, L., Serra, L., Sorrentino, G., Caltagirone, C., 2018. The neuroprotective effects of experience on cognitive functions: evidence from animal studies on the neurobiological bases of brain reserve. *Neuroscience* 370, 218–235.
- Georgescu, I., 2012. Quantum simulation: Toy model. *Nat. Phys.* 8, 444.
- Giagalone, A., Rocco, G., Ruberti, E., 2020. Physical health and psychosocial considerations during the coronavirus disease 2019 outbreak. *Psychosomatics* 61, 851–852.
- Hammond, F.M., Corrigan, J.D., Ketchum, J.M., Malec, J.F., Dams-O'Connor, K., Hart, T., Novack, T.A., Bogner, J., Dahdah, M.N., Whiteneck, G.G., 2019. Prevalence of medical and psychiatric comorbidities following traumatic brain injury. *J. Head Trauma Rehabil.* 34, E1–E10.
- Hare, B.D., Duman, R.S., 2020. Prefrontal cortex circuits in depression and anxiety: contribution of discrete neuronal populations and target regions. *Mol. Psychiatr.* 25, 2742–2758.
- Holz, N.E., Tost, H., Meyer-Lindenberg, A., 2020. Resilience and the brain: a key role for regulatory circuits linked to social stress and support. *Mol. Psychiatr.* 25, 379–396.
- Huang, C.M., Fan, Y.T., Lee, S.H., Liu, H.L., Chen, Y.L., Lin, C., Lee, T.M.C., 2019. Cognitive reserve-mediated neural modulation of emotional control and regulation in people with late-life depression. *Soc. Cognit. Affect Neurosci.* 14, 849–860.
- Huang, Y.Z., Lu, M.K., Antal, A., Classen, J., Nitsche, M., Ziemann, U., Ridding, M., Hamada, M., Ugawa, Y., Jaberzadeh, S., Suppa, A., Paulus, W., Rothwell, J., 2017. Plasticity induced by non-invasive transcranial brain stimulation: a position paper. *Clin. Neurophysiol.* 128, 2318–2329.
- Kimiskidis, V., 2019. The dynamics of TMS-induced seizures and epileptiform discharges. *Brain Stimul.* 12, 484.
- Kimiskidis, V.K., Tsimpliridis, A., Ryvlin, P., Kalviainen, R., Koutroumanidis, M., Valentin, A., Laskaris, N., Kugiumtzis, D., 2017. TMS combined with EEG in genetic generalized epilepsy: a phase II diagnostic accuracy study. *Clin. Neurophysiol.* 128, 367–381.
- Kroenke, K., Spitzer, R.L., Williams, J.B.W., Löwe, B., 2009. An ultra-brief screening scale for anxiety and depression: the PHQ-4. *Psychosomatics* 50, 613–621.
- Kybic, J., Clerc, M., Abboud, T., Faugeres, O., Keriven, R., Papadopoulos, T., 2005. A common formalism for the integral formulations of the forward EEG problem. *IEEE Trans. Med. Imag.* 24, 12–28.
- Lehmann, D., Skrandies, W., 1980. Reference-free identification of components of checkerboard-evoked multichannel potential fields. *Electroencephalogr. Clin. Neurophysiol.* 48, 609–621.
- Lioumis, P., Kicić, D., Savolainen, P., Mäkelä, J.P., Kähkönen, S., 2009. Reproducibility of TMS-evoked EEG responses. *Hum. Brain Mapp.* 30, 1387–1396.
- Livingston, G., Sommerlad, A., Orgeta, V., Costafreda, S.G., Huntley, J., Ames, D., Ballard, C.G., Banerjee, S., Burns, A., Cohen-Mansfield, J., Cooper, C., Fox, N., Gitlin, L.N., Howard, R., Kales, H.C., Larson, E., Ritchie, K., Rockwood, K., Sampson, E.L., Samus, Q.M., Schneider, L.S., Selbæk, G., Teri, L., Mukadam, N., 2017. The lancet international commission on dementia prevention and care. *Lancet* 390, 2673–2734.
- Lorant, V., Deliége, D., Eaton, W., Robert, A., Philippot, P., Anseau, M., 2003. Socioeconomic inequalities in depression: a meta-analysis. *Am. J. Epidemiol.* 157, 98–112.
- Maier, S.F., Watkins, L.R., 2010. Role of the medial prefrontal cortex in coping and resilience. *Brain Res.* 1355, 52–60.
- Mancuso, M., Sveva, V., Cruciani, A., Brown, K., Ibáñez, J., Rawji, V., Casula, E., Premoli, L., D'Ambrosio, S., Rothwell, J., Rocchi, L., 2021. Transcranial evoked potentials can be reliably recorded with active electrodes. *Brain Sci.* 11, 1–16.
- Marzuoli, A., 2008. Toy models in physics and the reasonable effectiveness of mathematics. In: *Deduction, Computation, Experiment*. Springer, pp. 49–64.
- Menon, V., 2015. Saliency Network, Brain Mapping: an Encyclopedic Reference. Elsevier Inc.
- Menon, V., Uddin, L.Q., 2010. Saliency, switching, attention and control: a network model of insula function. *Brain Struct. Funct.* 214, 655–667.
- Moore, T.M., White, L.K., Barzilay, R., Calkins, M.E., Jones, J.D., Young, J.F., Gur, R.C., Gur, R.E., 2020. Development of a scale battery for rapid assessment of risk and resilience. *Psychiatr. Res.* 288, 112996.
- Mutanen, T.P., Biabani, M., Sarvas, J., Ilmoniemi, R.J., Rogasch, N.C., 2020. Source-based artifact-rejection techniques available in TESA, an open-source TMS-EEG toolbox. *Brain Stimul.* 13, 1349–1351.
- Neitzel, J., Franzmeier, N., Rubinski, A., Ewers, M., 2019. Left frontal connectivity attenuates the adverse effect of entorhinal tau pathology on memory. *Neurology* 93, E347–E357.
- Ozdemir, R.A., Boucher, P., Fried, P.J., Momi, D., Jannati, A., Pascual-Leone, A., Santarnecchi, E., Shafi, M.M., 2021a. Reproducibility of cortical response modulation induced by intermittent and continuous theta-burst stimulation of the human motor cortex. *Brain Stimul.* 14, 949–964.
- Ozdemir, R.A., Tadayon, E., Boucher, P., Momi, D., Karakhanyan, K.A., Fox, M.D., Halko, M.A., Pascual-Leone, A., Shafi, M.M., Santarnecchi, E., 2020. Individualized perturbation of the human connectome reveals reproducible biomarkers of network dynamics relevant to cognition. *Proc. Natl. Acad. Sci. U. S. A.* 117, 8115–8125.
- Ozdemir, R.A., Tadayon, E., Boucher, P., Sun, H., Momi, D., Ganglberger, W., Westover, M.B., Pascual-Leone, A., Santarnecchi, E., Shafi, M.M., 2021b. Cortical responses to noninvasive perturbations enable individual brain fingerprinting. *Brain Stimul.* 14.
- Pascual-Leone, A., Bartres-Faz, D., 2021. Human brain resilience: a call to action. *Ann. Neurol.* 1–14.
- Pascual-Leone, A., Cattaneo, G., Macià, D., Solana, J., Tormos, J.M., Bartrés-Faz, D., 2021. Beware of optimism bias in the context of the COVID-19 pandemic. *Ann. Neurol.* 11, 2019–2021.
- Pfefferbaum, B., North, C.S., 2020. Mental health and the covid-19 pandemic. *N. Engl. J. Med.* 383, 510–512.
- Premoli, L., Biondi, A., Carlesso, S., Rivolta, D., Richardson, M.P., 2017. Lamotrigine and leveticetam exert a similar modulation of TMS-evoked EEG potentials. *Epilepsia* 58, 42–50.
- Redondo-Camós, M., Cattaneo, G., Perellón-Alfonso, R., Alviarez-Schulze, V., Morris, T.P., Solana-Sánchez, J., España-Irla, G., Delgado-Gallén, S., Pachón-García, C., Albu, S., Zetterberg, H., Tormos, J.M., Pascual-Leone, A., Bartres-Faz, D., 2022. Local prefrontal cortex TMS-induced reactivity is related to working memory and reasoning in middle-aged adults. *Front. Psychol.* 13, 1–10.
- Rocchi, L., Di Santo, A., Brown, K., Ibáñez, J., Casula, E., Rawji, V., Di Lazzaro, V., Koch, G., Rothwell, J., 2021. Disentangling EEG responses to TMS due to cortical and peripheral activations. *Brain Stimul.* 14, 4–18.
- Rocchi, L., Ibáñez, J., Benussi, A., Hannah, R., Rawji, V., Casula, E., Rothwell, J., 2018. Variability and predictors of response to continuous theta burst stimulation: a TMS-EEG study. *Front. Neurosci.* 12, 1–11.
- Rodman, A.M., Jenness, J.L., Weissman, D.G., Pine, D.S., McLaughlin, K.A., 2019. Neurobiological markers of resilience to depression following childhood maltreatment: the role of neural circuits supporting the cognitive control of emotion. *Biol. Psychiatr.* 86, 464–473.
- Rogasch, N.C., Sullivan, C., Thomson, R.H., Rose, N.S., Bailey, N.W., Fitzgerald, P.B., Farzan, F., Hernandez-pavon, J.C., 2017. NeuroImage Analysing concurrent transcranial magnetic stimulation and electroencephalographic data: a review and introduction to the open-source TESA software. *Neuroimage* 147, 934–951.
- Rossini, P.M., Burke, D., Chen, R., Cohen, L.G., Daskalakis, Z., Di Iorio, R., Di Lazzaro, V., Ferreri, F., Fitzgerald, P.B., George, M.S., Hallett, M., Lefaucheur, J.P., Langguth, B., Matsumoto, H., Miniussi, C., Nitsche, M.A., Pascual-leone, A., Paulus, W., Rossi, S., Rothwell, J.C., Siebner, H.R., Ugawa, Y., Walsh, V., Ziemann, U., Committee, I.F.C.N., Rossini, P.M., Burke, D., Chen, R., Cohen, L.G., Daskalakis, Z., Iorio, R., Di Lazzaro, V., Di Ferreri, F., Fitzgerald, P.B., George, M.S., Hallett, M., Lefaucheur, J.P., Langguth, B., Matsumoto, H., Miniussi, C., Nitsche, M.A., Pascual-leone, A., Paulus, W., Rossi, S., Rothwell, J.C., Siebner, H.R., 2015. Clinical Neurophysiology Non-invasive electrical and magnetic stimulation of the brain, spinal cord, and peripheral nerves: basic principles and procedures for routine clinical and research application. An updated report from an. *Clin. Neurophysiol.* 126, 1071–1107.
- RStudio Team, 2020. RStudio. Integrated Development Environment for R.
- Russo, S.J., Murrough, J.W., Han, M., Charney, D.S., Nestler, E.J., 2012. Review Neurobiology of resilience. *Nat. Neurosci.* 15, 1475–1484.
- Rutter, M., 2012. Resilience as a dynamic concept. *Dev. Psychopathol.* 24, 335–344.
- Salmelin, R., Baillet, S., 2009. Electromagnetic brain imaging. *Hum. Brain Mapp.* 30, 1753–1757.
- Santarnecchi, E., Rossi, S., Rossi, A., 2015. The smarter, the stronger: intelligence level correlates with brain resilience to systematic insults. *Cortex* 64, 293–309.
- Selemon, L.D., Zecevic, N., 2015. Schizophrenia: a tale of two critical periods for prefrontal cortical development. *Transl. Psychiatry* 5.
- Seligman, M.E., Rosellini, R.A., Kozak, M.J., 1975. Learned helplessness in the rat: time course, immunization, and reversibility. *J. Comp. Physiol. Psychol.* 88, 542–547.
- Shafi, M.M., Vernet, M., Klooster, D., Chu, C.J., Boric, K., Barnard, M.E., Romatoski, K., Westover, M.B., Christodoulou, J.A., Gabrieli, J.D.E., Whitfield-Gabrieli, S., Pascual-Leone, A., Chang, B.S., 2015. Physiological consequences of abnormal connectivity in a developmental epilepsy. *Ann. Neurol.* 77, 487–503.
- Stern, Y., Arenaza-Urquijo, E.M., Bartrés-Faz, D., Belleville, S., Cantillon, M., Chetelat, G., Ewers, M., Franzmeier, N., Kempermann, G., Kremen, W.S., Okonkwo, O., Scarmeas, N., Soldan, A., Udeh-Momoh, C., Valenzuela, M., Vemuri, P., Vuoksima, E., Arenaza Urquijo, E.M., Bartrés-Faz, D., Belleville, S., Cantillon, M., Chetelat, G., Clouston, S.A.P., Estanga, A., Ewers, M., Franzmeier, N., Gold, B., Habeck, C., Jones, R., Kempermann, G., Kochhann, R., Kremen, W., Lim, Y.Y., Martínez-Lage, P., Morbelli, S., Okonkwo, O., Ossenkoppele, R., Pettigrew, C., Rosen, A.C., Scarmeas, N., Soldan, A., Song, X., Udeh-Momoh, C., Stern, Y., Valenzuela, M., Van Loenhoud, A.C., Vemuri, P., Vuoksima, E., 2018. Whitepaper: defining and investigating cognitive reserve, brain reserve, and brain maintenance. *Alzheimer's Dementia* 1–7.
- Tadel, F., Bock, E., Niso, G., Mosher, J.C., Cousineau, M., Pantazis, D., Leahy, R.M., Baillet, S., 2019. MEG/EEG group analysis with brainstorm. *Front. Neurosci.* 13, 1–21.
- Thomas Yeo, B.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D., Hollinshead, M., Roffman, J.L., Smoller, J.W., Zölle, L., Polimeni, J.R., Fisch, B., Liu, H., Buckner, R.L., 2011. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J. Neurophysiol.* 106, 1125–1165.
- Torales, J., O'Higgins, M., Castaldelli-Maia, J.M., Ventriglio, A., 2020. The outbreak of COVID-19 coronavirus and its impact on global mental health. *Int. J. Soc. Psychiatr.* 66, 317–320.
- Tost, H., Champagne, F.A., Meyer-Lindenberg, A., 2015. Environmental influence in the brain, human welfare and mental health. *Nat. Neurosci.* 18, 4121–4131.
- WHO. WHO Executive Board stresses need for improved response to mental health impact of public health emergencies [WWW Document]. URL: <https://www.who.int/news/item/11-02-2021-who-executive-board-stresses-need-for-improved-response-to-mental-health-impact-of-public-health-emergencies>. (Accessed 23 February 2021).
- Wright, M.O., Masten, A.S., Narayan, A.J., 2013. Resilience processes in development: four waves of research on positive adaptation in the context of adversity. In: Goldstein, S., Brooks, R.B. (Eds.), *Handbook of Resilience in Children*. Springer US, Boston, MA, pp. 15–37.
- Xue, M., Atallah, B.V., Scanziani, M., 2014. Equalizing excitation-inhibition ratios across visual cortical neurons. *Nature* 511, 596–600.