

Preserflo™ MicroShunt implantation combined with Ologen™ in primary and secondary glaucoma patients in a clinical setting

María Rojo-Arno, José María Martínez-de-la-Casa^{1,2}, Oscar Albis-Donado³, Giovanni Yañez-Castro, Raquel Maroto-Cejudo, Jesús Téllez^{4,5}, Ricardo Menoyo-Calatayud

Purpose: This study aims to assess the effectiveness and safety of combining the Preserflo™ MicroShunt implant (MicroShunt) with a simultaneous Ologen™ implant in patients with glaucoma. **Methods:** We conducted a retrospective study on consecutive patients with medically uncontrolled glaucoma who underwent MicroShunt + Ologen implantation as a standalone procedure or in combination with phacoemulsification (combined procedure). Success was defined as achieving an intraocular pressure (IOP) of 6–15 mmHg at 18 months post surgery, with a preoperative IOP reduction of at least 20%, and without (complete success) or with (qualified success) the need for antiglaucoma medications. The primary endpoint was the success rate. **Results:** Forty-eight eyes from 47 patients were included, with 28 eyes (58.3%) undergoing the standalone procedure and 20 eyes (41.7%) undergoing the combined procedure. Overall, there was a significant reduction in preoperative IOP from 19.7 ± 5.8 mmHg to 11.4 ± 2.6 mmHg at 18 months ($P < 0.0001$). In the standalone procedure group, preoperative IOP decreased from 21.5 ± 5.2 mmHg to 11.7 ± 2.5 mmHg ($P < 0.0001$), and in the combined procedure group, preoperative IOP decreased from 17.1 ± 5.8 mmHg to 10.9 ± 2.7 mmHg ($P = 0.0002$), with no significant difference between the two groups regarding final IOP. The mean number of antiglaucoma medications significantly decreased from 3.2 ± 1.1 to 0.3 ± 0.7 in the overall study population ($P < 0.0001$). At 18 months, 40 eyes (83.3%) were classified as successful. Regarding safety, out of the total number of eyes, two (4.2%) experienced choroidal detachment without visual impairment, two (4.2%) had transient hyphema, one (2.1%) showed reactivation of a corneal herpetic ulcer, one (2.1%) had diplopia, and one (2.1%) exhibited a shallow anterior chamber during the first week. **Conclusion:** The combination of Ologen™ and Preserflo™ MicroShunt, either alone or in conjunction with phacoemulsification, demonstrated a favorable profile in terms of IOP reduction and safety.

Key words: Glaucoma, glaucoma surgery, intraocular pressure, MIGS, Ologen, Preserflo MicroShunt

Glaucoma is a progressive and often asymptomatic group of diseases that affect the optic nerve, leading to optic-nerve cupping, loss of retinal ganglion cells, and visual field loss, which can ultimately result in blindness. The primary modifiable risk factor for glaucoma is increased intraocular pressure (IOP), making IOP reduction a key goal in glaucoma treatment, including surgery.^[1,2]

Angle-based minimally invasive glaucoma surgeries (MIGS) and bleb-forming MIGS have emerged as viable

options for managing IOP and reducing medication dependence, particularly in cases of mild-to-moderate glaucoma damage.^[3]

Different ab interno and ab externo MIGS use filtration either into Schlemm's canal or the suprachoroidal space, whereas bleb-forming surgeries use filtration to the subconjunctival space, aiming to lower IOP in a safer way as compared with conventional and time-proven surgeries such as trabeculectomy and long-tube glaucoma implants.^[4,5]

However, most existing MIGS procedures offer only moderate IOP reduction, leaving a gap in the availability of a safer and more predictable option for eyes with severe damage or complicated secondary glaucoma that necessitate lower-target IOPs to prevent further optic nerve deterioration.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

Cite this article as: Rojo-Arno M, Martínez-de-la-Casa JM, Albis-Donado O, Yañez-Castro G, Maroto-Cejudo R, Téllez J, *et al.* Preserflo™ MicroShunt implantation combined with Ologen™ in primary and secondary glaucoma patients in a clinical setting. *Indian J Ophthalmol* 2024;72:417-26.

Access this article online

Website:

<https://journals.lww.com/ijjo>

DOI:

10.4103/IJO.IJO_1502_23

Quick Response Code:



Glaucoma Department, Hospital Virgen de la Luz, Cuenca, Spain, ¹Department of Ophthalmology and ORL, Faculty of Medicine, Ophthalmology Unit, Hospital Clínico San-Carlos, Universidad Complutense de Madrid, Instituto de Investigación Sanitaria del Hospital Clínico San-Carlos (IdISSC), Madrid, Spain, ²Department of Immunology, Ophthalmology and ORL, School of Medicine, Instituto de Investigaciones Oftalmológicas Ramon Castroviejo, Universidad Complutense de Madrid, Madrid, Spain, ³Glaucoma Specialist, Private Practice, Visual Sense, Mexico City, Mexico, ⁴Department of Ophthalmology, Hospital de la Santa Creu i de Sant Pau, Autònoma University of Barcelona, Barcelona, Spain, ⁵Department of Ophthalmology, Grupo Admiración, Barcelona, Spain

Correspondence to: Dr. María Rojo-Arno, Glaucoma Department, Hospital Virgen de la Luz, Hermandad de Donantes de Sangre, 1, 16002 Cuenca, Spain. E-mail: mariarojoarno@gmail.com

Received: 08-Jun-2023

Revision: 24-Sep-2023

Accepted: 18-Oct-2023

Published: 26-Dec-2023

Especially during the COVID-19 pandemic, conventional and bleb-forming MIGS devices have gained popularity due to their reduced surgical duration and lower risk of sight-threatening complications, resulting in a reduced number of postoperative visits.^[6] In this context, the Preserflo® MicroShunt (Preserflo MicroShunt, Santen Pharmaceutical Co., Osaka, Japan) has emerged as a novel bleb-forming device. The MicroShunt is made of poly (styrene-block-isobutylene-block-styrene) or SIBS, a biocompatible material that exhibits resistance to biodegradation, biologically inert properties, softness, and flexibility, thereby minimizing inflammation and scar tissue formation.^[7] Clinical reports have demonstrated promising mid- and long-term IOP control with few self-limiting complications.^[7]

Ologen™ (Aeon Astron Europe B. V., Netherlands) is a porous matrix implant composed of collagen-glycosaminoglycan. It serves as both a spacer and a wound modulator in glaucoma surgeries.^[8] Although it has been primarily used in trabeculectomy procedures, it has also been employed in combination with other glaucoma implants for both adult and pediatric patients.^[9,10] A recent publication examined the combination of Ologen™ and 0.2 mg/mL Mitomycin C (MMC) in Preserflo™ surgery and found no significant additional benefit compared to the manufacturer's recommended 0.2 mg/mL MMC alone after a 6-month follow-up in eyes with primary open-angle glaucoma (POAG).^[11] We have hypothesized that a thinner implant would induce less tension on the filtration bleb, which would translate into a lower risk of fibrosis. In addition, a larger surface area of the implant would mean greater filtration and, therefore, lower IOP.

This study aimed to assess the impact of combining a thinner and larger version of Ologen™ with Preserflo™ after an 18-month follow-up period. To the best of our knowledge, this is the first report investigating the use of this specific Ologen™ model in different types of glaucoma.

Methods

Study design

This retrospective, single-center study was conducted on patients who underwent Preserflo™ MicroShunt implant surgery combined with simultaneous Ologen™ implant. The follow-up period for the study was 18 months.

Ethical considerations

All patients provided written informed consent for their surgery, in accordance with the requirements of our institution Hospital Virgen de la Luz de Cuenca. The study complied with local and European legislation and ensured the use of clinical data without any identifiers. Furthermore, the study adhered to the principles outlined in the Declaration of Helsinki.

Participants

The study included consecutive patients aged 18 years or older, with glaucoma, who had insufficiently medically controlled IOP, poor treatment adherence, or intolerance to topical hypotensive medication. Patients with conjunctival pathologies (except for previous glaucoma surgery) that could potentially compromise the outcomes of the procedure, as determined by the surgeon, were excluded from the study.

Medication withdrawal

Prior to surgery, patients were instructed to discontinue the use of topical and systemic ocular hypotensive medications.

Surgical technique

All surgeries were performed by a single surgeon MRA under local anesthesia by using a combination of topical and sub-Tenon's bupivacaine 0.5% and mupivacaine 2%.

The Preserflo™ MicroShunt implantation technique followed the manufacturer's recommendation, with the additional implantation of the Ologen™ model 862051. This model, chosen due to its availability at our institution, features a round shape with a diameter of 12 mm and a height of 1 mm, placed under Tenon's capsule. The intention behind using this model was to achieve a larger extension area and a lower height of bleb filtration.

The surgical procedure involved a 70–90° fornix-based peritomy in the selected quadrant, followed by the use of light bipolar cautery and careful elevation of Tenon's capsule to expose the episclera adequately. Topical MMC at concentrations of either 0.2 or 0.4 mg/mL was applied using the provided 3 LASIK shields for a duration of 2–4 min. The choice of MMC concentration and exposure time was based on the surgeon's preference and patient characteristics. To prevent excessively elevated limbus bleb and reduce the chances of bleb leakage, the shields were placed as far away from the limbus as possible.

A marker was used to measure 3 mm from the limbus, and a small 1-mm-wide shallow scleral pocket was dissected using a triangular-shaped knife. Subsequently, a 27-G needle tract was created to enter the anterior chamber (AC) just above the iris, away from the corneal endothelium. The MicroShunt device was then inserted through the needle tract until the fins of the implant fixated it to the wider part of the scleral pocket. This ensures a predictable length inside the AC and prevents rotation, allowing the bevel to remain with the opening away from the iris.^[12]

Visual confirmation of adequate flow through the device was obtained, and if necessary, flow was promoted by increasing pressure through corneal indentation or by injecting balanced saline solution (BSS) into the anterior chamber. The Ologen™ implant was inserted above the outer tip of the MicroShunt and under Tenon's capsule, taking care to prevent it from occluding the distal portion of the MicroShunt. Conjunctival closure was achieved using nylon 10-0 sutures with a running mattress technique until no aqueous leaks were observed. Postoperative medication included a combination of topical tobramycin and dexamethasone for the first 2 weeks, followed by a switch to prednisolone acetate 10 mg/mL with a gradually reducing regimen over the next 8–12 weeks.

Study groups

The study sample was divided into two different groups: the MicroShunt solo procedure group, consisting of eyes that underwent Preserflo MicroShunt + Ologen standalone surgery, and the MicroShunt + Phaco group, comprising eyes that underwent combined surgery (MicroShunt + Ologen + phacoemulsification).

Definitions

Success was defined as achieving an IOP of 6–15 mmHg, with an IOP reduction of at least 20% relative to the preoperative value, either without any hypotensive medication (complete success) or with the use of hypotensive medication (qualified success) at the month-18 visit.

Patients with an IOP of <6 mmHg for more than two consecutive visits, those requiring further glaucoma surgery, or those undergoing surgery due to complications were considered treatment failures.

Outcomes

The primary endpoint of the study was the proportion of patients classified as successful at the month-18 visit.

Secondary endpoints included the mean IOP and mean IOP reduction at the month-18 visit, the mean number of antiglaucoma medications used and changes from baseline, the identification of predictive factors associated with success, and the incidence of adverse events.

Statistical analysis

Standard statistical analysis was conducted using MedCalc® Statistical Software version 20.2.18 (MedCalc Software Ltd, Ostend, Belgium; <https://www.medcalc.org>; 2023).

Descriptive statistics were reported as mean \pm standard deviation (SD), mean \pm standard error of the mean (SE), median (interquartile range) (IQR), or number (percentage) as appropriate.

The normal distribution of data was assessed using the Shapiro–Wilk test.

Patients with more than 10% missing data were excluded from the analysis, and case-wise deletion was applied to the remaining dataset. Patients with $\leq 10\%$ missing data were not withdrawn from the study. In these cases, missing values were allocated using an algorithm of multiple imputation.^[13]

Changes in IOP and the number of antiglaucoma medications were assessed using repeated measures analysis of variance (ANOVA) or Friedman's two-way analysis test, as appropriate. Post-hoc analysis for pairwise comparisons was conducted using Scheffé's method (ANOVA) or the Conover method (Friedman).

Repeated analysis of covariance (MANCOVA) was performed to evaluate changes in IOP between the study groups. The model included "type of surgery" (MicroShunt solo or MicroShunt + Phaco) as a factor, with age, preoperative IOP, pachymetry, and previous glaucoma surgery as covariates.

To assess preoperative differences between cohorts, the Mann–Whitney test was used for continuous variables.

Survival rates of success were analyzed using Kaplan–Meier analysis for the MicroShunt-solo and MicroShunt + Phaco groups, with comparison conducted using the log-rank test.

A logistic regression model was utilized to estimate and test factors associated with success. A backward strategy was adopted, with a statistically significant cutoff of 0.05 for variable screening. Factors demonstrating an association with progression in the univariate analysis at $P \leq 0.1$ were included in the multivariate analysis.

Categorical variables were compared using the Chi-square test and Fisher's exact test, as necessary. A P value of less than 0.05 was considered statistically significant.

Results

A total of 48 eyes from 47 patients were included in the study, with 28 (58.3%) eyes undergoing a standalone procedure (MicroShunt solo) and 20 (41.7%) eyes undergoing a combined procedure (MicroShunt + Phaco).

Preoperative demographic and clinical characteristics:

In the overall study population, the mean age was 72.5 ± 10.2 years, and 32 (66.7%) patients were women. The most frequent diagnoses were POAG and chronic angle-closure glaucoma (CACG), accounting for 33 (68.7%) and 9 (18.8%) eyes, respectively.

Fourteen eyes (29.2%) had previously undergone antiglaucomatous surgery. A majority of the eyes, 38 (79.2%), received MMC at a concentration of 0.4 mg/mL.

The mean preoperative IOP was significantly higher in the MicroShunt-solo group (21.5 ± 5.2 mmHg) compared to the MicroShunt + Phaco group (17.1 ± 5.8 mmHg), with a median difference of 4.0 mmHg (95% confidence interval, 95% CI: 4.0–7.0 mmHg, $P = 0.0025$).

Eyes that had previously undergone a surgical procedure underwent a MicroShunt standalone procedure.

The main demographic and clinical characteristics of the study sample are summarized in Table 1.

Success rate

The overall success rate in the study population was 83.3% (40/48) eyes. Specifically, the success rate was 92.9% (26/28) eyes in the MicroShunt-solo group and 70.0% (14/20) eyes in the MicroShunt + Phaco group ($P = 0.0532$). Kaplan–Meier survival analysis demonstrated significantly greater survival in the MicroShunt-solo group compared to the MicroShunt + Phaco group (Mean hazard ratio: 4.56; 95% CI: 1.06–19.59; $P = 0.0414$) [Fig. 1].

Intraocular pressure

In the overall study sample, there was a significant reduction in mean IOP from 19.7 ± 5.8 mmHg to 11.4 ± 2.6 mmHg [Fig. 2a], with a mean difference of -8.3 ± 5.7 mmHg (95% CI: -10.0 to -6.7 mmHg; $P < 0.0001$, repeated measures ANOVA).

Both the MicroShunt-solo and MicroShunt + Phaco groups showed a significant decrease in mean preoperative IOP from 21.5 ± 5.2 mmHg and 17.1 ± 5.8 mmHg to 11.7 ± 2.5 mmHg and 10.9 ± 2.7 mmHg, respectively ($P < 0.0001$ for each, repeated measures ANOVA).

No significant differences in mean IOP were observed between the two study groups at any of the measured time points [Fig. 2b].

Although unadjusted IOP lowering was significantly greater in the MicroShunt-solo group compared to the MicroShunt + Phaco group [Fig. 3], after adjusting for different covariates (age, preoperative IOP, pachymetry, and previous glaucoma surgery), there were no significant differences in IOP lowering between the groups [Table 2].

Table 1: Baseline demographic and clinical characteristics of the study population

Characteristic	Total population (n=48)	MicroShunt Solo* (n=28)	MicroShunt* + Phaco (n=20)	P
Age, years				
Mean (SD)	72.5±10.2	72.5±10.4	72.4±10.3	0.9913 ^a
Median (IqR)	73.0 (63.3–81.0)	73.0 (67.0–81.0)	78.0 (63.8–81.0)	
Sex, n (%)				0.7626 ^b
Female	32 (66.7)	18 (64.3)	14 (70.0)	
Male	16 (33.3)	10 (35.7)	6 (30.0)	
Eye, n (%)				1.0000 ^b
Right	28 (58.3)	16 (57.1)	12 (60.0)	
Left	20 (41.7)	12 (42.9)	8 (40.0)	
BCVA, LogMar				0.0008 ^a
Mean (SD)	0.30±0.47	0.23±0.57	0.39±0.26	
Median (IqR)	0.15 (0.03–0.40)	0.08 (0.0–0.23)	0.40 (0.25–0.50)	
Pachymetry, µm				0.6529 ^a
Mean (SD)	544.9±40.1	551.0±36.0	536.5±44.8	
Median (IqR)	542.5 (522.5–563.5)	542.5 (522.5–563.5)	542.5 (518.5–565.0)	
Diagnosis, n (%)				0.1378 ^c
POAG	33 (68.7)	21 (75.0)	12 (60.0)	
CACG	9 (18.8)	6 (21.4)	3 (15.0)	
NTG	3 (6.2)	0 (0.0)	3 (15.0)	
PG	1 (2.1)	0 (0.0)	1 (5.0)	
Other	2 (4.2)	1 (3.6)	1 (5.0)	
Previous Glaucoma surgery, n (%)				0.0001 ^b
Yes	14 (29.2)	14 (50.0)	0 (0.0)	
No	34 (70.8)	14 (50.0)	20 (100.0)	
MMC dose, n (%)				0.7209 ^b
0.02%	10 (20.8)	5 (17.9)	5 (25.0)	
0.04%	38 (79.2)	23 (82.1)	15 (75.0)	
Initial** IOP, mmHg				0.0855 ^a
Mean (SD)	28.5±7.3	30.0±7.3	26.5±6.9	
Median (IqR)	27.5 (25.0–29.5)	28.0 (26.0–31.0)	25.5 (22.5–28.5)	
Preop IOP, mmHg				0.0025 ^a
Mean (SD)	19.7±5.8	21.5±5.2	17.1±5.8	
Median (IqR)	18.0 (16.0–22.0)	20.0 (18.0–23.5)	16.0 (13.5–20.0)	
NOHM				0.0902 ^a
Mean (SD)	3.2±1.1	3.4±1.0	2.9±1.1	
Median (IqR)	3.0 (3.0–4.0)	3.5 (3.0–4.0)	3.0 (2.0–4.0)	
NOHM, n (%)				<0.1024 ^c
0	2 (4.2)	1 (3.6)	1 (5.0)	
1	1 (2.1)	0 (0.0)	1 (5.0)	
2	7 (14.6)	3 (10.7)	4 (20.0)	
3	18 (37.5)	10 (35.7)	8 (40.0)	
4	18 (37.5)	12 (42.9)	6 (30.0)	
5	2 (4.2)	2 (7.1)	0 (0.0)	

^aMann–Whitney test. ^bFisher's exact test. ^cChi-squared test for trend. *Preserflo™ MicroShunt + Ologen™ implant. **Untreated intraocular pressure, when the patient arrived at the clinic. SD: Standard deviation; CI: Confidence interval; POAG: Primary open-angle glaucoma; CACG: Chronic angle-closure glaucoma; NTG: Normal-tension glaucoma; PG: Pigmentary glaucoma; MMC: Mitomycin-C; IOP: Intraocular pressure; NOHM: Number of ocular hypotensive medications; Phaco: Phacoemulsification

Ocular hypotensive medications

The mean number of ocular hypotensive medications significantly decreased from 3.2 ± 1.1 drugs preoperatively to 0.3 ± 0.8 drugs at the month-18 visit. However, there were no statistically significant differences in the mean reduction of ocular hypotensive medications between the MicroShunt-solo (3.1 ± 1.1 drugs) and MicroShunt + Phaco (2.5 ± 1.3 drugs) groups (Hodges–Lehmann median difference: -1.0 ; 95% CI: -1.0 to 0.0 ; $P = 0.1197$) [Fig. 4].

Predictive factors for success

In logistic regression analysis, no factors were found to be significantly related to success in the univariate analysis.

However, there was a tendency for performing cataract surgery and having an IOP of >8 mmHg on postoperative day 1 to reduce the probability of success (odds ratio = 0.18 ; $P = 0.0513$ and odds ratio = 0.123 ; $P = 0.0682$, respectively), although these did not reach statistical significance [Table 3].

Visual acuity

In the overall study sample, there were no significant changes in LogMAR mean best-corrected visual acuity (BCVA) from preoperative to month-18 visit (0.30 ± 0.47 vs. 0.24 ± 0.61 ; $P = 0.2147$). However, in the MicroShunt + Phaco group, LogMAR BCVA significantly improved from 0.39 ± 0.26 preoperatively to 0.17 ± 0.19 at the month-18 visit (Hodges–

Lehmann median difference: -0.21 ; 95% CI: -0.35 to -0.13 ; $P = 0.0003$). In contrast, in the MicroShunt-solo group, BCVA remained stable (Hodges–Lehmann median difference: -0.00 ; 95% CI: -0.03 to 0.05 ; $P = 0.5693$).

Eyes with previous glaucoma surgery

Fourteen eyes (19.2%) had undergone previous glaucoma surgery, with 12 eyes undergoing one antiglaucomatous surgical procedure and two eyes undergoing two glaucoma surgeries [Table 4]. All of these eyes underwent a MicroShunt standalone procedure. The mean age was 72.6 ± 11.4 years (range: 39–86 years), with seven (50.0%) women, and 13 (92.9%) underwent surgery with MMC 0.04%. Among these eyes, 12 (85.7%) were classified as a success, with 10 (71.4%) as complete success and 2 (14.3%) as qualified success. The mean preoperative IOP significantly reduced from

21.5 ± 6.5 mmHg to 11.9 ± 2.7 mmHg at month 18 (Hodges–Lehmann median difference: -9.0 ; 95% CI: -13.0 to -6.0 mmHg; $P = 0.0001$). Similarly, the mean preoperative number of ocular hypotensive medications significantly reduced from 3.3 ± 1.3 to 0.4 ± 0.8 at month 18 (Hodges–Lehmann median difference: -3.0 ; 95% CI: -3.5 to -2.0 ; $P = 0.0002$). There were no significant differences in either preoperative BCVA (Hodges–Lehmann median difference: 0.05 ; 95% CI: -0.05 to 0.15 ; $P = 0.3825$) or month-18 BCVA (Hodges–Lehmann median difference: 0.0 ; 95% CI: -0.1 to 0.1 ; $P = 0.9435$) between the two time points.

Eyes without previous surgery

To minimize potential biases, a subanalysis was performed on eyes that had not undergone previous surgery. Except for preoperative IOP, which was significantly higher in the MicroShunt-solo group, no significant differences were observed at any of the different time points measured throughout the study follow-up [Fig. 5a]. However, the mean IOP lowering was significantly greater in the MicroShunt-solo group than in the MicroShunt + Phaco group [Fig. 5b]. In the overall study population, the preoperative mean IOP significantly reduced from 18.9 ± 5.4 mmHg to 11.2 ± 2.6 mmHg (mean difference: -7.8 ± 5.8 mmHg; 95% CI: -9.8 to -5.8 mmHg; $P < 0.0001$, Friedman test). In the MicroShunt-solo group, preoperative IOP significantly reduced from 21.6 ± 3.6 mmHg to 11.6 ± 2.4 mmHg at month 18 ($P < 0.0001$) [Fig. 5a]. Similarly, there was a significant IOP reduction in the MicroShunt + Phaco group from preoperative (17.1 ± 5.8 mmHg) to month 18 (10.9 ± 2.7 mmHg) [Fig. 5a]. Unadjusted IOP lowering was significantly greater in the MicroShunt-solo group compared to the MicroShunt + Phaco group at all the different time points measured [Fig. 5b].

Safety

In terms of surgical complications, two (4.2%) eyes experienced choroidal detachment without visual impairment, two (4.2%)




Figure 1: Kaplan–Meier survival analysis. The mean hazard ratio was 4.56 (95% CI: 1.06–19.59, $P = 0.0414$)




Figure 2: Mean intraocular pressure (IOP) over the course of follow-up. The vertical bars represent the 95% confidence interval. (a) Overall study population. $*P < 0.0001$ as compared to preoperative value (repeated measures ANOVA and the Greenhouse-Geisser correction). (b) Comparison between eyes that underwent MicroShunt-solo and those that underwent MicroShunt + phacoemulsification (MicroShunt + Phaco) surgery. The preoperative mean IOP was significantly lower in the MicroShunt + Phaco group ($P = 0.0025$). No significant differences were observed at any other measured IOP time points (Mann–Whitney U test). Compared to preoperative values, IOP was significantly lowered at all different time points. $**P < 0.0001$ as compared to preoperative value (Friedman test with the Conover method). IOP: Intraocular pressure; MicroShunt solo: Preserflo™ MicroShunt + Ologen™; MicroShunt + Phaco: Preserflo™ MicroShunt + Ologen™ + Phacoemulsification; Preop: Preoperative; D: Day; W: Week; M: Month; ns: not significant

Table 2: Adjusted Mean changes in intraocular pressure (IOP) over the course of follow-up in eyes that underwent Preserflo™ MicroShunt + Ologen™ (MicroShunt solo) versus those who underwent Preserflo™ MicroShunt + Ologen™ + Phacoemulsification (MicroShunt + Phaco)

Mean change in IOP, mmHg	MicroShunt solo		MicroShunt + Phaco		Difference		
	Mean	SE	Mean	SE	Mean (SE)	95% CI	P
Day 1	-12.4	0.7	-10.7	0.8	-1.7 (1.2)	-4.1 to 0.7	0.1573
Week 1	-10.4	0.8	-11.1	1.1	0.7 (1.5)	-2.3 to 3.7	0.6649
Month 1	-10.6	0.8	-8.2	0.9	-2.5 (1.3)	-0.4 to 5.0	0.0860
Month 3	-9.0	0.6	-9.5	0.8	0.5 (1.1)	-1.6 to 2.7	0.6341
Month 6	-8.5	0.6	-9.1	0.7	0.6 (1.0)	-1.4 to 2.5	0.5418
Month 12	-8.3	0.4	-8.8	0.6	0.5 (0.8)	-1.0 to 2.1	0.4890
Month 18	-8.2	0.4	-8.4	0.5	0.2 (0.8)	-1.4 to 1.8	0.7860
Significance*	P<0.0001		P<0.0001				

*Bonferroni corrected. *Repeated measures ANCOVA and the Greenhouse–Geisser correction. *Repeated measures analysis of covariance (MANCOVA). The model included "Type of surgery" (MicroShunt-solo vs. MicroShunt + Phaco) as a factor and age, preoperative IOP, pachymetry; and previous glaucoma surgery as covariates. IOP=Intraocular pressure; SE=Standard error; CI=Confidence interval




Figure 3: Comparison of the mean intraocular pressure (IOP) lowering from preoperative values in Preserflo™ MicroShunt + Ologen™ (MicroShunt-solo) and Preserflo™ MicroShunt + Ologen™ + Phacoemulsification (MicroShunt + Phaco) groups. * $p < 0.0001$, ** $p < 0.05$. Statistical significance was determined using the Mann–Whitney test. IOP: Intraocular pressure

eyes had transient hyphema, one (2.1%) eye had a reactivation of a corneal herpetic ulcer, one (2.1%) eye had diplopia, and one (2.1%) eye had a shallow anterior chamber during the first week. All cases of postoperative surgical complications were successfully managed with medical therapy.

No eyes required needling, and only one (2.1%) eye (diagnosed with chronic angle-closure glaucoma) needed repositioning of the implant due to the tip being too close to the corneal endothelium. After 18 months of follow-up, this eye had an IOP of 10 mmHg with the use of two medications.

Discussion

In recent years, significant advancements have been made in the field of glaucoma surgery. MIGS devices have emerged as safer and less traumatic alternatives for lowering IOP in glaucoma patients.^[14] Among the various MIGS devices available, the Preserflo™ MicroShunt is an ab-externo and subconjunctival device that has demonstrated favorable efficacy and safety profiles.^[15-25]




Figure 4: Mean number of ocular hypotensive medications over the course of follow-up. Statistical significance was determined using the Mann–Whitney test. Preop: Preoperative; ns: Not-significant; MicroShunt-solo: Preserflo™ MicroShunt + Ologen™; MicroShunt + Phaco: Preserflo™ MicroShunt + Ologen™ + Phacoemulsification

In this study, the overall success rate in the study population was 83.3%. The MicroShunt-solo group achieved a success rate of 92.9%, whereas the MicroShunt + Phaco group had a success rate of 70.0%. Although there were no significant differences between the two groups ($P = 0.0532$), both the standalone use of the MicroShunt and its combination with phacoemulsification demonstrated significant reductions in IOP and the number of ocular hypotensive medications in glaucoma patients treated in a real clinical setting.

Initially, the unadjusted analysis showed significantly greater IOP lowering in the MicroShunt-solo group at all measured time points. However, after adjusting for various covariates, there were no significant differences in IOP reduction between the standalone MicroShunt implant and its combination with cataract surgery at any of the measured time points.

Despite the increasing evidence supporting the use of the MicroShunt implant in glaucoma patients,^[15-25] there are limited studies analyzing its effectiveness in combination with the collagen–glycosaminoglycan porous matrix implant, Ologen™.

Table 3: Univariate and multivariate analysis of the 48 eyes included in the study, to evaluate potential factors for success. Factors associated with success in the univariate analysis at $P < 0.1$ were included in the multivariate analysis

Variable	Success			
	Univariate		Multivariate	
	OR (95% CI)	P	OR (95% CI)	P
Age*				
>73 years	1.67 (0.35–7.93)	0.5209		
Sex				
Ref: Female	0.80 (0.17–3.88)	0.7845		
Male				
Eye				
Ref: Right	0.67 (0.14–3.05)	0.6019		
Left				
Device implantation				
Ref: ST	1.80 (0.19–16.98)	0.6077		
SN	0.20 (0.01–3.66)	0.2779		
IN				
Previous glaucoma surgery				
Ref: No	1.29 (0.23–7.31)	0.7768		
Yes				
Cataract surgery				
Ref: No	0.18 (0.04–1.01)	0.0513	0.19 (0.03–1.13)	0.0679
Yes				
Pachymetry				
>542.5 μm	1.00 (0.22–4.56)	1.0000		
MMC dose				
Ref: 0.02%	1.33 (0.23–7.89)	0.7511		
0.04%				
Initial IOP				
>27.5 mmHg	0.54 (0.11–2.58)	0.4428		
Preop IOP				
>18 mmHg	3.32 (0.60–18.45)	0.1711		
Day 1				
IOP >8 mmHg	0.23 (0.05–1.12)	0.0682	0.24 (0.05–1.29)	0.0973
NOHM				
Per medication	0.64 (0.27–1.52)	0.3070		

*Reference group \leq Median. ST: Superotemporal; SN: Superonasal; IN: Inferonasal; MMC: Mitomycin-C; IOP: Intraocular pressure; NOHM: Number of ocular hypotensive medications

To the best of our knowledge, only one study has evaluated the clinical outcomes of the MicroShunt in pseudophakic eyes with moderate to advanced POAG.^[11]

Vastardis *et al.*^[11] reported a 6-month success rate of 91.8% using the Preserflo device with a success criterion of IOP ≥ 5 and ≤ 16 mmHg. In our study, we observed similar success rates of 92.9% in eyes that underwent the solo procedure, although our success criteria were stricter (IOP ≥ 6 and ≤ 15 mmHg) and our follow-up was three times longer (18 months compared to Vastardis *et al.*'s 6 months).

In terms of IOP reduction, Vastardis *et al.* reported a significant decrease from 26.04 ± 8.76 mmHg preoperatively to 11.75 ± 3.37 mmHg at 6 months. In our study, the mean preoperative IOP in the MicroShunt-solo group significantly reduced from 21.5 ± 5.2 mmHg to 10.9 ± 3.0 mmHg at 6 months and 11.7 ± 2.5 mmHg at 18 months.

Regarding the reduction of ocular hypotensive medications, our study yielded comparable results to Vastardis *et al.*,^[11] despite our much longer follow-up duration.

It is worth noting that unlike Vastardis *et al.*,^[11] who performed surgical revision in three eyes in the Ologen™ group, our study did not encounter any cases of failed or flat blebs. Although we did have two failures due to elevated IOP, which might have benefited from needling, this procedure was not included as part of our post-surgical management protocol. This disparity can be partially explained by differences in the use of MMC as we used a concentration of 0.04% in 82.1% of the eyes that underwent the solo procedure.

Another potential explanation for the differences is the Ologen™ model itself. In our study, we used a 12-mm diameter \times 1-mm thickness implant, which could increase the total filtering area and place less stress on the conjunctiva, whereas Vastardis *et al.*^[11] used a 6-mm diameter \times 2-mm thickness implant.

Direct comparisons with previous studies are challenging as they did not utilize the Ologen™ implant.^[15-25] Nevertheless, the observed good IOP lowering effect in our study was in line with those previous reports and did not significantly differ [Table 5].

Table 4: Overview of the main demographic and clinical characteristics of the eyes that had undergone previous glaucoma surgery

n	Age	Sex	Eye	Dx	Previous Surgery	Surgery	BCVA		IOP			NOHM		Complications*
							Preop	Month-18	Preop	Month-18	Preop	Month-18		
													Initial	
1	39	F	R	POAG	NPDS	Standalone	0.5	0.3	50	34	15	4	0	No
2	81	F	L	POAG	CP	Standalone	0.05	0.05	26	22	12	3	0	No
3	63	M	L	POAG	AhV	Standalone	0.4	1.5	50	35	17	5	2	No
4	81	M	L	POAG	XEN	Standalone	0	0	28	20	8	4	0	CD
5	66	M	R	CACG	TRAB	Standalone	0	0	26	18	14	4	0	SAC
6	72	F	L	POAG	CP	Standalone	0.1	0.1	25	16	9	3	0	No
7	81	F	R	POAG	CP	Standalone	0.15	0.1	26	19	11	2	0	Fibrosis
8	73	F	R	POAG	NPDS & AhV	Standalone	0.05	0.2	28	20	12	4	0	No
9	86	F	R	CACG	NPDS	Standalone	0.5	0.4	34	30	11	4	2	No
10	77	F	R	POAG	NPDS	Standalone	0	0.15	28	18	8	4	0	Diplopia
11	75	M	R	POAG	NPDS & AhV	Standalone	0	0	28	16	14	4	0	No
12	73	M	L	POAG	CP	Standalone	0.05	0	28	17	14	0	0	No
13	77	F	R	POAG	TRAB	Standalone	0	0.15	24	18	10	3	1	Hyphema
14	72	M	R	CACG	XEN	Standalone	0	0	25	18	11	2	0	No

*MicroShunt post-operative complications. N: Case number; IOP: Intraocular pressure; F: Female; M: Male; Dx: Diagnosis; POAG: Primary open-angle glaucoma; CACG: Chronic angle-closure glaucoma; NPDS: Non-penetrating deep sclerectomy; AhV: Ahmed valve; CP: Canaloplasty; TRAB: Trabeculectomy; Preop: Preoperative; BCVA: Best corrected visual acuity; IOP: Intraocular pressure; Initial: Baseline, untreated IOP, when the patient arrived to the clinic; NOHM: Number of ocular hypotensive medications; SAC: Shallow anterior chamber; CD: Choroidal detachment




Figure 5: Mean intraocular pressure (IOP) (a) and mean IOP lowering (b) over the course of follow-up in the eyes that did not undergo previous glaucoma surgery. The vertical bars represent the 95% confidence interval. (a) A comparison of mean IOP between the eyes that underwent MicroShunt-solo and those that underwent MicroShunt + phacoemulsification (MicroShunt + Phaco) surgery. The preoperative mean IOP was significantly lower in the MicroShunt + Phaco group. No significant differences were observed at any other measured IOP time points (Mann–Whitney U test). IOP was significantly lowered at all different time points compared to preoperative values (* $p < 0.0001$, Friedman test with the Conover method). (b) Comparison of the mean IOP lowering from preoperative values in MicroShunt-solo and MicroShunt + Phaco groups. Statistical significance was determined using the Mann–Whitney test ($\dagger p < 0.0001$, $\ddagger p < 0.05$). IOP: Intraocular pressure; MicroShunt-solo: Preserflo™ MicroShunt + Ologen™; MicroShunt + Phaco: Preserflo™ MicroShunt + Ologen™ + Phacoemulsification; Preop: Preoperative; D: Day; W: Week; M: Month; ns: Not Significant

Our study did not identify any factors significantly associated with surgery success. However, we observed a trend suggesting that a day-1 IOP of > 8 mmHg or the performance of combined surgery (MicroShunt + Phacoemulsification) may reduce the probability of success.

The preoperative IOP of 11 (5.0%) eyes who underwent combined surgery was ≤ 16 mmHg. This means that the main indication in these cases was cataract surgery and, secondarily,

the elimination/reduction of ocular hypotensive medication instead of lowering IOP.

Regarding eyes with previous glaucoma surgery, among the 28 eyes that underwent the solo procedure, 14 (50.0%) had undergone a previous glaucoma surgery, whereas 14 (50.0%) had not. The median preoperative IOP was slightly higher in eyes that did not undergo previous glaucoma surgery, but the difference was not statistically significant (Hodges–Lehmann

Table 5: Overview of the intraocular pressure (IOP) and number of hypotensive medications in eyes that underwent Preserflo™ MicroShunt implant surgery in different studies

Study	n	Follow-up	Preoperative IOP, mmHg	Final IOP, mmHg	Mean preoperative medications	Mean postoperative medications
Vastardis <i>et al.</i> ^[11]						
MMC 0.02%	25	6M	23.5±5.8*	11.6±3.1*	2.5±0.9*	0.04±0.2*
MMC 0.02% + OlogenTM	25	6M	26.0±8.8*	11.8±3.4*	2.6±0.8*	0.2±0.8*
Riss <i>et al.</i> ^[15]						
MMC 0.02%	31	12 M	27.9±6.7*	13.3±3.3*	2.5±1.4*	0.5±1.0*
MMC 0.04%	23	12 M	23.8±5.3*	10.7±2.8*	2.4±0.9*	0.3±0.8*
Batlle <i>et al.</i> ^[16]						
	23	12M	23.8±5.3*	10.7±2.8*	2.4±0.9*	0.3±0.8*
	22	24M	23.8±5.3*	11.9±3.7*	2.4±0.9*	0.4±1.0*
	22	36M	23.8±5.3*	10.7±3.5*	2.4±0.9*	0.7±1.1*
Scheres <i>et al.</i> ^[17]	41	18M	20.1±5.0*	12.5±3.3*	2.3±1.5*	0.6±0.9*
Batlle <i>et al.</i> ^[18]						
	21	48M	23.8±5.3*	12.8±5.6*	2.4±1.0*	0.5±1.1*
	21	60M	23.8±5.3*	12.4±6.5*	2.4±1.0*	0.8±1.3*
Martínez-de-la-Casa <i>et al.</i> ^[20]						
MicroShunt	35	12M	21.3±3.2*	14.4±3.4*	2.3±0.5*	-2.1±0.7**
MicroShunt + Phaco	23	12M	21.7±3.5*	14.9±3.6*	2.4±0.5*	-2.1±0.6**
Quaranta <i>et al.</i> ^[21]	31	12M	24.1±3.1*	12.6±2.6*	3.3±0.6*	0.5±0.8*
Fea <i>et al.</i> ^[22]	104	12M	25.1±6.5*	14.1±3.4*	3.0±1.0*	0.8±1.0*
Beckers <i>et al.</i> ^[23]						
	81	12M	21.7±3.4*	14.5±4.6*	2.1±1.3*	N.A.
	81	24M	21.7±3.4*	14.1±3.2*	2.1±1.3*	0.5±0.9*
Ibarz-Barberá <i>et al.</i> ^[24]						
MicroShunt	51	9M	22.5±0.9*	12.8±0.4*†	2.7±0.7*‡	0.19±0.5*‡
MicroShunt + Phaco	13	9M	20.1±1.3*	12.5±0.6*†		
Bhayani <i>et al.</i> ^[25]	100	12M	21.5 (19–28)□	13 (11–16)□	3 (2–3)□	0 (0–1)□
Current study						
Overall	48	18M	19.7±5.8*	11.4±2.6*	3.2±1.1*	0.3±0.8*
MicroShunt	28	18M	21.5±5.2*	11.7±2.5*	3.4±1.0*	0.3±0.7*
MicroShunt + Phaco	20	18M	17.1±5.8*	10.9±2.7*	2.9±1.1*	0.4±0.9*

*Mean±Standard deviation. **Mean±Standard deviation reduction. †"Last follow-up visit," without specifying follow-up. ‡Overall study sample. □Median (Interquartile range). IOP: Intraocular pressure; M: Month; NA: Not available

median difference: 2.0 mmHg; 95% CI: -2.0 to 4.0 mmHg, $P = 0.2856$). Similarly, there were no significant differences in month-18 IOP between the two groups (Hodges–Lehmann median difference: 0.0 mmHg; 95% CI: -2.0 to 2.0 mmHg, $P = 0.9443$) or in the median IOP lowering from preoperative values (Hodges–Lehmann median difference: 0.0 mmHg; 95% CI: -4.0 to 3.0 mmHg, $P = 0.7636$).

In terms of success rates, 14 (100%) eyes without previous glaucoma surgery and 12 (85.7%) eyes with previous glaucoma surgery were classified as successful (either complete or qualified) ($P = 0.1493$). In addition, no significant differences were observed in either preoperative or month-18 best-corrected visual acuity (BCVA) between both groups.

These results are consistent with those published by Quaranta *et al.*^[21] who reported that the Preserflo™ MicroShunt can be effectively and safely used in eyes with a single failed trabeculectomy. It is important to note that our study included patients with different types of glaucoma, including 11 eyes with POAG and three eyes with chronic angle-closure glaucoma (CAG), whereas Quaranta *et al.*'s study focused on patients with POAG.

These findings support the hypothesis that MicroShunt + Ologen implantation may serve as a safe and effective alternative in patients who have previously undergone glaucoma surgery, presenting new possibilities for its application

in clinical practice. However, despite the promising results, further studies are necessary to thoroughly analyze this option.

Previous studies have indicated that the dose and use of MMC can have a positive effect on surgical success.^[16-19] Higher MMC doses have been associated with a lower risk of MicroShunt failure.^[19] In our study, however, MMC dose was not found to be a predictor of surgery success. Furthermore, there were no significant differences in IOP lowering between MMC 0.02% and MMC 0.04% at 18 months (Hodges–Lehmann median difference: 0.0 mmHg; 95% CI: -3.0 to 4.0 mmHg, $P = 0.8986$). This could be due to variability in the application time or the fact that only 10 eyes received MMC 0.02%.

Study limitations

The study has certain limitations. First, its retrospective design introduces the possibility of selection bias and confounding factors, which are inherent to such studies. Nevertheless, the use of strict inclusion and exclusion criteria, as well as a large number of eyes, helps mitigate these concerns. The absence of a control group is another limitation. However, various subanalyses were performed (solo procedure vs. combined procedure and previous glaucoma surgery vs. primary surgery), providing valuable clinical information. It is also important to note that nine (18.8%) have been diagnosed with CAG. Among them, 6 eyes underwent a solo procedure because they had previously undergone cataract surgery, while

3 eyes underwent combined surgery (MicroShunt + Phaco) to facilitate the opening of the chamber angle.^[26]

The lack of difference in success rates and IOP lowering between different MMC doses may be attributed to the heterogeneity of our study sample or the limited number of eyes in the 0.02% MMC group. Therefore, further studies will be necessary to elucidate the role of MMC dose and application time on clinical outcomes.

Conclusions

The findings of this study suggest that the use of the Preserflo™ MicroShunt implant, either alone or in combination with cataract surgery, in conjunction with the Ologen™ implant, leads to a significant reduction in preoperative IOP and a decreased reliance on ocular hypotensive drugs in patients with glaucoma. In addition, this technique has shown favorable clinical outcomes in eyes with a history of previous glaucoma surgery. Although no factors were found to be significantly associated with surgery success in our study, further research is needed to validate or revise this finding. Importantly, no unexpected adverse events were observed in our study.

Ethical approval

“All procedures performed in studies involving human participants were in accordance with the ethical standards of the Ethics Committee of the Hospital Virgen de la Luz and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards”.

Informed consent

“Due to the characteristics of the study, the Ethics Committee of the Hospital Virgen de la Luz waived the need for written informed consent.” Nevertheless, all the patients signed a written informed consent for their surgery.

Data sharing statement

Data not published here are obtainable on reasonable request from the corresponding author.

Acknowledgments

Medical writing and Editorial assistant services have been provided by Antonio Martínez (MD) of Ciencia y Deporte S.L.

Financial support and sponsorship: Support for Editorial assistance was funded by Santen. It should be noted that Santen was not involved in the preparation of the manuscript nor did the company influence in any way the scientific conclusions reached.

Conflicts of interest: There are no conflicts of interest.

References

- European Glaucoma Society Terminology and Guidelines for Glaucoma, 4th Edition—Chapter 3: Treatment principles and options. Supported by the EGS Foundation: Part 1: Foreword; Introduction; Glossary; Chapter 3 Treatment principles and options. *Br J Ophthalmol* 2017;101:130–95.
- Joshi AB, Parrish RK 2nd, Feuer WF. 2002 survey of the American Glaucoma Society: Practice preferences for glaucoma surgery and antifibrotic use. *J Glaucoma* 2005;14:172–4.
- Saheb H, Ahmed II. Micro-invasive glaucoma surgery: current perspectives and future directions. *Curr Opin Ophthalmol* 2012;23:96–104.
- Jampel HD, Musch DC, Gillespie BW, Lichter PR, Wright MM, Guire KE, *et al*. Perioperative complications of trabeculectomy in the collaborative initial glaucoma treatment study (CIGTS). *Am J Ophthalmol* 2005;140:16–22.
- Gedde SJ, Feuer WJ, Lim KS, Barton K, Goyal S, Ahmed IIK, *et al*. Treatment outcomes in the primary tube versus trabeculectomy study after 3 years of follow-up. *Ophthalmology* 2020;127:333–45.

- Holland LJ, Kirwan JF, Mercieca KJ. Effect of COVID-19 pandemic on glaucoma surgical practices in the UK. *Br J Ophthalmol* 2022;106:1406–10.
- Sadruddin O, Pinchuk L, Angeles R, Palmberg P. Ab externo implantation of the MicroShunt, a poly (styrene- block-isobutylene- block-styrene) surgical device for the treatment of primary open-angle glaucoma: A review. *Eye Vis (Lond)* 2019;6:36.
- He M, Wang W, Zhang X, Huang W. Ologen implant versus mitomycin C for trabeculectomy: A systematic review and meta-analysis. *PLoS One* 2014;9:e85782.
- Sastre-Ibáñez M, Cabarga C, Canut MI, Pérez-Bartolomé F, Urcelay-Segura JL, Cordero-Ros R, *et al*. Efficacy of Ologen matrix implant in Ahmed glaucoma valve implantation. *Sci Rep* 2019;9:3178.
- Jacobson A, Rojas C, Bohnsack BL. Ologen augmentation of Ahmed glaucoma drainage devices in pediatric glaucomas. *BMC Ophthalmol* 2021;21:72.
- Vastardis I, Fili S, Perdikakis G, Kontopoulou K, Balidis M, Gatziofias Z, *et al*. Preliminary results of Preserflo Microshunt versus Preserflo Microshunt and Ologen implantation. *Eye Vis (Lond)* 2021;8:33.
- Ibarz-Barberá M, Morales-Fernández L, Corroto-Cuadrado A, Martínez-Galdón F, Tañá-Rivero P, Gómez de Liaño R, *et al*. Corneal endothelial cell loss after PRESERFLO™ MicroShunt implantation in the anterior chamber: Anterior segment OCT tube location as a risk factor. *Ophthalmol Ther* 2022;11:293–310.
- Lang KM, Little TD. Principled missing data treatments. *Prev Sci* 2018;19:284–94.
- Bar-David L, Blumenthal EZ. Evolution of glaucoma surgery in the last 25 years. *Rambam Maimonides Med J* 2018;9:e0024. doi: 10.5041/RMMJ.10345.
- Riss I, Batlle J, Pinchuk L, Kato YP, Weber BA, Parel JM. Résultats à un an de l'efficacité et de l'innocuité du MicroShunt InnFocus selon l'emplacement et la concentration de MMC [One-year results on the safety and efficacy of the InnFocus MicroShunt™ depending on placement and concentration of mitomycin C]. *J Fr Ophtalmol* 2015;38:855–60.
- Batlle JF, Fantes F, Riss I, Pinchuk L, Albuquerque R, Kato YP, *et al*. Three-Year follow-up of a novel aqueous humor MicroShunt. *J Glaucoma* 2016;25:e58–65.
- Scheres LMJ, Kujovic-Aleksov S, Ramdas WD, de Crom RMP, Roelofs LCG, Berendschot TTJM, *et al*. XEN[®]Gel Stent compared to PRESERFLO™ MicroShunt implantation for primary open-angle glaucoma: Two-year results. *Acta Ophthalmol* 2021;99:e433–40.
- Batlle JF, Corona A, Albuquerque R. Long-term results of the PRESERFLO MicroShunt in patients with primary open-angle glaucoma from a single-center nonrandomized study. *J Glaucoma* 2021;30:281–6.
- Schlenker MB, Durr GM, Michaelov E, Ahmed IIK. Intermediate outcomes of a novel standalone Ab Externo SIBS microshunt with mitomycin C. *Am J Ophthalmol* 2020;215:141–53.
- Martínez-de-la-Casa JM, Saenz-Francés F, Morales-Fernández L, Perucho L, Mendez C, Fernández-Vidal A, *et al*. Clinical outcomes of combined Preserflo Microshunt implantation and cataract surgery in open-angle glaucoma patients. *Sci Rep* 2021;11:15600.
- Quaranta L, Micheletti E, Carassa R, Bruttini C, Fausto R, Katsanos A, *et al*. Efficacy and safety of PreserFlo® MicroShunt after a failed trabeculectomy in eyes with primary open-angle glaucoma: A retrospective study. *Adv Ther* 2021;38:4403–12.
- Fea AM, Laffi GL, Martini E, Economou MA, Caselgrandi P, Sacchi M, *et al*. Effectiveness of MicroShunt in patients with primary open-angle and pseudoexfoliative glaucoma: A Retrospective European multicenter study. *Ophthalmol Glaucoma* 2022;5:210–8.
- Beckers HJM, Aptel F, Webers CAB, Bluwol E, Martínez-de-la-Casa JM, García-Feijóo J, *et al*. Safety and effectiveness of the PRESERFLO® MicroShunt in primary open-angle glaucoma: Results from a 2-year multicenter study. *Ophthalmol Glaucoma* 2022;5:195–209.
- Ibarz Barberá M, Martínez-Galdón F, Caballero-Magro E, Rodríguez-Piñero M, Tañá-Rivero P. Efficacy and safety of the Preserflo microshunt with mitomycin C for the treatment of open angle glaucoma. *J Glaucoma* 2022;31:557–66.
- Bhayani R, Martínez de la Casa JM, Figus M, Klabe K, Rabiolo A, Mercieca K. Short-term safety and efficacy of Preserflo™ Microshunt in glaucoma patients: A multicentre retrospective cohort study. *Eye (Lond)* 2023;37:644–9.
- Azuara-Blanco A, Burr J, Ramsay C, Cooper D, Foster PJ, Friedman DS, *et al*. Effectiveness of early lens extraction for the treatment of primary angle-closure glaucoma (EAGLE): A randomised controlled trial. *Lancet* 2016;388:1389–97.