

---

This is the **accepted version** of the journal article:

Ullmo, Johana [et al.]. «Cardiovascular risk assessment in women at high risk for pre-eclampsia in first half of pregnancy». *Ultrasound in obstetrics & gynecology*, 2025.

---

This version is available at <https://ddd.uab.cat/record/310453>

under the terms of the  <sup>IN</sup>COPYRIGHT license.

## **TITLE PAGE**

### **Cardiovascular risk assessment in women at high risk for pre-eclampsia in first half of pregnancy**

Johana Ullmo, MD<sup>1,2</sup>; Mónica Cruz-Lemini, MD PhD<sup>1,2,3</sup>; Lidia Bos-Real, MD<sup>4</sup>; Mireia Padilla, MD<sup>4</sup>; Marina Coma-Barbarà, MD<sup>1</sup>; Silvia Martorell, MD<sup>1</sup>; Josefina Mora, MD PhD<sup>2,5</sup>; Jordi Ordoñez-Llanos, MD PhD<sup>5,6</sup>; Elisa Llurba, MD PhD<sup>1,2,3</sup>, on behalf of the AngioCor working group.

<sup>1</sup>Department of Obstetrics and Gynecology, Hospital de la Santa Creu i Sant Pau, Sant Antoni Maria Claret 167, 08025 Barcelona, Spain. Universitat Autònoma de Barcelona, Barcelona, Spain.

<sup>2</sup>Women and Perinatal Health Research Group, Institut de Recerca (IR SANT PAU), Sant Quintí 77-79, 08041 Barcelona, Spain.

<sup>3</sup>Primary Care Interventions to Prevent Maternal and Child Chronic Diseases of Perinatal and Developmental Origin Network (RICORS-SAMID) (RD21/0012/0001) and Maternal and Child Health Development Network (SAMID, RD16/0022/0015), Instituto de Salud Carlos III, Madrid, Spain.

<sup>4</sup>Cardiology Department, Hospital de la Santa Creu i Sant Pau, IIB-SantPau, Universidad Autónoma de Barcelona, Barcelona, Spain.

<sup>5</sup>Department of Clinical Biochemistry, Hospital de la Santa Creu i Sant Pau, Institut d'Investigació Biomèdica Sant Pau (IIB Sant Pau), 08041 Barcelona, Spain.

<sup>6</sup>Fundació per la Bioquímica i la Patologia Molecular, Biochemistry Department, Santa Creu i Sant Pau University Hospital & Universitat Autònoma, Barcelona, Spain.

#### **Corresponding author:**

Elisa Llurba, MD PhD.

Director of the Obstetrics and Gynecology Department

Professor in Obstetrics and Gynecology, Universitat Autònoma de Barcelona

Hospital de la Santa Creu i Sant Pau – Institut de Recerca Sant Pau

Sant Quintí 89, 08025, Barcelona, Spain.

Tel: +34 93 553 70 48 / +34 93 553 70 41

email: ellurba@santpau.cat

ORCID: 000-0003-4048-6407

## **KEYWORDS**

Pre-eclampsia, angiogenic factors, echocardiography, cardiovascular risk, cardiac biomarkers.

## 1 **ABSTRACT**

2 **Background:** Women diagnosed with pre-eclampsia (PE), especially those with early-  
3 onset and severe forms, are at heightened risk of future cardiovascular disease (CVD). It  
4 is uncertain if women at high-risk for PE present cardiovascular impairment at the  
5 beginning of pregnancy. The aim of this study was to assess cardiovascular biomarkers  
6 and cardiac function in the first half of pregnancy in women at high-risk for PE.

7 **Methods:** This was a prospective cohort study including women attending first trimester  
8 screening for PE. Risk of PE in the first trimester was calculated using a previously  
9 validated model, which includes maternal factors and biophysical markers. A cut-off of  
10 1 in 250 was used to classify high-risk for developing PE. Maternal echocardiographic  
11 function and morphometry were evaluated, and peripheral blood samples were obtained  
12 to assess hematological and biochemical variables, including lipid profile and  
13 cardiovascular and angiogenic factors.

14 **Results:** A total of 349 pregnant individuals were included; 257 at high-risk and 92 at  
15 low-risk for PE. Maternal echocardiography was performed at a mean  $\pm$  SD gestational  
16 age of  $14.9 \pm 1.3$  weeks and blood sampling at  $16.3 \pm 1.9$  weeks. The high-risk PE group  
17 exhibited a significantly worse lipid profile (lower levels of high-density lipoproteins  $69.7$   
18  $\pm 14.2$  vs  $74.0 \pm 14.6$  mg/dL,  $p=0.020$ ), lower placental growth factor ( $0.89 \pm 0.63$  vs  $1.15$   
19  $\pm 0.58$  MoM,  $p<0.001$ ), and significantly lower levels of N-terminal pro-brain natriuretic  
20 peptide ( $51.0 \pm 2.1$  vs  $52.1 \pm 2.0$  ng/L,  $p=0.008$ ). Echocardiographic evaluation indicated  
21 concentric remodeling, along with an altered diastolic pattern in the high-risk group.

22 **Conclusions:** Women at high risk of developing PE exhibit differences in lipid profile,  
23 angiogenic factors, and echocardiographic parameters, indicative of cardiac  
24 maladaptation or impairment during the first half of pregnancy. These findings suggest a  
25 link between subclinical cardiovascular dysfunction, PE development and later CVD risk.

## **INTRODUCTION**

Heart disease and stroke cause 1 in 3 deaths among women each year - more than all cancers combined<sup>1</sup>. Although a reduction in mortality from cardiovascular disease (CVD) has been seen recently, the effect is less pronounced in women. Pregnancy complications such as preeclampsia (PE), are risk factors that are independently associated with CVD in women<sup>2</sup>. Women also have a significant window of opportunity after pregnancy to take proactive measures upon identifying their increased risk.

PE is a complication occurring in 2 to 8% of all pregnancies and remains a major cause of maternal morbidity and mortality<sup>3,4</sup>. PE is commonly associated with abnormal placental pathology and placenta-mediated complications such as fetal growth restriction<sup>5</sup>. Severe PE is directly related to cardiac injury, characterized by cardiac remodeling, decreased myocardial relaxation and global left ventricular diastolic dysfunction<sup>6</sup>. Previous studies have shown that women with PE have an increased risk of developing CVD in the future; the risk being higher in those with early-onset and severe PE<sup>7</sup>. Multiple epidemiological studies have shown that PE is associated with development of cardiovascular (CV) risk factors, and outcomes like arterial hypertension, myocardial infarction or stroke, renal complications, vascular and metabolic dysfunction, during and in the subsequent years after the affected pregnancy<sup>8</sup>. Some studies show a 2-to-4-fold higher risk of developing CV disease in the short- and long-term in women with PE<sup>8,9</sup>.

It is uncertain if women at high-risk of PE present CV impairment at the beginning of pregnancy or alternatively, CV changes observed are secondary to endothelial dysfunction and cardiac injury due to the condition<sup>10,11</sup>. Knowing the existing mechanisms behind cardiac remodeling in pregnant women at high-risk of PE is

important to identify those at increased risk of long-term CV disease, and to develop strategies to prevent or treat both early and late CV complications.

In this study, we assessed echocardiographic and CV biomarkers in the first half of pregnancy in women at high and at low risk for PE, to identify whether CV changes precede development of clinical PE.

## **METHODS**

### **Study population**

This was a prospective cohort study conducted at the Maternal-Fetal Medicine Department of Hospital Sant Pau, Barcelona, Spain as part of the AngioCor prospective cohort study<sup>12</sup>. Women were recruited from July 2020 to April 2023 upon first trimester screening for PE. Inclusion criteria were a singleton pregnancy in women between 18 and 45 years old, able to understand study requirements, and provide written informed consent. Women diagnosed with Type-1 diabetes mellitus, hyperthyroidism, renal disease or with history of thromboembolic events were excluded from the study. Likewise, fetal malformations and/or fetal chromosomal abnormalities were not included.

The AngioCor study is registered on the ClinicalTrials.gov website under NCT04162236. The Hospital's Ethics Committee approved the study protocol (September 27<sup>th</sup>, 2019; approval number EC/19/297/5698), and written informed consent was obtained from all women to participate in the study.

PE risk in the first trimester (between 11<sup>+0</sup> to 13<sup>+6</sup> weeks) was calculated using a multivariate Gaussian distribution model previously validated in our population<sup>4</sup>. Maternal factors (ethnicity, weight, height, waist/hip ratio, smoking status, parity, history

of PE, pre-existing diabetes, hypertension, thrombophilia, renal or autoimmune conditions), biophysical markers [mean arterial blood pressure (MAP) and mean uterine artery pulsatility index (UtA-PI)], and pregnancy-associated plasma protein A (PAPP-A) were included in the model, giving a numerical risk calculation. A cutoff point of 1 in 250 was considered as high-risk for developing PE<sup>4,13</sup>. In a subset of patients with intermediate risk for PE, placental growth factor (PlGF) was analyzed and included in the model, to refine the risk calculation. A detailed description of our risk calculation model can be found in **Appendix S1**.

### **Plasma biomarker analysis**

Peripheral blood samples were obtained upon inclusion in the study. Routine hematological and biochemical variables, including lipid profile, were measured by standard, fully automated methods. Angiogenic factors (soluble fms-like tyrosine kinase-1 (sFLT-1); PlGF), high-sensitivity cardiac troponin T (hs-cTnT) and N-terminal pro-brain natriuretic peptide (NT-proBNP) were measured using automated electrochemiluminescence immunoassays on a Roche 711 cobas platform (Roche Diagnostics, Basel, Switzerland). Copeptin was measured using an automated immunofluorescent assay in a Kryptor instrument using two polyclonal antibodies directed to the 132.164 aminoacids of proAVP (BRAHMS Copeptin proAVP; ThermoFisher Scientific, Henningsdorf, Germany). All immunoassays showed appropriate accuracy and within- and between-batch precision in the internal and external quality control assessments.

### **Maternal echocardiography**

Ultrasound examinations were performed at rest in the left lateral decubitus position using Affiniti 70G and EPIQ 7W (Philips Healthcare, Andover, MA, USA)

ultrasound systems. A 9 MHz sector probe (E10-2; Philips Healthcare,) was used. Images were acquired with duration of 5 cardiac consecutive cycles and frame rate of  $\geq 60$  Hz. Clip acquisition was performed during maternal apnea. High-resolution images were acquired by an experienced obstetrician and post-processed by a cardiologist using dedicated software (Xcelera; Philips Healthcare, Andover, MA, USA). Cardiologists were blinded to the study group at the time of the analysis.

Two-dimensional (2D), M-mode and Doppler echocardiographic measurements were acquired according to standard protocols<sup>14</sup>. Standard echocardiographic measurements included left ventricle (LV) end-diastolic diameter (LVEDd), LV end-systolic diameter, LV end-diastolic volume, as well as end-diastolic wall thickness of the interventricular septum (IVS) and LV posterior wall thickness (PWT), LV outflow tract diameter, right ventricular diameter at the midlevel in the apical 4-chamber view, and 2D left atrial (LA) volume. LV mass was calculated using the formula  $0.8 \times (1.04 \times [(IVS + LVEDd + PWT)^3 - LVEDd^3] + 0.6$ .

LV function was assessed by ejection fraction obtained by the Simpson method. Mitral annular peak systolic excursion (MAPSE) and tricuspid annular peak systolic excursion (TAPSE) were also measured. LV diastolic function was evaluated by transmitral flow velocities using pulsed Doppler, measuring early transmitral diastolic flow velocity (E) and flow velocity during atrial contraction (A), and calculating E/A ratio. LV filling pressure was predicted using average (septal and lateral) mitral E/E' ratio, obtained from pulsed tissue Doppler. Cardiac output was calculated using the formula: velocity-time integral (VTI)  $\times$  cross-sectional area (CSA)  $\times$  heart rate, in which the VTI is obtained through pulsed wave Doppler in the LV outflow tract, the CSA is of the LV outflow tract and heart rate (beats per min) is determined through automatic measurement of the R-R interval in the electrocardiogram<sup>14,15</sup>.

Measurement of 2D global longitudinal strain (GLS), was conducted with automated tracking of myocardial motion, and was manually adjusted by correcting the endocardial borders, if deemed necessary.

## **Statistical analysis**

Data distributions involved application of standard statistical tests. Variables manifesting non-Gaussian distributions underwent natural logarithm transformation, followed by reassessment of Gaussianity. Data are expressed as proportions, mean  $\pm$  SD, or median and interquartile range. Disparities in proportions observed within high- and low-risk subgroups were subjected to analysis using the Z-test for large-sized samples or the chi-square test for small-sized samples. Comparisons between numerical variables were executed using either parametric (Student's *t*-test) or non-parametric (Mann-Whitney *U*-test) tests, contingent upon variable distribution. Correlation analyses were investigated using the Pearson method and confirmed by Spearman. To adjust for gestational age, angiogenic markers were transformed into multiples of the median (MoM) using a subset of control/low-risk patients.

Statistical significance was attributed to a *P*-value less than 0.05. Analysis was performed using IBM SPSS Statistics 25 (2017) statistical package (IBM Corp., Armonk, NY, USA).

## **RESULTS**

A total of 4196 pregnant patients were screened during the study period, identifying and inviting 433 high-risk patients to participate. In addition, 102 low-risk patients were also identified. This means that 10.3% of the screened population was deemed as high-risk for PE according to the model previously described<sup>4</sup>. A total of 349 women were included, of which 257 were classified as having a high risk and 92 as low

145 risk for PE according to their first trimester screening. A flowchart detailing patient  
146 inclusion in the study is shown in **Figure 1**.

147 **Maternal characteristics**

148 Maternal characteristics of high- and low-risk subgroups are shown in  
149 **Table 1**. There were no significant differences in terms of age or parity between the two  
150 groups. As expected, other maternal characteristics such as weight, body mass index  
151 (BMI), body surface area and systolic and diastolic blood pressures were significantly  
152 higher in the high-risk PE group, since these variables contribute directly to the calculated  
153 PE risk. After first trimester screening, aspirin prophylaxis was started in 97.3% of high-  
154 risk women.

155 **Plasma biomarkers**

156 **Table 2** presents the plasma biomarkers associated with high and low-risk of PE.  
157 Blood samples were obtained at  $16 \pm 2$  vs  $15.9 \pm 2$  weeks' gestation in the high and low-  
158 risk groups, respectively. Platelet count was significantly higher in patients at high-risk  
159 for PE, with no other differences in the general blood count. Women at high-risk for PE  
160 had higher triglyceride and lower high-density lipoprotein concentrations. Copeptin and  
161 hs-cTnT showed significantly higher concentrations in the high-risk compared with the  
162 low-risk group. Of note, hs-cTnT was detectable ( $>3.0$  ng/L) in 50.2% of women at high-  
163 risk for PE and in 40.9% of those at low-risk. In contrast, NT-proBNP concentration was  
164 lower in the high-risk group. Finally, PIGF was significantly lower in high-risk women  
165 compared with the control group, and the sFlt-1/PIGF ratio had a non-significant trend  
166 towards higher values in the high-risk group.

167 **Maternal echocardiography**

The results of the maternal echocardiographic measures are presented in **Table 3**. Maternal echocardiography was performed just after 1<sup>st</sup> trimester screening, at a mean  $\pm$  SD gestational age of  $14.9 \pm 1.3$  weeks. Measures were corrected for ethnicity and BMI or body surface area, considering disparity in both variables between the high- and low-risk groups. Regarding morphometric parameters, there were significantly thicker ventricular and posterior walls, as well as increased ventricular diastolic volumes in the high-risk group, while there were no differences in cardiac index. Significant differences were found in the high-risk group compared with the low-risk group for LV mass and LV mass index. Diastolic function parameters were significantly different between groups, with high-risk women showing an increased A wave velocity, decreased E/A ratio, and decreased tissue Doppler E' velocity in both lateral and septal mitral annuli, with increased E/E' ratio. LV ejection fraction and ventricular strain were not different between the high- and low-risk groups.

Correlation coefficients between biochemical markers (angiogenic factors, NT-proBNP, hs-cTnT, copeptin), and echocardiographic parameters are detailed in **Table S1**. Analysis revealed that there was no significant correlation between angiogenic factors and most maternal echocardiographic parameters. However, the sFlt-1/PlGF ratio demonstrated a positive correlation with LV mass and LV mass index, suggesting that a higher ratio is associated with increased LV mass. Notably, the sFlt-1/PlGF ratio also showed a significant positive correlation with strain parameters. Moreover, NT-proBNP displayed a significant positive correlation with tissue Doppler E' velocity in the LV lateral wall. This finding is consistent with our hypothesis that higher levels of NT-proBNP might offer a protective effect in hypertensive disorders during pregnancy. Copeptin also demonstrated a correlation with echocardiographic parameters analyzed, such as septal thickness and LV mass.

193

## 194 **DISCUSSION**

195       To our knowledge, this study is the first to prospectively compare CV function  
196 and biomarkers in early pregnancy between women at high-risk and at low-risk for PE.  
197 High-risk women showed echocardiographic differences, suggesting pre-existing or  
198 pregnancy-induced CV dysfunction. We observed expected differences regarding  
199 maternal characteristics, such as significantly higher blood pressure and BMI in the high-  
200 risk group, consistent with previous studies that have identified these factors as important  
201 contributors to the development of PE<sup>16,17</sup>, which are associated with increased CV  
202 risk<sup>16,18</sup>. Our findings in the high-risk group support the idea that these factors may induce  
203 early cardiac changes even in early pregnancy.

204       This study confirms that women at high-risk for PE showed a more atherogenic  
205 lipid profile (lower high-density lipoprotein, higher triglycerides), aligning with other  
206 studies that support the role of lipid metabolism in early placental dysfunction and cardiac  
207 remodeling in PE pathogenesis<sup>19,20</sup>. The high-risk group also showed a significantly  
208 higher platelet count, potentially indicating increased inflammation or subclinical  
209 endothelial dysfunction, as elevated first-trimester platelet volume has been linked to PE,  
210 and may reflect endothelial function<sup>21</sup>.

211       Regarding angiogenic factors, PlGF was lower in the high-risk group, with the  
212 sFlt-1/PlGF ratio showing a non-significant trend towards higher values. This may be  
213 explained because sFlt-1 rises until the third trimester in cases of PE. Previous studies  
214 have shown that vascular endothelial growth factor and PlGF contribute to cardiac  
215 remodeling during pregnancy<sup>22</sup>. Both factors are reduced in women with clinical PE and  
216 linked to adverse cardiac remodeling<sup>23</sup>. Other studies suggest that the angiogenic/anti-

angiogenic imbalance in PE may drive cardiac remodeling by disrupting endothelial growth and cell death. This may change heart structure and function, increasing ventricular mass and reducing diastolic function<sup>24</sup>, as observed in this study.

Regarding cardiac enzymes, troponin is released from damaged heart cells and used as marker for acute myocardial infarction. Previous studies show elevated levels of troponin in pregnant women with *established* PE, associated with impaired cardiac function and adverse CV outcomes<sup>25</sup>. Women with PE had higher hs-cTnT levels than those with a normal pregnancy<sup>25,26</sup>. We observed similar hs-cTnT patterns in early pregnancy, suggesting its inclusion in PE prediction models may improve accuracy, though further studies are needed. Brain natriuretic peptide (BNP) is produced in the left atrium and ventricle in response to myocyte stretching due to volume expansion or pressure overload<sup>27,28</sup>. While BNP levels are typically normal throughout pregnancy, studies have shown higher concentrations in the first trimester compared with the third<sup>29</sup>. Elevated NT-proBNP levels in the first trimester correlate with lower risk for PE over a 2-7-year follow-up<sup>30</sup>. In this study, women at high-risk for PE had significantly lower NT-proBNP levels, aligning with recent studies linking higher NT-proBNP to a reduced risk of hypertensive disorders, even after adjusting for age, ethnicity, BMI, and aspirin use<sup>30,31</sup>. Finally, a metaanalysis of 16 studies showed higher copeptin concentrations in PE women across all pregnancy trimesters<sup>32</sup>, results that are in line with our findings. Cardiac enzymes may offer valuable insights into cardiac remodeling in women at high-risk for PE during early pregnancy, with ongoing research examining their progression into the third trimester.

Regarding echocardiography, Melchiorre et al.<sup>33,34</sup> reported elevated indices of cardiac effort and LV mass, which may indicate LV remodeling as a compensatory mechanism to sustain myocardial contractility in cases of established PE. However, no

studies have been conducted in early pregnancy in patients at high-risk for PE. One study reports significant differences in LV mass between those at high-risk and those at low risk for PE, but without laboratory parameters or additional echocardiographic measures evaluated<sup>35</sup>. In this study, both groups showed increased cardiac output, a normal hemodynamic change during pregnancy. We did not find differences in systolic LV functions, including ejection fraction and strain parameters. However, we observed significant differences in LV and LA dimensions, with larger volumes in the high-risk group. LA performance is impacted by LV loading changes, leading to reduced filling during diastole<sup>36</sup>. Increased LA volume in the high-risk group may be an early indicator of subclinical diastolic dysfunction. This suggests disrupted CV adaptation to pregnancy, causing atrial remodeling; LA enlargement is linked to poor CV outcomes in non-pregnant populations<sup>37</sup>. Worse Doppler patterns (i.e. LV A wave, E/A ratio, and E' velocities) were seen in the high-risk group, although still within normal ranges. Consistently, significant differences were found in LV mass and relative wall thickness (RWT). LV mass increase is attributed to adaptation of the ventricle to higher blood volume and peripheral vascular resistance. This adaptation may be pronounced in the high-risk group due to maladaptation and relative peripheral vasoconstriction, as demonstrated by higher blood pressure and lower cardiac adaptation, indicated by lower NT-proBNP levels. Notably, increased LV mass and RWT, are indicative of concentric remodeling rather than the usual eccentric remodeling<sup>38</sup>. In pregnancy, clinical significance of this finding may be magnified due to increased hemodynamic stress and CV changes. There is a clear trend to reduced cardiac adaptation in individuals at high-risk for PE, despite no evident pathological findings. Lack of pre-pregnancy data to assess whether changes were present before pregnancy, or develop during early pregnancy, is a limitation of this study.

267 Correlation coefficients between biochemical markers and echocardiographic  
268 parameters show interesting trends, however our sample size limits drawing conclusions.  
269 Another limitation is that PlGF was not included in the initial first-trimester screening in  
270 all patients, but measured later in a two-step process for intermediate-risk patients. While  
271 this method performs similarly to universal screening, the delayed measurement of PlGF  
272 could reduce external validation in settings where it is routinely included in first-trimester  
273 screening for PE.

274 In conclusion, women at high-risk for PE show early pregnancy differences in lipid  
275 profile, angiogenic factors, cardiac biomarkers, and echocardiography, suggesting  
276 subclinical CV dysfunction. Findings include low NT-proBNP, increased LV mass,  
277 concentric remodeling, and impaired diastolic function, highlighting CV adaptation in PE  
278 pathogenesis. Further research should assess persistence of these changes, their evolution  
279 with aspirin prophylaxis, and impact on PE and long-term maternal CV health.

280 **TABLES**

281 **Table 1** Baseline characteristics of study population, according to risk for pre-eclampsia  
282 (PE)

| Characteristic                       | PE risk                   |                         | <i>P</i> |
|--------------------------------------|---------------------------|-------------------------|----------|
|                                      | High<br>( <i>n</i> = 257) | Low<br>( <i>n</i> = 92) |          |
| GA at scan (weeks)                   | 15 ± 1                    | 14 ± 1                  | 0.001    |
| Maternal age (years)*                | 34 ± 5                    | 34 ± 4                  | 0.712    |
| Nulliparous                          | 171/256 (66.8%)           | 4/91 (4.4%)             | 0.967    |
| Ethnicity                            |                           |                         |          |
| Caucasian                            | 164/255 (64.3%)           | 75/91 (82.4%)           | 0.001    |
| Latin-American                       | 80/255 (31.4%)            | 15/91 (16.5%)           | 0.001    |
| Asian                                | 7/255 (2.7%)              | 0 (0)                   | —        |
| Black                                | 4/255 (1.6%)              | 1/91 (1.1%)             | —        |
| Smoker                               | 5 (1.9%)                  | 3/89 (3.4%)             | 0.469    |
| Chronic hypertension                 | 21 (8.2%)                 | 0 (0%)                  | 0.005    |
| Pregestational diabetes              | 22/253 (8.7%)             | 3/89 (3.4%)             | 0.097    |
| Previous PE                          | 32/222 (14.2%)            | 2/82 (2.4%)             | 0.004    |
| BMI (kg/m <sup>2</sup> )†            | 28.4 ± 5.8                | 24.4 ± 3.6              | < 0.001  |
| Body surface area (m <sup>2</sup> )† | 1.81 ± 0.17               | 1.71 ± 0.14             | < 0.001  |
| Waist/hip ratio‡                     | 0.86 ± 0.07               | 0.85 ± 0.07             | 0.154    |
| Heart rate (bpm)‡                    | 83 ± 12                   | 81 ± 12                 | 0.099    |
| Systolic BP (mmHg)†                  | 118 ± 12                  | 111 ± 9                 | < 0.001  |
| Diastolic BP (mmHg)†                 | 71 ± 8                    | 65 ± 8                  | < 0.001  |
| Aspirin prophylaxis (%)              | 218/219 (99.5%)           | 1/219 (0.5%)            | < 0.001  |

283 Data are given as mean ± SD, *n/N* (%) or *n* (%). \**n* = 254 for high-risk and *n* = 91 for low-risk  
284 group. †*n* = 255 for high-risk and *n* = 91 for low-risk group. ‡*n* = 250 for high-risk and *n* = 91 for  
285 low-risk group. BMI, body mass index; bpm, beats per minute; BP, Blood Pressure.

**Table 2** Plasma biomarker profile of study population, according to risk for pre-eclampsia (PE)

| Parameter                          | PE risk       |               | <i>P</i> |
|------------------------------------|---------------|---------------|----------|
|                                    | High          | Low           |          |
| GA at blood draw (weeks)*          | 16 ± 2        | 15.9 ± 2      | 0.020    |
| Hemoglobin (g/dL)*                 | 12.1 ± 1.0    | 12.1 ± 0.9    | 0.540    |
| Hematocrit (%)*                    | 35 ± 2        | 35 ± 3        | 0.446    |
| Platelets ( $\times 10^9/L$ )†     | 264 ± 56      | 233 ± 52      | < 0.001  |
| Protein/creatinine ratio (mg/dL)*  | 201.7 ± 22.9  | 199.9 ± 0.0   | 1.000    |
| Total cholesterol (mg/dL)‡         | 203.3 ± 34.6  | 203.5 ± 37.1  | 0.951    |
| Low-density lipoproteins (mg/dL)§  | 107.3 ± 28.6  | 108.7 ± 31.4  | 0.714    |
| High-density lipoproteins (mg/dL)‡ | 69.7 ± 14.2   | 74.0 ± 14.6   | 0.020    |
| Triglycerides (mg/dL)‡             | 121 ± 1.46    | 98 ± 1.45     | < 0.001  |
| NT-proBNP (ng/L)¶                  | 51.0 ± 2.1    | 52 ± 2.0      | 0.008    |
| hs-cTnT (ng/L)**                   | 3.1 (3.0–4.2) | 3.0 (3.0–3.4) | 0.010    |
| Copeptin (pmol/L)††                | 3.4 ± 1.9     | 2.9 ± 1.4     | 0.044    |
| sFlt-1 (MoM)*                      | 0.95 ± 0.43   | 1.17 ± 0.59   | < 0.001  |
| PlGF (MoM)*                        | 0.89 ± 0.63   | 1.15 ± 0.58   | < 0.001  |
| sFlt-1/PlGF ratio (MoM)*           | 1.32 ± 1.19   | 1.14 ± 0.60   | 0.086    |

Data are given as mean ± SD or median (interquartile range). \**n* = 249 for high-risk and *n* = 86 for low-risk group. †*n* = 248 for high-risk and *n* = 86 for low-risk group. ‡*n* = 248 for high-risk and *n* = 85 for low-risk group. §*n* = 248 for high-risk and *n* = 84 for low-risk group. ¶*n* = 227 for high-risk and *n* = 82 for low-risk group. \*\**n* = 227 for high-risk and *n* = 83 for low-risk group. ††*n* = 144 for high-risk and *n* = 65 for low risk group. GA, gestational age; NT-proBNP, N-terminal pro-brain natriuretic peptide; sFlt-1, soluble fms-like tyrosine kinase-1; PlGF, placental growth factor; MoM, multiples of the median; hs-cTnT, high-sensitivity cardiac troponin T.

**Table 3** Maternal echocardiographic measures of study population, according to risk for pre-eclampsia (PE)

| Parameter                                 | High PE risk    |          | Low PE risk     |          | <i>P</i> |
|-------------------------------------------|-----------------|----------|-----------------|----------|----------|
|                                           | Mean $\pm$ SD   | <i>n</i> | Mean $\pm$ SD   | <i>n</i> |          |
| Electrocardiogram heart rate (bpm)        | 80 $\pm$ 12     | 174      | 76 $\pm$ 12     | 60       | 0.017    |
| Septal thickness (cm)                     | 0.90 $\pm$ 0.14 | 222      | 0.81 $\pm$ 0.12 | 87       | < 0.001  |
| LV posterior wall (cm)                    | 0.85 $\pm$ 0.18 | 221      | 0.77 $\pm$ 0.13 | 87       | < 0.001  |
| LV end-diastolic diameter (cm)            | 4.4 $\pm$ 0.4   | 221      | 4.3 $\pm$ 0.4   | 87       | 0.02     |
| LV end-diastolic volume (mL)              | 110 $\pm$ 23    | 215      | 102 $\pm$ 20    | 80       | 0.006    |
| LV ejection fraction (%)                  | 65.1 $\pm$ 4.8  | 216      | 64.2 $\pm$ 5.2  | 81       | 0.186    |
| LV mass (g)                               | 122 $\pm$ 32    | 223      | 104 $\pm$ 25    | 87       | < 0.001  |
| LV mass index (g/m <sup>2</sup> )         | 67 $\pm$ 15     | 223      | 61 $\pm$ 12     | 87       | < 0.001  |
| Relative wall thickness                   | 0.39 $\pm$ 0.08 | 220      | 0.36 $\pm$ 0.07 | 87       | 0.008    |
| Cardiac output (L/min)                    | 5.7 $\pm$ 1.2   | 199      | 5.2 $\pm$ 1.2   | 74       | 0.018    |
| Cardiac output index                      | 3.15 $\pm$ 0.73 | 201      | 3.08 $\pm$ 0.69 | 75       | 0.459    |
| LV E wave velocity (cm/s)                 | 94.3 $\pm$ 16.7 | 222      | 97.5 $\pm$ 19.7 | 87       | 0.182    |
| LV A wave velocity (cm/s)                 | 67.1 $\pm$ 15.1 | 221      | 59.6 $\pm$ 13.1 | 87       | < 0.001  |
| LV E/A ratio                              | 1.5 $\pm$ 0.4   | 220      | 1.7 $\pm$ 0.5   | 87       | < 0.001  |
| Lateral mitral annular velocity E' (cm/s) | 16.7 $\pm$ 3.4  | 224      | 17.7 $\pm$ 3.3  | 83       | 0.025    |
| Septal mitral annular velocity E' (cm/s)  | 10.9 $\pm$ 4.5  | 222      | 11.8 $\pm$ 2.5  | 84       | 0.020    |
| E/E' ratio                                | 9.2 $\pm$ 2.4   | 218      | 8.6 $\pm$ 2.2   | 84       | 0.024    |
| Left atrial biplanar volume (mL)          | 47 $\pm$ 13     | 161      | 43 $\pm$ 11     | 52       | 0.023    |
| RV end-diastolic diameter (cm)            | 3.1 $\pm$ 0.4   | 149      | 3.0 $\pm$ 0.4   | 54       | 0.690    |
| TAPSE (cm)                                | 2.39 $\pm$ 0.36 | 228      | 2.38 $\pm$ 0.34 | 85       | 0.620    |
| MAPSE (cm)                                | 1.68 $\pm$ 0.25 | 223      | 1.69 $\pm$ 0.28 | 84       | 0.882    |
| Global longitudinal strain (%)            | -21.7 $\pm$ 2.8 | 173      | -21.9 $\pm$ 3.1 | 63       | 0.782    |

LV: Left ventricle; RV: Right Ventricle; TAPSE, tricuspid annular plane systolic excursion; MAPSE, mitral annular plane systolic excursion; bpm, beats per minute.

## FIGURE LEGEND

**Figure 1** Flowchart summarizing inclusion of patients at high and at low risk for pre-eclampsia (PE) in study. ECG, echocardiographic.

## COLLABORATORS

Members of the AngioCor working group: Carla Domínguez-Gallardo, MD PhD; Carmen Garrido-Gimenez, MD PhD; Cristina Trilla, MD PhD; and Judit Platero, MD (Obstetrics and Gynecology Department, Hospital de la Santa Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain); Francesc Carreras-Costa, MD PhD; Alessandro Sionis, MD PhD; and David Belmar Clivillé, MD (Cardiology Department, Hospital de la Santa Santa Creu i Sant Pau) and Olga Sánchez-García, PhD (Women and Perinatal Health Research Group, Institut de Recerca (IR SANT PAU)); and Álvaro García-Osuna, MD PhD (Department of Clinical Biochemistry, Hospital de la Santa Creu i Sant Pau, Institut d'Investigació Biomèdica Sant Pau).

## ACKNOWLEDGMENTS

This study was conducted as part of doctoral research in Pediatrics, Obstetrics, and Gynecology at the Universitat Autònoma de Barcelona, Barcelona, Spain. The authors thank all the professionals who participated in the development of this project. Finally, we extend our gratitude to all the patients that gave us the opportunity to perform this project. Dr Cruz-Lemini was supported by Juan Rodés contract JR19/00047, Instituto de Salud Carlos III - Spanish Ministry of Health. This work was supported by public funds obtained in competitive calls with peer review (grant PI19/00702, Instituto de Salud Carlos III, Madrid, Spain), the Maternal and Child Health and Development Network (SAMID, RD16/0022/0015, Instituto de Salud Carlos III, Madrid, Spain), the Spanish Clinical Research and Clinical Trials Platform, SCReN (Spanish Clinical Research Network), funded by the ISCIII-General Subdirectorate for Evaluation and Promotion of Research, through project PT13/0002/0028, integrated in the 2013-2016 R + D + I State Plan and co-financed by and the European Regional Development Fund (FEDER); and by the Primary Care Interventions to Prevent Maternal and Child Chronic Diseases of Perinatal and Developmental Origin Network (RICORS, RD21/0012/0001, Instituto de

Salud Carlos III, Madrid, Spain), funded by the Recovery, Transformation and Resilience Plan 2017-2020, ISCIII, and by the European Union – Next Generation EU. Funding sources were not involved in study design, collection, analysis and interpretation of data, or in the writing of this manuscript.

## DISCLOSURES

Elisa Llurba declares lecture fees from Roche Diagnostics. The remaining authors have nothing to disclose.

## REFERENCES

1. Wang H, Abbas KM, Abbasifard M, et al. Global age-sex-specific fertility, mortality, healthy life expectancy (HALE), and population estimates in 204 countries and territories, 1950–2019: a comprehensive demographic analysis for the Global Burden of Disease Study 2019. *The Lancet*. 2020;396(10258):1160-1203. doi:10.1016/S0140-6736(20)30977-6
2. Vogel B, Acevedo M, Appelman Y, et al. The Lancet women and cardiovascular disease Commission: reducing the global burden by 2030. *The Lancet*. 2021;397(10292):2385-2438. doi:10.1016/S0140-6736(21)00684-X
3. Cong J, Fan T, Yang X, Shen J, Cheng G, Zhang Z. Maternal cardiac remodeling and dysfunction in preeclampsia: a three-dimensional speckle-tracking echocardiography study. *Int J Cardiovasc Imaging*. 2015;31(7):1361-1368. doi:10.1007/s10554-015-0694-y
4. Trilla C, Luna C, De León Socorro S, et al. First-Trimester Sequential Screening for Preeclampsia Using Angiogenic Factors: Study Protocol for a Prospective, Multicenter, Real Clinical Setting Study. *Front Cardiovasc Med*. 2022;9. doi:10.3389/fcvm.2022.931943
5. Opichka MA, Rappelt MW, Gutterman DD, Grobe JL, McIntosh JJ. Vascular Dysfunction in Preeclampsia. *Cells*. 2021;10(11):3055. doi:10.3390/cells10113055
6. Mulder E, de Haas S, Mohseni Z, et al. Cardiac output and peripheral vascular resistance during normotensive and hypertensive pregnancy – a

- systematic review and meta-analysis. *BJOG*. 2022;129(5):696-707. doi:10.1111/1471-0528.16678
7. Borges VTM, Zanati SG, Peraçoli MTS, et al. Maternal left ventricular hypertrophy and diastolic dysfunction and brain natriuretic peptide concentration in early- and late-onset pre-eclampsia. *Ultrasound in Obstetrics & Gynecology*. 2018;51(4):519-523. doi:10.1002/uog.17495
  8. Chen CW, Jaffe IZ, Karumanchi SA. Pre-eclampsia and cardiovascular disease. *Cardiovasc Res*. 2014;101(4):579-586. doi:10.1093/cvr/cvu018
  9. Zhang J, Klebanoff MA, Roberts JM. Prediction of adverse outcomes by common definitions of hypertension in pregnancy. *Obstetrics & Gynecology*. 2001;97(2):261-267. doi:10.1016/S0029-7844(00)01125-X
  10. De Haas S, Ghossein-Doha C, Geerts L, van Kuijk SMJ, van Drongelen J, Spaanderman MEA. Cardiac remodeling in normotensive pregnancy and in pregnancy complicated by hypertension: systematic review and meta-analysis. *Ultrasound in Obstetrics & Gynecology*. 2017;50(6):683-696. doi:10.1002/uog.17410
  11. Reddy M, Wright L, Rolnik DL, et al. Evaluation of Cardiac Function in Women With a History of Preeclampsia: A Systematic Review and Meta-Analysis. *J Am Heart Assoc*. 2019;8(22). doi:10.1161/JAHA.119.013545
  12. Ullmo J, Cruz-Lemini M, Sánchez-García O, et al. Cardiac dysfunction and remodeling regulated by anti-angiogenic environment in patients with preeclampsia: the ANGIOCOR prospective cohort study protocol. *BMC Pregnancy Childbirth*. 2021;21(1):816. doi:10.1186/s12884-021-04263-w
  13. Trilla C, Mora J, Ginjaume N, et al. Reduction in Preterm Preeclampsia after Contingent First-Trimester Screening and Aspirin Prophylaxis in a Routine Care Setting. *Diagnostics*. 2022;12(8):1814. doi:10.3390/diagnostics12081814
  14. Lang RM, Badano LP, Mor-Avi V, et al. Recommendations for Cardiac Chamber Quantification by Echocardiography in Adults: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *Journal of the American Society of Echocardiography*. 2015;28(1):1-39.e14. doi:10.1016/j.echo.2014.10.003
  15. Nagueh SF, Smiseth OA, Appleton CP, et al. Recommendations for the Evaluation of Left Ventricular Diastolic Function by Echocardiography: An Update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging*. 2016;17(12):1321-1360. doi:10.1093/ehjci/jew082

16. Spradley FT. Metabolic abnormalities and obesity's impact on the risk for developing preeclampsia. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2017;312(1):R5-R12. doi:10.1152/ajpregu.00440.2016
17. Persson M, Cnattingius S, Wikström AK, Johansson S. Maternal overweight and obesity and risk of pre-eclampsia in women with type 1 diabetes or type 2 diabetes. *Diabetologia*. 2016;59(10):2099-2105.
18. Alston MC, Redman LM, Sones JL. An Overview of Obesity, Cholesterol, and Systemic Inflammation in Preeclampsia. *Nutrients*. 2022;14(10):2087. doi:10.3390/nu14102087
19. Winkler K, Wetzka B, Hoffmann MM, et al. Triglyceride-Rich Lipoproteins Are Associated with Hypertension in Preeclampsia. *J Clin Endocrinol Metab*. 2003;88(3):1162-1166. doi:10.1210/jc.2002-021160
20. Stadler JT, Scharnagl H, Wadsack C, Marsche G. Preeclampsia Affects Lipid Metabolism and HDL Function in Mothers and Their Offspring. *Antioxidants*. 2023;12(4):795. doi:10.3390/antiox12040795
21. Oğlak SC, Tunç Ş, Ölmez F. First Trimester Mean Platelet Volume, Neutrophil to Lymphocyte Ratio, and Platelet to Lymphocyte Ratio Values Are Useful Markers for Predicting Preeclampsia. *Ochsner Journal*. 2021;21(4):364-370. doi:10.31486/toj.21.0026
22. Verlohren S, Herraiz I, Lapaire O, et al. New Gestational Phase-Specific Cutoff Values for the Use of the Soluble fms-Like Tyrosine Kinase-1/Placental Growth Factor Ratio as a Diagnostic Test for Preeclampsia. *Hypertension*. 2014;63(2):346-352. doi:10.1161/HYPERTENSIONAHA.113.01787
23. Bates DO. An unexpected tail of VEGF and PlGF in pre-eclampsia. *Biochem Soc Trans*. 2011;39(6):1576-1582. doi:10.1042/BST20110671
24. Garrido-Gimenez C, Mendoza M, Cruz-Lemini M, et al. Angiogenic Factors and Long-Term Cardiovascular Risk in Women That Developed Preeclampsia During Pregnancy. *Hypertension*. 2020;76(6):1808-1816. doi:10.1161/HYPERTENSIONAHA.120.15830
25. Pasupathi P, Manivannan U, Manivannan P, Deepa M. Cardiac troponins and oxidative stress markers in non-pregnant, pregnant and preeclampsia women. *Bangladesh Med Res Counc Bull*. 2010;36(1):4-9. doi:10.3329/bmrcb.v36i1.4806

26. Bacmeister L, Goßling A, Buellesbach A, et al. High-Sensitivity Cardiac Troponin I Enhances Preeclampsia Prediction Beyond Maternal Factors and the sFlt-1/PlGF Ratio. *Circulation*. 2024;149(2):95-106. doi:10.1161/CIRCULATIONAHA.123.066199
27. Ouzounian JG, Elkayam U. Physiologic Changes During Normal Pregnancy and Delivery. *Cardiol Clin*. 2012;30(3):317-329. doi:10.1016/j.ccl.2012.05.004
28. Maalouf R, Bailey S. A review on B-type natriuretic peptide monitoring: assays and biosensors. *Heart Fail Rev*. 2016;21(5):567-578. doi:10.1007/s10741-016-9544-9
29. Furenäs E, Eriksson P, Wennerholm UB, Dellborg M. Pregnancy in a healthy population: dynamics of NTproBNP and hs-cTroponin T. *Open Heart*. 2020;7(2):e001293. doi:10.1136/openhrt-2020-001293
30. Hauspurg A, Marsh DJ, McNeil RB, et al. Association of N-Terminal Pro–Brain Natriuretic Peptide Concentration in Early Pregnancy With Development of Hypertensive Disorders of Pregnancy and Future Hypertension. *JAMA Cardiol*. 2022;7(3):268. doi:10.1001/jamacardio.2021.5617
31. Takahashi M, Suzuki L, Takahashi N, et al. Early-pregnancy N-terminal pro-brain natriuretic peptide level is inversely associated with hypertensive disorders of pregnancy. *Res Sq*. Published online March 11, 2024:1-19.
32. Bellos I, Pergialiotis V, Papapanagiotou A, Loutradis D, Daskalakis G. Association between serum copeptin levels and preeclampsia risk: A meta-analysis. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2020;250:66-73. doi:10.1016/j.ejogrb.2020.04.051
33. Melchiorre K, Sutherland GR, Baltabaeva A, Liberati M, Thilaganathan B. Maternal Cardiac Dysfunction and Remodeling in Women With Preeclampsia at Term. *Hypertension*. 2011;57(1):85-93. doi:10.1161/HYPERTENSIONAHA.110.162321
34. Melchiorre K, Sharma R, Thilaganathan B. Cardiovascular Implications in Preeclampsia. *Circulation*. 2014;130(8):703-714. doi:10.1161/CIRCULATIONAHA.113.003664
35. Ridder A, O'Driscoll J, Khalil A, Thilaganathan B. Routine first-trimester pre-eclampsia screening and maternal left ventricular geometry. *Ultrasound in Obstetrics & Gynecology*. 2024;63(1):75-80. doi:10.1002/uog.26306

36. Valensise H, Novelli GP, Vasapollo B, et al. Maternal Diastolic Dysfunction and Left Ventricular Geometry in Gestational Hypertension. *Hypertension*. 2001;37(5):1209-1215. doi:10.1161/01.HYP.37.5.1209
37. Pritchett AM, Mahoney DW, Jacobsen SJ, Rodeheffer RJ, Karon BL, Redfield MM. Diastolic dysfunction and left atrial volume. *J Am Coll Cardiol*. 2005;45(1):87-92. doi:10.1016/j.jacc.2004.09.054
38. Afari HA, Davis EF, Sarma AA. Echocardiography for the Pregnant Heart. *Curr Treat Options Cardiovasc Med*. 2021;23(8):55. doi:10.1007/s11936-021-00930-5

## **SUPPORTING INFORMATION ON THE INTERNET**

The following supporting information may be found in the online version of this article:

**Appendix S1** Sample size estimation and multivariate distribution model for estimation of risk of pre-eclampsia (PE)

**Table S1** Correlations between maternal cardiovascular biomarkers and echocardiographic measures evaluated in study population