

ORIGINAL RESEARCH

Alignment between the research question, design, and terminology is required in manual therapy trials: a methodological study

Sonia Roura^a, Gerard Alvarez^{b,c,*}, David Hohenschurz-Schmidt^d, Ivan Solà^{b,e,f},
Rodrigo Núñez-Cortés^g, Javier Bracchiglione^{b,e,h}, Carles Fernández-Jané^{c,i,j}, Jules Phalip^{k,l},
Ignasi Gich^{e,m}, Mercè Sitjà-Rabert^{c,i}, Gerard Urrútia^{b,e,f}

^aPhD Student in Biomedical Research Methodology and Public Health in the Medical Department of the Universitat Autònoma de Barcelona, Barcelona, Spain

^bIberoamerican Cochrane Centre, Institut de Recerca Sant Pau (IR SANT PAU), Barcelona, Spain

^cFacultat de Ciències de la Salut Blanquerna, Universitat Ramon Llull, Barcelona, Spain

^dImperial Clinical Trials Unit, School of Public Health, Imperial College, London, United Kingdom

^eCentro de Investigación Biomédica en Red de Epidemiología y Salud Pública (CIBERESP), Madrid, Spain

^fUniversitat Autònoma de Barcelona, Bellaterra, Spain

^gFaculty of Medicine, Department of Physical Therapy, University of Chile, Santiago, Chile

^hInterdisciplinary Centre for Health Studies (CIESAL), Universidad de Valparaíso, Viña del Mar, Chile

ⁱGlobal Research on Wellbeing (GRoW) Research Group, Universitat Ramon Llull, Barcelona, Spain

^jTecnocampus, Universitat Pompeu Fabra, Mataró-Maresme, Barcelona, Spain

^kFaculté de médecine, Fondation ANALGESIA, Clermont-Ferrand, France

^lCHU Clermont-Ferrand, Inserm 1107 Neuro-Dol, Service de pharmacologie médicale, Université Clermont Auvergne, Clermont-Ferrand, France

^mInstitut de Recerca Sant Pau (IR SANT PAU), Barcelona, Spain

Accepted 30 April 2025; Published online 8 May 2025

Abstract

Objectives: The study aimed to assess the design, reporting, and risk of bias in effectiveness trials (ETs) in manual therapy (MT), comparing pragmatic with nonpragmatic trials and trials with and without placebo controls.

Study Design and Setting: We searched MEDLINE and the Cochrane Central Register of Controlled Trials for randomized controlled trials with the term “effectiveness” in the title or abstract in the field of MT from inception to January 2024. Two independent reviewers extracted data on specific study characteristics, their reporting, and risk of bias and assessed them using the PRagmatic Explanatory Continuum Indicatory Summary-2 tool (PRECIS-2). Descriptive analysis using frequencies and percentages and a relation analysis between PRECIS-2 scores and specific study characteristics were performed.

Results: Of the 113 trials, 39 were self-labeled as pragmatic, 39 used usual care or other interventions, and 35 used placebo controls. ETs have increased in recent years. They are moderately pragmatic, whether they are self-described as pragmatic or not, and whether they use a placebo control group or not. Despite their aim to resemble clinical practice, the pragmatic features of these trials are often unclear. Pragmatic features are common in trials’ eligibility, recruitment, and outcome domains, but intervention and setting are rated as very explanatory. Compared to self-labeled pragmatic trials, “effectiveness” trials are less likely to follow reporting guidelines.

Conclusion: The term effectiveness is used in MT trials independently of its pragmatic connotations. Using a placebo or active control does not modify the pragmatic attitude of the ETs. © 2025 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Effectiveness clinical trials; Musculoskeletal manipulations; Randomized controlled trials; Manual therapy; Generalizability; PRECIS-2

1. Introduction

Both efficacy and effectiveness randomized clinical trials (RCTs) are essential for evaluating interventions [1,2]. Efficacy trials are conducted under controlled conditions to explore the causal relationship between interventions

* Corresponding author. Iberoamerican Cochrane Center, C. Sant Antoni Maria Claret 167, Pavelló 18, Planta 008025, Catalunya, Barcelona, Spain.

E-mail address: gerardalv@gmail.com (G. Alvarez).

What is new?

Key findings

- An increasing proportion of trials in manual therapy (MT) are aimed at evaluating effectiveness, as a primary aim.
- ETs in MT have a moderately pragmatic attitude regardless of whether they self-label themselves as pragmatic or not and whether they use a placebo control group or not. Most ETs aim to resemble clinical practice, but pragmatic features are not shown within the trial.
- Reporting ETs is often inadequate, exacerbated when studies do not call themselves pragmatic. It is also poorer when “usual care” or other interventions are used as a control intervention rather than a placebo control.
- Although selection bias is lower when studies call themselves pragmatic, detection bias is lower when using a placebo control than another intervention or “usual care”.

What this adds to what is known?

- The external validity of ETs is often compromised.
- The use of the term efficacy/effectiveness seems unclear among the MT research community, which uses the term effectiveness in efficacy designs.

What is the implication and what should change now?

- A thorough justification for the trial’s aim, rationale, and design can improve study quality.
- Modifications to trial methodologies are essential to enhance their clinical applicability in the MT field.
- Clear and comprehensive reporting is vital for understanding trial quality, mitigating biases, and improving the applicability of findings for clinical practice.

and their physiological effects (explanatory attitude) [3]. Conversely, effectiveness trials (ETs) aim to inform health care professionals by comparing interventions under real-world conditions (pragmatic attitude) [3,4]. The terms commonly used in the literature to designate an article as explanatory are efficacy trials and “pragmatic”, or “naturalistic”, to designate them as pragmatic, or referring to the “effectiveness” of an intervention [3,5–9]. These trial types are better understood as a continuum rather than a strict dichotomy, as they often share overlapping

methodologies and conceptual frameworks [4,10–13]. Furthermore, the terms “attitude” and “study intent” must be differentiated. The chosen “attitude” is the overarching orientation of the research, either choosing a more pragmatic or a more efficacy-focused orientation. In contrast, the “study intent” is the goal pursued within that orientation, in other words, how the author states and justifies the specific objective or research question [3,10], such as “determining if a specific manual therapy (MT) technique is correlated with a specific heart rate variability change (explanatory attitude)” vs “evaluating if adding MT to the standard treatment for obstructive pulmonary infection outperforms the standard treatment alone in real-world conditions (pragmatic attitude)”. ETs are necessary because they focus on increasing research’s relevance and applicability to clinical practice and policy decision-making. Thus, they might ensure external validity but also use methods that enhance internal trial validity, such as randomization and treatment fidelity monitoring.

In MT, efficacy and ETs are well-documented [1,14]. However, concerns have been raised regarding their overall methodological rigor due to internal validity issues and reporting deficiencies [14–22]. Although the term “effectiveness” is commonly used in MT trials [23–26], a systematic review revealed a lack of pragmatism in many cases [1]. This has led to concerns about whether these trials accurately simulate clinical practice and deliver meaningful insights for health care decisions [1,27,28]. The inconsistent use of terminology has implications for research, potentially affecting the interpretation of trial outcomes and influencing clinical guideline recommendations [29,30].

Research shows that MT trials often lean toward explanatory designs rather than pragmatic approaches [1]. Previous findings from the author group confirmed that self-labeled MT pragmatic RCTs (note that to facilitate readability, those will be named pragmatic trials (PTs) throughout the manuscript) have not increased in recent years, unlike in other disciplines [1,10,31–33]. Furthermore, their pragmatic attitude remains moderate, and better internal validity and reporting are needed [33]. The misuse of “efficacy” and “effectiveness” as interchangeable terms has resulted in research designs often inconsistent with their stated objectives [4,30,34,35]. This issue is particularly prevalent in studies focused on the clinical effects of interventions with limited efficacy evidence but widespread clinical use, such as MT [31,36].

Given the conceptual uncertainty regarding terminology, this study aimed to compare effectiveness MT trials self-labeled as pragmatic with those that are not. A substantial body of research exists that neither applies well in real-world settings nor is designed with the internal validity required to be considered efficacy studies. By assessing the methodological differences, the researchers sought to clarify whether there are critical design disparities and promote more accurate reporting and understanding of trial

characteristics. This research underscores the need for trial design and terminology clarity to strengthen academic inquiry and clinical guideline development. Establishing consistent definitions and methodological practices will help improve the quality and reliability of MT trials, ultimately benefiting clinical practice and health care decision-making.

2. Methods

2.1. Protocol registration

We conducted a methodological review of ETs on MT interventions, following the guidelines of the Cochrane Handbook for systematic reviews of interventions [37] and reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines (PRISMA) [38] ([Supplementary file 1](#)). The study was prospectively registered on the Open Science Framework (DOI 10.17605/OSF.IO/WKEPZ).

2.2. Eligibility criteria

We included published RCTs with the terms “effectiveness” (related to an intervention), “pragmatic” (methodological design), or “naturalistic” (methodological design) in either the title or abstract [6,7]. Eligible references had to include either a manual technique or a combination of manual techniques (soft tissue techniques, joint mobilization or manipulation, massage, myofascial release, nerve manipulation, strain/counterstrain, and acupressure). There were no restrictions regarding population, comparator groups or outcome measures. Exclusion criteria included experimental interventions using tools, devices (electrotherapy, kinesio taping, dry needling, and acupuncture), drugs, active exercises, or a combination of therapies without MT intervention. We also excluded non-English articles, protocols, and poster/conference presentations.

2.3. Search strategy

We conducted a comprehensive search of MEDLINE and the Cochrane Central Register of Controlled Trials from inception to January 2024, using a search strategy that combined controlled vocabulary with relevant Medical Subject Headings terms in the field of MT and the design of interest (see [Supplementary file 2](#)). The strategy was developed by an expert methodologist on our research team (I.S.).

2.4. Study selection

The records were uploaded onto the Rayyan software (www.rayyan.ai) [39]. On deduplication, the references were screened by two independent reviewers (S.R. and G.A.) based on the title and summary, resolving disagreements through discussion.

The references were initially classified into two groups: group 1 consisting of ETs labeled as “pragmatic” in the title and/or abstract (PTs), and group 2 comprising ETs that were not self-labeled as pragmatic. For further analysis, we split the ET group into studies that used a placebo/sham control group (effectiveness Randomised Controlled Trial using a placebo/sham comparator [ETpl])—GROUP 2A—and those that used an accepted or established comparator (effectiveness Randomised Controlled Trial using an active comparator [ETac])—GROUP 2B—based on title/abstract information, resulting in three groups (groups 1, 2A, and 2B). We determined the sample of studies to be included in each group according to the PTs found. As we found many more ETs than PTs, we randomized the ETs by matching their number to the number of PTs found.

2.5. Data collection process

Two reviewers, S.R. and an additional member from G.A., R.N., J.B., D.H., C.F., and J.P., gathered data from the included studies and resolved disagreements through consensus. The involvement of a third party was not required to resolve any discrepancies.

Before the review process, the team met with Dr Kirsty Loudon, the developer of the pragmatic-explanatory continuum indicatory summary (PRECIS)-2 tool [40]. It was deemed suitable to involve a member of the PRECIS-2 tool authors, given that PRECIS-2 was conceptualized as a tool for trial designers rather than as a method for retrospective assessment of pragmatism. The objective was to ensure that the method was used in the most reliable manner. Three articles were examined, compared, and discussed against Dr Loudon’s criteria.

2.6. Data items

Three authors (S.R., G.A., and G.U.) designed a data extraction form. It included bibliometric identification elements, the intent of the trial, the rationale of the intervention given by the authors, the experimental and control interventions, the limitations reported by the authors, the PRECIS-2 tool assessment, the Consolidated Standards of Reporting Trials (CONSORT) reporting assessment, and the risk of bias (RoB) assessment. Information about the data extraction form and the methods used to assess each item are included in [Supplementary file 3](#) and reported elsewhere [33].

The data extraction form was piloted to assess reviewers’ discrepancies. Subsequently, all the suggestions were incorporated. Finally, a guideline for reviewers was provided to the team ([Supplementary file 3](#)). When information about a PRECIS-2 domain was not found within the trial publication, it was rated as “blank”

2.7. Data analysis

A descriptive analysis of the categorical variables was conducted, wherein results were presented as relative and

absolute frequencies. Likewise, means and SDs were used to describe quantitative variables. Median and IQRs were applied for ordinal variables. Meanwhile, the PRECIS-2 wheel was used to present the pragmatic attitude of each article graphically. Furthermore, variables were compared using analysis of variance for quantitative variables, Chi-square for categorical variables and Kruskal–Wallis for ordinal variables. According to previous research [41–43], we used a 9–45 point scale to evaluate the total score PRECIS-2, apart from the PRECIS average. Ratings of 9–22 were described as slightly pragmatic, 23–34 moderately pragmatic, and ≥ 35 very pragmatic. We conducted post hoc analysis using the Scheffé test for analysis of variance and the Mann–Whitney test for Kruskal–Wallis. The Spearman test was performed to analyze the correlation between PRECIS-2 and some descriptive characteristics. The significance level was set at 5%

($\alpha = 0.05$). All data were analyzed using IBM-SPSS software (V26.0).

3. Results

3.1. Study selection

The search identified 3553 articles. After screening the title, abstract, and full-text availability, the final sample comprised 306 studies. Among these, 39 were identified as self-labeled pragmatic in the title and/or abstract, whereas 267 were ETs not self-labeled as pragmatic. We did not identify any articles in MT with the term naturalistic. Of the 267 ETs, 35 were ETpl, whereas the remaining 232 used other accepted or established interventions (ETac). To balance and compare the three groups, the 232 ETac were randomized using the Microsoft Excel random generator to get a group of




Figure 1. PRISMA flow diagram. MT manual therapy, PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses, RCT, randomized controlled trial.

39 trials (the same number of trials as in the PTs group), resulting in a total sample for this review of 113 studies. The PRISMA diagram is shown in [Figure 1](#).

3.2. Study characteristics of the sample

[Table 1](#) summarizes the main characteristics of our sample. A detailed table with all included studies and some additional characteristics can be found in [Supplementary File 4](#). The average number of participants was significantly higher in PTs (mean = 169) compared to ETac (mean = 65) and ETpl (mean = 70). In PTs, the most frequent intervention studied was a nonprotocolized combination of techniques (54%; 21/39), whereas in ETac, it was a protocol of a combination of techniques (51%; 20/39) and a protocol of an isolated technique in 49% (17/35) of the ETpl. No studies blinded therapists. Blinding of external assessors was more common in ETpl (74%) compared to PTs (69%) and ETac (59%). Multicentric settings were more prevalent in PTs (56%), whereas ETs primarily used unicentric settings.

Specific characteristics of PTs were reported elsewhere [33,44]. Although “pragmatic” was not mentioned in titles or abstracts, 64% of ETac studies claimed a pragmatic approach, compared to 31% of ETpl. Common self-reported study limitations included participant blinding (14/17/12), generalizability (13/11/11), and sample size (11/12/14). Inadequate control was frequent in PTs (13) and ETac (8) but rare in ETpl (4). Study length limitations were prevalent in ETs. [Figure 2](#) illustrates these limitations across study types.

3.3. PRECIS assessment

[Table 2](#) presents mean PRECIS-2 scores, revealing that 60% of studies in each group had a moderately pragmatic attitude. [Figure 3](#) illustrates the PRECIS-2 wheel based on study group.

[Table 3](#) shows PRECIS-2 rates by study type. Most domains in PTs, except domain 6 (adherence), were pragmatic in over 50% of studies. For ETs, domains 1 (eligibility), 2 (recruitment), and 8 (outcome measure) were pragmatic in more than 50% of cases. Domain 3 (setting) was largely explanatory in ETs (82%), whereas domain 5 (flexibility) was pragmatic in over 50% of PTs but explanatory in most ETs.

3.4. CONSORT assessment

Almost one-third of the trials (37/113) were reported according to the CONSORT statement: 41% (16/39) of the PTs, 31% (12/39) of the ETac, and 26% (12/35) of the ETpl.

One in four trials reported details related to more than 70% of CONSORT items (13/39 of the PTs, 9/39 of the ETac, and 13/32 of the ETpl), between 50% and 70% in 41% of the trials (46/113) and lower than 50% of the

CONSORT items in 28% of the sample (32/113). [Figure 4](#) shows the percentage of compliance with CONSORT items depending on the study type.

[Supplementary file 5](#) shows item compliance across study types. The data analysis from the pre-CONSORT and post-CONSORT periods indicates that the reporting of some items improved after the publication of the CONSORT tool. However, this improvement does not impact each item's final analyzed reporting result ([Supplementary file 6](#)). The items better reported (more than 85% of the trials) in all study types were structured summary, background, objectives, eligibility criteria, outcomes, statistical methods, why the trial stopped, and baseline data table. Pragmatic RCTs had more items reported in more than 85% of the trials (sequence generation, participants flow, numbers analyzed, and effect size) compared with ETs that were not self-labeled pragmatic.

Intervention descriptions were adequate in over 80% of studies (95/113). Still, the authors did not sufficiently report details of the intervention, the standardization or individualization of the intervention and the additional sources to resemble clinical practice. Item 5 b (Description of the different components of the interventions and, when applicable, description of the procedure for tailoring the interventions to individual participants) was better reported in the ETpl group (77%) than the ETac (56%) and the PTs groups (54%). Moreover, less than 11% (12/113) of the trials reported whether and how the adherence of participants or care providers was assessed.

How sample size was determined, blinding, and participants' flow were reported less optimally in ETac (46%, 56%, 67%, respectively), compared to ETpl (66%, 83%, 83%, respectively) and PTs (74%, 77%, 87%, respectively) groups. All the items regarding blinding were better reported when using a placebo/sham control group. Items regarding randomization sequence generation, allocation concealment, implementation, and generalizability of the trial results were reported better in PTs than in ETs.

Furthermore, items “harms” and “trial registration number”, and “where the protocol can be assessed” were not reported in the 49% (55/113) of the sample.

3.5. Risk of bias

Selection bias (random sequence generation and allocation concealment) was lower in PTs than in ETs. Detection bias was higher in PTs and ETac than ETpl. A table comparing the RoB between types of studies is presented in [Supplementary file 7](#).

Please refer to [Supplementary file 8](#) to compare the studies with a low vs high/unclear RoB according to compliance with CONSORT items. The results showed that the percentage of compliance with CONSORT items was higher in the trials with low RoB in the following domains: random sequence generation, allocation

Table 1. Main characteristics of the sample

Characteristics of the sample	PTs (N = 39)	ETac (N = 39)	ETpl (N = 35)
Number of participants (Mean)	169 (SD = 151)	65 (SD = 58)	70 (SD = 44)
Number of participants	Percent (n ^b)		
Intervention			
Combination of nonprotocolized techniques	54% (21)	10% (4)	9% (3)
Protocol of a combination of techniques	13% (5)	51% (20)	23% (8)
Isolated nonprotocolized technique	15% (6)	8% (3)	8% (3)
Protocol of an isolated technique	5% (2)	18% (7)	49% (17)
Combination of nonprotocolized therapies	10% (4)	3% (1)	3% (1)
Protocol of a combination of therapies	3% (1)	10% (4)	9% (3)
Control intervention			
2 arms			
Test treatment vs other active intervention	31% (12)	46% (18)	0% (0)
Test treatment vs placebo	3% (1)	0% (0)	69% (24)
Test treatment vs usual care	33% (13)	38% (15)	0% (0)
Test treatment vs no intervention	13% (5)	5% (2)	0% (0)
3 arms			
Test treatment vs 2 other active interventions	5% (2)	3% (1)	0% (0)
Test treatment vs 1 other active intervention and 1 placebo	2% (1)	3% (1)	11% (4)
Test treatment vs 1 active intervention and 1 usual care	10% (4)	3% (1)	0% (0)
Test treatment vs 1 usual care and 1 placebo	2% (1)	0% (0)	17% (6)
Test treatment vs 1 other active intervention and 1 no intervention	2% (1)	0% (0)	0% (0)
4 arms			
Test treatment vs 1 other active intervention, 1 no intervention and 1 placebo	0% (0)	3% (1)	3% (1)
Blinding (yes)			
Participants	8% (3)	8% (3)	20% (7)
Therapists	0% (0)	0% (0)	0% (0)
External assessors to providers and patients	69% (27)	59% (23)	74% (26)
Statistician	43% (17)	18% (7)	20% (7)
Rationale			
Yes (rationale provided)	64% (25)	64% (25)	63% (22)
Comparative effectiveness data	61%	38%	31%
Mechanistic experiments	36%	54%	63%
Highly sham-controlled	3%	8%	6%
No (no rationale provided)	21% (8)	28% (11)	26% (9)
Unclear	13% (5)	8% (3)	11% (4)
Funding			
Yes	67% (26)	21% (8)	37% (13)
Follow-up			
No follow-up	18% (7)	23% (9)	28% (9)
<2 wk	0% (0)	13% (5)	3% (1)

(Continued)

Table 1. Continued

Characteristics of the sample	PTs (N = 39)	ETac (N = 39)	ETpl (N = 35)
2–4 wk	2% (1)	8% (3)	3% (1)
4–12 wk	10% (4)	20% (8)	34% (12)
3–6 mo	18% (7)	15% (6)	20% (7)
6–12 mo	20% (8)	15% (6)	9% (3)
> 1 y	26% (10)	5% (2)	3% (1)
Individualized ^a	5% (2)	0% (0)	0% (0)
Setting			
Multicentric	56% (22)	8% (3)	20% (7)
Unicentric	33% (13)	87% (34)	77% (27)
Unclear reported	10% (4)	5% (2)	3% (1)

ETac, effectiveness trials active comparator; ETpl, effectiveness trials placebo comparator; PTs, pragmatic trials.

^a Trials assessing how many weeks patients remained pain-free.

^b Number of studies = 113.

concealment, incomplete outcome data, and selective reporting.

4. Discussion

This review compared the design, reporting, and bias risk of MT effectiveness RCTs, comparing trials self-labeled as pragmatic or not by the study authors, and using a placebo control or an active comparator. The results

showed a moderate pragmatic attitude regardless of labeling or placebo use. Although many trials aim to emulate clinical practice, designs often lack pragmatic methodology. The review also found that reporting was generally inadequate, especially when not self-labeled pragmatic or when using usual care/other interventions instead of placebo/sham controls as comparators. The findings highlight areas for improvement in designing and reporting MT effectiveness RCTs to better align with clinical practice and enhance methodological rigor.




Figure 2. Comparison of limitations reported between studies. ETac, effectiveness trials active comparator; ETpl, effectiveness trials placebo comparator; PTs, pragmatic trials.

Table 2. Mean PRECIS-2 scores between studies and pragmatic attitude

Type of study	Mean PRECIS score (SD)	Slightly pragmatic	Moderate pragmatic	Highly pragmatic
PTs	3,5 (0,6)	3% (1)	61% (24)	36% (14)
ETac	2,8 (0,6)	33% (13)	59% (23)	8% (3)
ETpl	2,8 (0,6)	34% (12)	63% (22)	3% (1)

ETac, effectiveness trials active comparator; ETpl, effectiveness trials placebo/sham comparator; PTs, pragmatic trials.

4.1. Designing MT effectiveness RCTs and correct use of the terminology

Although the growing interest in ETs designed to reflect clinical practice, probably a response to the demand for real-world assessments of interventions, the distinction between efficacy and ETs designs remains unclear [4]. Two key factors define a trial's position on the efficacy-effectiveness continuum: the study's intention [10] and the rationale for its design [4]. The intention determines the research question, affects the design, and influences the outcomes and applicability of the study [4,10]. When giving a study's rationale, authors should determine if sufficient efficacy trials justify evaluation in more realistic environments [4]. Our analysis revealed that 25% of trials lacked this rationale, and those that did often relied on mechanistic experiments rather than controlled placebo/sham trials. As previous studies suggest, our findings reveal a misalignment between trial objectives and methodologies, warranting further investigation into better design practices [1].

4.2. Pragmatic attitude and methodological characteristics of effectiveness MT RCTs

Effectiveness and PTs are terms that can be used interchangeably in many contexts to refer to clinical trials designed to answer whether an intervention works under routine health care conditions [9]. However, ETs scored lower on PRECIS-2 (2.8) than self-labeled PTs (3.5),

suggesting a greater emphasis on pragmatic design when explicitly labeled. In our sample, in contrast to PTs, ETs were primarily explanatory in the setting and intervention domains and had smaller sample sizes.

4.2.1. Experimental intervention

The most common interventions in ETs were protocolized combinations of techniques (ETac) and isolated techniques (ETpl), placing these studies closer to the efficacy pole of the continuum. Adopting nonprotocolized, patient-centered interventions could better reflect clinical practice [36,45,46]. In nonpharmacological research, such approaches are essential for generating relevant and actionable clinical results [1]. However, the lack of standardized interventions poses a significant challenge. Guidelines such as Template for Intervention Description and Replication [47] are essential for effective reporting and replicability, including detailed annexes to avoid interpretative gaps.

In addition, the use of placebo/sham comparators, uncentric settings, and blinding, which is common in explanatory studies, has led to a debate on their pragmatism [45–49]. Whether the same discussion should pertain to nonlabeled pragmatic ETs is debatable, as “effectiveness” is often deemed synonymous with “pragmatic” in research contexts [5–8].

4.2.2. Placebo/sham control

Including placebo/sham control interventions in MT RCTs may clarify therapeutic effects, which are




Figure 3. PRECIS-2 wheels of each study type. ETac, effectiveness trials active comparator; ETpl, effectiveness trials placebo/sham comparator; PRECIS, PRagmatic EXplanatory Continuum Indicator Summary; PTs, pragmatic trials.

Table 3. PRECIS-2 rates for each type of study

Type of study	PRECIS-2 score	Eligibility	Recruitment	Setting	Organization	Flexibility intervention	Flexibility adherence	Follow-up	Primary outcome	Primary analysis
PTs (<i>n</i> = 39)										
	4/5	54% (21)	54% (20)	55% (21)	51% (20)	65% (25)	39% (12)	62% (24)	95% (37)	71% (27)
	1/2	13% (5)	35% (13)	39% (15)	23% (9)	8% (3)	19% (6)	10% (4)	3% (1)	21% (8)
	3	33% (13)	11% (4)	5% (2)	26% (10)	28% (11)	42% (13)	28% (11)	3% (1)	8% (3)
	Blank	0% (0)	5% (2)	3% (1)	0% (0)	0% (0)	26% (8)	0% (0)	0% (0)	3% (1)
ETac (<i>n</i> = 39)										
	4/5	51% (20)	56% (22)	5% (2)	41% (16)	18% (7)	15% (6)	15% (6)	92% (36)	41% (16)
	1/2	28% (11)	18% (7)	82% (32)	15% (6)	51% (20)	15% (6)	41% (16)	3% (1)	28% (11)
	3	18% (7)	8% (3)	5% (2)	33% (13)	31% (12)	23% (9)	44% (17)	5% (2)	15% (6)
	Blank	3% (1)	18% (7)	8% (3)	10% (4)	0% (0)	46% (18)	0% (0)	0% (0)	15% (6)
ETpl (<i>n</i> = 35)										
	4/5	31% (11)	46% (16)	20% (7)	23% (8)	11% (4)	6% (2)	17% (6)	83% (29)	74% (26)
	1/2	31% (11)	31% (11)	74% (26)	43% (15)	60% (21)	34% (12)	23% (8)	9% (3)	26% (9)
	3	34% (12)	6% (4)	3% (1)	31% (11)	26% (10)	31% (11)	60% (21)	9% (3)	0% (0)
	Blank	3% (1)	11% (4)	3% (1)	3% (1)	0% (0)	29% (10)	0% (0)	0% (0)	0% (0)

ETac, effectiveness trials active comparator; ETpl, effectiveness trials placebo/sham comparator; PRECIS, PRagmatic EXplanatory Continuum Indicatory Summary; PTs, pragmatic trials.

Blank: percentage of studies without information about the domain within the study.

challenging to isolate in efficacy studies. The ongoing PRECIS-3 update aims to assess control groups more effectively, highlighting the need for further research on the role of placebos/shams in PTs, particularly in MT studies where efficacy trials struggle to pinpoint mechanisms despite demonstrating clinical outcomes.

Nonpharmacological interventions differ from pharmacological ones in development and implementation, often lacking prior efficacy studies. Our review found no significant variations in the PRECIS-2 scores between trials using placebo/sham or other comparators, indicating a moderately pragmatic approach. This finding is consistent




Figure 4. Compliance with CONSORT items. CONSORT, Consolidated Standards of Reporting Trials; ETac, effectiveness trials active comparator; ETpl, effectiveness trials placebo/sham comparator; PTs, pragmatic trials.

with the analysis conducted by Devos et al on nursing interventions [41]. Dal-Re et al argue that placebo-controlled trials cannot replicate clinical practice and should not be considered pragmatic. However, the PRECIS-2 authors believe that placebo/sham trials can still inform clinical decisions [12,48–52]. Dal-Re et al also found PRECIS-2 insufficiently sensitive to differentiate between masked and open-label trials [50]. The placebo effects in “touch” interventions are considered part of MT treatments’ therapeutic outcomes [53,54]. Although high-quality efficacy trials are required to clarify whether a MT intervention is more than a placebo/sham [55], including a placebo/sham control in MT RCTs (although it has explanatory connotations) could aid in understanding the real effect of MT interventions that are challenging to achieve in an efficacy study under less realistic conditions. The ongoing PRECIS-3 update aims to assess control groups more effectively, highlighting the need for further research on the role of placebos in PTs, which will be particularly interesting in MT [50–52,56].

4.2.3. Setting

Our results showed that many ETs were conducted in a single center, limiting results’ generalizability [48]. However, Zwarenstein et al argue that detailed reporting of the environment, health system, and patient types can still provide valuable clinical insights [49]. Although multi-centre trials are preferable for broader applicability, they are costly and challenging for MT researchers, who typically lack adequate public and private funding, a significant barrier to more extensive research.

4.2.4. Blinding

PRECIS-2 authors acknowledge that although blinding poses challenges, it remains beneficial in pragmatic studies by significantly enhancing decision-making data quality through the control of the placebo effect and subjectivity [49,56]. In MT, where mechanistic evidence and placebo-controlled studies are scarce, maintaining blinding in trials might help to preserve the rigor of RCTs.

4.3. Reporting and quality of effectiveness MT RCTs

Accurate trial reporting is essential for the applicability and quality of results [4,21]. Our review found that ETac were reported less comprehensively than PTs and ETpl. It is crucial to include adequate details on trial settings, participants, and interventions to facilitate external implementation and validation [4]. In MT, biases are common due to challenges in blinding therapists and patients, especially when patient-reported outcomes are used. This supports a pragmatic approach but poses a high RoB, reducing internal validity.

Our findings indicate that the external validity of ETs is often compromised. A thorough justification for the trial’s

aim, rationale, and design can improve study quality and interpretation. Modifications to trial methodologies are essential to enhance their clinical applicability in the MT field. Clear and comprehensive reporting is vital for understanding trial quality, mitigating biases, and improving the applicability of findings for clinical practice.

4.3. Strengths and limitations

Although the small number of studies in each review group might limit the generalizability of our findings, we intentionally split the sample to compare “effectiveness” studies with self-labeled PTs—ensuring that the groups were matched by the number of PTs. In addition, we acknowledge that not all MTs may be equally represented and that the inclusion of diverse reviewers could have introduced bias into the PRECIS-2 scores. To mitigate subjectivity, we implemented extensive training, thorough piloting of the data, and comprehensive discussions with the principal investigator for precise data extraction. We also recognize that using trial design as a proxy for the underlying intent is a limitation; ideally, direct consultation with the research teams would yield a more accurate assessment. However, this approach highlights the importance of researchers explicitly stating the primary objective of their trials in the published reports. Furthermore, although we acknowledge the limitations of retrospectively applying PRECIS-2, including the caution from its original developers against simply summing the domain scores, we performed a comprehensive evaluation of pragmatism through a meticulous analysis of the PRECIS-2 scores, the authors’ rationale, and the summary score. Our decision was motivated by the need for an objective, systematic comparison of study designs despite the known limitations of the tool.

5. Conclusions

In MT trials, “effectiveness” is used regardless of its pragmatic connotations. These trials are pragmatic regarding participant eligibility, recruitment, and outcome measures but explanatory regarding intervention and setting. The control type (placebo/sham or active) does not affect pragmatism. Compared to self-labeled PTs, ETs show lower compliance with reporting guidelines items.

CRediT authorship contribution statement

Sonia Roura: Writing – review & editing, Writing – original draft, Methodology, Investigation, Formal analysis, Data curation, Conceptualization. **Gerard Alvarez:** Writing – review & editing, Supervision, Methodology, Investigation, Conceptualization. **David Hohenschurz-**

Schmidt: Writing – review & editing, Investigation. **Ivan Solà:** Methodology. **Rodrigo Núñez-Cortés:** Investigation, Formal analysis. **Javier Bracchiglione:** Methodology, Investigation. **Carles Fernández-Jané:** Investigation. **Jules Phalip:** Investigation. **Ignasi Gich:** Formal analysis. **Mercè Sitjà-Rabert:** Supervision. **Gerard Urrútia:** Writing – review & editing, Supervision, Methodology, Conceptualization.

Declaration of competing interest

There are no competing interests for any authors.

Acknowledgments

Sònia Roura is a PhD student in Biomedical Research Methodology and Public Health in the Medical Department of the Universitat Autònoma de Barcelona, Barcelona, Spain. The authors appreciate the support and dedication of Dr Kirsty Loudon in the early stages of the project, teaching and guiding us with the PRECIS-2 tool. The authors would like to thank Roberto Acosta for his support in the first stages of the project.

Data availability

Data will be made available on request.

References

- [1] Daniel MC, Subialka JA, Young JL, Rhon DI. Over half of clinical trials of mobilization and manipulation for patients with low back pain may have limited real-world applicability. *A Systematic Review of 132 Clinical Trials*. *J Orthop Sports Phys Ther* 2022;52:532–45. <https://doi.org/10.1177/01410768231207536>.
- [2] Glasziou P, Matthews R, Boutron I, Chalmers I, Armitage P. The differences and overlaps between “explanatory” and “pragmatic” controlled trials: a historical perspective. *J R Soc Med* 2023;116:425–32. <https://doi.org/10.1177/01410768231207536>.
- [3] Schwartz D, Lellouch J. Explanatory and pragmatic attitudes in therapeutic trials. *J Chronic Dis* 1967;20:637–48. [https://doi.org/10.1016/0021-9681\(67\)90041-0](https://doi.org/10.1016/0021-9681(67)90041-0).
- [4] Singal AG, Higgins PDR, Waljee AK. A primer on effectiveness and efficacy trials. *Clin Transl Gastroenterol* 2014;5:e45. <https://doi.org/10.1038/ctg.2013.13>.
- [5] Janiaud P, Dal-Ré R, Ioannidis JPA. Assessment of pragmatism in recently published randomized clinical trials. *JAMA Intern Med* 2018;178:1278. <https://doi.org/10.1001/jamainternmed.2018.3321>.
- [6] Dal-Ré R, Janiaud P, Ioannidis JPA. Real-world evidence: how pragmatic are randomized controlled trials labeled as pragmatic? *BMC Med* 2018;16:49. <https://doi.org/10.1186/s12916-018-1038-2>.
- [7] Sacristán JA, Dilla T. Pragmatic trials revisited: applicability is about individualization. *J Clin Epidemiol* 2018;99:164–6. <https://doi.org/10.1016/j.jclinepi.2018.02.003>.
- [8] Ford I, Norrie J. Pragmatic trials. *New Engl J Med* 2016;375:454–63. <https://doi.org/10.1056/nejmra1510059>.
- [9] Fortney JC, Curran GM, Lyon AR, Check DK, Flum DR. Similarities and differences between pragmatic trials and hybrid effectiveness-implementation trials. *J Gen Intern Med* 2024;39:1735–43. <https://doi.org/10.1007/s11606-024-08747-1>.
- [10] Nicholls SG, Zwarenstein M, Hey SP, Giraudeau B, Campbell MK, Taljaard M. The importance of decision intent within descriptions of pragmatic trials. *J Clin Epidemiol* 2020;125:30–7. <https://doi.org/10.1016/j.jclinepi.2020.04.030>.
- [11] Treweek S, Zwarenstein M. Making trials matter: pragmatic and explanatory trials and the problem of applicability. *Trials* 2009;10:37. <https://doi.org/10.1186/1745-6215-10-37>.
- [12] Thorpe KE, Zwarenstein M, Oxman AD, Treweek S, Furberg CD, Altman DG, et al. A pragmatic–explanatory continuum indicator summary (PRECIS): a tool to help trial designers. *J Clin Epidemiol* 2009;62:464–75. <https://doi.org/10.1016/j.jclinepi.2008.12.011>.
- [13] Zwarenstein M, Treweek S, Gagnier JJ, Altman DG, Tunis S, Haynes B, et al. Improving the reporting of pragmatic trials: an extension of the CONSORT statement. *BMJ* 2008;337:a2390. <https://doi.org/10.1136/bmj.a2390>.
- [14] Alvarez G, Solà I, Sitjà-Rabert M, Fort-Vanmeerhaeghe A, Gich I, Fernández C, et al. A methodological review revealed that reporting of trials in manual therapy has not improved over time. *J Clin Epidemiol* 2020;121:32–44. <https://doi.org/10.1016/j.jclinepi.2020.01.006>.
- [15] Riley SP, Swanson BT, Sawyer SF, Brismée J-M. Is research quality in orthopedic manual therapy trials stagnating? Reflections and pathways for improving research quality and advance our profession. *J Man Manip Ther* 2016;24:239–40. <https://doi.org/10.1080/10669817.2016.1253561>.
- [16] Gonzalez GZ, Moseley AM, Maher CG, Nascimento DP, Costa L, da CM, et al. Methodologic quality and statistical reporting of physical therapy randomized controlled trials relevant to musculoskeletal conditions. *Arch Phys Med Rehabil* 2018;99:129–36. <https://doi.org/10.1016/j.apmr.2017.08.485>.
- [17] Koes BW. How to evaluate manual therapy: value and pitfalls of randomized clinical trials. *Man Ther* 2004;9:183–4. <https://doi.org/10.1016/j.math.2004.04.002>.
- [18] Cashin AG, Lee H, Bagg MK, O’Hagan E, Traeger AC, Kamper SJ, et al. A systematic review highlights the need to improve the quality and applicability of trials of physical therapy interventions for low back pain. *J Clin Epidemiol* 2020;126:106–15. <https://doi.org/10.1016/j.jclinepi.2020.06.025>.
- [19] Clar C, Tsertsvadze A, Court R, Hundt GL, Clarke A, Sutcliffe P. Clinical effectiveness of manual therapy for the management of musculoskeletal and non-musculoskeletal conditions: systematic review and update of UK evidence report. *Chiropr Man Therap* 2014;22:12. <https://doi.org/10.1186/2045-709X-22-12>.
- [20] Riley SP, Swanson B, Brismée J-M, Sawyer SF. A systematic review of orthopaedic manual therapy randomized clinical trials quality. *J Man Manipulative Ther* 2016;24:241–52. <https://doi.org/10.1080/10669817.2015.1119372>.
- [21] Innocenti T, Giagio S, Salvioli S, Feller D, Minnucci S, Brindisino F, et al. Completeness of reporting is suboptimal in randomized controlled trials published in rehabilitation journals, with trials with low risk of bias displaying better reporting: a meta-research study. *Arch Phys Med Rehabil* 2022;103:1839–47. <https://doi.org/10.1016/j.apmr.2022.01.156>.
- [22] Fregni F, Imamura M, Chien HF, Lew HL, Boggio P, Kaptchuk TJ, et al. Challenges and recommendations for placebo controls in randomized trials in physical and rehabilitation medicine: a report of the international placebo symposium working group. *Am J Phys Med Rehabil* 2010;89:160–72. <https://doi.org/10.1097/PHM.0b013e3181bc0bbd>.
- [23] Simoni G, Bozzolan M, Bonnini S, Grassi A, Zucchini A, Mazzanti C, et al. Effectiveness of standard cervical physiotherapy plus diaphragm manual therapy on pain in patients with chronic neck pain: a randomized controlled trial. *J Bodyw Mov Ther* 2021;26:481–91. <https://doi.org/10.1016/j.jbmt.2020.12.032>.
- [24] Scafoglieri A, Van den Broeck J, Willems S, Tamminga R, van der Hoeven H, Engelsma Y, et al. Effectiveness of local exercise therapy versus spinal manual therapy in patients with patellofemoral pain syndrome: medium term follow-up results of a randomized controlled trial. *BMC Musculoskelet Disord* 2021;22:446. <https://doi.org/10.1186/s12891-021-04310-9>.

- [25] Muñoz-Gómez E, Inglés M, Serra-Añó P, Espí-López GV. Effectiveness of a manual therapy protocol based on articulatory techniques in migraine patients. A randomized controlled trial. *Musculoskelet Sci Pract* 2021;54:102386. <https://doi.org/10.1016/j.msksp.2021.102386>.
- [26] Izaola-Azkona L, Vicenzino B, Olabarrieta-Eguia I, Saez M, Lascurain-Aguirrebeña I. Effectiveness of mobilization of the talus and distal fibula in the management of acute lateral ankle sprain. *Phys Ther* 2021;101:pzab111. <https://doi.org/10.1093/ptj/pzab111>.
- [27] Gamerman V, Cai T, Elsässer A. Pragmatic randomized clinical trials: best practices and statistical guidance. *Health Serv Outcomes Res Methodol* 2019;19:23–35. <https://doi.org/10.1007/s10742-018-0192-5>.
- [28] Wieland LS, Berman BM, Altman DG, Barth J, Bouter LM, D'Adamo CR, et al. Rating of Included Trials on the Efficacy-Effectiveness Spectrum: development of a new tool for systematic reviews. *J Clin Epidemiol* 2017;84:95–104. <https://doi.org/10.1016/j.jclinepi.2017.01.010>.
- [29] Deaton A, Cartwright N. Understanding and misunderstanding randomized controlled trials. *Soc Sci Med* 2018;210:2–21. <https://doi.org/10.1016/j.socscimed.2017.12.005>.
- [30] Porzolt F, Wiedemann F, Philippen M, Weiss C, Weiss M, Schmalting K, et al. The terminology conflict on efficacy and effectiveness in healthcare. *J Comp Eff Res* 2020;9:1171–8. <https://doi.org/10.2217/ce-2020-0149>.
- [31] Hohenschurz-Schmidt D, Kleykamp BA, Draper-Rodi J, Vollert J, Chan J, Ferguson M, et al. Pragmatic trials of pain therapies: a systematic review of methods. *Pain* 2021;163:21–46.
- [32] Palakshappa JA, Gibbs KW, Lannan MT, Cranford AR, Taylor SP. Systematic review of the “pragmatism” of pragmatic critical care trials. *Crit Care Explorations* 2022;4:e0738. <https://doi.org/10.1097/CCE.0000000000000738>.
- [33] Roura S, Alvarez G, Hohenschurz-Schmidt D, Solà I, Núñez-Cortés R, Bracchiglione J, et al. Lack of pragmatic attitude of self-labelled pragmatic trials on manual therapy: a methodological review. *BMC Med Res Methodol* 2024;24. <https://doi.org/10.1186/s12874-024-02393-1>.
- [34] Fritz JM, Cleland J. Effectiveness versus efficacy: more than a debate over language. *J Orthop Sports Phys Ther* 2003;33:163–5. <https://doi.org/10.2519/jospt.2003.33.4.163>.
- [35] Witt CM. Efficacy, effectiveness, pragmatic trials—guidance on terminology and the advantages of pragmatic trials. *Forsch Komplementmed* 2009;16:292–4. <https://doi.org/10.1159/000234904>.
- [36] Gordon KS, Peduzzi P, Kerns RD. Designing trials with purpose: pragmatic clinical trials of nonpharmacological approaches for pain management. *Pain Med* 2020;21:S7–12. <https://doi.org/10.1093/pm/pnaa347>.
- [37] Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al, editors. *Cochrane handbook systematic reviews interventions*. 2nd edition. Chichester, UK: John Wiley & Sons; 2019.
- [38] Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Rev Esp Cardiol* 2021;74:790–9. <https://doi.org/10.1016/j.rec.2021.07.010>.
- [39] Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan-a web and mobile app for systematic reviews. *Syst Rev* 2016;5:210. <https://doi.org/10.1186/s13643-016-0384-4>.
- [40] Loudon K, Treweek S, Sullivan F, Donnan P, Thorpe KE, Zwarenstein M. The PRECIS-2 tool: designing trials that are fit for purpose. *BMJ* 2015;350:h2147. <https://doi.org/10.1136/bmj.h2147>.
- [41] Devos F, Foissac F, Bouazza N, Ancel P-Y, Tréluyer J-M, Chappuy H. Study characteristics impacted the pragmatism of randomized controlled trial published in nursing: a meta-epidemiological study. *J Clin Epidemiol* 2019;116:18–25. <https://doi.org/10.1016/j.jclinepi.2019.07.017>.
- [42] Elder WG, Munk N. Using the pragmatic-explanatory continuum indicator summary (PRECIS) model in clinical research: application to refine a practice-based research network (PBRN) study. *The J Am Board Fam Med* 2014;27:846–54. <https://doi.org/10.3122/jabfm.2014.06.140042>.
- [43] Rosas LG, Lv N, Azar K, Xiao L, Yank V, Ma J. Applying the pragmatic—explanatory continuum indicator summary model in a primary care—based lifestyle intervention trial. *Am J Prev Med* 2015;49:S208–14. <https://doi.org/10.1016/j.amepre.2015.05.011>.
- [44] Alvarez G, Solà I, Sitjà-Rabert M, Fort-Vanmeerhaeghe A, Gich I, Fernández V, et al. A call for improving the Internal validity and the reporting of manual therapy trials self-labelled as pragmatic: a methodological review. *Int J Osteopath Med* 2025;55:100754. <https://doi.org/10.1016/j.ijosm.2025.100754>.
- [45] Bishop MD, Torres-Cueco R, Gay CW, Lluch-Girbés E, Beneciuk JM, Bialosky JE. What effect can manual therapy have on a patient's pain experience? *Pain Manag* 2015;5:455–64. <https://doi.org/10.2217/pmt.15.39>.
- [46] Lin I, Wiles L, Waller R, Goucke R, Nagree Y, Gibberd M, et al. What does best practice care for musculoskeletal pain look like? Eleven consistent recommendations from high-quality clinical practice guidelines: systematic review. *Br J Sports Med* 2020;54:79–86. <https://doi.org/10.1136/bjsports-2018-099878>.
- [47] Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014;348:g1687. <https://doi.org/10.1136/bmj.g1687>.
- [48] Dal-Ré R, de Boer A, James SK. The design can limit PRECIS-2 retrospective assessment of the clinical trial explanatory/pragmatic features. *J Clin Epidemiol* 2020;126:193–201. <https://doi.org/10.1016/j.jclinepi.2020.03.027>.
- [49] Zwarenstein M, Thorpe K, Treweek S, Loudon K. PRECIS-2 for retrospective assessment of RCTs in systematic reviews. *J Clin Epidemiol* 2020;126:202–6. <https://doi.org/10.1016/j.jclinepi.2020.06.023>.
- [50] Dal-Ré R. The PRECIS-2 tool seems not to be useful to discriminate the degree of pragmatism of medicine masked trials from that of open-label trials. *Eur J Clin Pharmacol* 2021;77:539–46. <https://doi.org/10.1007/s00228-020-03030-8>.
- [51] Dal-Ré R. Pragmatic trials, blinding, placebos, and the usefulness of the PRECIS-2 tool. *Eur J Clin Pharmacol* 2021;77:1071–2. <https://doi.org/10.1007/s00228-020-03079-5>.
- [52] Zwarenstein M, Howie A. Blinding, pragmatism, and the PRECIS-2 tool for designing and assessing randomized trials. *Eur J Clin Pharmacol* 2021;77:1069–70. <https://doi.org/10.1007/s00228-020-03078-6>.
- [53] Rossetini G, Carlino E, Testa M. Clinical relevance of contextual factors as triggers of placebo and nocebo effects in musculoskeletal pain. *BMC Musculoskelet Disord* 2018;19:27. <https://doi.org/10.1186/s12891-018-1943-8>.
- [54] Bialosky JE, Bishop MD, Penza CW. Placebo mechanisms of manual therapy: a sheep in wolf's clothing? *J Orthop Sports Phys Ther* 2017;47:301–4. <https://doi.org/10.2519/jospt.2017.0604>.
- [55] Hohenschurz-Schmidt D, Liem T. Placebo effects in osteopathy and other manual therapies — what they are and why they matter to clinical practice, education, and research. *Int J Osteopath Med* 2025;56:100762. <https://doi.org/10.1016/j.ijosm.2025.100762>.
- [56] Willis A, Shiely F, Treweek S, Taljaard M, Loudon K, Howie A, et al. Comments, suggestions and criticisms of the PRECIS-2 design tool: a citation analysis. *J Clin Epidemiol* 2024;176:111534. <https://doi.org/10.1016/j.jclinepi.2024.111534>.