

Hot Sulfur on the Rocks: The Reaction of Electronically Excited Sulfur Atoms with Water in an Ice-Surface Model

Published as part of ACS Earth and Space Chemistry *special issue* “Harold Linnartz Festschrift”.

Gabriella Di Genova,* Jessica Perrero, Marzio Rosi, Cecilia Ceccarelli, Albert Rimola,* and Nadia Balucani*



Cite This: *ACS Earth Space Chem.* 2025, 9, 844–855



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: In this contribution, we present a theoretical investigation of the reaction involving atomic sulfur in its first electronically excited state, $S(^1D)$, and H_2O on an ice-surface model. This study is motivated by the work of Giustini et al. (*ACS Earth Space Chem.*, 2024, 8, 2318), which indicated a strong effect of the presence of four additional water molecules in the $S(^1D) + H_2O$ reaction compared to the pure gas-phase case. Our simulation treats the long-range interactions (H-bonds and dispersion forces) with the ice water molecules in a much more realistic way being based on the use of a cluster of 18 water molecules, thus overcoming the limits of the small cluster used by Giustini et al. According to our results, $S(^1D)$ reacts via two possible reaction mechanisms: (1) addition to the O atom of a water molecule with the formation of H_2OS or (2) insertion into one of the O–H bonds of a water molecule with the formation of $HOSH$. Both H_2OS and $HOSH$ are stabilized on ice by energy dissipation rather than isomerizing or dissociating into two products as seen in the gas-phase reaction. The interaction with surrounding water molecules affects the entire reaction pathway by stabilizing intermediate species, reducing some barriers, and impeding the only two-product open channel of the gas-phase reaction. $S(^1D)$ can be produced by UV-induced photodissociation of various precursor molecules on the surface of interstellar or cometary ice or by other high-energy processes induced by electrons or cosmic rays also in the ice bulk. Therefore, our results can be of help in elucidating the mysterious sulfur chemistry occurring in the icy mantles of interstellar grains or in cometary nuclei. Furthermore, this study demonstrates that the product branching ratios of gas-phase reactions should not be uncritically used in modeling interstellar ice chemistry.

KEYWORDS: astrochemistry, DFT, interstellar medium, grain-surface chemistry, potential energy surfaces, kinetics



1. INTRODUCTION

Sulfur is the tenth most abundant element in the Solar System and local interstellar medium (its abundance with respect to hydrogen¹ is $1.4 \cdot 10^{-5}$) and is ubiquitous in space, with S-bearing species detected in interstellar and intergalactic regions,^{2,3} on planet surfaces and atmospheres,^{4–6} on moons,^{7,8} and small bodies like comets, asteroids, and meteorites.^{9–14} It is also one of the six essential elements (HCNOPS) of life, being present (mainly in its reduced forms) in numerous biomolecules, including proteins, sugars, nucleic acids, vitamin cofactors, and metabolites.¹⁵ Therefore, it is of interest to understand the carriers of reduced sulfur to early Earth. In the interstellar medium (ISM), sulfur can be found in molecules - either in the gas phase (e.g., SO , H_2S , CS , H_2CS , and C_3S ¹⁶) or adsorbed in the icy mantles of interstellar dust grains (OCS and, possibly, SO_2 ^{17–19}) - and in the refractory material of grains (as, for instance, iron sulfide²⁰) that eventually form planetesimals. According to the Cologne Database for Molecular Spectroscopy,² 39 S-bearing molecules

have been detected so far in ISM. Nevertheless, in molecular clouds and star-forming regions sulfur is significantly depleted from the gas phase (the “sulfur depletion problem”) with only 1% of the initial S-budget observed in its gaseous species.^{16,21–25} In these regions, S is mostly present in its ionized form (S^+), which largely increases the sulfur binding energy to the grain surface as for all the other charged species.²⁶ In addition, dust grains typically carry a negative charge²⁷ so that their electrostatic attraction causes an even larger sticking coefficient on dust grains. As a consequence, it has been suggested that the missing sulfur is present in the ice covering the dust grains. Yet, the only two sulfur species

Received: November 24, 2024

Revised: March 14, 2025

Accepted: March 14, 2025

Published: March 25, 2025



detected so far on the icy mantles of interstellar grains account for ca. 5% of the S elemental budget^{18,19} and, contrary to what initially suggested, H₂S is not the main S carrier on ice.^{17,28} The main carriers of sulfur in dense clouds remain unidentified. Refractory sulfur species, such as sulfur allotropes (S_n, *n* = 3–8), iron sulfide, or ammonium sulfide have been invoked as the main sulfur reservoirs.^{21,29–31} Some clues on the sulfur carriers in star-forming regions are offered by pristine small bodies of our solar system, such as asteroids, meteorites, and comets. Formed in the outer protosolar disc, cometary nuclei are believed to be formed by pristine material^{32,33} and their composition is alleged to resemble that of interstellar dust grains (and their icy mantles).⁹ Cometary comae have been investigated by remote spectroscopy for some years and, after the first detection of atomic S and CS in the comet C/1975 West,³⁴ other simple sulfur species (H₂S, OCS, SO, SO₂, S₂, CS₂, H₂CS, and NS) have been identified.^{9,35–42} The Rosetta orbiter Spectrometer for Ion and Neutral Analysis (ROSINA) allowed the identification of other S-species in the quiet coma of the Jupiter family comet 67P/Churyumov-Gerasimenko.⁴³ In addition to the usual S-bearing species, S₃, S₄, CH₃SH, and C₂H₅SH/CH₃SCH₃^{10,43,44} were identified for the first time. Even more complex S-bearing species were detected during two episodes of enhanced dust emission (e.g., CH₃OS, CH₄OS, C₂H₄OS, C₃H₈OS)^{45,46} including those with gross formula H₂SO and HSO.⁴⁶ Finally, a rich inventory of S-compounds was compiled after the analysis of the surface material of Ryugu,^{11,47,48} similar to the one obtained by the analysis of meteorites like Murchinson and Allende.¹² Clearly, the more complex S-species found in pristine small objects like comets and asteroids/meteorites are the result of some unidentified chemistry involving the simple S-molecules found in star-forming regions.

In this work, we consider the possible role of atomic sulfur in its first electronically excited state, ¹D, in the chemical evolution of interstellar or cometary ice. The ¹D state of S atoms has a large internal energy content (it lies 110.5 kJ mol^{−1} above the ground ³P state) and is a metastable state with a radiative lifetime of ca. 30 s.⁴⁹ In extraterrestrial environments, it is produced in the gas phase by UV-induced photodissociation or electron impact-induced dissociation of several precursor molecules like OCS and CS₂.⁵⁰ Other precursors of S(¹D) are H₂S and SO₂ via their photodissociation at $\lambda \leq 155$ nm (including the Lyman- α wavelength).^{51–53} While gas-phase S(¹D) has been spectroscopically identified in the coma of the comet 153P/Ikeya-Zhang,⁵⁴ the production of S(¹D) is also expected when photodissociation occurs for the same molecules residing on the ice surface as recently demonstrated for the analogous case of atomic oxygen in its ¹D state by Bergner et al.^{55,56} who investigated the O(¹D) reactions on methane ice, ethane ice, and acetylene ice. Methanol and formaldehyde were observed in the case of the reaction with solid methane, ethylene oxide, and acetaldehyde in the case of the reaction with ethane, and ketene in the case of the reaction with acetylene.⁵⁶ Recently, models on interstellar ice chemistry^{57,58} and ice chemistry in cometary nuclei⁵⁹ have started considering explicitly so-called photodissociation-induced (PDI) reactions, in which transient species are produced by UV photodissociation (or radiolysis induced by galactic cosmic rays (GCRs)) and immediately react with an adjacent molecule of the ice matrix. This approach, originally introduced to account for nondiffusive ice chemistry,⁵⁷ has been extended to the case of O(¹D) and also suggested for

other species of the same kind, such as C(¹D)), by Carder et al.⁵⁸

Several gas-phase reactions of S(¹D) have been recently investigated and revealed its ability to insert into σ bonds (thus activating molecules that are not much reactive otherwise^{49,60}) in addition to adding to the π -system of multiple bonds^{61,62} or lone pairs.⁶³ This is significant because, unlike atomic sulfur in its ground electronic ³P state, which has low reactivity with closed-shell species, S(¹D) reactive collisions can be crucial in forming organosulfur and other sulfur-bearing compounds in various environments.^{60–62} Since the main component of cometary and interstellar ice is water, according to the PDI scheme, once formed, S(¹D) will react with the surrounding species, and water is by far the most probable candidate. Our simulation could also contribute to understanding sulfur chemistry in atmospheric water aerosols containing S(¹D) precursors like OCS, the most abundant atmospheric sulfur species that photodissociates in the upper troposphere by absorbing photons in the window between the O₂/O₃ absorption bands and produce both S(³P) and S(¹D)).⁶⁴

To assess the possible role of the S(¹D) reaction with the water molecules of ice, we have simulated the reaction occurring on amorphous ice by considering a cluster of 18 water molecules holding several dangling oxygen and hydrogen atoms (the dangling OH ice feature was recently observed toward the dense cloud Chamaeleon I⁶⁵) and characterized the potential energy surface (PES) of the S(¹D) + H₂O reaction on the cluster model to highlight the differences with the same reaction in the gas phase. The results are compared with those previously obtained by Giustini et al.,⁶³ who investigated the gas-phase and S(¹D) reactions with a cluster of five water molecules. In that study, Giustini et al.⁶³ noted significant differences in the reaction mechanism caused by the presence of four additional water molecules with respect to the case of gas-phase reaction (see below). Those differences motivated the present study since an improved simulation of the amorphous water ice will allow for a more accurate assessment of the influence of the ice on the reaction outcome.

2. METHODOLOGY

2.1. Ice Model. Interstellar icy mantles were simulated with an 18-water cluster model (*W*_{ice}), adopted in previous studies^{66–69} mimicking a compact, amorphous ice surface that presents flatter regions. The size of *W*_{ice} is a compromise between a realistic model (relatively large and amorphous) and the accuracy of the employed methodology, enabling the use of accurate, cost-effective, state-of-the-art theoretical methods without incurring high computational costs. The *W*_{ice} cluster model presents several adsorption sites, consisting of 8 dangling H atoms (dH, labeled as *a–f* in Figure 1) and 9 dangling O atoms (dO, labeled as 1–9 in Figure 1), showing a significant binding site variability, as expected for real amorphous water ice surfaces.

2.2. Electronic Structure Calculations. We employed GAUSSIAN16⁷⁰ and ORCA 6.0.1⁷¹ software packages to carry out electronic structure calculations of the PESs for the S(¹D)+H₂O reaction in the gas phase and on *W*_{ice}.

For the gas-phase reaction, given its reduced size, we performed geometry optimizations of the PES stationary points using the single- and double-electronic excitation coupled-cluster method with perturbative description of triple excitations, CCSD(T),^{72–74} in conjunction with the correlation consistent valence polarized basis set,^{75–77} augmented




Figure 1. ω B97XD/ma-def2-TZVP-optimized structure for the water ice cluster model consisting of 18 water molecules (W_{ice}).

with a tight d function for the sulfur atoms to correct for the core polarization effects⁷⁸ (aug-cc-pV(Q+d)Z). The energy of each stationary point was also computed at the CCSD(T)-F12 level of theory to compare them with previously published work. The F12 approximation allows for the computation of electronic correlation energies close to the limit of a complete basis set (CBS).⁷⁹ The calculation requires a complementary auxiliary basis set (CABS) to factorize the many-electron integrals resulting from the use of explicitly correlated wave functions. This allows using medium-sized standard basis sets without compromising the quality of the results.^{80,81} Additionally, the PES was also computed at the domain-based local pair natural orbital coupled cluster theory with single-, double-, and perturbative triple-excitations DLPNO-CCSD(T) level of theory, in prevision of applying this methodology to W_{ice} .⁸² When considering W_{ice} , given the larger size of the system, geometry optimizations were performed at the DFT ω B97XD level of theory,⁸³ in conjunction with the minimally augmented Karlsruhe basis set ma-def2-TZVP.⁸⁴ The selection of the ω B97XD method was based on a preliminary benchmarking analysis, which allowed us to identify a suitable DFT functional

that properly describes the electronic structure of the systems and reactions to study. Three DFT functionals were employed: B3LYP-D3(BJ), M062X-D3, and ω B97XD. The evaluation was based on comparing the DFT energy barriers against the values computed with single-point energy calculations at CCSD(T) level of theory on the optimized DFT geometries. This analysis was performed by considering the reactivity of $S(^1D)$ with a water molecule in the presence of a second water molecule assisting the reaction. The ω B97XD/ma-def2-TZVP methodology was used to compute the harmonic vibrational frequencies with the aim of checking the nature of the stationary points and deriving the vibrational zero point energy (ZPE) corrections. Minima, such as reactants, intermediates, and products, are characterized by positive eigenvalues of the Hessian matrix, while first-order saddle points, such as transition state structures, are identified by the presence of a single negative eigenvalue. Every energy accounts for the thermochemistry at 0 K, obtained by adding the ω B97XD-ZPE corrections onto the potential energies calculated at the different theory levels, this way deriving enthalpies at 0 K.

Intrinsic reaction coordinate (IRC)^{85,86} calculations were performed to link each identified saddle point with the respective reactant and product.

All the calculations involving one radical species were run as open-shell systems based on an unrestricted formalism. For those involving two radical species, we first optimized the structures as triplet states, followed by an open-shell singlet optimization to describe the biradical system. Singlet biradical systems were treated using the unrestricted broken symmetry approach,⁸⁷ in which the most stable initial wave function was found using the *guess = mix* and *stable = opt* keywords in GAUSSIAN16.

At all the levels of calculation, the energy of the $S(^1D)$ electronic state was estimated by adding the experimental $S(^3P)$ - $S(^1D)$ separation, corresponding to 110.5 kJ mol⁻¹,⁴⁹ to the computed energy of the $S(^3P)$ state.




Figure 2. ZPE-corrected PES for the gas-phase $S(^1D) + H_2O$ reaction channels. Geometries and energies are calculated at the CCSD(T)/aug-cc-pV(Q+d)Z level of theory. The values obtained at the CBS level of theory on optimized ω B97XD structures by Giustini et al.⁶³ are also shown in parentheses for comparison.

Table 1. Relative Energies (in kJ mol⁻¹) with Respect to the S(¹D) + H₂O Asymptote of the Stationary Points along the PES Shown in Figure 2^a

	CCSD(T)/aug-cc-pV(Q+d)Z this work	DLPNO-CCSD(T) this work	CCSD(T)-F12 this work	CBS Giustini et al. ⁶³	CCSD(T)/aug-cc-pV(T+d)Z Giustini et al. ⁶³
H ₂ OS	-93	-93	-94	-98	-86
HOSH	-249	-248	-256	-258	-244
H ₂ SO	-176	-176	-188	-188	-171
TS1	-44	-42	-46	-50	-39
TS2	-4	-5	-9	-13	+2
TS3	-3	-3	-8	-10	+5
TS4	+153	+154	+148	+145	+153
³ OS + H ₂	-138	-139	-142	-141	-152
¹ OS + H ₂	-40	-40	-45	-47	-35
OH + SH	+33	+33	+31	+32	+31
HSO + H	+58	+57	+49	+51	+60
HOS + H	+71	+72	+64	+65	+73
H ₂ S + O(¹ D)	+291	+291	+278	+270	+265

^aThe data at the CCSD(T)/aug-cc-pV(Q+d)Z level obtained in this work are shown in the first column, while the single energy point computed at the DLPNO-CCSD(T) and CCSD(T)-F12 level of theory are presented, respectively, in the second and third column. The fourth and fifth columns list the data obtained by Giustini et al.⁶³ at the CBS and CCSD(T)/aug-cc-pV(T+d)Z levels on optimized ω B97XD structures for comparison.

2.3. Kinetic Calculations. After calculating the energy barriers for the isomerization reactions (see Section 3.2) in the W_{ice} cluster model, we performed kinetic calculations to estimate the reactive lifetimes of the addition and insertion products. We used the Ramsperger-Rice-Kassel-Marcus (RRKM) theory⁸⁸ to compute the reaction rate constant for the isomerization steps as a function of the total energy, $k(E)$, using an in-house code^{61,89,90} adapted to the case of reactions on large clusters of water molecules.^{67,68,91,92} The $k(E)$ for a certain reaction at a specific total energy is given by

$$k(E) = \frac{N_{TS}(E)}{h\rho_T(E)} \quad (1)$$

where $N_{TS}(E)$ is the sum of states of the transition state at the total energy E , $\rho_T(E)$ is the reactant density of states at the total energy E , and h is the Planck's constant. $N_{TS}(E)$ is obtained by integrating the relevant density of states up to energy E , assuming a rigid rotor/harmonic oscillator model. The density of states is symmetrized with respect to the number of identical configurations of the reactants and/or transition states. We treated each step as unimolecular and considered the whole cluster as isolated from its surroundings. Inside the isolated cluster, intramolecular vibrational redistribution is assumed to be fast. This approximation was verified to be good enough in other cases^{67,91} once we consider the time scale of energy randomization.^{93,94} From the unimolecular rate constants, we derived the lifetime of the reaction intermediates as the inverse of $k(E)$ to evaluate if isomerization is competitive with the energy dissipation toward the entire ice matrix.

3. RESULTS

3.1. Reaction S(¹D) + H₂O in the Gas Phase. The PES of the gas-phase S(¹D) + H₂O reaction, optimized at CCSD(T)/aug-cc-pV(Q+d)Z, is summarized in Figure 2. The energy values are reported rounded to the nearest kJ/mol considering the accuracy of the calculations. As already indicated by the calculations of Giustini et al.,⁶³ S(¹D) can either interact with one of the lone pairs of the oxygen atom, forming the addition

intermediate H₂OS (MIN1), or insert into one of the O–H bonds of the water molecule, forming the insertion intermediate HOSH (MIN2). Both entrance channels are barrierless. The H₂OS intermediate is formed by electron charge transfer, mainly from the O atom of H₂O to S, as indicated by the natural population analysis calculation (see Supporting Information, SI). This charge transfer is possible because S(¹D) has two pure electronic microstates that present an empty 3p orbital to accommodate electron density. Once formed, H₂OS can either isomerize to HOSH through TS1, by overcoming an intrinsic energy barrier of 49 kJ mol⁻¹ (TS1 of Figure 2), or decompose into SO + H₂ (since this is a singlet PES, the SO molecule is formed in its excited state a¹Δ) by overcoming a very high intrinsic barrier of 246 kJ mol⁻¹ (associated with a three-center elimination transition state, TS4 of Figure 2), or decompose into HOS + H in an endothermic channel. HOSH (MIN2) can either isomerize to MIN1 through the TS1 barrier, now of 205 kJ mol⁻¹, or isomerize to H₂SO (MIN3) by overcoming an intrinsic barrier of 245 kJ mol⁻¹ (TS2), or fragment into OH + SH, HSO + H and HOS + H (all these dissociation channels are endothermic). MIN3 can isomerize back to MIN2 (now with a barrier of 174 kJ mol⁻¹, TS2), or decompose into HSO + H (an endothermic channel) or into SO(a¹Δ) + H₂ by overcoming a barrier (TS3) located at -3 kJ mol⁻¹ with respect to the reactants asymptote. As already noted by Giustini et al. 2024, this is the only two-product open channel since it is both exothermic and characterized by submerged barriers along the energy path.

Table 1 compares the relative energies of the stationary points for the S(¹D) + H₂O PES with the values obtained by Giustini et al.⁶³ at the CBS and CCSD(T)/aug-cc-pV(T+d)Z levels of theory on optimized ω B97XD structures. The results of the present work are in good agreement with the CBS data, the two methods lying within the chemical accuracy (4 kJ mol⁻¹), with the only significant discrepancy associated with the energy of the O(¹D) + H₂S product channel. The agreement is less good with respect to the CCSD(T)/aug-cc-pV(T+d)Z calculations.⁶³ Our data also nicely compare with the $\Delta H_{0,r}^\circ = +30.7$ kJ/mol for the OH + SH channel (derived




Figure 3. DLPNO–CCSD(T)/aug-cc-pVTZ// ω B97XD/ma-def2-TZVP PES, including ZPE corrections (at ω B97XD/ma-def2-TZVP), of the surface reaction of $S(^1D) + H_2O$. Energies are in kJ mol^{-1} .




Figure 4. Optimized structures and relative energies (at the corresponding energy levels) of the stationary points for the $S(^1D) + H_2O$ addition reaction in the gas phase (A1) and on W_{ice} (A2).

by using the accepted values of ΔH_{0f}° for OH and SH, see Appendix A5 of Atkinson et al.⁹⁵). The agreement is also quite good when considering the difference in energy with respect to the SO product in its ground triplet state ($X^3\Sigma^-$) which is -143.75 kJ/mol (when using the values by Atkinson et al.⁹⁵). However, the internal energy content of the excited $SO(a^1\Delta)$ corresponds to 98 kJ/mol to be compared with the accepted value of 77.11 kJ/mol ($T_0 = 5861 \text{ cm}^{-1}$, see Borin and Ornellas⁹⁶). Therefore, our method, as well as those used by Giustini et al.,⁶³ confirm their problems in dealing with electronic excitation.

3.2. Reaction of $S(^1D)$ with a Cluster of 18 Water Molecules. According to our benchmarking analysis, the method yielding the smallest error on the energy barriers was ω B97XD (15%), followed by M062X-D3 (21%), and BHLYP-D3(BJ) (23%) (see SI). Therefore, ω B97XD was selected to

perform our calculations on W_{ice} . $S(^1D)$ was adsorbed on the 9 different dO positions of the W_{ice} cluster (see Figure 1), the dO4 binding site being the one providing the most favorable interaction energy (-219 kJ mol^{-1}). The geometries and interaction energies of each binding site (dO1–9) are shown in SI. Figure 3 shows the PES of the $S(^1D) + H_2O$ reaction on W_{ice} , the geometries of the stationary points optimized at the ω B97XD level of theory and the energetics refined at DLPNO–CCSD(T). Bond distances for each stationary point are shown in Figures 4 and 5. As in the gas phase, the $S(^1D)$ atom can either add to one of the lone pairs of dO4 forming H_2OS_{ice} (MIN1_{ice}) or directly insert into one of the O–H bonds protruding from the cluster forming $HOSH_{ice}$ (MIN2_{ice}). The formation of H_2OS_{ice} and $HOSH_{ice}$ is highly exothermic (-219 kJ mol^{-1} and -274 kJ mol^{-1} , respectively). $HOSH_{ice}$ can also be formed via the isomerization of H_2OS_{ice}




Figure 5. Optimized structures and relative energies (at the corresponding energy levels) of the stationary points for the $S(^1D) + H_2O$ insertion reaction in the gas phase (B1) and on W_{ice} (B2).

via a barrier of 20 kJ mol^{−1}. Finally, H_2SO_{ice} can be formed by the isomerization of $HOSH_{ice}$ but a barrier of 115 kJ mol^{−1} needs to be overcome.

The gas-phase and the ice-surface reaction pathways starting with the $S(^1D)$ addition to H_2O forming H_2SO and continuing with its isomerization into $HOSH$ are compared in Figure 4, while the gas-phase and the ice-surface reaction pathways starting with the $S(^1D)$ insertion forming $HOSH$ and continuing with its isomerization into H_2SO are compared in Figure 5. The presence of the ice water molecules significantly stabilizes the addition product, with H_2OS_{ice} being located at −219 and H_2OS at −93 kJ mol^{−1} with respect to their respective reactants asymptotes. Such a large stabilization is due to the H-bonds that H_2OS establishes with the water molecules of the ice surfaces. H-bond cooperativity in H_2OS_{ice} is such that the S–O bond length decreases from 1.89 Å in H_2OS to 1.75 Å in H_2OS_{ice} . $HOSH_{ice}$ and H_2SO_{ice} are also somewhat stabilized with respect to their gas phase counterparts by establishing H bonds with ice water molecules, but to a smaller extent (−274 vs −249 kJ mol^{−1} for $HOSH_{ice}$ and $HOSH$, respectively; −197 vs −176 kJ mol^{−1} for H_2SO_{ice} and H_2SO , respectively). Therefore, H_2OS_{ice} is the primary beneficiary of the presence of water ice, being highly stabilized compared to its gas-phase analog. The presence of the water ice surface also has important effects on energy barriers. Indeed, significant reduction of the energy barriers associated with the $TS1_{ice}$ and $TS2_{ice}$ (20 and 115 kJ mol^{−1}) with respect to the $TS1$ and $TS2$ (48 and 246 kJ mol^{−1}) is observed, which can be explained by the fact that the presence of the ice water molecules assists the H-transfer involved in the molecular rearrangement. Indeed, in $TS1_{ice}$, three water molecules participate in the H-transfer mechanism, while in $TS2_{ice}$, six water molecules are involved. In the gas phase, the $TS1$ and $TS2$ have a highly strained three-membered ring structure. In contrast, in W_{ice} the H-transfer mechanism assisted by water molecules relies on a TS structure less strained with larger rings corresponding to a lower barrier. The imaginary frequencies of each transition state are also reduced when moving from the gas-phase to W_{ice} . The imaginary frequencies for $TS1$ and $TS1_{ice}$ are 1565.32i cm^{−1} and 1001.10i cm^{−1},

respectively, and for $TS2$ and $TS2_{ice}$ are 1786.60i cm^{−1} and 701.89i cm^{−1}. The displacement vectors for $TS1_{ice}$ and $TS2_{ice}$ are shown in Figure 1 of the SI. These decreases confirm the participation of more atoms in the TS_{ice} structures, with the lowering caused by the larger total mass of the atoms involved. Since the curvature along a specific direction on the PES is proportional to the eigenvalue of the Hessian matrix in that direction, which corresponds to the vibrational frequency of a mode, such decreases indicate that the degree of curvature of the potential barrier highest points becomes smoother on W_{ice} , in line with the less-strained TS geometries.

Another significant effect is that we could not identify a TS connecting $MIN3_{ice}$ to the $SO(a^1\Delta) + H_2$ products. This can be easily justified considering that, for the gas-phase reaction, the H_2 elimination occurs via a 3-center elimination mechanism that requires a large distortion of the structure of $MIN3$. The presence of H-bonds with the additional water molecules contrasts the necessary distortion of $MIN3_{ice}$ and the only two-product exothermic channel leading to $SO(a^1\Delta) + H_2$ channel is not open in this case. Also, the channel associated with the S–O bond cleavage from $MIN2$ does not seem to be open starting from $MIN2_{ice}$ because of the additional resistance of the network of H-bonds to which $HOSH$ belongs.

3.3. Kinetics of the Reaction. The rates of the unimolecular isomerization processes occurring on W_{ice} shown in Figure 3 were computed via the RRKM theory to estimate the average reactive lifetime of each species formed on the water ice. As commented above, these estimates are made by considering the cluster as an isolated system that cannot exchange energy or matter with the environment. Once the atom is adsorbed on the cluster, the reaction energy generated is retained as internal energy by the molecule. If the lifetime of this newly formed species is longer than 1 ps, the time in which almost all the reaction energy is dissipated throughout the ice surfaces,^{93,94,97} we can conclude that the isomerization process cannot take place. In contrast, if the lifetime is shorter, the internal energy can be used to overcome the isomerization energy barrier.

Figure 6 shows the unimolecular rate coefficient for the conversion of $\text{H}_2\text{OS}_{\text{ice}}$ to HOSH_{ice} as a function of the energy




Figure 6. Unimolecular rate constants (s^{-1}) for the conversion of HOSH_{ice} to $\text{H}_2\text{OS}_{\text{ice}}$ and HOSH_{ice} to $\text{H}_2\text{SO}_{\text{ice}}$ (upper panel) and $\text{H}_2\text{OS}_{\text{ice}}$ to HOSH_{ice} (bottom panel) as a function of the energy (kJ mol^{-1}).

available to the system. The lifetime of the $\text{H}_2\text{OS}_{\text{ice}}$ species is the inverse of the unimolecular rate constant. The latter corresponds to the initial energy content deriving from the interaction of $\text{S}(^1\text{D})$ with the water molecule of the ice surface (219 kJ mol^{-1}). The calculated lifetime of $\text{H}_2\text{OS}_{\text{ice}}$ with respect to its isomerization is 64.2 ps. In the case of HOSH_{ice} , the lifetime was calculated as the inverse of the sum of the unimolecular rate constants of the processes $\text{HOSH}_{\text{ice}} \rightarrow \text{H}_2\text{SO}_{\text{ice}}$ and $\text{HOSH}_{\text{ice}} \rightarrow \text{H}_2\text{OS}_{\text{ice}}$. Considering an initial energy content of 274 kJ mol^{-1} , the lifetime of HOSH_{ice} is 0.19 ms.

Therefore, the estimated lifetimes of the $\text{H}_2\text{OS}_{\text{ice}}$ and especially of HOSH_{ice} (with respect to their isomerization processes) are much longer than the expected time scale in which the energy released by their formation is dissipated throughout the ice. According to *ab initio* molecular dynamics simulations (aimed at studying the fate of the energy released in exothermic reactions occurring in the presence of water ice), in 1 ps: (1) two-thirds of the energy liberated in the $\text{H} + \text{H}$ recombination reaction (ca. 300 kJ mol^{-1}) is absorbed by the

ice,⁹⁴ (2) 90% of the energy liberated in the $\text{H} + \text{CO} \rightarrow \text{HCO}$ reaction (ca. 130 kJ mol^{-1}) is absorbed by the ice,⁹⁴ and (3) about 60–90% of the energy released in the H additions to N up to the formation of NH_3 is absorbed by the ice.⁹⁷ Considering the large vibrational coupling with the water molecules of the present system, the energy dissipation is expected to occur within the same time scale, if not faster. Therefore, we speculate that isomerization does not occur appreciably, especially in the case of HOSH .

4. DISCUSSION AND ASTROPHYSICAL IMPLICATIONS

The main motivation for the present study was the fact that by comparing the reaction mechanism of $\text{S}(^1\text{D}) + \text{H}_2\text{O}$ in the gas-phase and in the presence of four additional water molecules significant differences were noted.⁶³ However, the use of a minimal cluster like that provided by 4 water molecules (hereafter referred to as 4WMC, see Giustini et al.⁶³) is rather simplistic and important surface effects could have been neglected. Our work aims at gaining deeper insights into the title reaction occurring on ice using a more realistic model of the water ice surface. As a reminder, the 4WMC PES was derived at the $\text{CCSD(T)}/\omega\text{B97XD}$ level, while our results were obtained using the W_{ice} model at the $\text{CCSD(T)}/\omega\text{B97X-D3}$ level. Therefore, the comparison is highly consistent as far as the methodology is concerned.

Based on our results, the effects arising from W_{ice} are even more significant than those using the minimal 4WMC. More specifically: (1) the relative energies (with respect to their asymptotes) of H_2OS and $\text{H}_2\text{OS}_{\text{ice}}$ are -93 and -219 kJ mol^{-1} , respectively, while in the 4WMC model they changed only from -86 to -161 kJ mol^{-1} . This implies a significant extra-stabilization when long-range interactions (i.e., H-bond cooperativity and dispersion forces) are better taken into account. (2) the intrinsic energy barriers are 20 kJ mol^{-1} for TS1_{ice} vs 111 kJ mol^{-1} for $\text{TS1}_{4\text{WMC}}$, and 115 kJ mol^{-1} for TS2_{ice} vs 244 kJ mol^{-1} for $\text{TS2}_{4\text{WMC}}$. Clearly, the 4WMC model, due to its small size, cannot take into account all the stabilizing H-bonds in the TS structures. Moreover, the number of water molecules involved in the water-assisted H-transfers is limited to one on the 4WMC model for both TSs, while on W_{ice} more water molecules are involved. As already commented on, this has the important consequence of strongly reducing the geometrical strain of the TSs, and hence an enhanced decrease of the associated energy barriers. In fact, with 4WMC, only one water molecule assists the proton transfer process, forming a six-member ring, whereas in the presence of W_{ice} for TS1_{ice} , three water molecules assist the process, forming a 10-member ring. Minor differences are instead noted for HOSH and H_2SO , the relative energies of which are -249 for HOSH vs -274 kJ mol^{-1} for HOSH_{ice} (to be compared with the Giustini et al. values of -244 for HOSH and -250 kJ mol^{-1} for $\text{HOSH}_{4\text{WMC}}$), and -176 for H_2SO vs -197 kJ mol^{-1} for $\text{H}_2\text{SO}_{\text{ice}}$ (to be compared with the Giustini et al. values of -171 for H_2SO and -180 kJ mol^{-1} for $\text{H}_2\text{SO}_{4\text{WMC}}$). This means that the stabilization of these two species is similar irrespectively of the water ice cluster model, as both water clusters can hold similar H-bond interactions with them. Finally, the only open two-product channel for the gas-phase reaction is not active on ice since $\text{H}_2\text{SO}_{\text{ice}}$ is never formed and the critical geometry of the 3-center elimination transition state could not be reached because the H-bonds that

$\text{H}_2\text{SO}_{\text{ice}}$ forms with the surrounding water molecules will impede it.

Despite these differences, our work corroborates the suggestion put forward by Giustini et al. We verified that, once formed by the reaction of $\text{S}(^1\text{D})$ with the water ice molecules, $\text{H}_2\text{OS}_{\text{ice}}$ and HOSH_{ice} will be stabilized by the ice. By comparing the reaction lifetimes in the most favorable scenarios (all the energy retained among the cluster molecules) with the energy dissipation time scale consistently derived in three different cases by using ab initio molecular dynamics, we can conclude that isomerization will not be significant.

$\text{H}_2\text{OS}_{\text{ice}}$ and HOSH_{ice} can accumulate on the ice surface or in the ice bulk, especially in the case of HOSH (hydrogen thioperoxide), the most stable H_2SO isomer. $\text{H}_2\text{OS}_{\text{ice}}$ can react on the icy surface of interstellar grains with H atoms ultimately being transformed into HOSH (for a similar process, see Góbi et al.⁹⁸). HOSH could undergo the typical interstellar ice reactions (namely, hydrogenation), but for this species, we expect a significant barrier, as already observed for the similar species H_2O_2 , whose reaction with H is characterized by a very large barrier (ca. 27 kJ mol⁻¹), which increases when adding 3 water molecules.⁹⁹ Even considering the tunneling effect at the very low temperature of interstellar ice, a significant portion of HOSH is expected to remain unchanged, as demonstrated by ice models in the case of H_2O_2 .

Our work also supports the modeling approach by Garrod⁵⁹ and Carder et al.,⁵⁸ who were the first to explicitly consider PDI reactions on ice. Once formed by the photodissociation (or GCR radiolysis) of their precursors, very reactive species like $\text{S}(^1\text{D})$ or $\text{O}(^1\text{D})$ will immediately react with the surrounding molecules. Water is the best candidate, being the most abundant species in interstellar or cometary ice by far. The radiative lifetime of $\text{O}(^1\text{D})$ was estimated to be 1 s in solid matrices. Considering that the lifetime of $\text{O}(^1\text{D})$ is ca. 3 times larger than that of $\text{S}(^1\text{D})$ in the gas phase, we expect that the lifetime of $\text{S}(^1\text{D})$ in ice will be a fraction of one second. During this time, it reacts with water and forms either H_2OS or HOSH. Notably, HOSH and H_2SO can also form by the reaction of $\text{O}(^1\text{D})$ if produced by UV photodissociation of $\text{H}_2\text{O}/\text{CO}_2/\text{CO}$ in the vicinity of one H_2S molecule.^{100,101} To the best of our knowledge, HOSH has never been detected so far in the interstellar medium, even though its presence has been already speculated.^{102,103} A related species, the HSO radical, was recently detected toward several cold dark clouds by using the Yebes 40 m telescope.¹⁰⁴ HSO resulted to be a widespread molecule in cold regions, but its formation mechanism is still obscure.¹⁰⁴ The gas-phase formation routes suggested so far are either endothermic (like the reaction between OH and SH, see Figure 2) or characterized by a significant barrier (like the $\text{O} + \text{H}_2\text{S}$ reaction¹⁰¹) while no formation routes from ion–molecule reactions have been envisaged.¹⁰⁴

Our results can also contribute to explaining the presence of a peak associated with H_2SO (and HSO) in the mass spectrum recorded by ROSINA on September 5, 2016 after a chunk of ice and dust hit the instrument. In that occasion, sublimation of less volatile species occurred inside the equilibrium chamber of ROSINA at a temperature of 273 K, so increasing the signal of all the sulfur species already detected and causing the appearance of new species in the mass spectrum. It is unclear if molecules with this formula were present in the ice or if the signal at these masses originated by dissociative ionization of larger molecules in the mass spectrometer. However, the signal

recorded at $m/z = 50$ attributed by Mahjoub et al.⁴⁶ to H_2SO^+ can plausibly be associated with the electron impact ionization of HSOH. The ice reaction investigated in this manuscript can be responsible for the formation and accumulation over time of HOSH that, being involved in strong H-bonds with the water molecules of ice, is not easily released in the gas phase. Notably, the similar species H_2O_2 , rarely identified in the ISM, has been detected in the coma of the 67P comet.¹⁰⁵

To conclude, a more general remark is in order. In the ice models by Carder et al.,⁵⁸ the branching ratios of the gas-phase reactions have been adopted also for the $\text{O}(^1\text{D})$ reactions on ice (except for the reaction with CH_4 for which the value of Bergner et al.⁵⁵ was used). This is a common approach when the data necessary to model ice chemistry are missing. However, our results clearly indicate that this approximation can cause serious mistakes. The presence of water molecules can impede the reaching of critical configurations for the reaction to occur while the formation of two products can be seriously disfavored by the presence of H-bonds. In most cases, if a closed-shell polar molecule is formed by the reaction of the electronically excited singlet species, it will easily be stabilized by energy dissipation.

5. CONCLUSIONS

In this work, we have investigated the $\text{S}(^1\text{D}) + \text{H}_2\text{O}$ reaction occurring on an ice surface model of 18 water molecules using electronic structure and kinetics calculations. As in the gas-phase reaction, the initial approach of $\text{S}(^1\text{D})$ can occur via the addition to the lone pair of an O atom of water (H_2OS is formed), or via insertion into the O–H σ bonds of water (HOSH is formed). The initial approaches are the same as seen for the gas-phase reaction. However, on the water ice cluster, the reaction stops at the formation of H_2OS and HSOH. For this specific reaction, we can clearly observe all the effects that water ice can exert: as a concentrator/supplier of reactants, as a third body that stabilizes the products formed, and as a chemical catalyst that provides alternative routes with lower energy barriers. Indeed, the ice surface stabilizes the newly formed products by acting as an energy dissipator thus preventing them from dissociating into two products or isomerizing. The catalytic effect exerted by the ice surface is also very clear: the energy barriers along the minimum energy path are significantly lowered because of the stabilization of the TS structures by H-bonding within the structures, and the occurrence of water-assisted H-transfer mechanisms.

Our results can be of help in elucidating the mysterious sulfur chemistry occurring in the icy mantles of interstellar grains or in cometary nuclei. We suggest that HOSH accumulates in interstellar ice. Furthermore, this study demonstrates that the product branching ratios of gas-phase reactions should not be uncritically used in modeling interstellar ice chemistry.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsearthspacechem.4c00351>.

Natural population analysis; attack sites; displacement vectors for TS1_{ice} and TS2_{ice} ; energetics and errors of the benchmarking study; and optimized Cartesian coordinates (PDF)

AUTHOR INFORMATION

Corresponding Authors

Gabriella Di Genova – Dipartimento di Chimica, Biologia e Biotecnologie, Università degli Studi di Perugia, 06123 Perugia, Italy; Email: gabriella.digenova@dottorandi.unipg.it

Albert Rimola – Departament de Química, Universitat Autònoma de Barcelona, 08193 Catalonia, Spain; orcid.org/0000-0002-9637-4554; Email: albert.rimola@uab.cat

Nadia Balucani – Dipartimento di Chimica, Biologia e Biotecnologie, Università degli Studi di Perugia, 06123 Perugia, Italy; orcid.org/0000-0001-5121-5683; Email: nadia.balucani@unipg.it

Authors

Jessica Perrero – Departament de Química, Universitat Autònoma de Barcelona, 08193 Catalonia, Spain; Dipartimento di Chimica and Nanostructured Interfaces and Surfaces (NIS) Centre, Università degli Studi di Torino, 10125 Torino, Italy

Marzio Rosi – Dipartimento di Ingegneria Civile e Ambientale, Università degli Studi di Perugia, 06125 Perugia, Italy; orcid.org/0000-0002-1264-3877

Cecilia Ceccarelli – Univ. Grenoble Alpes, CNRS, Institut de Planétologie et d'Astrophysique de Grenoble (IPAG), 38100 Grenoble, France; orcid.org/0000-0001-9664-6292

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acsearthspacechem.4c00351>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

G.D.G., M.R., C.C., and N.B. thank the Italian MUR PRIN2020 (2020AFB3FX-Astrochemistry beyond the second period elements) and the European Union's Horizon 2020 research and innovation program from the European Research Council (ERC)(project 'The Dawn of Organic Chemistry' DOC, grant agreement No 741002) for support. M.R. acknowledges financial support under the National Recovery and Resilience Plan (NRRP), Mission 4, Component 2, Investment 1.1, Call for tender No. 104 published on 2.2.2022 by the Italian Ministry of University and Research (MUR), funded by the European Union—NextGenerationEU—Project Title 2022JC2Y93 ChemicalOrigins: linking the fossil composition of the Solar System with the chemistry of protoplanetary disks—CUP J53D23001600006—Grant Assignment Decree No. 962 adopted on 30.06.2023 by the Italian Ministry of University and Research (MUR). Support from the Italian Space Agency (Bando ASI Prot. n. DC-DSR-UVS-2022-231, Grant no. 2023-10-U.0 MIGLIORA) is also acknowledged (J.P. and N.B.). This project has received funding within the European Union's Horizon 2020 research and innovation program from the European Research Council (ERC) for the project "Quantum Chemistry on Interstellar Grains" (QUANTUMGRAIN), grant agreement no. 865657 (A.R.). MICINN is also acknowledged for funding the projects PID2021-126427NB-I00 and CNS2023-144902 (A.R.). A.R. gratefully acknowledges support through 2023 ICREA Award.

REFERENCES

- (1) Lodders, K. Relative atomic solar system abundances, mass fractions, and atomic masses of the elements and their isotopes, composition of the solar photosphere, and compositions of the major chondritic meteorite groups. *Space Sci. Rev.* **2021**, 217, 44.
- (2) Endres, C. P.; Schlemmer, S.; Schilke, P.; Stutzki, J.; Müller, H. S. The Cologne database for molecular spectroscopy, CDMS, in the virtual atomic and molecular data centre, VAMDC. *J. Mol. Spectrosc.* **2016**, 327, 95–104.
- (3) McGuire, B. A. 2021 Census of Interstellar, Circumstellar, Extragalactic, Protoplanetary Disk, and Exoplanetary Molecules. *Astrophysical Journal Supplement Series* **2022**, 259, 30.
- (4) Gaillard, F.; Michalski, J.; Berger, G.; McLennan, S. M.; Scaillet, B. Geochemical reservoirs and timing of sulfur cycling on Mars. *Space Science Reviews* **2013**, 174, 251–300.
- (5) Wong, M. H.; Mahaffy, P. R.; Atreya, S. K.; Niemann, H. B.; Owen, T. C. Updated Galileo probe mass spectrometer measurements of carbon, oxygen, nitrogen, and sulfur on Jupiter. *Icarus* **2004**, 171, 153–170.
- (6) Esposito, L. W. Planetary science: Rising sulphur on Venus. *Nature Geoscience* **2013**, 6, 20–21.
- (7) Steenstra, E.; Seegers, A.; Eising, J.; Tomassen, B.; Webers, F.; Berndt, J.; Klemme, S.; Matveev, S.; van Westrenen, W. Evidence for a sulfur-undersaturated lunar interior from the solubility of sulfur in lunar melts and sulfide-silicate partitioning of siderophile elements. *Geochim. Cosmochim. Acta* **2018**, 231, 130–156.
- (8) Feaga, L. M.; McGrath, M.; Feldman, P. D. Io's dayside SO₂ atm. *Icarus* **2009**, 201, 570–584.
- (9) Mumma, M. J.; Charnley, S. B. The chemical composition of comets—emerging taxonomies and natal heritage. *Annual Review of Astronomy and Astrophysics* **2011**, 49, 471–524.
- (10) Altwegg, K.; Balsiger, H.; Fuselier, S. A. Cometary chemistry and the origin of icy solar system bodies: the view after Rosetta. *Annual Review of Astronomy and Astrophysics* **2019**, 57, 113–155.
- (11) Yoshimura, T.; et al. Chemical evolution of primordial salts and organic sulfur molecules in the asteroid 162173 Ryugu. *Nat. Commun.* **2023**, 14, 5284.
- (12) Zhrebekker, A.; Kostyukovich, Y.; Volkov, D. S.; Chumakov, R. G.; Friederici, L.; Rüger, C. P.; Kononikhin, A.; Kharybin, O.; Korochantsev, A.; Zimmermann, R.; Perminova, I. V.; Nikolaev, E. Speciation of organosulfur compounds in carbonaceous chondrites. *Sci. Rep.* **2021**, 11, 7410.
- (13) Bradley, J. P.; Ishii, H. A.; Bustillo, K.; Ciston, J.; Ogliore, R.; Stephan, T.; Brownlee, D. E.; Joswiak, D. J. On the provenance of GEMS, a quarter century post discovery. *Geochim. Cosmochim. Acta* **2022**, 335, 323–338.
- (14) Alexander, C. M. O.; Wynn, J. G.; Bowden, R. Sulfur abundances and isotopic compositions in bulk carbonaceous chondrites and insoluble organic material: Clues to elemental and isotopic fractionations of volatile chalcophiles. *Meteoritics & Planetary Science* **2022**, 57, 334–351.
- (15) Biology and brimstone. *Nat. Chem. Biol.* **2006**, 2, 169.
- (16) Rivière-Marichalar, P.; Fuente, A.; Goicoechea, J.; Pety, J.; Le Gal, R.; Gratier, P.; Guzmán, V.; Roueff, E.; Loison, J.; Wakelam, V.; et al. Abundances of sulphur molecules in the Horsehead nebula-First NS⁺ detection in a photodissociation region. *Astronomy & Astrophysics* **2019**, 628, A16.
- (17) Boogert, A.; Gerakines, P.; Whittet, D. Observations of the icy universe. *Annu. Rev. Astron. Astrophys.* **2015**, 53, 541–581.
- (18) McClure, M. K.; Rocha, W.; Pontoppidan, K.; Crouzet, N.; Chu, L. E.; Dartois, E.; Lamberts, T.; Noble, J.; Pendleton, Y.; Perotti, G.; et al. An Ice Age JWST inventory of dense molecular cloud ices. *Nature astronomy* **2023**, 7, 431–443.
- (19) Rocha, W. R. M.; et al. JWST Observations of Young protoStars (JOYS+): Detecting icy complex organic molecules and ions - I. CH₄⁺, SO₂⁺, HCOO⁺, OCN⁺, H₂CO, HCOOH, CH₃CH₂OH, CH₃CHO, CH₃OCHO, and CH₃COOH. *Astron. Astrophys.* **2024**, 683, A124.

- (20) Keller, L.; Hony, S.; Bradley, J.; Molster, F.; Waters, L.; Bouwman, J.; De Koter, A.; Brownlee, D.; Flynn, G.; Henning, T.; et al. Identification of iron sulphide grains in protoplanetary disks. *Nature* **2002**, *417*, 148–150.
- (21) Wakelam, V.; Caselli, P.; Ceccarelli, C.; Herbst, E.; Castets, A. Resetting chemical clocks of hot cores based on S-bearing molecules. *Astronomy & Astrophysics* **2004**, *422*, 159–169.
- (22) Vastel, C.; Quénard, D.; Le Gal, R.; Wakelam, V.; Andrianasolo, A.; Caselli, P.; Vidal, T.; Ceccarelli, C.; Lefloch, B.; Bachiller, R. Sulphur chemistry in the L1544 pre-stellar core. *Mon. Not. R. Astron. Soc.* **2018**, *478*, S514–S532.
- (23) Laas, J. C.; Caselli, P. Modeling sulfur depletion in interstellar clouds. *Astronomy & Astrophysics* **2019**, *624*, A108.
- (24) Fuente, A.; Navarro, D.; Caselli, P.; Gerin, M.; Kramer, C.; Roueff, E.; Alonso-Albi, T.; Bachiller, R.; Cazaux, S.; Commerçon, B.; et al. Gas phase Elemental abundances in Molecular clouds (GEMS)-I. The prototypical dark cloud TMC 1. *Astron. Astrophys.* **2019**, *624*, A105.
- (25) Fuente, A.; et al. Gas phase Elemental abundances in Molecular clouds (GEMS). VII. Sulfur elemental abundance. *Astron. Astrophys.* **2023**, *670*, A114.
- (26) Perrero, J.; Beitia-Antero, L.; Fuente, A.; Ugliengo, P.; Rimola, A. Theoretical modelling of the adsorption of neutral and charged sulphur-bearing species on to olivine nanoclusters. *Mon. Not. R. Astron. Soc.* **2023**, *527*, 10697–10704.
- (27) Ruffle, D.; Hartquist, T.; Caselli, P.; Williams, D. The sulphur depletion problem. *Mon. Not. R. Astron. Soc.* **1999**, *306*, 691–695.
- (28) Bariosco, V.; Pantaleone, S.; Ceccarelli, C.; Rimola, A.; Balucani, N.; Corno, M.; Ugliengo, P. The binding energy distribution of H₂S: why it is not the major sulphur reservoir of the interstellar ices. *Mon. Not. R. Astron. Soc.* **2024**, *531*, 1371–1384.
- (29) Jiménez-Escobar, A.; Caro, G. M. Sulfur depletion in dense clouds and circumstellar regions-I. H₂S ice abundance and UV-photochemical reactions in the H₂O-matrix. *Astron. Astrophys.* **2011**, *536*, A91.
- (30) Shingledecker, C. N.; Lamberts, T.; Laas, J. C.; Vasyunin, A.; Herbst, E.; Kästner, J.; Caselli, P. Efficient production of S₈ in interstellar ices: The effects of cosmic-ray-driven radiation chemistry and nondiffusive bulk reactions. *Astrophysical Journal* **2020**, *888*, 52.
- (31) Cazaux, S.; Carrascosa, H.; Caro, G. M.; Caselli, P.; Fuente, A.; Navarro-Almada, D.; Riviere-Marichalar, P. Photoprocessing of H₂S on dust grains-Building S chains in translucent clouds and comets. *Astron. Astrophys.* **2022**, *657*, A100.
- (32) Capria, M. T.; Capaccioni, F.; Filacchione, G.; Tosi, F.; De Sanctis, M. C.; Mottola, S.; Ciarniello, S.; Formisano, M.; Longobardo, A.; Migliorini, A.; et al. How pristine is the interior of the comet 67P/Churyumov–Gerasimenko? *Mon. Not. R. Astron. Soc.* **2017**, *469*, S685–S694.
- (33) Bockelée-Morvan, D.; Biver, N. The composition of cometary ices. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **2017**, *375*, 20160252.
- (34) Smith, A. M.; Stecher, T. P.; Casswell, L. Production of carbon, sulfur, and CS in Comet West. *Astrophys. J.* **1980**, *242*, 402–410.
- (35) Crovisier, J.; Bockelée-Morvan, D. Remote Observations of the Composition of Cometary Volatiles. *Space Science Reviews* **1999**, *90*, 19–32.
- (36) Bockelée-Morvan, D.; Colom, P.; Crovisier, J.; Despois, D.; Paubert, G. Microwave detection of hydrogen sulphide and methanol in comet Austin (1989c1). *Nature* **1991**, *350*, 318–320.
- (37) Woodney, L.; McMullin, J.; A'Hearn, M. Detection of OCS in comet Hyakutake (C/1996 B2). *Planetary and space science* **1997**, *45*, 717–719.
- (38) Bockelée-Morvan, D.; et al. New molecules found in comet C/1995 O1 (Hale-Bopp). Investigating the link between cometary and interstellar material. *Astron. Astrophys.* **2000**, *353*, 1101–1114.
- (39) Jackson, W. M.; Scodinu, A.; Xu, D.; Cochran, A. L. Using the Ultraviolet and Visible spectrum of Comet 122P/de Vico to Identify the Parent Molecule CS₂. *Astrophysical Journal, Letters* **2004**, *607*, L139–L141.
- (40) A'Hearn, M. F.; Schleicher, D. G.; Feldman, P. D. The discovery of S₂ in comet IRAS-Araki-Alcock 1983d. *Astrophys. J., Lett.* **1983**, *274*, L99–L103.
- (41) Woodney, L.; A'Hearn, M.; McMullin, J.; Samarasingha, N. Sulfur chemistry at millimeter wavelengths in C/Hale-Bopp. *Earth, Moon, and Planets* **1997**, *78*, 69.
- (42) Irvine, W. M.; Senay, M.; Lovell, A. J.; Matthews, H. E.; McGonagle, D.; Meier, R. NOTE: Detection of Nitrogen Sulfide in Comet Hale-Bopp. *Icarus* **2000**, *143*, 412–414.
- (43) Calmonte, U.; Altwegg, K.; Balsiger, H.; Berthelier, J.-J.; Bieler, A.; Cessateur, G.; Dhooghe, F.; Van Dishoeck, E.; Fiethe, B.; Fuselier, S.; et al. Sulphur-bearing species in the coma of comet 67P/Churyumov–Gerasimenko. *Mon. Not. R. Astron. Soc.* **2016**, *462*, S253–S273.
- (44) Le Roy, L.; Altwegg, K.; Balsiger, H.; Berthelier, J.-J.; Bieler, A.; Brioso, C.; Calmonte, U.; Combi, M. R.; De Keyser, J.; Dhooghe, F.; et al. Inventory of the volatiles on comet 67P/Churyumov–Gerasimenko from Rosetta/ROSINA. *Astronomy & Astrophysics* **2015**, *583*, A1.
- (45) Hänni, N.; Altwegg, K.; Combi, M.; Fuselier, S.; De Keyser, J.; Rubin, M.; Wampfler, S. Identification and characterization of a new ensemble of cometary organic molecules. *Nat. Commun.* **2022**, *13*, 3639.
- (46) Mahjoub, A.; Altwegg, K.; Poston, M. J.; Rubin, M.; Hodyss, R.; Choukroun, M.; Ehlmann, B. L.; Hänni, N.; Brown, M. E.; Blacksborg, J.; Eiler, J. M.; Hand, K. P. Complex organosulfur molecules on comet 67P: Evidence from the ROSINA measurements and insights from laboratory simulations. *Sci. Adv.* **2023**, *9*, No. eadh0394.
- (47) Schmitt-Kopplin, P.; Hertkorn, N.; Harir, M.; Moritz, F.; Lucio, M.; Bonal, L.; Quirico, E.; Takano, Y.; Dworkin, J. P.; Naraoka, H.; et al. Soluble organic matter Molecular atlas of Ryugu reveals cold hydrothermalism on C-type asteroid parent body. *Nat. Commun.* **2023**, *14*, 6525.
- (48) Naraoka, H.; et al. Soluble organic molecules in samples of the carbonaceous asteroid (162173) Ryugu. *Science* **2023**, *379*, No. eabn9033.
- (49) Schofield, K. Critically evaluated rate constants for gaseous reactions of several electronically excited species. *Journal of physical and chemical reference data* **1979**, *8*, 723–798.
- (50) Meier, R.; A'Hearn, M. F. Atomic Sulfur in Cometary Comae Based on UV Spectra of the SITriplet near 1814 Å. *Icarus* **1997**, *125*, 164–194.
- (51) Cook, P. A.; Langford, S. R.; Dixon, R. N.; Ashfold, M. N. An experimental and ab initio reinvestigation of the Lyman- α photodissociation of H₂S and D₂S. *J. Chem. Phys.* **2001**, *114*, 1672–1684.
- (52) Zhao, Y.; Luo, Z.; Chang, Y.; Wu, Y.; Zhang, S.-E.; Li, Z.; Ding, H.; Wu, G.; Campbell, J. S.; Hansen, C. S.; et al. Rotational and nuclear-spin level dependent photodissociation dynamics of H₂S. *Nat. Commun.* **2021**, *12*, 4459.
- (53) Chang, Y.; Fu, Y.; Chen, Z.; Luo, Z.; Zhao, Y.; Li, Z.; Zhang, W.; Wu, G.; Fu, B.; Zhang, D. H.; et al. Vacuum ultraviolet photodissociation of sulfur dioxide and its implications for oxygen production in the early Earth's atmosphere. *Chemical Science* **2023**, *14*, 8255–8261.
- (54) Lupu, R. E.; Feldman, P. D.; Weaver, H. A.; Tozzi, G.-P. The Fourth Positive System of Carbon Monoxide in the Hubble Space Telescope Spectra of Comets. *Astrophysical Journal* **2007**, *670*, 1473.
- (55) Bergner, J. B.; Öberg, K. I.; Rajappan, M. Methanol formation via oxygen insertion chemistry in ices. *Astrophysical Journal* **2017**, *845*, 29.
- (56) Bergner, J. B.; Öberg, K. I.; Rajappan, M. Oxygen atom reactions with C₂H₆, C₂H₄, and C₂H₂ in ices. *Astrophysical Journal* **2019**, *874*, 115.
- (57) Jin, M.; Garrod, R. T. Formation of Complex Organic Molecules in Cold Interstellar Environments through Nondiffusive Grain-surface and Ice-mantle Chemistry. *Astrophysical Journal Supplement Series* **2020**, *249*, 26.

- (58) Carder, J. T.; Ochs, W.; Herbst, E. Modelling the insertion of O (^1D) into methane on the surface of interstellar ice mantles. *Mon. Not. R. Astron. Soc.* **2021**, *508*, 1526–1532.
- (59) Garrod, R. T. Simulations of Ice Chemistry in Cometary Nuclei. *Astrophysical Journal* **2019**, *884*, 69.
- (60) Berteloite, C.; Le Picard, S. D.; Sims, I. R.; Rosi, M.; Leonori, F.; Petrucci, R.; Balucani, N.; Wang, X.; Casavecchia, P. Low temperature kinetics, crossed beam dynamics and theoretical studies of the reaction $S(^1D) + CH_4$ and low temperature kinetics of $S(^1D) + C_2H_2$. *Phys. Chem. Chem. Phys.* **2011**, *13*, 8485–8501.
- (61) Leonori, F.; Petrucci, R.; Balucani, N.; Casavecchia, P.; Rosi, M.; Skouteris, D.; Berteloite, C.; Le Picard, S. D.; Canosa, A.; Sims, I. R. Crossed-beam dynamics, low-temperature kinetics, and theoretical studies of the reaction $S(^1D) + C_2H_4$. *J. Phys. Chem. A* **2009**, *113*, 15328–15345.
- (62) Leonori, F.; Petrucci, R.; Balucani, N.; Hickson, K. M.; Hamberg, M.; Geppert, W. D.; Casavecchia, P.; Rosi, M. Crossed-beam and theoretical studies of the $S(^1D) + C_2H_2$ reaction. *J. Phys. Chem. A* **2009**, *113*, 4330–4339.
- (63) Giustini, A.; Di Genova, G.; Skouteris, D.; Ceccarelli, C.; Rosi, M.; Balucani, N. Gas-Phase and Model Ice-Surface Reactions of S (^1D) with Water and Methanol: A Computational Investigation and Implications for Cosmochemistry/Astrochemistry. *ACS Earth and Space Chemistry* **2024**, *8*, 2318–2333.
- (64) Karu, E.; Li, M.; Ernle, L.; Brenninkmeijer, C. A. M.; Lelieveld, J.; Williams, J. Carbonyl Sulfide (OCS) in the Upper Troposphere/Lowermost Stratosphere (UT/LMS) Region: Estimates of Lifetimes and Fluxes. *Geophys. Res. Lett.* **2023**, *50*, No. e2023GL105826.
- (65) Noble, J. A.; Fraser, H.; Smith, Z.; Dartois, E.; Boogert, A.; Cuppen, H.; Dickinson, H.; Dulieu, F.; Egami, E.; Erkal, J.; et al. Detection of the elusive dangling OH ice features at $\sim 2.7 \mu m$ in Chamaeleon I with JWST NIRC2. *Nat. Astron.* **2024**, *8*, 1169–1180.
- (66) Rimola, A.; Taquet, V.; Ugliengo, P.; Balucani, N.; Ceccarelli, C. Combined quantum chemical and modeling study of CO hydrogenation on water ice. *Astronomy & Astrophysics* **2014**, *572*, A70.
- (67) Rimola, A.; Skouteris, D.; Balucani, N.; Ceccarelli, C.; Enrique-Romero, J.; Taquet, V.; Ugliengo, P. Can formamide be formed on interstellar ice? An atomistic perspective. *ACS Earth and Space Chemistry* **2018**, *2*, 720–734.
- (68) Enrique-Romero, J.; Rimola, A.; Ceccarelli, C.; Ugliengo, P.; Balucani, N.; Skouteris, D. Reactivity of HCO with CH_3 and NH_2 on water ice surfaces. a comprehensive accurate quantum chemistry study. *ACS Earth and Space Chemistry* **2019**, *3*, 2158–2170.
- (69) Perrero, J.; Enrique-Romero, J.; Martínez-Bachs, B.; Ceccarelli, C.; Balucani, N.; Ugliengo, P.; Rimola, A. Non-energetic formation of ethanol via CCH reaction with interstellar H_2O ices. A computational chemistry study. *ACS Earth and Space Chemistry* **2022**, *6*, 496–511.
- (70) Frisch, M.; Trucks, G.; Schlegel, H.; Scuseria, G.; Robb, M.; Cheeseman, J.; Montgomery, Jr, J.; Vreven, T.; Kudin, K.; Burant, J. et al. *Gaussian 03, Revision C.02*; Gaussian, Inc.: Wallingford, CT, 2004. Google Scholar There is no corresponding record for this reference 2013.
- (71) Neese, F. Software update: The ORCA program system—Version 5.0. *Wiley Interdiscip. Rev.: Comput. Mol. Sci.* **2022**, *12*, No. e1606.
- (72) Bartlett, R. J. Many-body perturbation theory and coupled cluster theory for electron correlation in molecules. *Annu. Rev. Phys. Chem.* **1981**, *32*, 359–401.
- (73) Olsen, J.; Koch, H.; Balkova, A.; Bartlett, R. J.; et al. Full configuration–interaction and state of the art correlation calculations on water in a valence double-zeta basis with polarization functions. *J. Chem. Phys.* **1996**, *104*, 8007–8015.
- (74) Raghavachari, K.; Trucks, G. W.; Pople, J. A.; Head-Gordon, M. A fifth-order perturbation comparison of electron correlation theories. *Chem. Phys. Lett.* **1989**, *157*, 479–483.
- (75) Kendall, R. A.; Dunning, T. H.; Harrison, R. J. Electron affinities of the first-row atoms revisited. Systematic basis sets and wave functions. *J. Chem. Phys.* **1992**, *96*, 6796–6806.
- (76) Woon, D. E.; Dunning, T. H., Jr Gaussian basis sets for use in correlated molecular calculations. III. The atoms aluminum through argon. *J. Chem. Phys.* **1993**, *98*, 1358–1371.
- (77) Dunning, T. H., Jr Gaussian basis sets for use in correlated molecular calculations. I. The atoms boron through neon and hydrogen. *J. Chem. Phys.* **1989**, *90*, 1007–1023.
- (78) Dunning, T. H., Jr; Peterson, K. A.; Wilson, A. K. Gaussian basis sets for use in correlated molecular calculations. X. The atoms aluminum through argon revisited. *J. Chem. Phys.* **2001**, *114*, 9244–9253.
- (79) Werner, H.-J.; Adler, T. B.; Manby, F. R. General orbital invariant MP2-F12 theory. *J. Chem. Phys.* **2007**, *126*, 164102.
- (80) Valeev, E. F. Improving on the resolution of the identity in linear R12 ab initio theories. *Chem. Phys. Lett.* **2004**, *395*, 190–195.
- (81) Klopper, W.; Samson, C. C. M. Explicitly correlated second-order Møller-Plesset methods with auxiliary basis sets. *J. Chem. Phys.* **2002**, *116*, 6397–6410.
- (82) Riplinger, C.; Pinski, P.; Becker, U.; Valeev, E. F.; Neese, F. Sparse maps—A systematic infrastructure for reduced-scaling electronic structure methods. II. Linear scaling domain based pair natural orbital coupled cluster theory. *J. Chem. Phys.* **2016**, *144*, No. 024109.
- (83) Chai, J.-D.; Head-Gordon, M. Systematic optimization of long-range corrected hybrid density functionals. *J. Chem. Phys.* **2008**, *128*, No. 084106.
- (84) Zheng, J.; Xu, X.; Truhlar, D. G. Minimally augmented Karlsruhe basis sets. *Theor. Chem. Acc.* **2011**, *128*, 295–305.
- (85) Gonzalez, C.; Schlegel, H. B. An improved algorithm for reaction path following. *J. Chem. Phys.* **1989**, *90*, 2154–2161.
- (86) Gonzalez, C.; Schlegel, H. B. Reaction path following in mass-weighted internal coordinates. *J. Phys. Chem.* **1990**, *94*, 5523–5527.
- (87) Neese, F. Definition of corresponding orbitals and the diradical character in broken symmetry DFT calculations on spin coupled systems. *J. Phys. Chem. Solids* **2004**, *65*, 781–785.
- (88) Gilbert, R. G.; Smith, S. C. *Theory of unimolecular and recombination reactions*; Blackwell Scientific Publications, 1990.
- (89) Leonori, F.; Skouteris, D.; Petrucci, R.; Casavecchia, P.; Rosi, M.; Balucani, N. Combined crossed beam and theoretical studies of the $C(^1D) + CH_4$ reaction. *J. Chem. Phys.* **2013**, *138*, No. 024311.
- (90) Balucani, N.; Caracciolo, A.; Vanuzzo, G.; Skouteris, D.; Rosi, M.; Pacifici, L.; Casavecchia, P.; Hickson, K. M.; Loison, J.-C.; Dobrijevic, M. An experimental and theoretical investigation of the $N(^2D) + C_6H_6$ (benzene) reaction with implications for the photochemical models of Titan. *Faraday Discuss.* **2023**, *245*, 327–351.
- (91) Enrique-Romero, J.; Ceccarelli, C.; Rimola, A.; Skouteris, D.; Balucani, N.; Ugliengo, P. Theoretical computations on the efficiency of acetaldehyde formation on interstellar icy grains. *A&A* **2021**, *655*, A9.
- (92) Enrique-Romero, J.; Rimola, A.; Ceccarelli, C.; Ugliengo, P.; Balucani, N.; Skouteris, D. Quantum Mechanical Simulations of the Radical–Radical Chemistry on Icy Surfaces. *Astrophysical Journal Supplement Series* **2022**, *259*, 39.
- (93) Pantaleone, S.; Enrique-Romero, J.; Ceccarelli, C.; Ugliengo, P.; Balucani, N.; Rimola, A. Chemical desorption versus energy dissipation: insights from ab initio molecular dynamics of HCO-formation. *Astrophys. J.* **2020**, *897*, 56.
- (94) Pantaleone, S.; Enrique-Romero, J.; Ceccarelli, C.; Ferrero, S.; Balucani, N.; Rimola, A.; Ugliengo, P. H_2 Formation on Interstellar Grains and the Fate of Reaction Energy. *Astrophys. J.* **2021**, *917*, 49.
- (95) Atkinson, R.; Baulch, D. L.; Cox, R. A.; Crowley, J. N.; Hampson, R. F.; Hynes, R. G.; Jenkin, M. E.; Rossi, M. J.; Troe, J. Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I - gas phase reactions of O_x , HO_x , NO_x and SO_x species. *Atmospheric Chemistry and Physics* **2004**, *4*, 1461–1738.

- (96) Borin, A. C.; Ornellas, F. R. The lowest triplet and singlet electronic states of the molecule SO. *Chemical physics* **1999**, *247*, 351–364.
- (97) Ferrero, S.; Pantaleone, S.; Ceccarelli, C.; Ugliengo, P.; Sodupe, M.; Rimola, A. Where does the energy go during the interstellar NH₃ formation on water ice? A computational study. *Astrophysical Journal* **2023**, *944*, 142.
- (98) Góbi, S.; Csonka, I. P.; Bazsó, G.; Tarczay, G. Successive Hydrogenation of SO and SO₂ in Solid para-H₂: Formation of Elusive Small Oxoacids of Sulfur. *ACS Earth and Space Chemistry* **2021**, *5*, 1180–1195.
- (99) Lamberts, T.; Samanta, P. K.; Köhn, A.; Kästner, J. Quantum tunneling during interstellar surface-catalyzed formation of water: the reaction $\text{H} + \text{H}_2\text{O}_2 \rightarrow \text{H}_2\text{O} + \text{OH}$. *Phys. Chem. Chem. Phys.* **2016**, *18*, 33021–33030.
- (100) Balucani, N.; Beneventi, L.; Casavecchia, P.; Stranges, D.; Volpi, G. The effect of reagent electronic energy on the dynamics of chemical reactions: A high-resolution crossed beam study of O (³P,¹D) + H₂S. *J. Chem. Phys.* **1991**, *94*, 8611–8614.
- (101) Balucani, N.; Stranges, D.; Casavecchia, P.; Volpi, G. G. Crossed beam studies of the reactions of atomic oxygen in the ground ³P and first electronically excited ¹D states with hydrogen sulfide. *J. Chem. Phys.* **2004**, *120*, 9571–9582.
- (102) Palmer, C. Z.; Fortenberry, R. C.; Francisco, J. S. Spectral Signatures of Hydrogen Thioperoxide (HOSH) and Hydrogen Persulfide (HSSH): Possible Molecular Sulfur Sinks in the Dense ISM. *Molecules* **2022**, *27*, 3200.
- (103) Wang, J.; Marks, J. H.; Tuli, L. B.; Mebel, A. M.; Azyazov, V. N.; Kaiser, R. I. Formation of Thioformic Acid (HCOSH)-The Simplest Thioacid-in Interstellar Ice Analogues. *J. Phys. Chem. A* **2022**, *126*, 9699–9708. PMID: 36534075
- (104) Marcelino, N.; Puzzarini, C.; Agúndez, M.; Fuentetaja, R.; Tercero, B.; de Vicente, P.; Cernicharo, J. First detection of the HSO radical in space. *A&A* **2023**, *674*, L13.
- (105) Bieler, A.; Altwegg, K.; Balsiger, H.; Bar-Nun, A.; Berthelier, J.-J.; Bochsler, P.; Briois, C.; Calmonte, U.; Combi, M.; De Keyser, J.; et al. Abundant molecular oxygen in the coma of comet 67P/Churyumov–Gerasimenko. *Nature* **2015**, *526*, 678–681.



CAS BIOFINDER DISCOVERY PLATFORM™

**PRECISION DATA
FOR FASTER
DRUG
DISCOVERY**

CAS BioFinder helps you identify
targets, biomarkers, and pathways

Unlock insights

CAS
A division of the
American Chemical Society