



OPEN Assessing geometric microliths as cultural markers through an analysis of shape variation and projectile performance

Alfredo Cortell-Nicolau^{1,2✉}, Alastair Key² & Antoni Palomo³

European geometric microlith shape variation is often used as a marker of cultural differences between groups of Mesolithic hunter gatherers and/or Neolithic farmers. Indeed, the 2D plan-view shape of these lithics is known to vary in spatially and temporally systematic ways between archaeological sites. Such differences are well evidenced in the Iberian Peninsula between the 9th and 8th millennia BP. Here we test an alternative hypothesis for the structured variation observed in geometric microliths: whether their plan-view shape significantly impacts the force, energy and displacement experienced when they are used as projectile tips. If functional differences between groups help to explain the shape variation observed in the archaeological record, then any role for cultural (non-functional social) explanations is potentially reduced. We undertook controlled static penetration tests using an Instron materials tester and an assemblage of replicated Iberian geometric microliths hafted to standardised wooden shafts. Results indicate that the maximum force required, energy used, and displacement at maximum force experienced by these hafted geometric microliths when used as projectile armatures is not significantly influenced by their 2D plan-view shape. Rather, gross form attributes such as maximum thickness, distance from the tip of the microlith to the start of the shaft, and the maximum width of the hafting substrate/adhesive are the greatest determinants of penetration ease, along with the positioning of the microlith when hafted. Our data therefore supports past research that proposes a cultural role for geometric microlithic shape variation in the European Mesolithic and Early Neolithic. Moreover, it highlights the functional importance of maintaining relatively thin microblades during microlith production, along with taking care to minimise the size of hafting components and the necessity to haft them in the most efficient way.

Keywords Projectile penetration potential, Force, Energy, Geometric microliths, Experimental archaeology, Mesolithic, Neolithic

Geometric microliths are a specific type of lithic artefact most often inferred to have been used as part of composite arrowheads. They are usually trapezoid, triangular or lunate in shape. Their spatial and chronological distribution arguably dates from as early as 70,000 years before present (BP) in Africa^{1,2}, but some of the clearest early examples are found in Asia and date to older than 30,000 BP^{3–5}. In Europe, although some have been reported from the Solutrean period around 20,000 BP⁶, geometric microliths did not become predominant until the beginning of the Holocene. Indeed, from around 10,000 BP they slowly acquire omnipresence in the European archaeological record and are a dominant lithic artefact until the Early Neolithic, with reappearances throughout the Middle and Late Neolithic⁷. During this time in Europe, they show great size and shape variability, from (broadly speaking) Sauveterrian micro-triangles and micro-segments through to Castelnovian trapezes or larger Early Neolithic tranchettes. Often, an assemblage shows a wide array of intermixed forms which are commonly collated as part of named typologies. Due to this variation, they have often been considered one of the main cultural markers for the societies using them, and have been used to not only build periodisations for the groups involved—alone or in combination with other lithic proxies e.g.^{8–12}—but also to help understand other

¹Max Planck Institute for Evolutionary Anthropology, Deutscher Pl. 6, 04103 Leipzig, Germany.

²Department of Archaeology, University of Cambridge, Cambridge CB2 3DZ, UK. ³Department of Prehistory, Universitat Autònoma de Barcelona, Edifici B, C/Fortuna, 08193 Bellaterra, Barcelona, Spain. ✉email: alfredo_cortell_nicolau@eva.mpg.de

relevant categorizations and patterns such as, among other things, social networks¹³, stratigraphic sequences¹⁴ or aggregation sites¹⁵.

Hindering their use as a cultural proxy is the fact that morphological differences observed between and within assemblages may indeed represent variable past cultural identities, or alternatively, they may represent differences in the use of these composite tools. Although several studies have assessed the potential functional applications of geometric microliths, principally through the use of experimental reproductions coupled with studies of breakage patterns^{16–21} and microwear analysis^{22–27}, there remains an absence of work investigating how variation in their form influences their performance *during* a given task. That is, the potential utility of these lithic objects has been investigated and arguably demonstrated, but how utilitarian performance and any associated performance attributes relate to variation in their form remains unknown. This has major implications for studies that treat form variation in geometric microliths (on a stand-alone basis, or integrated within broader lithic assemblages) as a cultural proxy e.g.^{11,21,28–31}. That is, despite their use as arrowheads seemingly being consistently supported see references above, but also^{22,32}, their morphology—and any morphological differences observed within and between microlith assemblages—could be derived from variation in, or changes to, the use of these arrowheads within tasks or when different task-outcomes were intended; for example, hunting ground nesting birds relative to boars, or killing a rabbit relative to stunning it. If microlith morphometric variation was tightly linked to measures of utilitarian performance, and potentially therefore different functional tasks³⁴, then the shape of these arrowheads could not reliably be used as a proxy for cultural diversity. Instead, they may better represent the ecology, and in turn faunal elements, present at the time and place of the artefacts deposition into the archaeological record. Such a scenario would reduce the social (cultural) explanatory power that has traditionally been attributed to the morphological variation observed in geometric microliths. Artefacts could have held a double value, even combining optimal functionality *and* optimal cultural value, but we adopt here a dichotomic framework to aid discussions on, and interpretation of, the relative role of each.

The style versus function debate of the 1980s and 1990s is highly pertinent e.g.^{35–44}, as is more recent work built on these earlier ideas e.g.^{45–49}. We do not intend to reproduce or exhaustively cover the full debate, however, some comments on this topic are useful. First, regardless of how one defines stylistic or functional traits (for debates on stochastic stylistic variation see Hurt & Ratika⁴⁴), the cultural evolutionary mechanisms impacting each are governed by diverse selective pressures and, therefore, analytical approaches investigating each need to be tailored to one or the other. If the shape of any microlith artefact was conditioned by functional pressures, for example, then understanding the relevant selective environment (e.g. a specific type of prey or broader ecological niche, etc.) is essential to understanding why any variation (or lack thereof) exists. If traits are stylistic (even considering potential social functions *sensu* Sackett³⁷), we can feasibly use geometric microliths as markers of past cultural groupings, and therefore gain insight into the social structure of these prehistoric populations, but given we cannot directly observe these past groups, the security of this inference is dependent on other explanatory factors (e.g., functional pressures) not playing a significant role.

To investigate the relative contribution of cultural drift, the founder effect and other cultural evolutionary mechanisms, including functional selective pressures, on past artefact forms, researchers often rely on experiments. For example, it is possible to get a handle on the role of drift mechanisms through experiments that replicate the production and social transfer of past artefact forms e.g.^{50,51}. In this scenario, if changes to the morphometry of geometric microliths were identified as being governed by social dynamics internal and specific to the group (of modern participants) who produced the artefacts, these objects could feasibly continue to aid our understanding of social chrono-spatial patterns in the European Mesolithic-to-Neolithic period. Similarly, experiments that investigate the potential role of functional selective pressures on artefact forms are also common e.g.^{52–55}. In these investigations, experimentation allows for specific utilitarian hypotheses to be tested—including form-function relationships—within diverse conditions, with the internal and external validity of any results varying dependent on the specific experimental set-up⁵⁶. Of most relevance here are studies that investigate potential performance characteristic differences between arrowheads or atlatl dart armature forms, with examples ranging from mathematical modelling⁵⁷ and highly controlled machine-performed penetration tests^{58–62} through to those conducted by human participants using replica whole artefact-systems cf.⁶³ to propel projectiles into simulated or real animal carcasses^{17,21,64}.

Here, we investigate whether the form of geometric microliths can be confidently used as a proxy for cultural identity by experimentally investigating an alternative utilitarian hypothesis. Our approach relies on understanding whether microlithic morphometric variation has an effect on their performance (and in turn, inferred utilitarian efficiency) when hafted as arrowheads. If a form variable is demonstrated to have an effect on the performance, then it feasibly could have been a relevant functional factor for Mesolithic and Neolithic individuals. The rationale is that if shape does not alter microlith performance, then efficiency differences within 'real world' (i.e., prehistoric hunting) conditions would have been minimal and tool design choices would not have been linked to functional optimisation. We would like to clarify that our intention is not to understand which shapes may perform better than others but, rather, to assess whether shape affects performance at all. We do this through a combination of experimental replication, mechanical testing and statistical modelling. First, a replica assemblage of geometric microliths based on the shapes found in Eastern Iberia were produced with intentionally wide ranges of morphometric variability. Second, we undertake controlled static penetration tests following Sittin et al.⁶¹ and Mullen et al.⁶⁰ to investigate form-performance relationships. Finally, we statistically assess which factors lead to optimal levels of performance when geometric microliths are used as projectiles.

Methods and data

Ballistic analyses are fairly common in prehistoric archaeological research e.g.^{20,62,64–70}. To the best of our knowledge, while limited-in-scale actualistic experiments investigating potential types of utility for geometric microliths have been published e.g.^{17,18}, large-scale structured analyses exploring form optimisation and

performance variation are lacking. For the present study, we focus on the performance of geometric microliths and hence it is useful to define the term ‘performance’ in advance. One way to think about performance is via the optimisation of an artefact’s production costs (e.g., time, energy, risk cf.⁷¹). Neeley and Barton⁷² provide clear time-costs during the production of geometric microliths, with each taking an average of 2–3 min (also see below), and this being, along with their modularity and ease of substitution, key for their success see⁷³. Another relevant performance measure for microliths concerns the ability to track fauna see⁷⁴. In theory, after an arrow strikes an animal microliths can detach from the haft and contribute to internal bleeding, more easily facilitating the tracking of the wounded prey. In this context, performance could be further enhanced by the potential use of poison e.g.^{75–77}. These performance characteristics cf.⁴³ were potentially important to the design of geometric microliths, but of principal concern here, and with most prehistoric projectile research see⁷⁸, is the relative ability of the lithic armature to penetrate intended targets under low force and energy (incl. velocity) requirements. Indeed, if an arrowhead propelled at a given velocity cannot perforate an animal’s skin then its performance may be considered ineffective, if it perforates but only minimally penetrates performance would be considered inefficient, and if it both perforates and penetrates to a depth suitable enough to cause substantial injury performance can be considered efficient. Importantly, inefficient-to-efficient can be viewed as a scaled measure and recorded as such. Thus, in this work we specifically test the capacity of the geometric microliths to penetrate a substrate and how this relates to force and energy inputs, and we refer to performance as such.

According to these lines of thought, in order to assess whether geometric microlith morphometry/shape impacts efficiency we undertook three broad work packages, as follows.

Replication and geometric morphometric process

First, an experimental sample of 29 geometric microliths were created, based on 14 defined shapes found during the Iberian Late Mesolithic and Early Neolithic^{9,11,15,79}. It was our intention to replicate most of the typologies found in the area during this time, but we acknowledge outliers may exist in the archaeological record.

Each microlith was produced from small blades knapped through abdominal pressure by AP, following the experimental process shown in Fig. 1. Fine grained chert gathered from Normandy (France) and Aragon (Spain) was used, and our choice for blade selection was guided by the intended final geometric shape. Based on prior work the two highly-siliceous and homogenous raw materials likely display similar sharpness measures⁸⁰. In general, the replica blades presented transversal trapezoidal sections and longitudinal straight sections with slightly convex distal sides. All geometrics were segmented through flexion of the blade. Slim blades aid the segmentation process by allowing greater control over the fracture, with ‘clean’ fracture perpendicular to the long axis being more common. Once the blade was segmented we used a cervid antler tine with a sharp edge, along with a leather-protected table, to configure the geometric shape. Retouching (final shaping) typically did not last more than two or three minutes per geometric. Each geometric was hafted to the end of a 150 mm long wooden dowel, with the dowel used in place of an arrow shaft. We used a commercially available thermoplastic



Fig. 1. Experimental production process of the geometric microliths. 1—Production of blades through abdominal pressure; 2—Blade fracture through bending; 3—Blade fragments obtained through bending; 4—Blade fragments and deer antler tool used for shaping the geometric; 5, 6—Pressure retouch; 7—Comparison of the experimental versus the archaeological geometric. The individual in the image has given consent for publication in an online open-access journal.

adhesive (hot-melt adhesive) to attach the lithics as this allowed the glue to be remelted, and in turn, each geometric could be used multiple times; in this case, in three distinct positions (see below). For the experiment, every point was positioned at the tip of the shaft. This was not their only potential prehistoric positioning and, in some specific cases, not even the most likely one^{18,26,27}. We have proceeded this way because our intention was to assess the optimality of their penetrating capabilities, disregarding their lacerating properties. Each geometric was tested three times with three different positions on the tip of the shaft: obliquely pointing with the distal side, obliquely pointing with the proximal side and transversally/horizontally with the long base as potential cutting edge (see Fig. 4 below). For symmetric arrowheads we did the same for consistency, although we are aware that, in this case, there should be no significant differences between distal and proximal sides. This allowed us to investigate the impact of hafting position on penetration potential alongside the role of geometric shape/size variation. Regardless of their typology and morphometry, each geometric was hafted in the above mentioned three positions and using the same hafting method.

The shape of the geometric microliths was recorded using outline-based Geometric Morphometric Analysis (GMM), a technique widely utilised in modern archaeological practice. Details on the application of the method (outline or landmark-based) on lithic assemblages can be found elsewhere e.g.^{15,30,81–90}. We first performed a General Procrustes Analysis (GPA) followed by a Fourier analysis with seven harmonics and principal component analysis (PCA). The intention here was not to use GMM as an end goal, as is typically the case but, rather, to use it as a proxy for recording each geometric's morphology. With a cumulative variance of 88.2% for the first two components in our PCA (Fig. 2), we can interpret these components as broadly capturing the shape of each geometric. Therefore, instead of using linear measures to account for aspects of form variation, we use PC1 and PC2 (*M1* and *M2* respectively, see below) to describe the general outline shape of each microlith.

Mechanical tests

For the second part of our analysis we investigated the penetration potential of each geometric microlith in each of its three hafted positions. We follow Sitton et al.⁶¹ and Mullen et al.⁶⁰ and used an Instron materials tester (34SC-5 model) to perform machine-controlled static penetration tests. We secured the shaft of each hafted geometric into an upper grip on the Instron before lowering it at a controlled rate (50 mm/min) into a substrate. Throughout the force (newtons [N]), energy (joules [J]) and displacement (mm) experienced by the upper grip could be recorded, allowing us to observe how each metric varied depending on the replica lithic and hafting position used. Our specific interest was in recording peak force experienced (N), total energy (J) used, and displacement at the point of maximum force (mm), when pushing the microlith into the substrate. Each unique lithic and position combination was 'thrown' (i.e., pushed by the Instron into the substrate) three times, resulting in a total of 261 individual penetration tests (three times per geometric and hafting position) being conducted.




Fig. 2. PCA on the geometric microliths studied. PC1 and PC2 have been included as co-variables to define the shape of the geometric microliths in the final model.

It was essential for us to use an industrially produced material as the ‘target substrate’ to ensure testing conditions were identical for all samples. Following a large number of prior projectile experiments^{59,61,91,92} we used potters clay (terracotta [purchased from HobbyCraft in Cambridge, UK]) in place of meat, as the former has previously been demonstrated to display a reasonable proxy for the latter when investigating projectile penetration energy and forces in small armatures, although Key et al.⁹³ stress the two materials are not directly interchangeable and differences increase when assessing larger armatures (which is not the case here). The clay was secured within a 10 × 15 cm plastic box, with the central portion of the clay being aligned underneath the microlith. On the superior surface of the clay we positioned a taut sheet of thin polythene plastic to act as a ‘skin’ that may at first deform and be pushed into the clay under pressure from the microlith, before being perforated after five mm of deformation and the lithic continuing into clay. Prior to the test beginning all lithics were aligned with the surface of the polythene such that they were touching but no force was recorded by the Instron. The lithics were pushed into the clay to a depth of 30 mm before the test ceased. Data were collected at a rate of 20 points/second with the maximum force (N) experienced, the displacement (distance moved) at the point of maximum force (mm), and the area (energy [J]) beneath the resulting force–displacement curve being collected for analysis (Fig. 3).

Analytical framework

Variables and observations

We recorded the following three response variables:

- *E* (Energy): Total energy used by the geometric during the process of perforating the target and penetrating to a depth of 30 mm (measured in Joules).
- *F* (Maximum Force): The peak force experienced by the geometric when perforating the target (measured in Newtons).
- *D* (Displacement at Maximum Force): Displacement produced on the target by the penetrating arrowhead (measured in millimetres).

We considered a total of 10 potential predictors, as follows:

- *Sym* (Symmetry): Whether the gross shape of the geometric can be considered symmetric or not. Binomial.
- *I* (Inclination): Position of the geometric on the haft. Categorical with four categories: oblique distal, oblique proximal, horizontal, symmetric.
- *B* (Breakage): Whether the geometric breaks during the penetration testing. Binomial.
- *S.par* (Parallel width shaft): Width of the shaft measured parallel to the geometric’s positioning. Continuous in mm.
- *S.per* (Perpendicular width shaft): Width of the shaft measured perpendicular to the geometric’s positioning. Continuous in mm.
- *W* (Width). The maximum width of the geometric after hafting. Continuous in mm.
- *W.S.* (Width before shaft). Width of the geometric prior to the start of the shaft. Continuous in mm.
- *T* (Distance from tip to shaft). Distance from the tip of the geometric to the start of the shaft. Continuous in mm.
- *M1* (Morphometry 1). Plan-view shape of the geometric. PC1 result after the GMM analysis explained above.
- *M2* (Morphometry 2). Plan-view shape of the geometric. PC2 result after the GMM analysis explained above.

From these, *Sym* and *B* were discarded prior to the start of the analyses. For *Sym* this was because the symmetry of the geometric should be accounted for by its plan-view shape. As for *B*, no geometrics broke during the testing, so this co-variate became uninformative. The rest of the predictors were considered within the model. The low number of observations available from our analysis limited the complexity of our systematic component. Thus, to improve the model’s efficiency and reduce its complexity, two variables were modified: the variable *I* was converted to binomial with only two categories (oblique/horizontal), while the variables *S.par* and *S.per* were merged into one variable, *S*, the mean of the two (as in $S_i = \frac{S.par_i + S.per_i}{2}$ for $i = 1, 2, \dots, n$ with n being the number of observations) and accounting for the overall thickness of the shaft.

Our main criteria for co-variate inclusion in the final model was based on expert knowledge, usually advised over automatic and purely diagnostic procedures^{94–97}. We complemented this with a collinearity analysis and a PCA on the rest of the co-variables to see whether they could provide insight for further model simplification, with no success. Thus, for the final systematic component of the model we included the co-variables *W*, *I*, *S*, *M1*, *M2* and *T*, as shown in Fig. 4. As it can be seen, each geometric was positioned in three different ways, although only two (oblique and transversal) were considered for the analysis. The decision to leave *W.S* out was also reinforced by the model’s performance worsening (higher WAICs) when it was included. To check for potential outliers, we computed the group Mahalanobis distance including the co-variables considered for the final model. Our results identified two potential group outliers for observations nine and 29. Observation nine had a negative impact on the model’s overall fitting, but observation 29 did not. Thus, we removed observation nine from the analysis, resulting in 86 observations for each response.

Models

We developed a hierarchical Bayesian model for each response (*E*, *D* and *F*) using R and nimble. The systematic component remained the same for each model and consisted of the variables mentioned above, plus one random effect group to account for the fact that each geometric was tested three times. All of our responses were strictly positive but they did not follow the same distribution, with some of them displaying skewness. To assess which



Fig. 3. The sample of replica geometric microliths hafted to the wooden dowels (a), three pairs of geometrics prior to testing (b–d), the Instron device and wider experimental set-up, including an exemplar force–deformation curve (e), and a close-up of the experiment post-test (f).

specific type of model to use for the analysis of each response we tested a linear regression, linear regression with log-transformed data, gamma regression and lognormal regression. Our preferred model was chosen based on: (1) their standardised residuals and (2) their posterior predictive checks on mean and median, as adapted from Gelman^{98,99}. According to this, we decided to use a standard linear regression for *E*, a lognormal regression for *F* and a gamma regression for *D*. We selected weakly informative priors consisting of (1) normal distributions with




Fig. 4. Predictors used as they were gathered for each geometric. The subimage in the top right hand corner shows the variable *I* and how each geometric was positioned along the shaft. For simplicity, the variable *I* was reduced to only two categories: transversal and oblique. M1 and M2 (not in this image) are explained in the previous section.

large standard deviation (100) for the parameters of the systematic component ($\beta_1, \dots, \beta_n$) and (2) half-cauchy distributions (coded as Student-t distributions with 1 degree of freedom) for the distribution's parameters, such as the variances of the response and the random effects [see^{98,100}]. Furthermore, we censored the response *D*. This is because the machine was set to a maximum level of 30 mm, and thus a concentration of values were observed on the upper limit of this range [29, 30], indicating that displacement at maximum force values for these samples could theoretically have exceeded 30 mm. Hence, we censored this response at 29.5, which returned a much better performance of this model.

We estimated the parameters using an MCMC with 300,000 iterations, burn-in period of 3000 and sampling every 300 iterations to obtain 1,000 non-autocorrelated values for each parameter's posterior distribution. Chains converged properly (Fig. 5), with Gelman-Rubin equaling 1 for the vast majority of parameters (occasionally 1.01) and with number of effective sizes > 2000 in any case.

All data and code for the analyses mentioned above, along with exploratory analysis, plots and further model diagnostics are available in the supplementary material. Supplementary material has been prepared as a git repository, with its dedicated readme, and can be found at https://github.com/acortell3/Geo_performance. We have used the language R¹⁰¹ and the R packages nimble^{102,103}, MCMCvis¹⁰⁴, coda¹⁰⁵, psych¹⁰⁶ and Momocs¹⁰⁷.

Results

Our results focus on the statistical models developed to assess the effect of the different co-variables on the response variables. One of our main criteria when selecting the best model to use was whether it was capable of fully recapturing the mean and median through a posterior predictive check (Fig. 6) (see Gelman⁹⁸, and see Crema & Shoda¹⁰⁸ and Timpson et al.¹⁰⁹ for archaeological applications). In this regard, the models selected obtained practically identical means (Table 1) with very similar medians (Table 2); these data act to confirm the models worked as expected.

After checking the adequacy of the model our main interest was to understand how the parameters considered affected the responses for each model (*E*, *D* and *F*). Figure 7 demonstrates some consistent trends. When analysed individually, *T* (the distance from the tip of the geometric microlith to the beginning of the shaft) has a small effect on the penetration of the arrowhead, considering the responses *E* and *D*. In the first case, as *T* increases, less force is required and, in the second, for higher values of *T*, more displacement would be recorded. The small values of these effects (see Table 3) make them almost neglectable. Taking into account the response *F*, *T* does not differentiate from zero. On the other hand, the inclination of the geometric microlith, *I*, plays a key role in penetration potential, with arrowheads obliquely positioned resulting in lower energy being required to penetrate the target and displacement at maximum force being greater. For these two responses, and as shown in Table 3, the posterior distributions for this parameter are significantly different from zero, showing its potential effect. Although the upper credible interval on this co-variate on the response *F* is slightly above zero, its lower credible interval and mean suggest that an arrowhead obliquely positioned would require reduced force to penetrate the target, but this is not strongly supported by our results.

The maximum width of the arrowhead (*W*) and the shaft/hafting width (*S*) also play a key role in the penetrating capabilities of geometric microlith arrowheads, in all cases clearly differentiating from zero. For the former, higher values of *W* indicated that the arrowhead requires greater force to perforate the target, greater energy during penetration and deeper penetration at the point of maximum force. For *S*, we observed similar relationships, albeit with larger credible intervals. These, following Table 3 and Fig. 7, could be interpreted as thicker shafts requiring more energy and force to penetrate the target, along with greater forces being required at greater penetration depths. The effect of displacement at maximum force (0.03–0.29) is significantly larger than energy (0.03–0.07) and maximum force (0.03–0.1). Interestingly, for the variables representing the shape of the




Fig. 5. Convergence of the MCMCs. The pattern in the left hand side images, essentially random and without peaks, indicates good convergence. The same can be said for the right hand side images, which are approaching normal distributions, with Gelman-Rubin indices (rhat) = 1 and with number of effective sizes around 3000 (usually anything higher than 200 can be considered enough). Due to space constraints, only three parameters are shown. The rest can be observed in the code and plots provided as supplementary material.




Fig. 6. Posterior predictive checks on mean and medians of the models selected. Red is the observed mean/median and blue is the prediction from our parameters’ posterior distributions.

	Area	Maximum force	Displacement
Observed	0.5	2.78	22.34
Predicted	0.49	2.77	22.32

Table 1. Posterior predictive checks on means.

	Area	Maximum force	Displacement
Observed	0.44	2.55	21.82
Predicted	0.43	2.53	21.5

Table 2. Posterior predictive checks on medians.

arrowhead (*M1* and *M2*) neither diverged from zero (indicating that the shape or morphometry of the geometric does not influence the penetration of the throw), with in some cases (e.g. for responses *F* and *D*) zero being almost within one standard distribution from the posterior’s mean. For the case of *E* as a response, the credible interval of *M2* (which could be broadly representing slimmer arrowheads) did differ from zero, although some of its values were still negative. The exact impact of the different co-variates regarding each response has been summarised in Table 3.

In sum, these results demonstrate the position of the geometric (oblique vs. transversal; predictor *I*) to have the strongest effect on performance during each test, followed by the shaft’s width (*S*) and the maximum width of the geometric (*W*). The distance from the tip to the shaft (*T*) does also have an effect, but it is small and not for all responses. Finally, 2D plan-view shape (*M1* and *M2*) does not appear to improve or worsen the penetrating capabilities of geometric microliths.




Fig. 7. Posterior distributions of the parameters considered for each model (energy, maximum force and displacement). Effects can be considered relevant to the interpretation of geometric performance if their credible intervals deviate from 0.

	E (mean)	E (CI)	F (mean)	F (CI)	D (mean)	D (CI)
T (Distance)	−0.01	−0.02, −0.01	0.01	0, 0.02	0.05	0.04, 0.07
I (Inclination)	−0.11	−0.16, −0.07	−0.06	−0.13, 0.01	0.23	0.13, 0.33
W (Maximum width)	0.03	0.02, 0.03	0.04	0.04, 0.05	0.04	0.02, 0.05
M1 (PC1)	−0.05	−0.13, 0.01	−0.06	−0.17, 0.06	−0.05	−0.2, 0.09
M2 (PC2)	0.11	0.04, 0.19	0.05	−0.08, 0.18	0.06	−0.09, 0.24
S (Shaft)	0.05	0.03, 0.07	0.06	0.03, 0.1	0.24	0.19, 0.29

Table 3. Means and credible intervals of the posterior distributions of the parameters studied for each response.

Discussion and conclusion

Our main objective was to understand whether morphometric variation in geometric microliths impacts their performance when used as projectile tips (armatures). This allows us to assess whether their prehistoric design (form) was likely conditioned by utilitarian optimisation, or alternatively, if it was governed by other factors, including cultural constraints. As reported in the results, our analysis suggests the main factors affecting their performance as projectiles are their positioning on the shaft (oblique or transverse), the maximum thickness of the shaft (when hafted, including adhesive), the maximum width of the geometric and, to a lesser extent, the distance of the geometric's tip to the start of the shaft. Plan-view shape does not seem to play an important role. These results are consistent with the fundamental mechanics of projectiles as they penetrate substrate¹¹⁰. Oblique positioning on the haft and the lower shaft and geometric thickness measures, all contribute to reducing the surface area of the projectile perpendicular to the target as it penetrates, reducing the resistance experienced. Further, transversal arrowhead positioning, along with thicker shaft and arrowheads, result in increased displacement prior to maximum force being recorded meaning that once the arrow penetrates, the work experienced by the projectile appears to increase as it pushes deeper into the substrate, rather than being experienced closer to the point of initial penetration. Admittedly, this study has not considered other elements which might impact penetration potential, such as the varying use of bows, spears or atlatls. Given the poor preservation of wood from the European Mesolithic and Neolithic we do not know a great deal about the bows used at this time, but there are exceptions in water-logged environments (see Piqué et al.¹¹¹, Bailey et al.¹¹² and see Bertin et al.¹¹³ for non water-logged case), as well as experimental studies assessing the potential requirements of prehistoric bows^{114,115}. There appear to be differences in the wood used for bows during the Mesolithic and Neolithic, with the earlier hunter-

gatherers usually preferring elm (*Ulmus sp.*) and, with exceptions, farmers tending to use yew (*Taxus bacatta*). Archaeobotanical data in Northern Europe explains how yew expanded through Europe during the sixth millennium, which might explain how hunter-gatherers started to diversify their wood use around this time¹¹⁶. In any case, the technology for bow manufacture seems to be different between the two periods, although at present this difference does not seem to allow easy discrimination between potential different uses³³.

Some other technological elements such as the bow's size or the fletching of the arrow might also impact their performance^{115,117}, but we understand even less about the selection of these factors in prehistoric Europe. Nevertheless, in our case it is important to consider that we are not analysing the overall hunting performance, which would refer to the full hunting artefact-system⁶³, including the bow, but how the shape of the arrowhead improves this performance. Arrowhead shape has been reported to change depending on the method of hafting used in some contexts (see Bettinger & Eerkens¹¹⁷ for the case of Palaeoindian points). This is particularly relevant because it is known that microliths were not always located on the tip of the shaft, but could also be used modularly along the side of shafts to form barbs in a harpoon-like system^{18,27,118,119}. We have not considered this hafting disposition as we focus on the penetration capabilities of the arrowheads alone. Alternative methodological approaches that display the 'wound' track of the projectile e.g.^{60,119} would be required to investigate laceration capabilities. Further, it would require assumptions as to which microliths should be positioned on the tip and which ones should be positioned along the shaft see^{7,120}. While it is true that we might have used some microliths as *tip* arrowheads when their prehistoric position was more often as a *shaft* arrowhead, this does not affect our results. Indeed, we have theoretically assessed the tip-functionality of 'shaft' armatures, with our data demonstrating that if shape is a determinant of which microliths are hafted to the side of shafts and used for laceration, it is likely because of their performance during laceration and not penetration (although more detailed exploration of how side and tip armatures impact performance together is required¹²¹).

Our results are distinct to similar experimental data returned for Clovis points, with 2D plan-view shape (i.e., size having been controlled for) having been demonstrated to significantly impact their penetration depth (and in turn, likely force and energy required over the first 30 mm of penetration)⁵⁴. This difference likely stems from the overall size of the armatures, with microliths being substantially smaller and the relative mechanical impact of shape differences being subsequently reduced. We did not consider cross-sectional shape here, so it is difficult to assess how our data compare to studies that found a relationship between this parameter and penetration depth e.g.^{65,121,122}, but our measure of maximum thickness—along with its significant impact on performance—is a highly related metric with likely similar results.

Our main question here is, all else being equal, does the shape of geometric microliths affect their capacity to be used as a projectile? Our results point to form-attributes other than microlith 2D plan-view shape, including hafting sections and gross size attributes (e.g., thickness), being more relevant to determining their penetration performance. This supports past research proposing that geometric microlith shape can be used to help understand cultural differences between groups of Mesolithic hunter gatherers and Neolithic farmers^{11,15,28,29,124–126}. Indeed, our research suggests that their morphometry and shape could potentially be used to assess complex questions regarding group identity and cultural transmission patterns in Prehistory e.g.^{126–130}. Moreover, due to their role as a distinctive functional end product (as opposed to, for example, scrapers, blades or microburins), we believe that geometric microlith shape may be one of the most informative well-preserved cultural markers of this time e.g.^{9,11,79,131–135}. If and how geometric microlith shape was related to social functions, however, is another question altogether, and one beyond the scope of our experimental paper.

In sum, we have shown that the shape of these arrowheads does not appear to retain information relating to the penetration potential of projectiles, and instead they likely contain cultural information relevant to understanding the group dynamics and interactions of these past peoples. While additional studies are required to further support (or refute) our inferences, this work provides a novel utilitarian foundation from which to analyse geometric microlith morphometric variability across broad geographic and diachronic ranges.

Data availability

All the data and code generated for this article can be found in a reproducible format here https://github.com/acortell3/Geo_performance.

Received: 3 January 2025; Accepted: 18 March 2025

Published online: 02 April 2025

References

1. Wurz, S. & Lombard, M. 70,000-year-old geometric backed tools from the Howiesons Poort at Klasies River, South Africa: Were they used for hunting? *South. Afr. Humanit.* **19**, 1–16 (2007).
2. Brown, K. S. et al. An early and enduring advanced technology originating 71,000 years ago in South Africa. *Nature* **491**, 590–593 (2012).
3. Clarkson, C. et al. The oldest and longest enduring microlithic sequence in India: 35,000 years of modern human occupation and change at the Jwalapuram locality 9 rockshelter. *Antiquity* **83**, 326–348 (2009).
4. Roberts, P., Boivin, N. & Petraglia, M. The Sri Lankan 'Microlithic' Tradition c 38,000 to 3000 years ago: Tropical technologies and adaptations of homo sapiens at the southern edge of Asia. *J. World Prehistory* **28**, 69–112 (2015).
5. Basak, B. & Srivastava, P. Earliest dates of microlithic Industries (42–25 ka) from West Bengal, Eastern India: New light on modern human occupation in the Indian subcontinent. *Asian Perspect.* **56**, 237–259 (2017).
6. Elston, R. G. & Kuhn, S. L. *Thinking Small: Global Perspectives on Microlithization* (American Anthropological Association, 2002).
7. Fernández López de Pablo, J., Gibaja Bao, J. F. & Palomo, A. Geométricos y puntas usadas como proyectiles en contextos neolíticos de la fachada mediterránea. in *IV Congreso del Neolítico Peninsular: 27-30 de noviembre de 2006*, Vol. 2, 2008, ISBN 978-84-96979-14-7, 305–312 (Museo Arqueológico de Alicante - MARQ, 2008).
8. Rozoy, J.-G. Microburins et armatures microlithiques dans « le » Néolithique. *Bull. Société Préhistorique Fr.* **68**, 145–151 (1971).
9. Fortea, J. *Los Complejos Microlaminares y Geométricos Del Epipaleolítico Mediterráneo Español* (Universidad de Salamanca, 1973).

10. Binder, D. L. *Néolithique Ancien Provençal Typologie et Technologie Des Outillages Lithiques* (Éditions du CNRS, 1987).
11. García Puchol, O. *El Proceso de Neolitización En La Fachada Mediterránea de La Península Ibérica. Tecnología y Tipología de La Piedra Tallada* Vol. 1430 (BAR International Series, 2005).
12. Perrin, T. et al. The late Mesolithic of Western Europe: Origins and chronological stages. *Ann. Fond. Fyssen* **24**, 160–176 (2009).
13. Barrera-Cruz, M., García-Puchol, O., Jiménez-Puerto, J., Cortell-Nicolau, A. & Bernabeu-Aubán, J. Weaving social networks from cultural similarities on the neolithisation process in the Western Mediterranean: Evolutionary trajectories using projectile tools. *PLoS ONE* **19**, e0306027 (2024).
14. Perrin, T. & Defranould, E. The Montclus rock shelter (Gard) and the continuity hypothesis between 1st and 2nd Mesolithic in Southern France. *Quat. Int.* **423**, 230–241 (2016).
15. Cortell-Nicolau, A., García-Puchol, O. & Juan-Cabanilles, J. The geometric microliths of cueva de la cocina and their significance in the mesolithic of Eastern Iberia: A morphometric study. *Quat. Int.* <https://doi.org/10.1016/j.quaint.2023.05.014> (2023).
16. Fischer, A., Hansen, P. V. & Rasmussen, P. Macro and micro wear traces on lithic projectile points. Experimental results and prehistoric examples. *J. Dan. Archaeol.* **3**, 19–46 (1984).
17. Gibaja, J. F. & Palomo, A. Geométricos usados como proyectiles. Implicaciones económicas, sociales e ideológicas en sociedades neolíticas del VI-IV milenio CAL BC en el Noroeste de la Península Ibérica. *Trab. Prehist.* **61**, 81–97 (2004).
18. Yaroshevich, A., Kaufman, D., Nuzhnyj, D., Bar-Yosef, O. & Weinstein-Evron, M. Design and performance of microlith implemented projectiles during the Middle and the late Epipaleolithic of the Levant: Experimental and archaeological evidence. *J. Archaeol. Sci.* **37**, 368–388 (2010).
19. Chesnaux, L. Des microlithes sauveterriens, témoins de l'armement des derniers chasseurs-cueilleurs dans les Alpes du Nord. *Palethnol. Archéol. Sci. Hum.* <https://doi.org/10.4000/palethnologie.9400> (2009).
20. Hutchings, W. K. Measuring use-related fracture velocity in lithic armatures to identify spears, javelins, darts, and arrows. *J. Archaeol. Sci.* **38**, 1737–1746 (2011).
21. Philibert, S., Chesnaux, L., Costamagno, S. & Lacarriere, J. Something new in the quivers: experimental approach to functioning of the late Mesolithic and Early Neolithic geometric bitruncations. in (Toulouse, 2020).
22. Domingo Martínez, R. *La Funcionalidad de Los Microlitos Geométricos: Bases Experimentales Para Su Estudio* (Universidad de Zaragoza. Departamento de Ciencias de la Antigüedad, 2005).
23. Domingo Martínez, R. Caracterización funcional de los microlitos geométricos. El caso del valle del Ebro. In *El Mesolítico Geométrico en la Península Ibérica* (eds Utrilla Miranda, P. & Montes Ramírez, L.) 357–389 (Universidad de Zaragoza, Zaragoza, 2009).
24. Plisson, H., Dubreuil, L. & Guilbert, R. The functional significance of Sauveterrian microlithic assemblages: Broadening the focus of investigation. In *Prehistoric Technology' 40 Years Later: Functional Studies and the Russian Legacy: Proceedings of the International Congress Verona (Italy), 20-23 April 2005* (eds Longo, L. & Skakun, N.) 147–156 (BAR International Series, 2008).
25. Cristiani, E., Pedrotti, A. & Gialanella, S. Tradition and innovation between the Mesolithic and early Neolithic in the Adige Valley (northeast Italy). New data from a functional analysis of trapezes from the Gaban rock-shelter. *Doc. Praehist.* **36**, 191–205 (2009).
26. García Puchol, O., Mazzucco, N., Gibaja Bao, J. F. & Juan Cabanilles, J. Use-wear analysis of a set of geometric projectile points from the mesolithic context of Cocina cave (Eastern Spain). In *International Conference of Use-Wear* (eds Marreiros, J. et al.) 213–224 (Cambridge Scholar Publishing, 2014).
27. Philibert, S. Ist versus IInd Mesolithic in southern. France Functional approach of techno-economic behavior through the Castelnovian of Montclus rock shelter (Gard): First results. *Quat. Int.* **423**, 242–251 (2016).
28. Lo Vetro, D. & Martini, F. Mesolithic in Central-Southern Italy: Overview of lithic productions. *Quat. Int.* **423**, 279–302 (2016).
29. Marchand, G. & Perrin, T. Why this revolution? Explaining the major technical shift in Southwestern Europe during the 7th millennium cal. BC. *Quat. Int.* **428**, 73–85 (2017).
30. Cortell-Nicolau, A., García-Puchol, O., Barrera-Cruz, M. & García-Rivero, D. The spread of agriculture in Iberia through approximate Bayesian computation and neolithic projectile tools. *PLoS ONE* **16**, e0261813 (2021).
31. Defranould, E., Philibert, S. & Perrin, T. Evolutionary dynamics of armatures in southern France in the late Mesolithic and early Neolithic. *Open Archaeol.* **8**, 905–924 (2022).
32. Finlayson, B. & Mithen, S. The microwear and morphology of microliths from Gleann Mor. In *Projectile technology* (ed. Knecht, H.) 107–131 (Springer Science-Business Media, LLC, 1997).
33. Jukmanns, J. *Pfeil Und Bogen. Vonder Altsteinzeit Bis Zum Mittelatlter* (Verlag Angelika Hörnig, 2013).
34. Palomo Pérez, A. et al. La caza en el yacimiento lacustre de La Draga (Banyoles, Girona). In *Actas del III Congreso del Neolítico en la Península Ibérica: Santander, 5 a 8 de octubre de 2003, 2005* (eds Arias Cabal, P. et al.) 135–144 (Instituto Internacional de Investigaciones Prehistóricas de Cantabria, 2005).
35. Close, A. E. The identification of style in lithic artefacts. *World Archaeol.* **10**, 223–237 (1978).
36. Dunnell, R. C. Style and function: A fundamental dichotomy. *Am. Antiq.* **43**, 192–202 (1978).
37. Sackett, J. R. Approaches to style in lithic archaeology. *J. Anthropol. Archaeol.* **1**, 59–112 (1982).
38. Sackett, J. R. Style and ethnicity in the Kalahari: A reply to Wiessner. *Am. Antiq.* **50**, 154–159 (1985).
39. Wiessner, P. Style or Isochrestic Variation? A Reply to Sackett. *Am. Antiq.* **50**, 160–166 (1985).
40. Dibble, H. L. Interpreting typological variation of middle paleolithic scrapers: Function, style of sequence of reduction. *J. Field Archaeol.* **11**, 431–436 (1984).
41. Barton, C. M. Beyond style and function: A View from the middle paleolithic. *Am. Anthropol.* **92**, 57–72 (1990).
42. Bettinger, R. L., Boyd, R. & Richerson, P. J. Style, function, and cultural evolutionary processes. In *Darwinian Archaeologies* (ed. Maschner, H. D. G.) 133–164 (Springer, 1996). https://doi.org/10.1007/978-1-4757-9945-3_8.
43. Schiffer, M. B. & Skibo, J. M. The explanation of artifact variability. *Am. Antiq.* **62**, 27–50 (1997).
44. Hurt, T. D. & Rakita, G. F. M. *Style and Function: Conceptual Issues in Evolutionary Archaeology* (Bergin & Garvey, 2001).
45. Skibo, J. M. *Understanding Pottery Function* (Springer, 2013). <https://doi.org/10.1007/978-1-4614-4199-1>.
46. Lycett, S. J., Schillinger, K., Eren, M. I., von Cramon-Taubadel, N. & Mesoudi, A. Factors affecting Acheulean handaxe variation: Experimental insights, microevolutionary processes, and macroevolutionary outcomes. *Quat. Int.* **411**, 386–401 (2016).
47. Shennan, S. Style, function and cultural transmission. In *Culture History and Convergent Evolution: Can We Detect Populations in Prehistory?* (ed. Groucutt, H. S.) 291–298 (Springer, 2020).
48. Gowlett, J. A. J. Deep structure in the Acheulean adaptation: Technology, sociality and aesthetic emergence. *Adapt. Behav.* **29**, 197–216 (2021).
49. Baldino, J. et al. North American Clovis point form and performance V: An experimental assessment of spear thrusting penetration depth and entry wound size. *Lithic Technol.* **49**, 295–310 (2024).
50. Schillinger, K., Mesoudi, A. & Lycett, S. J. Copying error and the cultural evolution of 'additive' versus 'reductive' material traditions: An experimental assessment. *Am. Antiq.* **79**, 128–143 (2014).
51. Page, S. N. Cultural transmission of lithic artefact traditions: an experimental approach. *Doctoral thesis, UCL (University College London)*. 1–375 (UCL University College London, 2014).
52. Mesoudi, A. & O'Brien, M. J. The cultural transmission of great basin projectile-point technology I: An experimental simulation. *Am. Antiq.* **73**, 3–28 (2008).
53. Key, A. & Lycett, S. J. Form and function in the Lower Palaeolithic: History, progress, and continued relevance. *J. Anthropol. Sci. Riv. Antropol. JASS* **95**, 67–108 (2017).

54. Eren, M. I. et al. North American Clovis Point form and performance: An experimental assessment of penetration depth. *Lithic Technol.* **45**, 263–282 (2020).
55. Stileman, F., Shipton, C. & Ashton, N. Not just scraping by: Experimental evidence for large cutting tools in the high lodge non-handaxe industry. *J. Paleolit. Archaeol.* **7**, 8 (2024).
56. Eren, M. I. et al. Test, model, and method validation: The role of experimental stone Artifact replication in hypothesis-driven archaeology. *Ethnoarchaeology* **8**, 103–136 (2016).
57. Thomas, K. A. et al. Explaining the origin of fluting in North American Pleistocene weaponry. *J. Archaeol. Sci.* **81**, 23–30 (2017).
58. Bebbler, M. R. et al. Controlled experiments support the role of function in the evolution of the North American copper tool repertoire. *J. Archaeol. Sci. Rep.* **26**, 101917 (2019).
59. Mika, A. et al. Miniaturization optimized weapon killing power during the social stress of late pre-contact North America (AD 600–1600). *PLoS ONE* **15**, e0230348 (2020).
60. Mullen, D. et al. Comparison of four ballistic and thrusting target materials: An experimental and Bayesian approach using static testing of stone and steel arrow tips. *Archaeometry* **65**, 1108–1124 (2023).
61. Sitton, J., Stenzel, C., Buchanan, B., Eren, M. I. & Story, B. Static penetration assessment of stone weapon tip geometry metrics and comparison of static and dynamic penetration depths. *Archaeometry* **65**, 463–479 (2023).
62. Conrad, G. et al. Clovis bone versus stone weapon tip penetration: Thinking about relative costs and benefits, experimental assumptions, and archaeological unknowns at Sheriden Cave, Ohio. *U.S.A. J. Archaeol. Sci. Rep.* **52**, 104295 (2023).
63. Bleed, P. The optimal design of hunting weapons: Maintainability or reliability. *Am. Antiq.* **51**, 737–747 (1986).
64. Bebbler, M. R. et al. The gravity of Paleolithic hunting. *J. Archaeol. Sci. Rep.* **59**, 104785 (2024).
65. Sisk, M. L. & Shea, J. J. Experimental use and quantitative performance analysis of triangular flakes (Levallois points) used as arrowheads. *J. Archaeol. Sci.* **36**, 2039–2047 (2009).
66. Iovita, R., Schönekeß, H., Gaudzinski-Windheuser, S. & Jäger, F. Projectile impact fractures and launching mechanisms: results of a controlled ballistic experiment using replica Levallois points. *J. Archaeol. Sci.* **48**, 73–83 (2014).
67. Gaillard, Y. et al. Assessing hafting adhesive efficiency in the experimental shooting of projectile points: A new device for instrumented and ballistic experiments. *Archaeometry* **58**, 465–483 (2016).
68. Lowe, C. et al. Controlled ballistics tests of ground, percussion-flaked, and pressure-flaked projectile point impact durability: Implications for archaeological method and theory. *J. Archaeol. Sci. Rep.* **24**, 677–682 (2019).
69. Milks, A., Parker, D. & Pope, M. External ballistics of Pleistocene hand-thrown spears: Experimental performance data and implications for human evolution. *Sci. Rep.* **9**, 820 (2019).
70. Coppe, J., Lepers, C. & Rots, V. projectiles under a new angle: A ballistic analysis provides an important building block to grasp Paleolithic weapon technology. *J. Archaeol. Method Theory* **29**, 1131–1157 (2022).
71. Torrence, R. *Time, Energy and Stone Tools* (Cambridge University Press, 1989).
72. Neeley, M. P. & Barton, C. M. A new approach to interpreting late Pleistocene microlith industries in southwest Asia. *Antiquity* **68**, 275–288 (1994).
73. Kuhn, S. L. & Shimelmitz, R. From hafting to retooling: Miniaturization as tolerance control in Paleolithic and Neolithic blade production. *J. Archaeol. Method Theory* **30**, 678–701 (2023).
74. Villaverde Bonilla, V., Domingo Sanz, I. & López Montalvo, E. Las figuras levantinas del Abric i La Sarga: Aproximación a su estilo y composición. In *La Sarga Arte Rupestre y Territorio* (eds Mauro, H. P. & Josep Maria, S. M.) 101–126 (Ayuntamiento de Alcoy/ Ajuntament d'Alcoi, Alcoi, 2002).
75. Borgia, V., Carlin, M. G. & Crezzini, J. Poison, plants and Palaeolithic hunters. An analytical method to investigate the presence of plant poison on archaeological artefacts. *Quat. Int.* **427**, 94–103 (2017).
76. Martínez Varea, C. M. *La Recolección de Vegetales Por Los Grupos Humanos Durante el Paleolítico Superior. Frutos, Semillas, Tallos y Hojas de la Cova de les Cendres (Teulada-Moraira, Alicante)* (Universitat de València, 2019).
77. Riede, F. & Lombard, M. Hunting with poisoned arrows during the Terminal Pleistocene in Northern Europe? A tip cross-sectional area assessment and list of potential arrow poison ingredients. *J. Archaeol. Sci. Rep.* **59**, 104757 (2024).
78. Eren, M. I. & Meltzer, D. J. Controls, conceits, and aiming for robust inferences in experimental archaeology. *J. Archaeol. Sci. Rep.* **53**, 104411 (2024).
79. Juan Cabanilles, J. *El Utillaje de Piedra Tallada En La Prehistoria Reciente Valenciana: Aspectos Tipológicos, Estilísticos y Evolutivos* (Servicio de Investigación Prehistórica, 2008).
80. Key, A. et al. Quantifying edge sharpness on stone flakes: comparing mechanical and micro-geometric definitions across multiple raw materials from Olduvai gorge (Tanzania). *J. Archaeol. Method Theory* **31**, 51–74 (2024).
81. Lycett, S. J. Quantifying Transitions: Morphometric Approaches to Palaeolithic Variability and Technological Change. In *Sourcebook of Paleolithic Transitions: Methods, Theories, and Interpretations* (eds Camps, M. & Chauhan, P.) 79–92 (Springer, 2009). https://doi.org/10.1007/978-0-387-76487-0_5.
82. Buchanan, B. & Collard, M. A geometric morphometrics-based assessment of blade shape differences among Paleoindian projectile point types from western North America. *J. Archaeol. Sci.* **37**, 350–359 (2010).
83. Thulman, D. K. Discriminating Paleoindian point types from Florida using landmark geometric morphometrics. *J. Archaeol. Sci.* **39**, 1599–1607 (2012).
84. Buchanan, B., O'Brien, M. J. & Collard, M. Continent-wide or region-specific? A geometric morphometrics-based assessment of variation in Clovis point shape. *Archaeol. Anthropol. Sci.* **6**, 145–162 (2014).
85. Okumura, M. & Araujo, A. G. M. Archaeology, biology, and borrowing: A critical examination of Geometric Morphometrics in Archaeology. *J. Archaeol. Sci.* **101**, 149–158 (2019).
86. Cortell-Nicolau, A. L. *evolució de Las Puntas de Proyectoil en la Transición a Las Economías Productoras en el Este y Noreste de la Península Ibérica: Implicaciones Filogenéticas, Culturales y Sociales* (Universitat de València, 2020).
87. Timbrell, L. et al. Testing inter-observer error under a collaborative research framework for studying lithic shape variability. *Archaeol. Anthropol. Sci.* **14**, 209 (2022).
88. Way, A. M., de la Peña, P., de la Peña, E. & Wadley, L. Howiesons Poort backed artifacts provide evidence for social connectivity across southern Africa during the Final Pleistocene. *Sci. Rep.* **12**, 9227 (2022).
89. Flanders, E. & Key, A. The West Tofts handaxe: A remarkably average, structurally flawed, utilitarian biface. *J. Archaeol. Sci.* **160**, 105888 (2023).
90. Clark, J., Shipton, C., Moncel, M.-H., Nigst, P. R. & Foley, R. A. When is a handaxe a planned-axe? Exploring morphological variability in the Acheulean. *PLoS ONE* **19**, e0307081 (2024).
91. Bebbler, M. R. et al. The non-invention of the ceramic arrowhead in world archaeology. *J. Archaeol. Sci. Rep.* **31**, 102283 (2020).
92. Eren, M. I., Bebbler, M. R., Knell, E. J., Story, B. & Buchanan, B. Plains Paleoindian projectile point penetration potential. *J. Anthropol. Res.* **78**, 84–112 (2022).
93. Key, A. et al. Comparing the use of meat and clay during cutting and projectile research. *Eng. Fract. Mech.* **192**, 163–175 (2018).
94. Flom, L. & Cassell, D. Stopping stepwise: Why stepwise and similar selection methods are bad, and what you should use. *NorthEast SAS Users Group Inc 20th Annu. Conf. 11–14th Novemb. 2007 Baltim. Md.* (2007).
95. Heinze, G. & Dunkler, D. Five myths about variable selection. *Transpl. Int. Off. J. Eur. Soc. Organ Transplant.* **30**, 6–10 (2017).
96. Smith, G. Step away from stepwise. *J. Big Data* **5**, 32 (2018).
97. Ullmann, T. et al. Evaluating variable selection methods for multivariable regression models: A simulation study protocol. *PLoS ONE* **19**, e0308543 (2024).

98. Gelman, A. Prior distributions for variance parameters in hierarchical models (comment on article by Browne and Draper). *Bayesian Anal.* **1**, 515–534 (2006).
99. Gelman, A., Hwang, J. & Vehtari, A. Understanding predictive information criteria for Bayesian models. *Stat. Comput.* **24**, 997–1016 (2014).
100. Polson, N. G. & Scott, J. G. On the half-Cauchy prior for a global scale parameter. *Bayesian Anal.* **7**, 887–902 (2012).
101. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing (2023).
102. de Valpine, P. et al. Programming with models: Writing statistical algorithms for general model structures with NIMBLE. *J. Comput. Graph. Stat.* **26**, 403–413 (2017).
103. de Valpine, P. et al. *NIMBLE User Manual*. vol. R package manual version 1.2.1 (2024).
104. Youngflesh, C. MCMCvis: Tools to visualize, manipulate, and summarize MCMC output. *J. Open Source Softw.* **3**, 640 (2018).
105. Plummer, M., Best, N., Cowles, K. & Vines, K. CODA: Convergence diagnosis and output analysis for MCMC. *R News* **6**, 7–11 (2006).
106. Revelle, W. *psych: Procedures for Psychological, Psychometric, and Personality Research* (Northwestern University, 2024).
107. Bonhomme, V., Picq, S., Gaucherel, C. & Claude, J. Momocs: Outline analysis using R. *J. Stat. Softw.* **56**, 1–24 (2014).
108. Crema, E. R. & Shoda, S. A Bayesian approach for fitting and comparing demographic growth models of radiocarbon dates: A case study on the Jomon-Yayoi transition in Kyushu (Japan). *PLoS ONE* **16**, e0251695 (2021).
109. Timpon, A., Barberena, R., Thomas, M. G., Méndez, C. & Manning, K. Directly modelling population dynamics in the South American arid diagonal using 14C dates. *Philos. Trans. R. Soc. B Biol. Sci.* **376**, 20190723 (2021).
110. Atkins, T. *The Science and Engineering of Cutting* (Butterworth-Heinemann, 2009). <https://doi.org/10.1016/B978-0-7506-8531-3.00018-3>.
111. Piqué, R. et al. Characterizing prehistoric archery: technical and functional analyses of the Neolithic bows from La Draga (NE Iberian Peninsula). *J. Archaeol. Sci.* **55**, 166–173 (2015).
112. Bailey, G., Galanidou, N., Peeters, H., Jöns, H. & Mennenga, M. *The Archaeology of Europe's Drowned Landscapes* (Springer, 2020).
113. Bertin, I. et al. First evidence of early neolithic archery from Cueva de los Murciélagos (Albuñol, Granada) revealed through combined chemical and morphological analysis. *Sci. Rep.* **14**, 29247 (2024).
114. Bergman, C. A., McEwen, E. & Miller, R. Experimental archery: Projectile velocities and comparison of bow performances. *Antiquity* **62**, 658–670 (1988).
115. Lepers, C. & Rots, V. The important role of bow choice and arrow fletching in projectile experimentation. A ballistic approach. *J. Archaeol. Sci. Rep.* **34**, 102613 (2020).
116. Out, W. A. *Sowing the seed?: Human Impact and Plant Subsistence in Dutch Wetlands During the Late Mesolithic and Early and Middle Neolithic (5500–3400 cal BC)* (Leiden University, 2009).
117. Bettinger, R. L. & Eerkens, J. Point typologies, cultural transmission, and the spread of bow-and-arrow technology in the prehistoric great basin. *Am. Antiq.* **64**, 231–242 (1999).
118. Crombé, P., Perdaen, Y., Sergeant, J. & Caspar, J.-P. Wear analysis on early mesolithic microliths from the verrebroek site, east flanders, Belgium. *J. Field Archaeol.* **28**, 253–269 (2001).
119. Zhilin, M. G. & Ruev, V. L. Arrowheads of the Shan-Koba culture in Crimea. *L'Anthropologie* **128**, 103258 (2024).
120. Milks, A., Champion, S., Cowper, E., Pope, M. & Carr, D. Early spears as thrusting weapons: Isolating force and impact velocities in human performance trials. *J. Archaeol. Sci. Rep.* **10**, 191–203 (2016).
121. Pétillon, J.-M. et al. Hard core and cutting edge: Experimental manufacture and use of Magdalenian composite projectile tips. *J. Archaeol. Sci.* **38**, 1266–1283 (2011).
122. Sittin, J., Story, B., Buchanan, B. & Eren, M. I. Tip cross-sectional geometry predicts the penetration depth of stone-tipped projectiles. *Sci. Rep.* **10**, 13289 (2020).
123. Grady, J. H. & Churchill, S. E. Projectile point morphology and penetration performance. *J. Archaeol. Sci. Rep.* **48**, 103893 (2023).
124. Binder, D., Collina, C., Guilbert, R., Perrin, T. & GarcíaPuchol, O. Pressure-Knapping Blade Production in the North-Western Mediterranean Region During the Senventh Millenium cal B.C. In *The Emergence of Pressure Blade Making: From Origin to Modern Experimentation* (ed. Desrosiers, M.) 199–217 (Springer, 2012).
125. Perrin, T., Manen, C., Valdeyron, N. & Guislaine, J. Beyond the sea... The Neolithic transition in the southwest of France. *Quat. Int.* **470**, 318–332 (2018).
126. Soto, A., Domingo, R., García-Simón, L. M., Alday, A. & Montes, L. For a fistful of geometric microliths: Reflections on the Sauveterrian industries from the upper and middle Ebro Basin (Spain). *Quat. Int.* **564**, 61–74 (2020).
127. Eerkens, J. W. & Lipo, C. P. Cultural transmission theory and the archaeological record: Providing context to understanding variation and temporal changes in material culture. *J. Archaeol. Res.* **15**, 239–274 (2007).
128. Shennan, S. Evolution in archaeology. *Annu. Rev. Anthropol.* **37**, 75–91 (2008).
129. Lycett, S. J. Cultural evolutionary approaches to artifact variation over time and space: Basis, progress, and prospects. *J. Archaeol. Sci.* **56**, 21–31 (2015).
130. Crema, E. R., Kandler, A. & Shennan, S. Revealing patterns of cultural transmission from frequency data: equilibrium and non-equilibrium assumptions. *Sci. Rep.* **6**, 39122 (2016).
131. Crema, E. R., Bortolini, E. & Lake, M. How cultural transmission through objects impacts inferences about cultural evolution. *J. Archaeol. Method Theory* **31**, 202–226 (2024).
132. García Puchol, O. & La Aura Tortosa, J. E. dinámica secuencial del Mesolítico en la fachada mediterránea peninsular. In *El abrigo de la Falguera (Alcoi Alacant): 8.000 años de Ocupación Humana en la Cabecera del río de Alcoi* (eds GarcíaPuchol, O. & Aura Tortosa, J. E.) 137–157 (Museo Arqueológico de Alicante-MARQ, 2006).
133. Martí Oliver, B., Aura Tortosa, J. E., Juan Cabanilles, J., García Puchol, O. & Fernández López de Pablo, J. El Mesolítico Geométrico de tipo 'Cocina' en el País Valenciano. In *El Mesolítico Geométrico en la Península Ibérica* (eds Utrilla Miranda, P. & Montes Ramírez, L.) 205–258 (Universidad de Zaragoza, 2009).
134. Jover Maestre, F. J. El instrumental lítico tallado de Benàmer: Continuidad y ruptura en los procesos de producción lítica tallada entre el VII y el IV milenio cal BC. In *Benàmer (Muro d'Alcoi, Alicante)* (eds TorregrosaGiménez, P. et al.) 133–204 (Servicio de Investigación Prehistórica del Museo de Prehistoria de Valencia, 2011).
135. JuanCabanilles, J. & GarcíaPuchol, O. Rupture et continuité dans la néolithisation du versant méditerranéen de la péninsule Ibérique: Mise à l'épreuve du modèle de dualité culturelle. In *Transitions, Ruptures et Continuité en préhistoire* Vol. 1 (eds Jaubert, J. et al.) 405–417 (Société Préistorique Française, 2013).

Acknowledgements

The Instron materials tester used in this study was available due to a grant from UKRI Arts and Humanities Research Council (AHRC) Capability for Collections (CapCo) Fund for the Cambridge Heritage Science Hub (CHERISH) Initiative (AH/V011685/1). We want to thank Dr. Miguel Ángel Martínez-Beneito, from the Department of Statistics and Operational Research at the University of València, for his key advice on specific stats development. Any potential misinterpretations are exclusively our own. We finally want to thank two anonymous reviewers, whose comments greatly improved the quality of this paper.

Author contributions

ACN, AK and AP designed the overall research protocol equally. ACN did the statistical analysis and main writing. AK did the penetration tests and main writing. AP produced the experimental microliths. All authors added, commented, edited and reviewed the final manuscript.

Funding

Open Access funding enabled and organized by Projekt DEAL.

Declarations

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to A.C.-N.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025