



Intrinsic timed-Hausdorff convergence and its implications

Raquel Perales^{1,2} 

Received: 23 November 2025 / Accepted: 23 April 2026

© The Author(s) 2026

Abstract

Sakovich–Sormani introduced several notions of distance between certain classes of Lorentzian manifolds. These distances use the Hausdorff and Gromov–Hausdorff distances and, therefore, extend naturally to a broader class of spaces. Here we show that, for timed-metric-spaces, intrinsic timed–Hausdorff convergence implies (timeless) Gromov–Hausdorff convergence as well as big bang convergence, among other related implications for future-developed convergence.

Mathematics Subject Classification 53C23 53C50

Contents

1	Introduction	
2	Background	
2.1	Metric spaces and distances between them	
2.2	Timed-metric-spaces and distances between them	
3	Relationships between the notions of convergence	
3.1	Proof of Theorem 1.2	
3.2	Proof of first part of Theorem 1.3 and Theorem 1.4	
3.3	Proof of second part of Theorem 1.3 and Proof of Theorem 1.5	
	References	

1 Introduction

Sakovich and Sormani in [8] introduced several notions of distance between certain classes of Lorentzian manifolds. To do this, they gave a canonical method for converting smooth Lorentzian manifolds satisfying certain hypotheses—known as causally null compactifiable—into compact timed-metric-spaces. This conversion uses the cos-

R. Perales was funded by the Austrian Science Fund (FWF) [Grant DOI: 10.55776/EFP6].

✉ Raquel Perales
raquel.perales@cimat.mx

¹ Math Department, University of Vienna, Oskar-Morgenstern-Platz 1, 1090 Wien, Austria

² Centro de Investigación en Matemáticas, De Jalisco s/n, Valenciana, Guanajuato, Gto. Mexico 36023, Austria

mological time function of Anderson–Galloway–Howard [2] together with the null distance of Sormani–Vega [10].

Other approaches to convergence for Lorentzian manifolds were developed by Minguzzi–Suhr [6] and by Mondino–Sämman [7]. Minguzzi–Suhr established a compactness theorem for “Lorentzian metric spaces” (which are not metric spaces). In addition, Mondino–Sämman introduced the notion of Lorentzian Gromov–Hausdorff convergence for “Lorentzian pre-length spaces”, using causal diamonds and time separation functions, and proved a precompactness theorem.

In this short note, we prove several relationships between Sormani–Sakovich’s distances in the general setting of timed-metric-spaces, c.f. Conjectures 6.1–6.5 in [8]. No knowledge of Lorentzian geometry is required, but applications to the convergence of Lorentzian manifolds are expected. We note that we do not use the compactness and embedding theorem for the intrinsic timed-Hausdorff distance of Che–Perales–Sormani [5], although it could provide alternative proofs of some of the results.

We begin by defining timed-metric-spaces and stating our first result. All remaining relevant definitions can be found in Sect. 2.

Definition 1.1 A **timed-metric-space**, (X, d, τ) , is a metric space, (X, d) , endowed with a 1-Lipschitz function $\tau: X \rightarrow [0, \infty)$, that we call time function.

Theorem 1.2 (Part of Conjecture 6.1 in [8]) *If (X_j, d_j, τ_j) , $j \in \mathbb{N}$, and $(X_\infty, d_\infty, \tau_\infty)$ are compact timed-metric-spaces converging in the intrinsic timed-Hausdorff sense as in Definition 2.4,*

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0, \quad (1)$$

then we have (timeless) Gromov-Hausdorff convergence:

$$d_{GH}\left((X_j, d_j), (X_\infty, d_\infty)\right) \rightarrow 0. \quad (2)$$

There are special classes of timed-metric-spaces in which the time functions have explicit formulas. These include big bang spaces and future-developed spaces. For big bangs, Sormani–Vega [9] and Sakovich–Sormani [8] defined *BB–GH* convergence, while for future-developed spaces Sakovich–Sormani [8] introduced *FD–HH* convergence, which can be seen as a particular case of the GH convergence for metric pairs defined by Che, Galaz-García, Guijarro, and Membrillo Solis [4]. We now state some of the relationships between intrinsic timed–Hausdorff convergence and these other two notions of convergence.

Theorem 1.3 (Part of Conjectures 6.2 and 6.4 in [8]) *Let (X_j, d_j, τ_j) , $j \in \mathbb{N}$, and $(X_\infty, d_\infty, \tau_\infty)$ be compact timed-metric-spaces.*

If the (X_j, d_j, τ_j) , $j \in \mathbb{N}$, are big bang as in Definition 2.5 and converge in the intrinsic timed-Hausdorff sense as in Definition 2.4,

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0, \quad (3)$$

then $(X_\infty, d_\infty, \tau_\infty)$ is a big bang and we have convergence in the BB-GH sense as in Definition 2.7,

$$d_{BB-GH}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0. \quad (4)$$

Conversely, if all the (X_j, d_j, τ_j) , $j \in \mathbb{N} \cup \{\infty\}$, are big bang and converge in BB-GH sense,

$$d_{BB-GH}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0 \quad (5)$$

then they converge in the intrinsic timed-Hausdorff sense,

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0. \quad (6)$$

We conclude this section with two results on future-developed spaces. We recall that Conjecture 6.3 in [8] asks for a sequence of (smooth) future-developed spaces that converges in the intrinsic timed-Hausdorff sense to a big bang space. Thus, one should not expect a sequence of future-developed spaces that converges in the intrinsic timed-Hausdorff sense to also converge in the FD-HH sense. However, convergence does hold in the more general setting of metric pairs.

Theorem 1.4 *Let (X_j, d_j, τ_j) , $j \in \mathbb{N}$, and $(X_\infty, d_\infty, \tau_\infty)$ be compact timed-metric-spaces converging in the intrinsic timed-Hausdorff sense as in Definition 2.4,*

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0, \quad (7)$$

If the (X_j, d_j, τ_j) , $j \in \mathbb{N}$, are future-developed as in Definition 2.6, then $\tau_\infty^{-1}(0) \neq \emptyset$, $\tau_\infty = d_\infty(\tau_\infty^{-1}(0), \cdot)$, and the sequence $(X_j, \tau_j^{-1}(0))$ converges as metric pairs in the Gromov-Hausdorff sense to $(X_\infty, \tau_\infty^{-1}(0))$.

Theorem 1.5 (Part of Conjecture 6.5 in [8]) *Let (X_j, d_j, τ_j) , $j \in \mathbb{N}$, and $(X_\infty, d_\infty, \tau_\infty)$ be compact future-developed timed-metric-spaces as in Definition 2.6 with*

$$d_{FD-HH}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0, \quad (8)$$

then

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0. \quad (9)$$

The structure of the paper is as follows. In Sect. 2 we review basic notions and results on the Gromov-Hausdorff distance, followed by the convergence notions for timed-metric-spaces introduced by Sakovich-Sormani. In Sect. 3 we prove the results stated in the introduction.

2 Background

We review basic notions and results on the Gromov-Hausdorff distance, followed by the convergence notions for timed-metric-spaces introduced by Sakovich-Sormani,

closely following [8]. For further background on metric spaces, we refer the reader to [3].

2.1 Metric spaces and distances between them

Given a metric space (Z, d_Z) , the **Hausdorff distance in Z** between subsets, $X, Y \subset Z$, is defined as

$$d_H^Z(X, Y) = \inf \left\{ \varepsilon > 0 : \begin{array}{l} \forall x \in X, \exists y \in Y \text{ s.t. } d_Z(x, y) < \varepsilon \\ \forall y \in Y, \exists x \in X \text{ s.t. } d_Z(x, y) < \varepsilon \end{array} \right\} \quad (10)$$

For two compact metric spaces (X, d_X) and (Y, d_Y) , the **Gromov-Hausdorff distance** (GH) between them is defined by

$$d_{GH}((X, d_X), (Y, d_Y)) = \inf d_H^Z(\varphi(X), \psi(Y)), \quad (11)$$

where the infimum is over all distance-preserving maps, $\varphi : X \rightarrow Z$ and $\psi : Y \rightarrow Z$, into all possible metric spaces, Z .

We now focus in a particular class of distance-preserving maps. For that, let

$$\ell^\infty = \{(s_1, s_2, \dots) : s_i \in \mathbb{R}, d_{\ell^\infty}((s_1, s_2, \dots), (0, 0, \dots)) < \infty\}$$

where

$$d_{\ell^\infty}((s_1, s_2, \dots), (r_1, r_2, \dots)) = \sup\{|s_i - r_i| : i \in \mathbb{N}\}.$$

Definition 2.1 Given a compact metric space, (X, d) , and a countable and dense collection of points $\mathcal{N} = \{x_1, x_2, \dots\} \subset X$, the Fréchet map of X given by $\mathcal{N}$,

$$\kappa_X = \kappa_{X, \mathcal{N}} : (X, d) \rightarrow (\ell^\infty, d_{\ell^\infty}),$$

is defined as

$$\kappa_X(x) = (d(x_1, x), d(x_2, x), \dots). \quad (12)$$

Fréchet maps are distance-preserving, c.f. [8]. Following the notation of Sakovich-Sormani (see Section 2.6 in [8]), given two compact metric spaces (X, d_X) and (Y, d_Y) , let

$$d_{\kappa-GH}((X, d_X), (Y, d_Y)) := \inf d_H^{\ell^\infty}(\kappa_X(X), \kappa_Y(Y)) \quad (13)$$

where the infimum is taken over all pairs of Fréchet maps κ_X and κ_Y . With this notation, the following inequalities hold:

$$d_{GH} \leq d_{\kappa-GH} \leq 2d_{GH}. \quad (14)$$

The first inequality follows right away. The second follows from the next proposition, which is proven in [8].

Proposition 2.2 *Let (X, d_X) and (Y, d_Y) be two compact metric spaces. If*

$$d_{GH}(X, Y) < R \quad (15)$$

then there exist Fréchet maps $\kappa_X : X \rightarrow \ell^\infty$ and $\kappa_Y : Y \rightarrow \ell^\infty$ such that

$$d_H^{\ell^\infty}(\kappa_X(X), \kappa_Y(Y)) < 2R. \quad (16)$$

More explicitly, there exist countable dense sets

$$\{x_1, x_2, \dots\} \subset X \text{ and } \{y_1, y_2, \dots\} \subset Y$$

such that the Fréchet maps defined by them satisfy (16).

2.2 Timed-metric-spaces and distances between them

For this section, recall the definition of timed-metric space, Definition 1.1.

Definition 2.3 Given a compact timed-metric-space, (X, d, τ) , and a countably dense collection of points, $\mathcal{N} \subset X$, we define the timed-Fréchet map,

$$\kappa_{\tau, X} = \kappa_{\tau, X, \mathcal{N}} : X \rightarrow [0, \tau_{\max}] \times \ell^\infty \subset \ell^\infty,$$

by $\kappa_{\tau, X} = (\tau, \kappa_X)$, where $\kappa_X = \kappa_{X, \mathcal{N}}$ is as in Definition 2.1 and $\tau_{\max} = \max_{x \in X} \tau(x)$.

Timed-Fréchet maps are distance-preserving and restricting to them, one can define the intrinsic timed-Hausdorff distance.

Definition 2.4 The **intrinsic timed-Hausdorff distance** between two compact timed-metric-spaces, (X_1, d_1, τ_1) and (X_2, d_2, τ_2) , is defined as

$$d_{\tau-H}((X_1, d_1, \tau_1), (X_2, d_2, \tau_2)) = \inf d_H^{\ell^\infty}(\kappa_{\tau_1, X_1}(X_1), \kappa_{\tau_2, X_2}(X_2)) \quad (17)$$

where the infimum is taken over all possible timed-Fréchet maps κ_{τ_1, X_1} and κ_{τ_2, X_2} .

We now define two special cases of timed-metric-spaces and corresponding intrinsic distances between them.

Definition 2.5 A compact **timed-metric-space** (X, d, τ) **has a big bang** if $\tau^{-1}(0)$ consists of a single point, that we denote as p_{BB} and call it the big bang point, and the time function τ equals the distance function to this point:

$$\tau(p) = d(p, p_{BB}) \quad \forall p \in X. \quad (18)$$

In this case, we say that (X, d, τ) is a big bang timed-metric-space or a BB space.

Equivalently, we could define (X, d, τ) to be a BB space if there exists a point $p_{BB} \in X$, called the big bang point, such that the time function τ is given as in (18).

Definition 2.6 A **future-developed timed-metric-space**,¹ (X, d, τ) , consists of a timed-metric-space (X, d, τ) and a non-empty set $Y \subset X$, of cardinality greater than or equal to 2, such that

$$Y = \tau^{-1}(0) \quad \text{and} \quad \tau(\cdot) = d(Y, \cdot).$$

In this case, we also say that (X, d, τ) is a FD space.

Definition 2.7 The **intrinsic $BB - GH$ distance** between two compact **big bang timed-metric-spaces** is defined as

$$d_{BB-GH}((X_1, d_1, \tau_1), (X_2, d_2, \tau_2)) = d_{pt-GH}((X_1, d_1, p_{BB,1}), (X_2, d_2, p_{BB,2})), \quad (19)$$

where $p_{BB,i} = \tau_i^{-1}(0)$. Here,

$$d_{pt-GH}((X_1, d_1, p_{BB,1}), (X_2, d_2, p_{BB,2}))$$

is defined as the infimum over all distance-preserving maps $\varphi_i : X_i \rightarrow Z$ and metric spaces Z of sums of the form:

$$d_H^Z(\varphi_1(X_1), \varphi_2(X_2)) + d_Z(\varphi_1(p_{BB,1}), \varphi_2(p_{BB,2})).$$

Definition 2.8 The **$FD - HH$ distance** between two compact future-developed space-times is defined as

$$d_{FD-HH}((X_1, d_1, \tau_1), (X_2, d_2, \tau_2)) = \quad (20)$$

$$= \inf \left(d_H^Z(\varphi_1(X_1), \varphi_2(X_2)) + d_H^Z(\varphi_1(Y_1), \varphi_2(Y_2)) \right), \quad (21)$$

where the infimum is over all distance-preserving maps $\varphi_i : X_i \rightarrow Z$ and over all metric spaces Z .

Finally, we recall that given compact metric spaces (X_j, d_j) and subsets $Y_j \subset X_j$, $j \in \mathbb{N} \cup \{\infty\}$, it is said that (X_j, Y_j) , $j \in \mathbb{N}$, converge as metric pairs in the Gromov-Hausdorff to (X_∞, d_∞) sense if

$$\inf \left(d_H^Z(\varphi_j(X_j), \varphi_\infty(X_\infty)) + d_H^Z(\varphi_j(Y_j), \varphi_\infty(Y_\infty)) \right) \rightarrow 0 \quad \text{as } j \rightarrow \infty, \quad (22)$$

where the infimum is taken over all metric spaces Z and distance-preserving maps $\varphi_j : X_j \rightarrow Z$. For additional details, see [1, 4].

¹ The current synthetic definition of a future-developed timed-metric-space is not yet fully satisfactory. In the smooth Lorentzian setting, $Y \subset X$ must be a hypersurface (satisfying suitable conditions). In particular, it cannot be a big bang space and, thus, this definition precisely rules out this case.

3 Relationships between the notions of convergence

Here, we prove the results stated in the introduction.

3.1 Proof of Theorem 1.2

Lemma 3.1 *If (X_1, d_1, τ_1) and (X_2, d_2, τ_2) are compact timed-metric-spaces such that their intrinsic timed-Hausdorff distance is bounded above,*

$$d_{\tau-H}\left((X_1, d_1, \tau_1), (X_2, d_2, \tau_2)\right) < \varepsilon, \quad (23)$$

then their (timeless) Gromov-Hausdorff distance is also bounded above:

$$d_{GH}\left((X_1, d_1), (X_2, d_2)\right) < \varepsilon. \quad (24)$$

Proof By definition of intrinsic timed-Hausdorff distance, Definition 2.4, there exist timed-Fréchet maps $\varphi = \kappa_{\tau_1, X_1}$ and $\psi = \kappa_{\tau_2, X_2}$ such that setting $Z = \ell^\infty$, we have

$$d_H^Z(\varphi(X_1), \psi(X_2)) < \varepsilon. \quad (25)$$

Denote by $F : Z \rightarrow \ell^\infty$ the map that deletes the first coordinate, i.e.

$$F(x_1, x_2, x_3, \dots) = (x_2, x_3, \dots).$$

Then, for the Fréchet maps $\kappa_{X_1} = F \circ \varphi$ and $\kappa_{X_2} = F \circ \psi$ by (25), we have

$$d_H^{\ell^\infty}(\kappa_{X_1}(X_1), \kappa_{X_2}(X_2)) < \varepsilon.$$

This implies, recalling the definition of Gromov-Hausdorff distance, that

$$d_{GH}\left((X_1, d_1), (X_2, d_2)\right) < \varepsilon,$$

which is what we wanted to prove. $\square$

With the previous lemma at hand we establish our first main result.

Proof of Theorem 1.2 Assume

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0.$$

Then, for all $\varepsilon > 0$ there exists $N(\varepsilon) \in \mathbb{N}$ such that for all $j \geq N(\varepsilon)$, we have

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) < \varepsilon.$$

Thus, by applying Lemma 3.1 we obtain

$$d_{GH}\left((X_j, d_j), (X_\infty, d_\infty)\right) < \varepsilon,$$

which implies (2) and concludes the proof. $\square$

3.2 Proof of first part of Theorem 1.3 and Theorem 1.4

In this subsection, we give the proofs of two results stated in the introduction, which are very similar.

Proof of first part of Theorem 1.3 Assume

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0.$$

Thus, by definition, for all $\varepsilon > 0$ there exists $N(\varepsilon) \in \mathbb{N}$ such that for all $j \geq N(\varepsilon)$, we have

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) < \varepsilon.$$

By definition of intrinsic timed-Hausdorff distance, Definition 2.4, there exist timed-Fréchet maps $\varphi_j = \kappa_{\tau_j, X_j}$ and $\psi_j = \kappa_{\tau_\infty, X_\infty}$ such that for $j \geq N(\varepsilon)$ and setting $Z = \ell^\infty$, we have

$$d_H^Z(\varphi_j(X_j), \psi_j(X_\infty)) < \varepsilon. \quad (26)$$

Note that φ_j and ψ_j also depend on ε , when necessary and to avoid confusion, we will add a superscript and denote them as φ_j^ε and ψ_j^ε , respectively.

First we show that $\tau_\infty^{-1}(0) \neq \emptyset$. Fix $j \geq N(\varepsilon)$. By hypotheses, $\tau_j^{-1}(0)$ consists of only one point, which we denote by p_j for brevity. By (26) there exist $y_j^\varepsilon \in X_\infty$ such that

$$d_Z(\varphi_j^\varepsilon(p_j), \psi_j^\varepsilon(y_j^\varepsilon)) < \varepsilon. \quad (27)$$

By the previous inequality, and since $\varphi_j^\varepsilon(p_j) - \psi_j^\varepsilon(y_j^\varepsilon)$ has as first entry $\tau_j(p_j) - \tau_\infty(y_j^\varepsilon)$, we see that

$$\tau_\infty(y_j^\varepsilon) = |\tau_j(p_j) - \tau_\infty(y_j^\varepsilon)| < \varepsilon. \quad (28)$$

Hence, taking a sequence of positive numbers $\varepsilon_j \rightarrow 0$, we can construct a subsequence of points, that we do not relabel, y^{ε_j} contained in X_∞ such that $\tau_\infty(y^{\varepsilon_j}) < \varepsilon_j$. Since X_∞ is compact, there exists a point $y \in X_\infty$ such that, up to a subsequence, $y^{\varepsilon_j} \rightarrow y$, and by continuity of τ_∞ , we get $\tau_\infty(y) = 0$.

Now we show that $\tau_\infty^{-1}(0)$ consists of only one point. Assume that $y_1, y_2 \in \tau_\infty^{-1}(0)$. Fix $j \geq N(\varepsilon)$. By (26) there exist $x_{j1}, x_{j2} \in X_j$ such that

$$d_Z(\varphi_j(x_{ji}), \psi_j(y_i)) < \varepsilon, \quad i = 1, 2. \quad (29)$$

Since $y_1, y_2 \in \tau_\infty^{-1}(0)$, by the previous inequality and given that $\varphi_j(x_{ji}) - \psi_j(y_i)$ has as first entry $\tau_j(x_{ji}) - \tau_\infty(y_i)$, we see that

$$\tau_j(x_{ji}) = |\tau_j(x_{ji}) - \tau_\infty(y_i)| < \varepsilon \quad i = 1, 2. \quad (30)$$

By hypotheses $d_j(x_{ji}, p_j) = \tau_j(x_{ji})$. Hence, by the triangle inequality, (29) and (30):

$$\begin{aligned} d_\infty(y_1, y_2) &= d_Z(\psi_j(y_1), \psi_j(y_2)) \\ &\leq d_Z(\psi_j(y_1), \varphi_j(x_{j1})) + d_Z(\varphi_j(x_{j1}), \varphi_j(p_j)) \\ &\quad + d_Z(\varphi_j(p_j), \varphi_j(x_{j2})) + d_Z(\varphi_j(x_{j2}), \psi_j(y_2)) \\ &< 4\varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, this shows that $y_1 = y_2$. Denote by p_∞ the only point in $\tau_\infty^{-1}(0)$.

Now we show that $\tau_\infty = d_\infty(p_\infty, \cdot)$. Fix $j \geq N(\varepsilon)$. First note that by (29) and (30), we have

$$\begin{aligned} d_Z(\psi_j(p_\infty), \varphi_j(p_j)) &\leq d_Z(\psi_j(p_\infty), \varphi_j(x_{ji})) + d_Z(\varphi_j(x_{ji}), \varphi_j(p_j)) \\ &= d_Z(\psi_j(y_i), \varphi_j(x_{ji})) + \tau_j(x_{ji}) < 2\varepsilon. \end{aligned} \quad (31)$$

Let $y \in X_\infty$. Take $x_y \in X_j$ such that

$$d_Z(\varphi_j(x_y), \psi_j(y)) < \varepsilon. \quad (32)$$

By considering the first coordinate function of φ_j and ψ_j , we know that $|\tau_\infty(y) - \tau_j(x_y)| < \varepsilon$. Furthermore,

$$\begin{aligned} \tau_j(x_y) &= d_Z(\varphi_j(x_y), \varphi_j(p_j)) \\ &\leq d_Z(\varphi_j(x_y), \psi_j(y)) + d_Z(\psi_j(y), \psi_j(p_\infty)) + d_Z(\psi_j(p_\infty), \varphi_j(p_j)) \\ &< d_Z(\psi_j(y), \psi_j(p_\infty)) + 3\varepsilon, \end{aligned}$$

where we used the triangle inequality, (32) and (31). We can obtain a similar inequality and conclude that

$$|\tau_\infty(y) - d_Z(\psi_j(y), \psi_j(p_\infty))| < 3\varepsilon.$$

Since $\varepsilon > 0$ is arbitrary this proves that $\tau_\infty = d_\infty(p_\infty, \cdot)$. Hence, $(X_\infty, d_\infty, \tau_\infty)$ is a BB space.

We finally show BB-GH convergence. Denote by $F : \ell^\infty \rightarrow \ell^\infty$ the map given by $F(x_1, x_2, x_3, \dots) = (x_2, x_3, \dots)$. For the Fréchet maps $\kappa_{X_j} = F \circ \varphi_j$ and $\kappa_{X_\infty} = F \circ \psi_j$ by (26) and (31), we have

$$d_H^{\ell^\infty}(\kappa_{X_j}(X_j), \kappa_{X_\infty}(X_\infty)) + d^{\ell^\infty}(\kappa_{X_j}(p_j), \kappa_{X_\infty}(p_\infty)) < 3\varepsilon. \quad (33)$$

Concluding that

$$d_{BB-GH}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) < 3\varepsilon. \quad (34)$$

□

Proof of Theorem 1.4 Assume

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0.$$

Thus, by definition, for all $\varepsilon > 0$ there exists $N(\varepsilon) \in \mathbb{N}$ such that for all $j \geq N(\varepsilon)$, we have

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) < \varepsilon.$$

By definition of intrinsic timed-Hausdorff distance, Definition 2.4, there exist timed-Fréchet maps $\varphi_j = \kappa_{\tau_j, X_j}$ and $\psi_j = \kappa_{\tau_\infty, X_\infty}$ such that for $j \geq N(\varepsilon)$ and setting $Z = \ell^\infty$, we have

$$d_H^Z(\varphi_j(X_j), \psi_j(X_\infty)) < \varepsilon. \quad (35)$$

Note that φ_j and ψ_j also depend on ε , when necessary and to avoid confusion, we will add a superscript and denote them as φ_j^ε and ψ_j^ε , respectively.

First we show that $\tau_\infty^{-1}(0) \neq \emptyset$. Fix $j \geq N(\varepsilon)$. By hypotheses, there exists $p_j \in \tau_j^{-1}(0)$. By (35) there exists $y_j^\varepsilon \in X_\infty$ such that

$$d_Z(\varphi_j^\varepsilon(p_j), \psi_j^\varepsilon(y_j^\varepsilon)) < \varepsilon. \quad (36)$$

By the previous inequality, and since $\varphi_j^\varepsilon(p_j) - \psi_j^\varepsilon(y_j^\varepsilon)$ has as first entry $\tau_j(p_j) - \tau_\infty(y_j^\varepsilon)$, we see that

$$\tau_\infty(y_j^\varepsilon) = |\tau_j(p_j) - \tau_\infty(y_j^\varepsilon)| < \varepsilon. \quad (37)$$

Hence, taking a sequence of positive numbers $\varepsilon_j \rightarrow 0$, we can construct a subsequence of points, that we do not relabel, y^{ε_j} contained in X_∞ such that $\tau_\infty(y^{\varepsilon_j}) < \varepsilon_j$. Since X_∞ is compact, there exists a point $y \in X_\infty$ such that, up to a subsequence, $y^{\varepsilon_j} \rightarrow y$, and by continuity of τ_∞ , we get $\tau_\infty(y) = 0$.

We now show that $\tau_\infty = d_\infty(Y_\infty, \cdot)$. Fix $j \geq N(\varepsilon)$. Let $y \in X_\infty$. By (35) we can take $x_y \in X_j$ such that

$$d_Z(\varphi_j(x_y), \psi_j(y)) < \varepsilon. \quad (38)$$

Since X_j is a compact FD space, there exists $p \in Y_j$ such that

$$\tau_j(x_y) = d_j(Y_j, x_y) = d_j(p, x_y). \quad (39)$$

Furthermore, there exists $y_p \in X_\infty$ such that

$$d_Z(\varphi_j(p), \psi_j(y_p)) < \varepsilon. \quad (40)$$

Hence, recalling that τ_j and τ_∞ are the first coordinate functions of φ_j and ψ_j , respectively, we get

$$\tau_\infty(y_p) = |\tau_\infty(y_p) - \tau_j(p)| \quad (41)$$

$$\leq d_Z(\psi_j(y_p), \varphi_j(p)) < \varepsilon. \quad (42)$$

By (39), (38), the triangle inequality, (40) and (39), we get

$$\begin{aligned} \tau_\infty(y) &\leq \tau_j(x_y) + d_Z(\psi_j(y), \varphi_j(x_y)) \\ &\leq d_j(p, x_y) + \varepsilon \\ &\leq \{d_Z(\varphi_j(p), \psi_j(y_p)) + d_Z(\psi_j(y_p), \psi_j(y)) + d_Z(\psi_j(y), \varphi_j(x_y))\} + \varepsilon \\ &\leq d_\infty(y_p, y) + 3\varepsilon. \end{aligned}$$

Furthermore,

$$\begin{aligned} -\tau_\infty(y) &\leq -\tau_j(x_y) + d_Z(\psi_j(y), \varphi_j(x_y)) \\ &\leq -d_j(p, x_y) + \{d_Z(\psi_j(y), \psi_j(y_p)) + d_Z(\psi_j(y_p), \varphi_j(p)) + d_Z(\varphi_j(p), \psi_j(x_y))\} \\ &\leq d_\infty(y_p, y) + \varepsilon. \end{aligned}$$

Hence,

$$|\tau_\infty(y) - d_\infty(y_p, y)| \leq 3\varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, one can construct a sequence of points $y_i \in X_\infty$ such that $\tau_\infty(y_i) < \varepsilon_i$ and

$$|\tau_\infty(y) - d_\infty(y_i, y)| \leq 3\varepsilon_i.$$

for $\varepsilon_i \searrow 0$. Since X_∞ is compact, up to a subsequence, y_i converges to $q \in X_\infty$ and since τ_∞ is continuous, $q \in Y_\infty$. Hence, $\tau_\infty(y) = d_\infty(y, q) \geq d_\infty(Y, y)$. Since τ_∞ is 1-Lipschitz, the other inequality follows,

$$\tau_\infty(y) = \min_{z \in Y_\infty} |\tau_\infty(y) - \tau_\infty(z)| \leq \min_{z \in Y_\infty} d_\infty(y, z) = d_\infty(Y, y).$$

Hence, $\tau_\infty = d_\infty(Y_\infty, \cdot)$.

Denote by $F : \ell^\infty \rightarrow \ell^\infty$ the map given by $F(x_1, x_2, x_3, \dots) = (x_2, x_3, \dots)$. For $j \geq N(\varepsilon)$, and the Fréchet maps $\kappa_{X_j} = F \circ \varphi_j$ and $\kappa_{X_\infty} = F \circ \psi_j$, by (35), we have

$$d_H^{\ell^\infty}(\kappa_{X_j}(X_j), \kappa_{X_\infty}(X_\infty)) < \varepsilon. \quad (43)$$

Finally, to see that

$$d_H^{\ell^\infty}(\kappa_{X_j}(Y_j), \kappa_{X_\infty}(Y_\infty)) < 2\varepsilon, \quad (44)$$

we proceed as follows. Let $p \in Y_j$. By (35), we can take $y_p \in X_\infty$ such that

$$d_Z(\varphi_j(p), \psi_j(y_p)) < \varepsilon. \quad (45)$$

Hence, $\tau_\infty(y_p) = |\tau_\infty(y_p) - \tau_j(p)| < \varepsilon$. Since X_∞ is a compact FD space, there exists $q \in Y_\infty$ such that

$$\tau_\infty(y_p) = d_\infty(y_p, q).$$

Furthermore, there exists $y_p \in X_\infty$ such that

$$d_Z(\varphi_j(p), \psi_j(y_p)) < \varepsilon. \quad (46)$$

By the triangle inequality, we get

$$d_Z(\varphi_j(p), \psi_j(q)) < 2\varepsilon. \quad (47)$$

Similarly, for each $q \in Y_\infty$ there exists $p \in Y_j$ such that the previous inequality holds. This implies the required estimate and concludes the proof. $\square$

3.3 Proof of second part of Theorem 1.3 and Proof of Theorem 1.5

In this subsection, we give the proofs of two results stated in the introduction, which are very similar.

Lemma 3.2 *Let (X_j, d_j, τ_j) and $(X_\infty, d_\infty, \tau_\infty)$ be two compact big bang timed-metric-spaces. Assume that*

$$d_{BB-GH}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) < \varepsilon. \quad (48)$$

Then

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) < 2\varepsilon. \quad (49)$$

More explicitly, there exist countable dense sets $\mathcal{N}_{X_j}$ and $\mathcal{N}_{X_\infty}$ such that the timed-Fréchet maps κ_{τ_j, X_j} and $\kappa_{\tau_\infty, X_\infty}$ defined by them satisfy

$$d_H^{\ell_\infty}(\kappa_{\tau_j, X_j}(X_j), \kappa_{\tau_\infty, X_\infty}(X_\infty)) < 2\varepsilon. \quad (50)$$

The proof is very similar to the proof of Proposition 2.2, but we give all the details for completeness.

Proof Assume that

$$d_{BB-GH}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) < \varepsilon. \quad (51)$$

By definition of $BB-GH$ distance, there exist a compact metric space Z and isometric embeddings $\varphi_j : X_j \rightarrow Z$ and $\varphi_\infty : X_\infty \rightarrow Z$ such that

$$d_H^Z(\varphi_j(X_j), \varphi_\infty(X_\infty)) + d_Z(\varphi_j(p_{BB,j}), \varphi_\infty(p_{BB,\infty})) < \varepsilon. \quad (52)$$

So for all $x \in X_j$ there exists $y_x \in X_\infty$ such that

$$d_Z(\varphi_j(x), \varphi_\infty(y_x)) < \varepsilon, \quad (53)$$

and for all $y \in X_\infty$ there exists $x_y \in X_j$ such that

$$d_Z(\varphi_j(x_y), \varphi_\infty(y)) < \varepsilon. \quad (54)$$

Let $\{x'_1, x'_2, \dots\}$ and $\{y'_1, y'_2, \dots\}$ be two countably dense collection of points in X_j and X_∞ , respectively. Applying (53) and (54), we obtain larger countably and dense collections of points,

$$\mathcal{N}_j = \{x_1, x_2, x_3, \dots\} = \{x'_1, x_{y'_1}, x'_2, x_{y'_2}, x'_3, x_{y'_3}, \dots\} \subset X_j$$

and

$$\mathcal{N}_\infty = \{y_1, y_2, y_3, \dots\} = \{y_{x'_1}, y'_1, y_{x'_2}, y'_2, y_{x'_3}, y'_3, \dots\} \subset X_\infty.$$

We then define timed-Fréchet maps, $\kappa_{\tau_{X_j} X_j} = (\tau_j, \kappa_{X_j})$ and $\kappa_{\tau_\infty X_\infty} = (\tau_\infty, \kappa_{X_\infty})$, using these collections.

Given $x \in X_j$ and $y_x \in X_\infty$ that satisfy (53), we claim that

$$d_{\ell^\infty}(\kappa_{\tau_j X_j}(x), \kappa_{\tau_\infty X_\infty}(y_x)) < 2\varepsilon. \quad (55)$$

To prove the claim, recall that

$$\begin{aligned} \kappa_{\tau_j X_j}(x) &= (\tau_j(x), d_j(x'_1, x), d_j(x_{y'_1}, x), d_j(x'_2, x), d_j(x_{y'_2}, x), \dots) \\ \kappa_{\tau_\infty X_\infty}(y_x) &= (\tau_\infty(y_x), d_\infty(y_{x'_1}, y_x), d_\infty(y'_1, y_x), d_\infty(y_{x'_2}, y_x), d_\infty(y'_2, y_x), \dots). \end{aligned}$$

Then by the fact that φ_j and φ_∞ are distance-preserving, the triangle inequality, (53) and the second term to the left of (52), we have

$$\begin{aligned} |\tau_j(x) - \tau_\infty(y_x)| &= |d_j(x, p_{BBj}) - d_\infty(y_x, p_{BB\infty})| \\ &\leq d_Z(\varphi_j(x), \varphi_\infty(y_x)) + d_Z(\varphi_j(p_{BBj}), \varphi_\infty(p_{BB\infty})) \\ &< 2\varepsilon. \end{aligned} \quad (56)$$

Similarly, for any $i \in \mathbb{N}$,

$$\begin{aligned} |d_j(x'_i, x) - d_\infty(y_{x'_i}, y_x)| &= |d_Z(\varphi_j(x'_i), \varphi_j(x)) - d_Z(\varphi_\infty(y_{x'_i}), \varphi_\infty(y_x))| \\ &\leq d_Z(\varphi_j(x'_i), \varphi_\infty(y_{x'_i})) + d_Z(\varphi_j(x), \varphi_\infty(y_x)) \\ &< 2\varepsilon, \\ |d_j(x_{y'_i}, x) - d_\infty(y'_i, y_x)| &< 2\varepsilon. \end{aligned}$$

Hence, (55) holds. Analogously, one can show that for any $y \in X_\infty$ there exists $x_y \in X_j$ such that

$$d_{\ell^\infty}(\kappa_{\tau_j X_j}(x_y), \kappa_{\tau_\infty X_\infty}(y)) < 2\varepsilon. \quad (57)$$

From (55) and (57) we conclude that (50) holds. $\square$

Proof of second part of Theorem 1.3 Assume that

$$d_{BB-GH}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) \rightarrow 0, \quad (58)$$

Then, for all $\varepsilon > 0$ there exists $N(\varepsilon) \in \mathbb{N}$ such that for all $j \geq N(\varepsilon)$, we have

$$d_{BB-GH}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) < \varepsilon. \quad (59)$$

Now by Lemma 3.2, for all $j \geq N(\varepsilon)$ we have

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) < 2\varepsilon, \quad (60)$$

which concludes the proof of the theorem. $\square$

Lemma 3.3 *Let (X_j, d_j, τ_j) and $(X_\infty, d_\infty, \tau_\infty)$ be two compact future-developed timed-metric-spaces. Assume that*

$$d_{FD-HH}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) < \varepsilon, \quad (61)$$

then

$$d_{\tau-H}\left((X_j, d_j, \tau_j), (X_\infty, d_\infty, \tau_\infty)\right) < 2\varepsilon. \quad (62)$$

More explicitly, there exist countable dense sets $\mathcal{N}_{X_j}$ and $\mathcal{N}_{X_\infty}$ such that the timed-Fréchet maps κ_{τ_j, X_j} and $\kappa_{\tau_\infty, X_\infty}$ defined by them satisfy

$$d_H^{\ell^\infty}(\kappa_{\tau_j, X_j}(X_j), \kappa_{\tau_\infty, X_\infty}(X_\infty)) < 2\varepsilon. \quad (63)$$

Proof The proof is very similar to the one of Lemma 3.2. So, we use the same notation as in that proof and only focus in the parts where they differ.

By definition of $FF - DH$, there exist a compact metric space Z and isometric embeddings, $\varphi_j : X_j \rightarrow Z$ and $\varphi_\infty : X_\infty \rightarrow Z$, such that

$$d_H^Z(\varphi_j(X_j), \varphi_\infty(X_\infty)) + d_H^Z(\varphi_j(Y_j), \varphi_\infty(Y_\infty)) < \varepsilon. \quad (64)$$

Construct timed-Fréchet maps, $\kappa_{\tau_j, X_j} = (\tau_j, \kappa_{X_j})$ and $\kappa_{\tau_\infty, X_\infty} = (\tau_\infty, \kappa_{X_\infty})$, as in the proof of Lemma 3.2. Given $x \in X_j$ and $y_x \in X_\infty$ that satisfy

$$d_Z(\varphi_j(x), \varphi_\infty(y_x)) < \varepsilon,$$

we claim that

$$d_{\ell^\infty}(\kappa_{\tau_j, X_j}(x), \kappa_{\tau_\infty, X_\infty}(y_x)) < 2\varepsilon.$$

The proof of this claim proceeds as in the proof of Lemma 3.2, except for (56). For this, note that $Y_j = \tau_j^{-1}(0) \subset X_j$ is nonempty and compact, so there exists $p_x \in Y_j$ such that

$$\tau_j(x) = d_j(Y, x) = d_j(x, p_x).$$

By the second term to the left of (64) there exist $y_{p_x} \in Y_\infty$ such that

$$d_Z(\varphi_j(p_x), \varphi_\infty(y_{p_x})) < \varepsilon.$$

Hence,

$$\begin{aligned} \tau_\infty(y_x) - \tau_j(x) &= \tau_\infty(y_x) - d_j(x, p_x) \\ &\leq d_\infty(y_x, y_{p_x}) - d_j(x, p_x) \\ &\leq d_Z(\varphi_j(x), \varphi_\infty(y_x)) + d_Z(\varphi_j(p_x), \varphi_\infty(y_{p_x})) < 2\varepsilon, \end{aligned}$$

where we used the fact that $\tau_\infty = d(Y_\infty, \cdot)$. In a similar way, we get the other inequality and conclude that

$$|\tau_j(x) - \tau_\infty(y_x)| < 2\varepsilon.$$

This concludes the proof of the claim. The remaining part of the proof proceeds as in the proof of the Lemma 3.2. $\square$

Proof of Theorem 1.5 This follows applying Lemma 3.3. $\square$

Funding Open access funding provided by University of Vienna.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Ahumada Gómez, A., Che, M.: Gromov-Hausdorff convergence of metric pairs and metric tuples. *Differ. Geom. Appl.* **94**, 102135 (2024)
2. Andersson, L., Galloway, G.J., Howard, R.: The cosmological time function. *Class. Quantum Gravity* **15**(2), 309–322 (1998)
3. Burago, D., Burago, Y., Ivanov, S.: A Course in Metric Geometry, vol. 33. American Mathematical Society, Providence, RI (2001)
4. Che, M., Galaz-García, F., Guijarro, L., Solís, Membrillo Solís, I.: Metric geometry of spaces of persistence diagrams. *J. Appl. Comput. Topol.* **8**(8), 2197–2246 (2024)
5. Che, M., Perales, R., Sormani, C.: Gromov's compactness theorem for the intrinsic timed-Hausdorff distance (2025)
6. Minguzzi, E., Suhr, S.: Lorentzian metric spaces and their Gromov-Hausdorff convergence. *Lett. Math. Phys.* **114**(3), 73 (2024)
7. Mondino, A., Sämann, C.: Lorentzian Gromov-Hausdorff convergence and pre-compactness (2025)

8. Sakovich, A., Sormani, C.: Introducing various notions of distances between space-times (2025)
9. Sormani, C., Vega, C.: Big bang spacetimes. unpublished ideas (2014)
10. Sormani, C., Vega, C.: Null distance on a spacetime. *Class. Quantum Gravity* **33**(8), 085001 (2016)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.