

Is There a Role for Mifamurtide in Nonmetastatic High-Grade Osteosarcoma? Results From the Italian Sarcoma Group (ISG/OS-2) and Spanish Sarcoma Group (GEIS-33) Trials

Emanuela Palmerini, MD, PhD^{1,2} ; Cristina Meazza, MD³; Angela Tamburini, MD⁴; Catalina Márquez-Vega, MD⁵ ; Gianni Bisogno, MD⁶ ; Franca Fagioli, MD⁷ ; Virginia Ferraresi, MD⁸; Giuseppe Maria Milano, MD⁹; Luca Coccoli, MD¹⁰ ; Alba Rubio-San-Simón, MD¹¹ ; Oscar Gallego, MD¹² ; María Esther Llinares Riestra, MD¹³; Carla Manzitti, MD¹⁴; Jaume Mora, MD¹⁵; M^a Ángeles Vaz-Salgado, MD¹⁶; Roberto Luksch, MD³; Cristina Mata, MD¹⁷ ; Michela Pierini, BSc¹; Elisa Carretta, BSc¹⁸ ; Marilena Cesari, MD¹ ; Anna Paioli, MD¹ ; Andrea Marrari, MD¹; Katia Scotlandi, PhD, BSc¹⁹ ; Massimo Serra, BSc¹; Sebastian Dorin Asafei, MD⁷ ; Marco Gambarotti, MD²⁰ ; Piero Picci, MD, PhD¹⁹ ; Stefano Ferrari, MD¹; Claudia Valverde, MD²¹; Toni Ibrahim, MD, PhD¹ ; and Javier Martín Broto, MD^{22,23,24} 

DOI: <https://doi.org/10.1200/JCO-25-00210>

ABSTRACT

PURPOSE Outcome of patients with localized osteosarcoma is challenging. The role of mifamurtide is still a matter of debate. Two prospective trials were carried out in Italy (ISG/OS-2) and Spain (GEIS-33) with mifamurtide in ABCB1/P-glycoprotein (Pgp)-positive patients.

PATIENTS AND METHODS Patients age ≤40 years with localized extremity high-grade osteosarcoma were eligible. Analysis of Pgp expression from diagnostic biopsy was centralized. Patients received two cycles of preoperative methotrexate, doxorubicin, and cisplatin (MAP) before surgery. Postoperatively, in case of Pgp overexpression (Pgp-positive), mifamurtide was added, combined with doxorubicin (one cycle) and four consecutive cycles of high-dose ifosfamide (HDIFO) for patients with poor histologic response, or with MAP in case of good response. Patients who were Pgp-negative received MAP postoperatively. We present the merged analysis of ISG/OS-2 and GEIS-33 trial, an observational study with same inclusion criteria and treatment of ISG/OS-2. The primary endpoint was 5-year event-free survival (EFS) according to the use of mifamurtide. Secondary endpoint was overall survival (OS).

RESULTS From March 2013 to April 2018, 398 patients were analyzed. The median age was 14 years (range, 4–40), male/female: 238/160 (1.48/1.0); 211 of 398 (53%) tumors were Pgp-positive, and 204 of 398 (51.3%) patients received mifamurtide. With a median follow-up of 70 months (IQR, 49–90 months), the 5-year EFS and OS were 65.2% (95% CI, 60.1 to 69.8) and 74.8% (95% CI, 69.8 to 79.0), respectively, with superior EFS for patients undergoing mifamurtide and chemotherapy as compared with EFS of patients undergoing chemotherapy alone (5-year EFS 71.4% v 58.3%; $P = .0139$) not confirmed at multivariable analysis ($P = .0593$).

CONCLUSION In this merged analysis with a risk-adapted strategy for nonmetastatic osteosarcoma, the group with unfavorable prognoses, identified by Pgp expression, performed well when mifamurtide, combined with HDIFO in case of poor response, was administered after surgery.

ACCOMPANYING CONTENT

-  Appendix
-  Data Sharing Statement
-  Data Supplement
-  Protocol

Accepted June 13, 2025

Published August 18, 2025

J Clin Oncol 43:3113-3122

© 2025 by American Society of Clinical Oncology



[View Online Article](#)

Creative Commons Attribution
Non-Commercial No Derivatives
4.0 License

INTRODUCTION

High-grade osteosarcoma is the most frequent primary malignant bone tumor among children and adolescents and young adults (AYAs).^{1–3} In the past 50 years, survival has not improved, and novel therapeutic strategies are urgently needed.^{4,5}

Histologic response to neoadjuvant chemotherapy according to Huvo's has consistently shown a strong correlation with survival in patients with localized osteosarcoma.⁶ In addition, high levels of ABCB1/P-glycoprotein (Pgp), an efflux pump that reduces the intracellular concentration of doxorubicin, demonstrated an unfavorable prognostic factor in osteosarcoma in several published series, but controversy exists on its

CONTEXT

Key Objective

Is there a role for mifamurtide in patients with nonmetastatic extremity high-grade osteosarcoma?

Knowledge Generated

This analysis on about 400 patients with nonmetastatic osteosarcoma treated within two European prospective trials shows that high-risk patients, here defined by expression of P-glycoprotein, particularly in case of poor response to adjuvant chemotherapy, when ifosfamide was added, might benefit most from adjuvant mifamurtide.

Relevance (R.G. Maki)

Although it is unavailable in the United States, the post-chemotherapy period appears to be an opportune moment to impose other novel immunotherapeutic strategies in this patient population.*

*Relevance section written by JCO Associate Editor Robert G. Maki, MD, PhD, FACP, FASCO.

prognostic role.⁷⁻²⁴ Pgp mediates drug efflux through the plasma membrane and increases drug trapping within lysosomal drug safe houses^{13,20,24,25} (Appendix Fig A1, online only).

The fine interplay between tumor and immune cells is not fully understood, but M2-polarized macrophages have shown enhanced expression of Pgp.²⁶ It can be hypothesized that mifamurtide reverses the negative impact of M2 macrophage polarization in tumors overexpressing Pgp. Besides, COX2+ tumor-associated macrophages promote cancer cell proliferation and survival through different mechanisms, including by increasing Pgp.²⁷

Mifamurtide is a nonspecific immunomodulator binding to the extracellular Toll-like receptor-4 thus activating monocytes and macrophages, promoting antitumor activity.^{4,28,29} In 2009, the European Medicines Agency granted a centralized marketing authorization for muramyltripeptide (mifamurtide; MTP) for patients age between 2 and 30 years presenting a high-grade nonmetastatic resected osteosarcoma (EU/1/08/502/001). However, it is unclear which patients could potentially benefit from mifamurtide the most, and no approval has been obtained from the Food and Drug Administration. Moreover, in several European countries, the use of mifamurtide is still limited as the results of the INT-0133 trial have been a matter of debate between key-opinion leaders.²⁹

The ISG/OS-2 and GEIS-33 studies are prospective trials evaluating the role of a risk-modulated approach in patients with localized high-grade osteosarcoma of the extremities, stratifying patients on the basis of Pgp expression and tumor response to neoadjuvant chemotherapy.^{30,31}

The results of the ISG/OS2 study were recently published,³⁰ showing that adjuvant mifamurtide, combined with high-dose ifosfamide (HDIFO) in case of poor responder (PR) to

induction chemotherapy, could improve event-free survival (EFS) in Pgp-positive patients. This clinical trial also reported improved survival in comparison with previous studies.^{10,32}

This study is a secondary analysis of patients included in both ISG/OS-2 and GEIS-33 trial. All patients received methotrexate, doxorubicin, and cisplatin (MAP) for induction chemotherapy, but those with Pgp expression also received adjuvant mifamurtide. Moreover, in Pgp-positive patients with poor response to preoperative treatment, mifamurtide was combined with a cycle of doxorubicin + four consecutive cycles of HDIFO. Pgp expression might be measured by RNA and immunohistochemistry. Because differences in immunohistochemical Pgp assessment techniques might jeopardize the interpretation of results, Pgp assessment was performed in one institution for all of the patients.

The aim of this study was to analyze the outcome of patients in both studies, thus addressing the role of mifamurtide in osteosarcoma treatment in a larger series of patients.

PATIENTS AND METHODS

Study Design

This study was carried out on patients treated within the ISG/OS-2 and GEIS-33. ISG/OS2 (ClinicalTrials.gov identifier: [NCT01459484](#)) is an Italian, multicenter, uncontrolled phase II trial. GEIS-33 (ClinicalTrials.gov identifier: [NCT04383288](#)) is a Spanish phase IV, postauthorization, observational, multicenter study in patients age between 2 and 30 years, diagnosed with nonmetastatic high-grade osteosarcoma of the extremities.

For both trials, data collection was performed on electronic case report form and centralized.

Both studies evaluated the efficacy of risk-adapted chemotherapy regimens, with adjuvant mifamurtide and HDIFO in Pgp-positive patients PR to induction chemotherapy. A written informed consent was obtained from the adult patients or from the guardians in case of pediatric patients. The local Ethics Committees approved the protocols.

Patients' inclusion and exclusion criteria were presented elsewhere.³⁰ Main inclusion criteria were a diagnosis of primary, central, high-grade localized osteosarcoma of the extremities; age ≤ 40 years; and no prior surgery or chemotherapy for osteosarcoma; main exclusion criterion was metastatic disease at diagnosis (Data Supplement, online only).

Immunohistochemical Detection of Pgp

For ISG/OS2 and GEIS-33 studies, assessment of Pgp was centralized at the Istituto Rizzoli and the expression of Pgp was evaluated by immunohistochemistry as described.^{7,33}

Briefly, only formalin-fixed, paraffin-embedded biopsies were used for immunohistochemical analysis of Pgp. Expression of Pgp was assessed with three monoclonal antibodies, which react with different, mutually exclusive epitopes of this protein: JSB-1 (Sanbio, Uden, the Netherlands), MRK16 (Kamiya Biomedical, Thousand Oaks, CA), and C494 (Signet Laboratories, Dedham, MA). Immunohistochemistry was performed by using an avidin-biotin

peroxidase complex method (Vectastain ABC kit, Vector Laboratories, Burlingame, CA), and the final reaction product was revealed by incubation with diaminobenzidine (Sigma, St Louis, MO). For each specimen, a negative control was carried out by replacing the primary antibody with normal horse serum, whereas to check the antigenicity of the sample, one additional section was incubated with the V9 anti-vimentin monoclonal antibody (Roche Molecular Biochemicals, Mannheim, Germany). Sections of normal human kidney were used as reference control for Pgp immunostaining procedure because of its reported overexpression in proximal tubuli. Only specimens with a diffused immunostaining for Pgp were classified as positive.

From here onward, because the Pgp+ group is superimposable with the mifamurtide-receiving group, in this analysis, we will only refer to the Pgp+ group as the mifamurtide group.

Study Description

Methotrexate is given once a day, 4 hour infusion, while adriamycin and cisplatin are administered as a continuous infusion. Frequency of administration is illustrated in Figure 1. After surgery, chemotherapy-induced necrosis was expressed as a percentage. When the percentage of tumor necrosis was 90% or more, patients were classified as good responders (GRs); when the percentage of tumor necrosis was lower, patients were defined as PRs. In




FIG 1. Treatment scheme of the ISG/OS-2 study by Pgp expression. HDIFO, high-dose ifosfamide; Pgp, P-glycoprotein.

Pgp-positive patients, postoperative chemotherapy consisted of MAP and mifamurtide for GRs or one cycle of A 90 mg/m² intravenous (IV) continuous infusion (24h), every 3 weeks followed by four consecutive cycles of HDIFO (3 g/m²/d, day 1–5, continuous infusion) and mifamurtide (2 mg/m²) for PRs. MTP was administered twice a week for 3 months and then weekly for 6 months at the dose of 2 mg/m² (total duration 48 weeks). Patients who were Pgp-negative continued MAP postoperatively, regardless of the response to induction chemotherapy. The total cumulative dose of A was 420 mg/m² (75 mg/m²/day × two cycles in the neoadjuvant setting, combined with P, plus

90 mg/m² × three cycles when administered in monotherapy, in the adjuvant setting).

Statistical Analysis

Continuous data were summarized as median with interquartile range and categorical data as frequencies with percentage. Alkaline phosphatase (ALP) and lactate dehydrogenase levels were classified as high when they exceed the upper limit of the normal range according to laboratories ranges, whereas levels within the ranges are deemed low.

TABLE 1. Baseline Demographic and Clinical Characteristics According to Mifamurtide Treatment

Characteristic	All Sample (N = 398), No. (%)	Mifamurtide Yes (n = 204), No. (%)	Mifamurtide No (n = 194), No. (%)	P
Age, years				
0-14	218 (54.8)	126 (61.8)	92 (47.4)	.0041
15-40	180 (45.2)	78 (38.2)	102 (52.6)	
Sex				
Male	238 (59.8)	117 (57.4)	121 (62.4)	.3074
Female	160 (40.2)	87 (42.6)	73 (37.6)	
Serum ALP				
High	133 (33.4)	62 (30.4)	71 (36.6)	.0288
Normal	233 (58.6)	131 (64.2)	102 (52.6)	
Unknown	32 (8.0)	11 (5.4)	21 (10.8)	
LDH				
Normal	258 (64.8)	134 (65.7)	124 (63.9)	.5080
High	115 (28.9)	55 (27.0)	60 (30.9)	
Unknown	25 (6.3)	15 (7.3)	10 (5.2)	
Histologic response				
GR	164 (41.2)	87 (42.6)	77 (39.7)	.4301
PR	227 (57.0)	115 (56.4)	112 (57.7)	
Unknown	7 (1.8)	2 (1.0)	5 (2.6)	
Surgery				
Resection	375 (94.2)	195 (95.6)	180 (92.8)	.3495
Amputation	20 (5.0)	9 (4.4)	11 (5.7)	
Rotationplasty	1 (0.3)		1 (0.5)	
Unknown	2 (0.5)		2 (1.0)	
Margins				
Adequate	361 (90.7)	192 (94.1)	169 (87.1)	.0181
Inadequate	34 (8.5)	12 (5.9)	22 (11.3)	
Unknown	3 (0.8)		3 (1.6)	
Histotype				
Osteoblastic	287 (72.1)	147 (72.1)	140 (72.2)	.8652
Other	110 (27.6)	57 (27.9)	53 (27.3)	
Chondroblastic	45 (40.9)	19 (33.3)	26 (49.0)	
Fibroblastic	21 (19.1)	14 (24.6)	7 (13.2)	
Hemorrhagic	21 (19.1)	14 (24.6)	7 (13.2)	
Small cell	16 (14.5)	6 (10.5)	10 (18.9)	
Unspecified	7 (6.4)	4 (7.0)	3 (5.7)	
Unknown	1 (0.3)		1 (0.5)	

Abbreviations: ALP, alkaline phosphatase; GR, good response; LDH, lactate dehydrogenase; PR, poor response.

Baseline characteristics according to mifamurtide treatment were compared using the chi-square test or Fisher exact test.

EFS was calculated from the first day of chemotherapy to recurrence (local or distant), death from all causes, the appearance of secondary tumors, or the last follow-up examination. Overall survival (OS) was calculated from the first day of chemotherapy to death or the last follow-up examination. Survival curves were calculated using the Kaplan-Meier method and compared using the log-rank test. Univariable and multivariable hazard ratios with 95% CIs were estimated using the Cox proportional hazards model. The multivariable model was computed including covariates statistically significant in univariable analysis. The proportional hazards assumption was tested using Schoenfeld residuals.

Tests were performed two-sided, and results were reported with 95% CI. We considered a $P \leq .05$ as statistically significant. Statistical analyses were performed using SAS 9.4 (SAS Institute Inc, Cary, NC).

RESULTS

A total of $N = 401$ patients (from ISG/OS $n = 279$, and from GEIS $n = 122$) were included in this study. Three patients did not have survival data and therefore were excluded (Appendix Fig A2). Overall, 398 patients were analyzed: 204 (51.3%) treated with mifamurtide and 194 not treated with mifamurtide (48.7%).

The median age was 14 years (range, 4-40). Clinical characteristics according to mifamurtide use are reported in Table 1. As shown in Table 1, there was an imbalance at




FIG 2. Probability of EFS for (A) the entire cohort and (B) by mifamurtide administration. Probability of OS for (C) the entire cohort and (D) by mifamurtide administration. EFS, event-free survival; OS, overall survival.

baseline for some clinical potential prognostic factors favoring the mifamurtide arm. Toxicities and compliance to therapy were reported elsewhere.^{30,31}

Outcomes

EFS

Overall, the median follow-up was 70 months (IQR, 49–90 months). The 5-year EFS was 65.2% (95% CI, 60.1 to 69.8; Fig 2A).

Five-year EFS was 71.4% (95% CI, 64.5 to 77.1) for patients undergoing mifamurtide and MAP (HDIFO in case of PR) and 58.3% (95% CI, 50.6 to 65.3) for patients undergoing MAP only ($P = .0131$; Table 2; Fig 2B). This difference was also significant in the subgroup of patients with poor response ($P = .0359$), whereas no significant difference in the GR patients was detected (Table 3; Appendix Fig A3).

Poor histological response to induction chemotherapy and high baseline ALP levels were associated with significant inferior EFS (Table 2), whereas age ≥ 15 years was not a negative prognostic factor in this series.

In the multivariable analysis, poor response to induction chemotherapy and high baseline ALP level were

both confirmed significant prognostic factors for EFS (Table 4).

OS

The 5-year OS was 74.8% overall (95% CI, 69.8 to 79.0; Figs 2A and 2C). The 5-year OS was 65.8% (95% CI, 57.8 to 72.6) for MAP only and 75.4% (95% CI, 68.4 to 81.1) in case of chemotherapy plus mifamurtide ($P = .0457$; Table 2; Figs 2B and 2D). Poor histologic response to induction chemotherapy and high ALP levels at diagnosis were associated with significant inferior OS (Table 2).

In the multivariable analysis, poor response to induction chemotherapy and high baseline ALP level were both confirmed significant prognostic factors for OS (Table 4).

Survival Analysis on the Basis of Histologic Necrosis and Pgp

A subgroup of EFS and OS analysis on the basis of patients stratification according to histologic response in the Pgp– and Pgp+ subgroups was performed (Appendix Fig A4): Chemotherapy-induced necrosis after neoadjuvant chemotherapy was confirmed as a strong prognostic factor, in both Pgp– and Pgp+ subgroups, supporting the evidence reported in Table 2.

TABLE 2. Univariable Analysis of EFS and OS in Patients With Nonmetastatic Extremity Osteosarcoma Treated in ISG/OS-2 and GEIS-33 Trials

Characteristic	5-Year EFS, % (95% CI)	HR (95% CI)	<i>P</i>	5-Year OS, % (95% CI)	HR (95% CI)	<i>P</i>
Age, years						
0–14	66.8 (60.0 to 72.8)			74.3 (67.4 to 79.9)		
15–40	63.3 (55.4 to 70.2)	1.0 (0.7 to 1.5)	.8650	75.4 (67.9 to 81.5)	0.9 (0.6 to 1.4)	.6233
Sex						
Male	65.8 (59.1 to 71.7)			76.0 (69.5 to 81.3)		
Female	64.3 (56.2 to 71.4)	1.0 (0.7 to 1.5)	.7972	73.2 (65.0 to 79.7)	1.1 (0.7 to 1.7)	.6423
Serum ALP						
Normal	70.6 (64.0 to 76.2)			80.4 (74.2 to 85.3)		
High	56.3 (47.2 to 64.4)	1.8 (1.2 to 2.5)	.0017	63.7 (54.5 to 71.6)	2.1 (1.3 to 3.2)	.0008
LDH						
Normal	65.8 (59.4 to 71.5)			75.2 (69.0 to 80.3)		
High	62.6 (52.8 to 71.0)	1.2 (0.8 to 1.7)	.4168	69.9 (59.9 to 77.9)	1.2 (0.8 to 1.9)	.3286
Histologic response						
GR	79.0 (71.7 to 84.6)			85.8 (79.0 to 90.6)		
PR	54.9 (47.9 to 61.4)	2.5 (1.7 to 3.7)	<.0001	67.1 (60.0 to 73.2)	2.5 (1.5 to 4.1)	.0002
Mifamurtide						
Yes	71.4 (64.5 to 77.1)			75.4 (68.4 to 81.1)		
No	58.3 (50.6 to 65.3)	1.5 (1.1 to 2.2)	.0139	65.8 (57.8 to 72.6)	1.5 (1.0 to 2.3)	.0457
Histotype						
Osteoblastic	66.2 (60.2 to 71.5)			75.8 (70.1 to 80.7)		
Other	62.5 (52.2 to 71.1)	1.1 (0.8 to 1.7)	.4721	71.9 (61.5 to 78.0)	1.2 (0.8 to 1.9)	.4755

Abbreviations: ALP, alkaline phosphatase; EFS, event-free survival; GR, good response; HR, hazard ratio; LDH, lactate dehydrogenase; OS, overall survival; PR, poor response.

TABLE 3. Univariable EFS According to Histologic Response to Induction Chemotherapy and Mifamurtide

Characteristic	No.	5-Year EFS, % (95% CI)	HR (95% CI)	P
Good responder				
MAP + mifamurtide (Pgp+)	87	83.5 (73.7 to 89.9)		.1238
MAP (Pgp-)	77	73.5 (61.5 to 82.3)	1.7 (0.9 to 3.4)	
Poor responder				
MAP + HDIFO + mifamurtide (Pgp+)	115	62.5 (52.7 to 70.8)		.0454
MAP (Pgp-)	112	47.0 (36.9 to 56.4)	1.5 (1.0 to 2.2)	

NOTE. Seven patients with unknown necrosis were excluded from this analysis.

Abbreviations: EFS, event-free survival; HDIFO, high-dose ifosfamide; HR, hazard ratio; MAP, methotrexate, doxorubicin, and cisplatin; Pgp, P-glycoprotein.

DISCUSSION

This study demonstrated a significant superior EFS and OS in patients undergoing mifamurtide as compared with chemotherapy alone at univariable analysis. Despite the unbalance for age and ALP, the multivariable analysis showed that among these factors, only ALP had an independent prognostic role, whereas the *P* value for mifamurtide administration was .059.

Our study suggests that the anticipated poor prognosis for patients with Pgp+ tumors may have been mitigated by mifamurtide. Despite conflicting evidence regarding the prognostic effect of Pgp expression, we maintain that it holds significant prognostic value.

We believe that there are two different ways to interpret these findings, according with the potential role of Pgp as a prognostic factor.

If we assume that Pgp selects two groups with different prognosis, these results demonstrate that mifamurtide was able to improve EFS in the high-risk patient group. Intriguingly, Pgp-positive patients with PR to neoadjuvant chemotherapy had a statistically significant EFS improvement with this experimental approach (mifamurtide and HDIFO postoperatively), as compared with the Pgp-negative PR cohort (MAP only), suggesting that Pgp prognostic value might be outplayed by an integrated treatment and reinforces the hypothesis that ifosfamide might act

synergistically with mifamurtide. It is plausible that ifosfamide and mifamurtide are potentially synergistic. Two decades ago, it was reported that ifosfamide could be an immunomodulator upregulating the expansion of Th1 cells.³⁴

Unfortunately, because the Pgp-negative population did not receive mifamurtide, our study cannot exclude the potential benefit of mifamurtide for this subgroup.

On the other hand, if we are skeptical on the prognostic role of Pgp, these analysis shows that, in a cohort of unselected patients with localized osteosarcoma age ≤40 years, a strategy based on mifamurtide and chemotherapy with HDIFO in case of PRs is superior to chemotherapy alone in terms of EFS.

Particularly, INT0133 did not find a correlation between Pgp expression and worse EFS or OS. However, Pgp expression analysis was possible in only 139 (for C494 antibody) of 685 patients (20.3%), using tissues from either diagnostic biopsies or surgical specimens, and the staining technique differed from ours.¹⁰

In both cases, these data would support, 17 years later, the findings of the controlled study by Meyer et al³⁵. The former controlled study from the Children's Oncology Group, INT0133 trial, using a randomized two-by-two factorial design, already hypothesized an interaction between mifamurtide and ifosfamide. This study reported a 3-year EFS of 68% in the group that received a combination of mifamurtide and MAP and 78% for those who received

TABLE 4. Multivariable Analysis of EFS and OS in Patients With Nonmetastatic Extremity Osteosarcoma Treated in ISG/OS-2 and GEIS-33 Trials

Characteristic	Variable	5-Year EFS		5-Year OS	
		HR (95% CI)	P	HR (95% CI)	P
Serum ALP	High v normal	1.9 (1.3 to 2.7)	.0006	2.0 (1.3 to 3.2)	.0013
Histologic response	PRs v GRs	2.5 (1.7 to 3.8)	<.0001	2.6 (1.6 to 4.3)	.0002
Mifamurtide	No v yes	1.4 (1.0 to 2.0)	.0593	1.4 (0.9 to 2.1)	.1480

NOTE. Models were performed on patients with completed data (n = 359).

Abbreviations: ALP, alkaline phosphatase; EFS, event-free survival; GRs, good responses; HR, hazard ratio; OS, overall survival; PRs, poor responses.

mifamurtide combined with MAP and ifosfamide (9 g/m² continuous infusion per cycle). In a follow-up study, mifamurtide was associated with an OS benefit and with a positive EFS trend.³⁵ On the basis of these results, concerns whether INT0133's results met generally accepted standards for practice-changing conclusions were raised.

Is the improvement observed in our study because of mifamurtide? Or is it the HDIFO salvage therapy in case of poor response? The EURAMOS study showed that the addition of ifosfamide (2.8 g/m² per day combined with etoposide 100 mg/m² per day, on days 1–5, every 21 days), to MAP does not improve survival, increases toxicity, and should not be considered as a salvage therapy for high-risk patients.^{36,37} A recent Japanese phase II study has come to a similar conclusion, failing to demonstrate a survival benefit for ifosfamide given to patients with poor response to MAP.³⁸ As the most important benefit in terms of EFS in this study was obtained for Pgp-positive patients with PR to induction chemotherapy, treated with both mifamurtide and HDIFO, a positive interaction between these drugs could be hypothesized, and this combination might be proposed as salvage treatment of patients with unresponsive osteosarcoma.

Also, our study confirmed that poor histologic response to induction chemotherapy and high ALP at diagnosis predict worse outcome, as confirmed at multivariable analysis.

Interestingly, no relationship was found between the level of Pgp expression and histologic response after preoperative chemotherapy, as previously reported.^{9,10} Most likely, the two parameters identify two phenomena related to resistance to chemotherapy, only partly overlapping. In fact, although poor histologic response might be the result of a cumulative resistance to all drugs administered preoperatively, Pgp mediates a specific resistance to doxorubicin and might also be related to other tumor features, such as differential biological aggressiveness of tumor cells.

We observed that survival for pediatric patients and AYA was not different when treated within the same protocol. This conflicts with other studies²⁹ and reinforce the suggestion of the AYA Working Group of the European Society for Medical Oncology and the European Society for Pediatric Oncology, that multicenter cooperation, including pediatric and adult cooperation, should be encouraged.³

This is a retrospective analysis of data collected prospectively. The major limitation of these clinical trials is the lack of a control group because of inherent difficulties to run a randomized trial in the adjuvant setting of a rare disease. Unfortunately, this is a limitation that will remain

unaddressed for now because the design of most of the trials in osteosarcoma is not controlled (Appendix Table A1). Results of the UNIFRA study, which will test the hypothesis that high-risk patients, defined by metastases at diagnosis or poor response to induction chemotherapy, might benefit from mifamurtide, are awaited.³⁹

An observational study is ongoing in Italy since 2021 (ISG/AIEOP OS2 Oss) to collect real-world data on osteosarcoma; on the basis of these findings, the recommended approach, as per national guidelines, is to use mifamurtide in patients younger than 30 years, because of regulatory constraints, with a PR after neoadjuvant chemotherapy, and in combination with HDIFO. Similarly in Spain, this study facilitated the prescription of mifamurtide within its framework, promoting its rational use. The results of this study will be critical in establishing the use of high-dose ifosfamide in conjunction with mifamurtide as a national protocol, especially for patients younger than 30 years exhibiting a poor pathologic response to induction therapy with MAP.

Mifamurtide use will also be allowed according to investigator preferences in the future European FOSTER-Cabos trial. This study will randomize maintenance cabozantinib in patients with localized and metastatic osteosarcoma nonprogressing after up-front chemotherapy. A run-in phase will also be undertaken to assess the feasibility of mifamurtide in combination with cabozantinib.

Identifying predictive factors to select patients and stratify treatment on the basis of their likelihood of responding to mifamurtide could lead to increased cost-effectiveness, particularly relevant considering reimbursement and cost saving concerns.

A translational genomic sequencing study on pretreatment sample, to identify subgroup of responders, is ongoing (ClinicalTrials.gov identifier: [NCT03737435](https://clinicaltrials.gov/ct2/show/study?term=NCT03737435)).

In conclusion, key finding from this study including almost 400 patients with nonmetastatic osteosarcoma is the potential role of HDIFO combined with mifamurtide as salvage treatment for Pgp-positive patients with poor histologic response to neoadjuvant chemotherapy. These results might pave the path for future treatment strategies in selected high-risk osteosarcoma groups. No specific indications on the generalized use of mifamurtide should be derived from this study. Nonetheless, the results of this trial, in the absence of a randomized controlled trial to further explore the role of mifamurtide, might have a prescriptive role for mifamurtide along with HDIFO in patients exhibiting a poor response to neoadjuvant treatment with MAP.

AFFILIATIONS

¹Osteoncology, Bone and Soft Tissue Sarcomas, and Innovative Therapies Unit, IRCCS Istituto Ortopedico Rizzoli, Bologna, Italy

²Sylvester Comprehensive Cancer Center, Miller School of Medicine, University of Miami, Miami, FL

³Pediatric Oncology Unit, Fondazione IRCCS Istituto Nazionale dei Tumori, Milano, Italy

⁴Department of Pediatric Onco-Hematology, AOU Meyer—IRCCS, Florence, Italy

⁵Pediatric Oncology Unit, Hospital Infantil Universitario Virgen del Rocío, Sevilla, Spain

⁶Department of Women's and Children's Health, University of Padova, Padova, Italy

⁷S.C. Oncoematologia Pediatrica—Regina Margherita Children's Hospital, A.O.U. Città della Salute e della Scienza di Torino, Torino, Italy

⁸Sarcomas and Rare Tumors Departmental Unit, IRCCS Regina Elena National Cancer Institute, Rome, Italy

⁹Division of Pediatric Hematology and Oncology, Gene and Cellular Therapy, IRCCS Ospedale Pediatrico Bambino Gesù, Roma, Italy

¹⁰Pediatric Onco-Hematology Unit, S. Chiara Hospital, AOUP, Pisa, Italy

¹¹Paediatric Haematology-Oncology Department, Children's University Hospital Infantil Niño Jesús, Madrid, Spain

¹²Hospital de la Santa Creu i Sant Pau, Barcelona, Spain

¹³Pediatric Oncology Unit, Hospital Infantil Universitario Clínico Virgen de la Arrixca, Murcia, Spain

¹⁴IRCCS Istituto Giannina Gaslini, Genova, Italy

¹⁵Hospital Sant Joan de Déu, Barcelona, Spain

¹⁶Hospital Universitario Ramón y Cajal, Irycis, Madrid, Spain

¹⁷Pediatric and Adolescent Oncohematology Unit, Pediatric Sarcomas National Reference Center, Hospital General Universitario Gregorio Marañón, Madrid, Spain

¹⁸Department of Programming and Monitoring, IRCCS Istituto Ortopedico Rizzoli, Bologna, Italy

¹⁹Laboratory of Experimental Oncology, IRCCS Istituto Ortopedico Rizzoli, Bologna, Italy

²⁰Department of Pathology, IRCCS Istituto Ortopedico Rizzoli, Bologna, Italy

²¹Department of Medical Oncology, Hospital Universitari Vall d'Hebron, Vall d'Hebrón Institute of Oncology, Barcelona, Spain

²²Research Health Institute of Fundación Jiménez Díaz (IIS/FJD; UAM), Madrid, Spain

²³Department of Medical Oncology, Fundación Jiménez Díaz University Hospital, Madrid, Spain

²⁴University Hospital General de Villalba, Madrid, Spain

CORRESPONDING AUTHOR

Emanuela Palmerini, MD, PhD; Social Media Handle: @ManuPalmerini; e-mail: emanuela.palmerini@ior.it.

EQUAL CONTRIBUTION

T.I. and J.M.B. equally contributed.

PRIOR PRESENTATION

Presented at the Connective Tissue Oncology Society (CTOS) Annual Meeting, San Diego, CA, November 13-16, 2024.

SUPPORT

Supported by the Associazione Onlus Il Pensatore; Matteo Amitrano; the Associazione Mario Campanacci; the Carisbo Foundation Call for

REFERENCES

- Gitelis S: Bone and soft tissue tumors: Clinical features, imaging, pathology and treatment. 2nd ed. J Bone Joint Surg 83:801-802, 2001
- Baumhoer D, Böhlring TO, Cates JMM, et al: Osteosarcoma, in WHO Classification of Tumours Editorial Board (ed): Soft Tissue and Bone Tumours, Volume 3 (ed 5). Lyon, France, International Agency for Research on Cancer, 2020, pp 403-409
- Ferrari A, Stark D, Peccatori FA, et al: Adolescents and young adults (AYA) with cancer: A position paper from the AYA Working Group of the European Society for Medical Oncology (ESMO) and the European Society for Paediatric Oncology (SIOPE). ESMO Open 6:100096, 2021
- Bielack SS, Hecker-Nolting S, Blattmann C, et al: Advances in the management of osteosarcoma. F1000Res 5:2767, 2016
- Reed DR, Tulpule A, Metts J, et al: Pediatric leukemia roadmaps are a guide for positive metastatic bone sarcoma trials. J Clin Oncol 42:2955-2960, 2024

Translational and Clinical Medical Research; the 5 x mille contributions to the Istituto Ortopedico Rizzoli.

CLINICAL TRIAL INFORMATION

[NCT01459484](#); [NCT04383288](#)

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO-25-00210>.

DATA SHARING STATEMENT

A data sharing statement provided by the authors is available with this article at DOI <https://doi.org/10.1200/JCO-25-00210>.

AUTHOR CONTRIBUTIONS

Conception and design: Emanuela Palmerini, Angela Tamburini, Gianni Bisogno, Luca Coccoli, Roberto Luksch, Katia Scotlandi, Massimo Serra, Piero Picci, Stefano Ferrari

Provision of study materials or patients: Emanuela Palmerini, Angela Tamburini, Gianni Bisogno, Franca Fagioli, Luca Coccoli, Alba Rubio-San-Simón, Jaume Mora, M^a Ángeles Vaz-Salgado, Marilena Cesari, Andrea Marrari, Piero Picci, Stefano Ferrari, Claudia Valverde, Javier Martín Broto

Collection and assembly of data: Emanuela Palmerini, Catalina Márquez-Vega, Gianni Bisogno, Virginia Ferraresi, Giuseppe Maria Milano, Luca Coccoli, Alba Rubio-San-Simón, Oscar Gallego, María Esther Llinares Riestra, Carla Manzitti, Jaume Mora, M^a Ángeles Vaz-Salgado, Cristina Mata, Michela Pierini, Anna Paioli, Andrea Marrari, Massimo Serra, Sebastian Dorin Asaftei, Marco Garbarotti, Stefano Ferrari, Claudia Valverde, Javier Martín Broto

Data analysis and interpretation: Emanuela Palmerini, Cristina Meazza, Gianni Bisogno, Franca Fagioli, Virginia Ferraresi, Luca Coccoli, Oscar Gallego, Roberto Luksch, Elisa Carretta, Marilena Cesari, Anna Paioli, Andrea Marrari, Massimo Serra, Marco Garbarotti, Stefano Ferrari, Toni Ibrahim, Javier Martín Broto

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

ACKNOWLEDGMENT

The authors thank the participating patients and their family and the nurses. The authors also thank Michela Pierini, Valquiria Broll, and Antonio Carella for all support and editorial assistance; Stefano Zagnoni, Federica Sella, Francesca Michelacci, Marilù Fanelli, Serena Vella, Camilla Cristalli, Claudia Hattinger, Silvia Luppi, and Federica Magagnoli for immunohistochemical studies; and Maria Pia Cumani and Cristina Ghinelli for their help with the graphic design.

6. Strauss SJ, Frezza AM, Abecassis N, et al: Bone sarcomas: ESMO-EURACAN-GENTURIS-ERN PaedCan Clinical Practice Guideline for diagnosis, treatment and follow-up. *Ann Oncol* 32:1520-1536, 2021
7. Serra M, Scotlandi K, Reverter-Branchat G, et al: Value of P-glycoprotein and clinicopathologic factors as the basis for new treatment strategies in high-grade osteosarcoma of the extremities. *J Clin Oncol* 21:536-542, 2003
8. Zhao ZG, Ding F, Liu M, et al: Association between P-glycoprotein expression and response to chemotherapy in patients with osteosarcoma: A systematic and meta-analysis. *J Cancer Res Ther* 10: C206-C209, 2014
9. Pakos EE, Ioannidis JPA: The association of P-glycoprotein with response to chemotherapy and clinical outcome in patients with osteosarcoma: A meta-analysis. *Cancer* 98:581-589, 2003
10. Serra M, Picci P, Ferrari S, et al: Prognostic value of P-glycoprotein in high-grade osteosarcoma. *J Clin Oncol* 25:4858-4861, 2007
11. Liu T, Li Z, Zhang Q, et al: Targeting ABCB1 (MDR1) in multi-drug resistant osteosarcoma cells using the CRISPR-Cas9 system to reverse drug resistance. *Oncotarget* 7:83502-83513, 2016
12. Schwartz CL, Gorlick R, Teot L, et al: Multiple drug resistance in osteogenic sarcoma: INT0133 from the Children's Oncology Group. *J Clin Oncol* 25:2057-2062, 2007
13. Seebacher NA, Richardson DR, Jansson PJ: A mechanism for overcoming P-glycoprotein-mediated drug resistance: Novel combination therapy that releases stored doxorubicin from lysosomes via lysosomal permeabilization using Dp44mT or DpC. *Cell Death Dis* 7:e2510, 2016
14. Serra M, Pasello M, Manara MC, et al: May P-glycoprotein status be used to stratify high-grade osteosarcoma patients? Results from the Italian/Scandinavian Sarcoma Group 1 treatment protocol. *Int J Oncol* 29:1459-1468, 2006
15. Park YB, Kim HS, Oh JH, et al: The co-expression of p53 protein and P-glycoprotein is correlated to a poor prognosis in osteosarcoma. *Int Orthop* 24:307-310, 2001
16. Hornicek FJ, Gebhardt MC, Wolfe MW, et al: P-glycoprotein levels predict poor outcome in patients with osteosarcoma. *Clin Orthop Relat Res* 373:11-17, 2000
17. Baldini N, Scotlandi K, Serra M, et al: P-glycoprotein expression in osteosarcoma: A basis for risk-adapted adjuvant chemotherapy. *J Orthop Res* 17:629-632, 1999
18. Gorlick R, Huvos AG, Heller G, et al: Expression of HER2/erbB-2 correlates with survival in osteosarcoma. *J Clin Oncol* 17:2781-2788, 1999
19. Serra M, Maurici D, Scotlandi K, et al: Relationship between P-glycoprotein expression and p53 status in high-grade osteosarcoma. *Int J Oncol* 14:301-307, 1999
20. Shnyder SD, Hayes AJ, Pringle J, et al: P-glycoprotein and metallothionein expression and resistance to chemotherapy in osteosarcoma. *Br J Cancer* 78:757-759, 1998
21. Chan HS, Grogan TM, Haddad G, et al: P-glycoprotein expression: Critical determinant in the response to osteosarcoma chemotherapy. *J Natl Cancer Inst* 89:1706-1715, 1997
22. Radig K, Häckel C, Herting J, et al: Expression of P-glycoprotein in high grade osteosarcomas with special emphasis on chondroblastic subtype. *Gen Diagn Pathol* 142:139-145, 1997
23. Baldini N, Scotlandi K, Barbanti-Bròdano G, et al: Expression of P-glycoprotein in high-grade osteosarcomas in relation to clinical outcome. *N Engl J Med* 333:1380-1385, 1995
24. Bodey B, Taylor CR, Siegel SE, et al: Immunocytochemical observation of multidrug resistance (MDR) p170 glycoprotein expression in human osteosarcoma cells. The clinical significance of MDR protein overexpression. *Anticancer Res* 15:2461-2468, 1995
25. Serra M, Scotlandi K, Manara MC, et al: Analysis of P-glycoprotein expression in osteosarcoma. *Eur J Cancer* 31A:1998-2002, 1995
26. Robinson K, Tiriveedhi V: Perplexing role of P-glycoprotein in tumor microenvironment. *Front Oncol* 10:265, 2020
27. Li H, Yang B, Huang J, et al: Cyclooxygenase-2 in tumor-associated macrophages promotes breast cancer cell survival by triggering a positive-feedback loop between macrophages and cancer cells. *Oncotarget* 6:29637-29650, 2015
28. Kleinerman ES, Jia SF, Griffin J, et al: Phase II study of liposomal muramyl tripeptide in osteosarcoma: The cytokine cascade and monocyte activation following administration. *J Clin Oncol* 10: 1310-1316, 1992
29. Meyers PA, Schwartz CL, Krailo M, et al: Osteosarcoma: A randomized, prospective trial of the addition of ifosfamide and/or muramyl tripeptide to cisplatin, doxorubicin, and high-dose methotrexate. *J Clin Oncol* 23:2004-2011, 2005
30. Palmerini E, Meazza C, Tamburini A, et al: Phase 2 study for nonmetastatic extremity high-grade osteosarcoma in pediatric and adolescent and young adult patients with a risk-adapted strategy based on ABCB1/P-glycoprotein expression: An Italian Sarcoma Group trial (ISG/OS-2). *Cancer* 128:1958-1966, 2022
31. Salazar J, Arranz MJ, Martin-Broto J, et al: Pharmacogenetics of neoadjuvant MAP chemotherapy in localized osteosarcoma: A study based on data from the GEIS-33 protocol. *Pharmaceutics* 16: 1585, 2024
32. Smeland S, Bielack SS, Whelan J, et al: Survival and prognosis with osteosarcoma: Outcomes in more than 2000 patients in the EURAMOS-1 (European and American Osteosarcoma Study) cohort. *Eur J Cancer* 109:36-50, 2019
33. van der Valk P, van Kalken CK, Ketelaars H, et al: Distribution of multi-drug resistance-associated P-glycoprotein in normal and neoplastic human tissues: Analysis with 3 monoclonal antibodies recognizing different epitopes of the P-glycoprotein molecule. *Ann Oncol* 1:56-64, 1990
34. Binotto G, Trentin L, Semenzato G: Ifosfamide and cyclophosphamide: Effects on immunosurveillance. *Oncology* 65:17-20, 2003 (suppl 2)
35. Meyers PA, Schwartz CL, Krailo MD, et al: Osteosarcoma: The addition of muramyl tripeptide to chemotherapy improves overall survival—A report from the Children's Oncology Group. *J Clin Oncol* 26:633-638, 2008
36. Marina NM, Smeland S, Bielack SS, et al: Comparison of MAPIE versus MAP in patients with a poor response to preoperative chemotherapy for newly diagnosed high-grade osteosarcoma (EURAMOS-1): An open-label, international, randomised controlled trial. *Lancet Oncol* 17:1396-1408, 2016
37. Bielack SS, Werner M, Tunn PU, et al: Methotrexate, doxorubicin, and cisplatin (MAP) plus maintenance pegylated interferon alfa-2b versus MAP alone in patients with resectable high-grade osteosarcoma and good histologic response to preoperative MAP: First results of the EURAMOS-1 good response randomized controlled trial. *J Clin Oncol* 33:2279-2287, 2015
38. Hiraga H, Machida R, Kawai A, et al: Methotrexate, doxorubicin, and cisplatin versus methotrexate, doxorubicin, and cisplatin + ifosfamide in poor responders to preoperative chemotherapy for newly diagnosed high-grade osteosarcoma (JCOG0905): A multicenter, open-label, randomized trial. *J Clin Oncol* 43:1886-1897, 2025
39. Brard C, Piperno-Neumann S, Delaye J, et al: Sarcome-13/OS2016 trial protocol: A multicentre, randomised, open-label, phase II trial of mifamurtide combined with postoperative chemotherapy for patients with newly diagnosed high-risk osteosarcoma. *BMJ Open* 9:e025877, 2019

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Is There a Role for Mifamurtide in Nonmetastatic High-Grade Osteosarcoma? Results From the Italian Sarcoma Group (ISG/OS-2) and Spanish Sarcoma Group (GEIS-33) Trials

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/authors/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](#)).

Emanuela Palmerini

Consulting or Advisory Role: Daiichy Sankyo, Deciphera Pharmaceuticals, Eusa Pharma, SynOx Therapeutics, Ipsen Biopharmaceuticals and Servier
Travel, Accommodations, Expenses: Takeda

Gianni Bisogno

Consulting or Advisory Role: Bayer

Franca Fagioli

Research Funding: Jazz Pharmaceuticals (Inst), Novartis (Inst), EUSA Pharma (Inst), Gilead Sciences (Inst), Takeda (Inst), Medac Pharma (Inst), Merck (Inst), Novimmune (Inst), Roche (Inst)

Virginia Ferraresi

Consulting or Advisory Role: Bristol Myers Squibb, Novartis, GlaxoSmithKline, Serb
Speakers' Bureau: Bristol Myers Squibb, Novartis, Pierre Fabre, MSD Oncology, PharmaMar
Travel, Accommodations, Expenses: Pierre Fabre

Giuseppe Maria Milano

Consulting or Advisory Role: Recordati Rare Disease

Alba Rubio-San-Simón

Honoraria: Roche, Miltenyi Biotec, Ricordati Rare Disease
Speakers' Bureau: SERB
Travel, Accommodations, Expenses: EUSA Pharma, Roche

Jaume Mora

Consulting or Advisory Role: Ymabs Therapeutics Inc, Bristol Meyers
Research Funding: Roche/Genentech (Inst)

M^a Ángeles Vaz-Salgado

Consulting or Advisory Role: Servier, Novocure
Speakers' Bureau: Novocure

Roberto Luksch

Consulting or Advisory Role: Recordati, SERB Pharmaceuticals
Travel, Accommodations, Expenses: Recoati Rare Diseases

Cristina Mata

Consulting or Advisory Role: ITM Isotope Technologies Munich, Ipsen
Travel, Accommodations, Expenses: Novartis

Claudia Valverde

Consulting or Advisory Role: PharmaMar, Boehringer Ingelheim, Deciphera, Philogen, Bayer
Research Funding: Bayer (Inst), Boehringer Ingelheim (Inst)
Travel, Accommodations, Expenses: PharmaMar, Bayer, Deciphera
Uncompensated Relationships: GEIS (Spanish Sarcoma Group)

Toni Ibrahim

Speakers' Bureau: Istituto Gentili
Research Funding: Ipsen (Inst), PharmaMar (Inst)

Javier Martin Broto

Leadership: Sarcoma Research Solutions SL
Consulting or Advisory Role: PharmaMar, GlaxoSmithKline, Novartis, Amgen, Bayer, Roche, Lilly, Tecnofarma, Asofarma, Boehringer Ingelheim, Cogent Medicine
Speakers' Bureau: PharmaMar
Research Funding: GlaxoSmithKline (Inst), PharmaMar, Novartis (Inst), Eisai (Inst), Lilly (Inst), Bayer (Inst), Eisai, Novartis, Lixte Biotechnology (Inst), Karyopharm Therapeutics (Inst), Deciphera (Inst), Blueprint Medicines (Inst), Nektar (Inst), Forma Therapeutics (Inst), Amgen (Inst), Daiichi Sankyo (Inst), Immix BioPharma (Inst), Bristol Myers Squibb (Inst), Pfizer (Inst), Celgene (Inst), PharmaMar (Inst), Arog (Inst), Adaptimmune, Rain Therapeutics (Inst), InhibRx (Inst), Ayala Pharmaceuticals (Inst), Philogen (Inst), Cebiotex (Inst), PTC Therapeutics (Inst), SpringWorks Therapeutics (Inst)
Expert Testimony: PharmaMar, Novartis
Travel, Accommodations, Expenses: PharmaMar, Pfizer

No other potential conflicts of interest were reported.

APPENDIX

TABLE A1. Ongoing Studies With Mifamurtide in Patients With High-Grade Osteosarcoma

NCT Number	Study Title	Sponsor	Other ID
NCT00631631	Mifamurtide (L-MTP-PE) for High-Risk Osteosarcoma	Millennium Pharmaceuticals, Inc	MTP-OS-403
NCT01194284	Surveillance Study of Patients With Newly Diagnosed Osteosarcoma	Millennium Pharmaceuticals, Inc	C23003 2009-017204-89 (EudraCT Number)
NCT02441309	A Eurosarc Study of Mifamurtide in Advanced Osteosarcoma (MEMOS)	University of Oxford	OCTO_039
NCT04571229	Expanded Access Use of L-MTP-PE for the Treatment of Osteosarcoma	Memorial Sloan Kettering Cancer Center	20-324
NCT03643133	Mifamurtide Combined With Postoperative Chemotherapy for Newly Diagnosed High Risk Osteosarcoma Patients (SARCOME13)	UNICANCER	UC-0150/1704 2017-001165-24 (EudraCT_Number)
NCT01459484	ABCB1/P-glycoprotein Expression as Biologic Stratification Factor for Patients With Non Metastatic Osteosarcoma	Italian Sarcoma Group	ISG-OS2 2011-001659-36 (EudraCT_Number)
NCT04383288	ABCB1/P-glycoprotein Expression Influence on Nonmetastatic Osteosarcoma of the Extremities	Grupo Espanol de Investigacion en Sarcomas	Geis-33
NCT04890067	Observational Study in Localized Osteosarcoma	AIEOP/Italian Sarcoma Group	AIEOP/ISG OS2 Oss

Abbreviations: ID, identifier; L-MTP-PE, liposomal muramyl tripeptide phosphatidyl ethanolamine; NCT, noncontrolled trial.




FIG A1. A chronological roadmap for original manuscripts on P-glycoprotein role in osteosarcoma.




FIG A2. Study flow diagram of the combined analysis of the Italian ISG/OS-2 and Spanish GEIS-33 studies.




FIG A3. EFS according to induction chemotherapy and mifamurtide in (A) GRs and (B) PRs. EFS, event-free survival; GRs, good responses; HDIFO, high-dose ifosfamide; MAP, methotrexate, doxorubicin, and cisplatin; PRs, poor responses.




FIG A4. (A) EFS curves of Pgp-positive patients stratified according to tumor necrosis, (B) EFS curves of Pgp-negative patients stratified according to tumor necrosis. EFS, event-free survival; GR, good response; Pgp, P-glycoprotein; PR, poor response.