

Melatonin as a Possible Stimulus to Unmask an Oxytocin-Deficient State in Hypopituitarism and Hypothalamic Damage

Queralt Asla,^{1,2,3} Maite Garrido,⁴ Eulàlia Urgell,⁵ Sílvia Terzan,⁵ Alicia Santos,^{1,2,6,7} Nimmy Varghese,^{8,9} Cihan Atila,^{10,11} Betina Biagetti,^{6,12} Franziska Plessow,^{13,14} Mirjam Christ-Crain,^{10,11} Anne Eckert,^{8,9} Susan M. Webb,^{2,6,7} Elizabeth A. Lawson,^{13,14} and Anna Aulinas^{1,2,3,6}

¹Department of Endocrinology and Nutrition, Hospital de la Santa Creu i Sant Pau, ENDO-ERN, 08025 Barcelona, Spain

²Institut de Recerca Sant Pau (IR-SANT PAU), 08041 Barcelona, Spain

³Faculty of Medicine, University of Vic-Central University of Catalonia (UVic-UCC), 08500 Vic/Manresa, Catalonia, Spain

⁴Centre d'Investigació del Medicament (CIM), IR-SANT PAU, 08041 Barcelona, Spain

⁵Department of Biochemistry, Hospital de la Santa Creu i Sant Pau, 08025 Barcelona, Spain

⁶Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBER-ER, Unit 747), Instituto de Salud Carlos III, 28222 Majadahonda, Spain

⁷Department of Medicine, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

⁸Research Cluster Molecular and Cognitive Neurosciences, Universität Basel, 4055 Basel, Switzerland

⁹Neurobiology Lab for Brain Aging and Mental Health, University Psychiatric Clinics Basel, 4002 Basel, Switzerland

¹⁰Department of Endocrinology, Diabetology and Metabolism, Universitätsspital Basel, 4031 Basel, Switzerland

¹¹Department of Clinical Research, Universität Basel, 4031 Basel, Switzerland

¹²Department of Endocrinology and Nutrition, Hospital Universitari Vall d'Hebron, ENDO-ERN, 08035 Barcelona, Spain

¹³Neuroendocrine Unit, Massachusetts General Hospital, Boston, MA 02114, USA

¹⁴Department of Medicine, Harvard Medical School, Boston, MA 02115, USA

Correspondence: Anna Aulinas Masó, MD, PhD, Department of Endocrinology and Nutrition, Hospital de la Santa Creu i Sant Pau, Sant Antoni Maria Claret St, 167, 08025 Barcelona, Spain. Email: aaulinas@santpau.cat.

Abstract

Context: Increasing evidence supports the presence of oxytocin deficiency (OXT-D) in hypopituitarism and hypothalamic damage (HHD). Identifying an applicable and reliable test to diagnose OXT-D is an unmet need. Melatonin (MEL) might be a candidate for such a test as it regulates OXT release in animals.

Objective: This work aimed to examine the effects of melatonin on OXT release in HHD compared to healthy controls (HCs) and to describe the psychopathology, sexual function, and quality of life (QoL) and their associations with OXT.

Methods: This proof-of-concept study (NCT05319301) included 20 participants with HHD (11 women) and 20 HCs (11 women). Blood samples were collected over 120 minutes to assess plasma OXT. A linear mixed-effects regression model was used to evaluate the change in OXT in response to MEL in HHD compared to HCs.

Results: MEL significantly increased OXT at T90 vs T0 in HCs compared to the HHD group (difference 14.57 pg/mL 26% increase; 95% CI, 1.90–27.23; $P = .02$). The HHD group had more depression symptoms, alexithymia, impaired sexual function, and worse QoL compared to HCs. The mean percentage change in OXT from T0 to T90 was negatively associated with depressive and alexithymia symptoms in the HHD group and anxiety in both groups.

Conclusion: The reduced OXT response after MEL in HHD supports the existence of an impaired OXT response at least in a subset of patients with HHD. The associations between OXT changes and psychopathology suggest its role in mood and QoL. These findings support further investigation into MEL's role as a diagnostic tool to address OXT-D.

Key Words: hypopituitarism, melatonin, oxytocin, arginine-vasopressin deficiency, pineal gland, stimulation test

Abbreviations: ACTH, adrenocorticotropin; AUC, area under the curve; AVP, arginine-vasopressin; AVP-D, arginine-vasopressin deficiency; BIS, Barrat Impulsiveness Scale; BMI, body mass index; FSFI, Female Sexual Function Index; GH, growth hormone; HCs, healthy controls; HHD, hypopituitarism and hypothalamic damage; IIEF, International Index of Erectile Function; IIH, insulin-induced hypoglycemia; MDMA, 3,4-methylenedioxymethamphetamine; MEL, melatonin; OXT-D, oxytocin deficiency; QoL, quality of life; QoL-AGHDA, Quality of Life Assessment of Growth Hormone Deficiency in Adults; SF-36, Short-Form 36; STAI, State-Trait Anxiety Inventory; TAS-20, Toronto Alexithymia Score.

Received: 14 January 2025. Editorial Decision: 26 March 2025. Corrected and Typeset: 11 April 2025

© The Author(s) 2025. Published by Oxford University Press on behalf of the Endocrine Society.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com. See the journal About page for additional terms.

Hypopituitarism and hypothalamic damage (HHD) can be associated with multiple comorbidities and sequelae that substantially reduce quality of life (QoL) compared to the general population (1, 2). Increased rates of psychopathology (ie, anxiety, depression), alexithymia (ie, difficulty discerning and explaining one's own emotions), impaired prosocial functioning, and sexual dysfunction persist despite receiving standard hormone replacement, driving the need for additional therapeutic strategies to address the unmet needs of individuals with HHD (3, 4).

Importantly, the role of clinically relevant oxytocin deficiency (OXT-D) in patients with HHD has experienced a growing interest over the last decade, since OXT is involved in several physiological processes including psychological and social behaviors in both sexes (5-14). OXT, like arginine-vasopressin (AVP), is a neuropeptide mainly synthesized in the supraoptic and paraventricular nuclei of the hypothalamus and secreted into the peripheral circulation from the posterior lobe of the pituitary gland, as well as released to the central nervous system (3, 4). Therefore, an impaired OXT system could be present in some patients with HHD and might explain—at least partly—the observed psychopathology.

Initial research evaluating OXT secretion in HHD was limited to a single measurement of OXT (15-18). A more reliable approach using a 1-hour pool of fasting plasma OXT (19) evidenced significantly lower OXT levels in patients with AVP deficiency (AVP-D) who also had greater psychopathology compared to healthy controls (HCs). These initial findings suggested that an OXT deficient system might exist at least in a subset of patients with HHD; however, the difference compared to HCs was too small to be considered a reliable test (15-19). Going a step further searching for a stimulus to diagnose OXT-D, a few stimuli have been investigated in humans so far, including insulin-induced hypoglycemia (IIH) (20-23), hypertonic saline infusion (24), arginine (24), macimorelin (24), angiotensin II (25), 3,4-methylenedioxymethamphetamine (MDMA) (26), glucagon (27), corticotropin-releasing hormone (28), and kisspeptin (29), among others. Nonetheless, most of them were not found to be suitable as a provocative test to diagnose OXT-D, except for IIH, which variably increased OXT in HCs (20-23) but not in patients with AVP-D (20); hypertonic saline infusion, which increased the OXT response in healthy volunteers but with a high variance (24); and MDMA, which induced a remarkable and sustained increase in OXT concentrations in the HC group with no increase in patients with AVP-D (26). In addition, the lack of OXT increase in the AVP-D group was associated with reduced subjective prosocial and anxiolytic effects of MDMA compared to HCs, further supporting the central lack of OXT (26). Nevertheless, a safer and simpler stimulus to reveal an OXT deficient state in patients with HHD is still required since MDMA is not exempt from risks (ie, tachycardia, diaphoresis) and it is contraindicated in some clinical conditions.

Melatonin (MEL) is a neurohormone mainly produced by the pineal gland (30). Most of the physiological functions of MEL are controlled by the hypothalamus and the pituitary gland and mediated by MEL receptors (MT1) widely distributed in the hypothalamus, and colocalized with the oxytocinergic neurons in the paraventricular and supraoptic nuclei (30, 31). Several *in vitro* and *in vivo* studies have suggested that MEL might regulate the synthesis and release of OXT, evidencing an anatomic and functional interrelationship between the

pineal gland and the hypothalamic-neurohypophyseal axis (32). Pinelectomy in animal models was associated with a decrease in the storage of OXT in the neurohypophysis (33-36) and some *in vitro* and animal studies have shown a stimulating effect of MEL on OXT release (34, 37, 38). Preliminary studies in healthy volunteers showed an increase of OXT secretion following a low dose of oral MEL vs placebo (39). Likewise, MEL increased OXT-dependent myometrial contractions (40). Altogether these findings suggest that MEL, currently used to mitigate jet lag and to treat sleep disorders, may be a good stimulating agent for endogenous OXT secretion and ultimately be applicable as a potential diagnostic test to unmask an OXT deficient state in humans.

Thus, this study aimed to examine plasma OXT dynamics in response to a low dose of oral MEL (1.95 mg) in patients with HHD compared to HCs. To improve our understanding of OXT-D, we also explored the relationships between OXT levels and clinically relevant features of psychopathology, sexual function, and QoL in both groups. Based on prior evidence, we hypothesized that OXT release would be blunted in patients with HHD compared to HCs in response to MEL and, a blunted OXT response to MEL would be associated with greater psychopathology, impaired sexual function, and worse QoL in patients with HHD.

Materials and Methods

Trial Design

This is a proof-of-concept study with a quasi-experimental design conducted in accordance with the ethical principles of the Declaration of Helsinki and the Good Clinical Practice Guideline of the International Conference of Harmonisation (41, 42). The study protocol was approved by the institutional review board of the Hospital de la Santa Creu i Sant Pau (protocol code IIBSP-OXI-2021-112), and all participants provided written informed consent to a study physician before any study procedure. The study was registered at [ClinicalTrials.gov](https://clinicaltrials.gov) (NCT05319301).

Study Participants

Twenty-one adult patients with HHD and 20 HCs were included. HCs were of similar age and sex as the patients with HHD. Participants with HHD were primarily recruited from the Research Center for Pituitary Diseases at the Hospital de la Santa Creu i Sant Pau and Hospital Universitari Vall d'Hebron (Barcelona, Spain). HCs were recruited from the surrounding community and from the volunteers' list of the Research Center for Pituitary Diseases that had previously consented to be informed of future research projects.

The pituitary diagnoses were established as per routine clinical practice based on the standard protocol and clinical practice guidelines (2). We included patients with hypopituitarism (>1 anterior pituitary deficit) and at least one sign of hypothalamic damage (AVP-D and/or severe-onset obesity immediately after hypothalamic and pituitary damage and/or hyperphagia and/or brain tumors involving the hypothalamus [ie, craniopharyngioma, germinoma] and/or radiotherapy of the hypothalamic-pituitary area and/or hypothalamic damage visualized by neuroimaging). Patients with HHD were on stable hormone replacement for at least 3 months prior to the study visit (Supplementary Appendix 1 provides the main exclusion criteria (43)).

Procedures

This study comprised 2 visits: a screening visit to assess eligibility criteria and, if eligible, an outpatient visit.

The screening visit included personal medical history and, in those with HHD, a complete history of the hypothalamic and pituitary disease, a physical examination, and the assessment of baseline biochemical variables, including the hormone profile in patients with HHD to confirm appropriate hormone substitution. If a hormone imbalance was detected, dose adjustments were performed accordingly, and hormone levels were reassessed 6 to 8 weeks later. Before each study visit, urine pregnancy tests were performed on premenopausal female participants.

A 3-hour main visit was conducted at the Center for Drug Research of the Sant Pau's Biomedical Research Institute in a quiet room and was run by one investigator and a research nurse. The main visit started at 8 AM after an at least 8-hour fast, and ad libitum water intake was allowed in patients with AVP-D. We confirmed no alcohol intake nor vigorous physical activity 24 hours before each visit. Participants were asked to take their morning medication after finishing the main visit, except for desmopressin (in patients with AVP-D), as required. We scheduled the study visit of premenopausal female participants in the early to midfollicular phase of their menstrual cycle to minimize the effects of increasing estradiol on OXT levels (8). At the beginning of the visit, participants were asked about their current health status and changes in medication use, and a physical examination was performed (including body mass index [BMI], waist-to-hip ratio, blood pressure, heart rate, and temperature) to confirm eligibility and rule out any potential exclusion criteria not previously detected. Once eligibility was reconfirmed, an intravenous catheter was placed and participants were asked to complete validated questionnaires to assess mood (Beck Depression Inventory-2 [BDI-II] (44, 45) and State-Trait Anxiety Instrument [STAI] questionnaires (46-48)), alexithymia (Toronto Alexithymia Scale 20 [TAS-20] (49)), impulsivity (Barratt Impulsiveness Scale [BIS] (50)), sexual function (Female Sexual Function Index [FSFI] (51) in women and International Index of Erectile Function [IIEF] (52) in men), and QoL (with the Short Form 36 [SF-36] (53) and the QoL Assessment of Growth Hormone Deficiency when applicable [QoL-AGHDA] (54)). A more detailed description of the questionnaires used is provided in Supplementary Appendix 2 (43).

After completion of the questionnaires, a baseline blood sample was drawn (T0) for basic metabolic panel, lipid profile, complete blood count, anterior pituitary hormone assessment (thyrotropin, free thyroxine, luteinizing hormone, follicle-stimulating hormone, testosterone [men], estradiol [women], prolactin, adrenocorticotropin [ACTH], cortisol, growth hormone [GH], insulin-like growth factor-1), and OXT. Low-dose MEL, a compounded product considered a dietary supplement in Spain (1.95 mg), was administered sublingually. Blood samples for OXT measurements were collected after MEL administration over 120 minutes at the following time points: 30 (T30), 60 (T60), 90 (T90), and 120 (T120) minutes. Blood sampling and processing were performed by a research nurse and a study physician, with identical procedures used for the patients and the HCs. Participants were instructed to refrain from using cell phones, listening to the radio, watching TV, and engaging in conversations to avoid any potential increase in OXT concentrations due to social stimulation.

Safety Events

All participants were monitored by the investigator team for treatment-emergent adverse events for the entire duration of the visits. Vital signs (blood pressure, heart rate, and body temperature) were assessed at the beginning and the end of each main visit. All these data were collected in a specific case report form.

Biochemical Measurements

Biochemical and blood parameters were analyzed using routine laboratory methods, and all hormonal measurements were performed using immunoassay methods by the Hormones and Biochemistry laboratory in the Hospital de la Santa Creu i Sant Pau, except for OXT.

Equal aliquots of plasma from each time point for OXT measurement were immediately stored at -80°C and were run blinded in a single batch at the end of the study by the Neurobiology Laboratory for Brain Aging and Mental Health at Basel University (Switzerland), experienced in measuring OXT (24, 26, 27). The OXT concentration was measured using the Oxytocin ELISA kit (ENZO Life Sciences) (Arbor Assays catalog No. K048-H, RRID:AB_2924717) according to the manufacturer's protocol (55). Intra-assay coefficient of variation was 2.25% and interassay coefficient of variation was 4.45%. In brief, the plasma was diluted at a ratio of 2:1 in 1.5% TFA- H_2O and centrifuged at $17\,100 \times g$ for 15 minutes at 4°C . The supernatant was collected. In the next step, the supernatant was loaded onto an Oasis PRiME HLB 96-well plate, 30 mg sorbent (Waters Corporation) and washed with 0.10% TFA- H_2O . Afterwards, the sample was eluted with 95% acetonitrile + 5% of a 0.1% TFA- H_2O solution. Then, the elute was collected and dried in an Eppendorf centrifugal vacuum concentrator 5301. For the measurement, the residue was reconstituted in assay buffer. The reconstituted samples, standards, and controls were plated on the goat anti-rabbit immunoglobulin G microtiter plate and incubated with the OXT conjugate at 4°C overnight. Thereafter the plate was washed, and the substrate was added. The reaction was stopped after 1 hour and the plate was read at 405 nm. OXT concentration was calculated with a standard curve determined from 4 parameter logistics curve fit.

Statistical Analysis

Statistical analysis was performed using STATA software, version 17.0 (StataCorp LLC). For descriptive analysis, quantitative data were expressed as mean and SEM (gaussian distribution) or as median and interquartile range (25th percentile-75th percentile) (nongaussian distribution); and categorical data as absolute frequencies and percentages. Distribution of continuous data was tested for normality using the Shapiro-Wilk test. Means across groups (HHD vs HCs) were compared using the *t* test or Mann-Whitney *U* test accordingly, and the Fisher exact test was used to compare categorical variables between 2 or more groups.

The primary outcomes measure was the change in OXT concentrations from baseline (T0) to each time point evaluated (T30, T60, T90, T120) over 120 minutes following MEL administration. To analyze OXT level change relative to baseline at each time point post MEL administration, we used a linear mixed-effects regression model (mixed command), including time (5 time points as categorical variable: baseline, T30, T60, T90, and T120), and between-subject

factor diagnosis (patient with HHD vs HC), and the interactions (two-way) between both factors as fixed effects, and controlled for BMI. A random intercept for study participants was added to the model to account for the correlation between measures within the same participant, and a general unstructured variance-covariance structure was used to model the within-patient errors. The OXT area under the curve (AUC) was also calculated using the trapezoidal formula (56). All *P* values were 2-sided, and statistical significance was set at *P* less than .05.

We used Spearman rank correlation coefficients to explore the associations between questionnaire scores and fasting OXT levels, AUC of OXT, and relative change of OXT levels, only when differences in OXT levels were statistically significant between HHD and HCs. All *P* values were adjusted for multiple comparisons using the Holm-Bonferroni method.

Results

Sample Characteristics

Between April 2022 and July 2023, a total of 41 participants (21 patients with HHD and 20 HCs) participated in this quasi-experimental study. One woman with HHD voluntarily abandoned the study after the screening visit. Finally, 40 participants (20 patients with HHD and 20 HCs) completed the study.

Baseline characteristics across groups are summarized in Table 1. HHD and HC groups did not differ in age and sex distribution. The median BMI was higher in the HHD group compared to the HC group, 30.20 (3.79) vs 22.91 (7.34) kg/m², respectively (*P* = .02). The remaining demographic and baseline characteristics did not differ between both groups, except for a higher prevalence of psychiatric disorders (*P* = .02) and lower bone mineral density (*P* = .01) in the HHD group in comparison to HCs. No statistically significant differences in the prevalence of psychiatric disorders among patients with HHD with or without GH deficiency were observed as well as in the prevalence of psychiatric treatment used between patients with HHD and HCs.

The clinical history of the hypothalamic and pituitary disease of patients with HHD are summarized in Table 2.

The most frequent hypothalamic-pituitary disease was craniopharyngioma in 50% of the participants. Eighty-five percent of the patients with HHD (*n* = 17) presented with 3 or more anterior hormone pituitary deficits with a mean time since first anterior pituitary deficiency of 17.98 ± 2.37 years, ranging from 1.40 to 38.0 years. Eighty-five percent of participants (*n* = 17) had AVP-D at study entry for a median time of 14.52 (7.50) years, ranging from 0.52 to 38.0 years.

Ten (91%) women with HHD had central hypogonadism, whereas only 1 woman maintained gonadal axis function preserved but with irregular menses (see Tables 1 and 2). Among women with hypogonadism, 6 (60%) were menopausal and 4 (40%) had regular menses on estrogen/progestogen treatment (3 were treated with estrogen and progestogen at a substitutive dose and 1 was treated with oral contraceptive pills). Regarding healthy women, 4 (36.36%) were menopausal and the remaining had regular menses (of these, only 1 participant was treated with oral contraceptive pills). Mean days since the last menstrual period at the time of the study visit were similar between groups (HCs: 5.67 ± 1.28 days vs HHD: 3.60 ± 1.03 days; *P* = .25). Eight (89%) of the men with HHD presented with central hypogonadism; 4 (50%) were treated with daily transdermal testosterone and the other 4 (50%) received monthly to quarterly intramuscular testosterone. Median testosterone levels were similar between the groups (HCs: 18.30 [4.80] nmol/L vs HHD: 13.20 [12.70] nmol/L; *P* = .38).

Oxytocin Levels in Response to Melatonin

Table 3 shows the response of OXT concentrations over time (T0, T30, T60, T90, and T120) after MEL administration across groups (HHD vs HCs). Baseline OXT concentrations were similar between the groups (HHD 56.74 ± 4.13 vs HCs 57.05 ± 3.99 pg/mL; *P* = .86). In response to sublingual MEL administration, the change in plasma OXT concentrations at T90 vs T0 was greater in the HC group than the HHD group (OXT difference 14.57 pg/mL; 95% CI, 1.90-27.23; *P* = .02) (Fig. 1), with a significant increase in OXT levels at T90 (vs T0) only in the HC group (*P* = .003). Specifically, the mean percentage increase in OXT levels from T0 to T90 in HCs compared to the HHD group was

Table 1. Summary of baseline characteristics across groups (participants with hypopituitarism and hypothalamic damage and healthy controls)

	Participants with HHD (n = 20)	Healthy controls (n = 20)	<i>P</i>
Sex (male/female), n (%)	9 (45%)/ 11 (55%)	9 (45%)/ 11 (55%)	≥.99
Age at study entry, years	57.21 (18.46)	45.29 (28.43)	.12
Estrogen status in women, n (%)			.40
Menopausal	6 (54.55%)	4 (36.36%)	
Regular menses	4 (36.36%)	7 (63.64%)	
Irregular menses	1 (9.09%)	0 (0%)	
Body mass index, kg/m ²	30.20 (3.79)	22.91 (7.34)	.02
Waist-to-hip ratio, iliac	0.91 (0.13)	0.82 (0.20)	.06
Hypertension, n (%)	5 (25%)	2 (10.53%)	.41
Diabetes mellitus, n (%)	2 (10.53%)	2 (10%)	≥.99
Dyslipidemia, n (%)	10 (52.63%)	6 (30%)	.15
Psychiatric disorder, n (%)	8 (40%)	1 (5%)	.02

Numerical data are reported as median and interquartile range (25th percentile–75th percentile). Abbreviation: HHD, hypopituitarism and hypothalamic damage.

Table 2. Summary of the clinical history of hypothalamic-pituitary disease in participants with hypopituitarism and hypothalamic damage

Hypothalamic-pituitary disease	Participants with HHD (n = 20)
Diagnosis	
Age at diagnosis, years	36.46 ± 4.10
Hypothalamic pituitary disease, n (%)	
Craniopharyngioma	10 (50%)
NFPA	5 (25%)
Granular cell tumor of pituitary stalk	2 (10%)
Prolactinoma	1 (5%)
Other	2 (10%)
Tumor size, mm	26.25 ± 4.44
Treatment	
Surgery, n (%)	20 (100%)
Type of surgery, n (%)	
Transsphenoidal	10 (50%)
Craniotomy	9 (45%)
Both	1 (5%)
Radiotherapy, n (%)	7 (35%)
Current clinical situation	
≥3 anterior pituitary deficiencies, n (%)	17 (85%)
TSH deficiency, n (%)	19 (95%)
FSH/LH deficiency, n (%)	18 (90%) (10 women and 8 men)
ACTH deficiency, n (%)	15 (75%)
GH deficiency, n (%)	13 (65%)
AVP deficiency, n (%)	17 (85%)
Signs of hypothalamic damage, n (%)	
Suprasellar lesion or involving hypothalamus	19 (95%)
Multiple surgeries at hypothalamic-pituitary level	3 (15%)
Radiotherapy at hypothalamic-pituitary level	7 (35%)
Hypothalamic obesity	4 (20%)

Numerical data are reported as mean ± SEM (gaussian distribution), median, and interquartile range (nongaussian distribution). Abbreviations: ACTH, adrenocorticotropin; AVP, arginine-vasopressin; FSH, follicle-stimulating hormone; GH, growth hormone; HHD, hypopituitarism and hypothalamic damage; LH, luteinizing hormone; NFPA, nonfunctioning pituitary adenoma; TSH, thyrotropin.

25.85 ± 9.05 vs -1.39 ± 8.93%, respectively (95% CI, 1.30-53.18; $P = .04$). There were no changes in OXT levels in response to MEL at other time points. When analyzing the subgroup of participants with AVP-D ($n = 17$) in comparison to HCs, the overall results did not change and OXT concentrations at T90 vs T0 were higher in the HC group compared to the HHD group (OXT difference 13.29 pg/mL; 95% CI, 0.95-25.65; $P = .03$). Baseline sodium levels were within the normal range in all participants.

The median AUC of plasma OXT concentrations was lower in the HHD than in the HC group (7003.5 [1511.4] vs 7689.8 [1580.3] pg/mL, respectively; $P = .05$) (Fig. 2).

Adverse Events

MEL administration did not cause any adverse effects and did not affect vital signs.

Table 3. Adjusted mean plasma oxytocin concentrations (pg/mL) after melatonin administration across groups

Time points	Participants with HHD (n = 20)	HCs (n = 20)	HCs vs HHD
	OXT concentrations, pg/mL	OXT concentrations, pg/mL	Mean difference ^a (95% CI)
0	56.75 ± 4.13	57.05 ± 3.99	
30	57.62 ± 4.56	61.76 ± 4.66	3.8 (-7.2 to 14.9)
60	55.17 ± 4.56	62.91 ± 4.66	7.4 (-5.5 to 20.4)
90	54.07 ± 4.37	68.95 ± 4.37	14.6 (1.9 to 27.2)
120	60.50 ± 4.24	69.23 ± 4.13	8.4 (-5.6 to 22.5)

Abbreviations: HCs, healthy controls; HHD, hypopituitarism and hypothalamic damage; MEL, melatonin; OXT, oxytocin.
^aShows the mean adjusted difference in OXT concentrations (pg/mL) after MEL administration between HCs vs HHD across all postintervention time points (respect to baseline, T0) (two-way interaction between the 2 factors: participant and time point) and controlled for body mass index.

Psychopathology, Sexual Function, and Quality of Life Across Groups

Table 4 shows baseline scores of the questionnaires to assess psychopathology, sexual function, and QoL across groups.

In comparison to the HC group, patients with HHD scored higher on depression symptoms (BDI-II) (8.50 [11] vs 5 [7.50]; $P = .02$) (35% HHD vs 5% HC with mild-moderate depression and 5% HHD vs 0% HCs with severe depression). HHD individuals also had higher scores with regard to symptoms of alexithymia (TAS-20) in the subscales of difficulty identifying feelings (17.30 ± 1.59 vs 13 ± 1.02; $P = .03$) and externally oriented thinking (21.93 [7] vs 18 [4]; $P = .02$).

In addition, participants with HHD, when compared to HCs, had impaired sexual function (FSFI total score in women 15.10 [27.10] vs 28.10 [24.30]; $P = .05$; IIEF total score in men 25 ± 5.32 vs 47 ± 6.63; $P = .02$).

Finally, worse QoL was observed in the HHD group in comparison to the HC group (SF-36 physical functioning 85 [20] vs 97.50 [7.50]; $P < .01$; SF-36 physical health 90 [50] vs 100 [6.70]; $P = .01$; SF-36 bodily pain 66.70 [55.60] vs 88.90 [22.20]; $P = .01$; and SF-36 general health 45 [50] vs 82.5 [20]; $P < .001$).

Psychopathology, Sexual Function, and Quality of Life and Their Associations With Oxytocin

In both groups, the mean percentage change in OXT levels from T0 to T90 was associated with lower anxiety symptoms (HHD: $r = -0.58$; $P = .02$ and HCs: $r = -0.59$; $P = .01$). In the HHD group only, the mean percentage change in OXT levels from T0 to T90 was associated with lower depressive ($r = -0.59$; $P = .01$) and alexithymia total scale ($r = -0.59$; $P = .01$) symptoms. These relationships remained statistically significant after controlling for multiple comparisons. No significant associations were found between the mean percentage change in OXT levels from T0 to T90 and sexual function or QoL in either the HHD or HC groups. Overall baseline OXT concentrations and the AUC of OXT were not significantly associated with psychopathology, sexual function, or QoL in either patients with HHD or in the HCs (Supplementary Table S1 (43)).




Figure 1. Response of OXT levels over time (from T0 to T120) after MEL across groups (HHD vs HC). Dot represents the mean value of OXT and whiskers represent the SEM range. **P* less than .05 (statistically significant change from T0 to T90 between groups). HC, healthy controls; HHD, hypopituitarism and hypothalamic damage; MEL, melatonin; OXT, oxytocin; T, time point.




Figure 2. Dot plot comparing the AUC of plasma OXT concentrations after MEL across groups (HC and HHD). Dots represent the AUC of OXT after MEL for each participant, plus signs represent the median of AUC of OXT after MEL across groups (HC and HHD), and dashed lines represent the interquartile range for each median. HC, healthy controls; HHD, hypopituitarism and hypothalamic damage; MEL, melatonin; OXT, oxytocin.

Discussion

To the best of our knowledge, this is the first study that has investigated plasma OXT dynamics in response to low doses of sublingual MEL by comparing patients with HHD to healthy individuals, aimed at identifying a provocative test to unmask OXT-D and clinically characterize an OXT deficient state. We demonstrated lower plasma OXT concentrations after MEL administration in the HHD group in comparison to the control group, supporting the evidence of an OXT deficient state in HHD. However, further studies are required to identify the MEL dose regimen (eg, dose, time, route of administration) to more efficiently magnify the OXT differences from baseline to peak between HHD and HCs before suggesting MEL as a possible test to unmask OXT-D in HHD. The HHD group suffered more psychopathology, impaired sexual function, and worse QoL compared to HCs; notwithstanding, most of them were not in the clinically relevant range. Associations between greater percentage change in OXT levels from T0 to T90 and lower depressive and alexithymia symptoms in the HHD group and lower anxiety symptoms in both groups provide additional evidence for a clinically significant OXT deficient state.

Similar to our clinical observation, some but not all experimental studies found a stimulating effect of MEL on OXT release (34, 37, 38), while others showed either no effects (34, 57) or reduced OXT concentrations following MEL administration (33, 38, 58-63). Such variations could be explained by differences in study species (rodents, mammals vs humans and in vitro or in vivo), dose and regimen employed (pharmacological or supraphysiological vs physiological) (59), time of day when the research was performed (62), as well as in the route of administration (intravenous, oral vs sublingual) (34, 59, 64). Few human studies, mostly in healthy men, have evaluated circulating OXT release following MEL administration, evidencing a dose-dependent secretion of OXT, with high doses of MEL causing a decrease in OXT release, and low doses increasing OXT levels. In an initial study, 10 healthy men received 5 mg of MEL orally for 4 days in the late afternoon vs placebo, and a mild decrease of OXT concentrations after MEL was observed in comparison to the

placebo group (63). Similarly, the administration of higher doses (6 and 12 mg) of oral MEL vs placebo in basal conditions did not modify OXT release over 3 hours in healthy men (65). Thereafter, Forsling et al (39) explored the effect of oral MEL given in the afternoon at different doses (0.05, 0.5, and 5 mg) on OXT release over 150 minutes, showing dose-dependent changes in plasma OXT levels. While the lowest dose did not show a change in OXT levels, the dose of 0.5 mg showed a stimulatory effect and 5 mg an inhibitory effect on OXT levels. In particular, the 0.5 mg dose produced a peak of OXT of around 30% after 60 minutes compared to baseline.

According to the aforementioned studies, we chose a low dose of sublingual MEL (1.95 mg) to examine peripheral OXT secretion in HCs and in participants with HHD. In line with Forsling et al (39), we observed an increase of the AUC of plasma OXT concentrations in healthy individuals that peaked at 90 minutes, with a 26% increase of OXT compared to baseline, whereas, as expected, we did not observe any increase in OXT in participants with HHD. The difference in the time point when the peak was achieved –60 minutes in Forsling's vs 90 minutes in our study might be attributed to dose differences, pharmacokinetics (oral vs sublingual), the time of day when the study started (14:30 in Forsling's study vs 8:00 in the present study), as well as variability in OXT levels. It could be possible that MEL at doses above 2 mg as approved by drug agencies for sleep disorders would have had a greater effect on circulating OXT release. Nevertheless, a better understanding of the mechanisms involved (eg, receptor downregulation or desensitization, neurochemical interactions, feedback inhibition) in dose-dependent effects of MEL on OXT release is crucial to select the optimal dose regimen to be used as a provocative test for OXT-D.

The detailed mechanisms by which the pineal gland and MEL interact with the OXT system and increase OXT release remain unclear, although different mechanisms have been proposed: (i) OXT-containing fibers extend into the pineal gland, facilitating anterograde and retrograde axonal

Table 4. Baseline scores of questionnaires of study participants across groups

Questionnaires	Participants with HHD (n = 20)	HCs (n = 20)	P
Beck Depression Inventory-2	8.50 (11.0)	5 (7.50)	.02
STAI-State subscale^a	12.50 (11.50)	9 (7)	.13
STAI-Trait subscale	17 (23.50)	14 (10.50)	.40
TAS-20	49.22 ± 3.81	41.40 ± 2.33	.09
TAS-20 Difficulty describing feelings	13.20 ± 1.02	10.95 ± 0.88	.11
TAS-20 Difficulty identifying feelings	17.30 ± 1.59	13.0 ± 1.02	.03
TAS-20 Externally oriented thinking	21.93 (7)	18 (4)	.02
BIS	36.48 ± 3.07	33.25 ± 2.60	.43
BIS Cognitive impulsivity	12 ± 1.01	11.85 ± 1.06	.92
BIS Motor impulsivity	11.01 ± 1.44	11.50 ± 1.10	.79
BIS Nonplanning impulsivity	11 (10.50)	12 (7)	.71
FSFI^b	15.10 (27.10)	28.10 (24.30)	.05
FSFI Desire	2.84 ± 0.44	3.27 ± 0.32	.43
FSFI Arousal	2.40 (4.50)	4.50 (4.50)	.09
FSFI Lubrication	2.48 ± 0.75	3.71 ± 0.77	.27
FSFI Orgasm	2 (4.80)	4.80 (6)	.07
FSFI Satisfaction	2.69 ± 0.72	4.44 ± 0.42	.05
FSFI Pain	3.20 (4.80)	5.20 (6)	.16
IIEF^c	25 ± 5.32	47 ± 6.63	.02
IIEF Erectile function	8.89 ± 2.60	18.11 ± 3.70	.06
IIEF Intercourse satisfaction	0 (5)	9 (9)	.03
IIEF Orgasmic function	0 (6)	10 (3)	<.01
IIEF Sexual desire	5.33 ± 0.87	6.22 ± 0.87	.40
IIEF Overall satisfaction	6 (2)	8 (2)	.17
SF-36 score			
SF-36 Physical functioning	85 (20)	97.50 (7.50)	.003
SF-36 Role limitations due to physical health	90 (50)	100 (6.70)	.01
SF-36 Bodily pain	66.70 (55.60)	88.90 (22.20)	.01
SF-36 General health	45 (50)	82.50 (20)	<.01
SF-36 Vitality	59.40 (25)	68.80 (25)	.13
SF-36 Social	87.50 (43.75)	100 (12.50)	.06
SF-36 Emotional	95.85 (50)	100 (12.50)	.13
SF-36 Mental health	77.50 (32.50)	82.50 (15)	.70
QoL-AGHDA (n = 13)	8.77 ± 2.19	—	—

Data are reported as mean ± SEM (gaussian distribution) and as median (25th percentile–75th percentile) (nongaussian distribution).

Abbreviations: BIS, Barrat Impulsiveness Scale; FSFI, Female Sexual Function Index; HCs, healthy controls; HHD, hypopituitarism and hypothalamic damage; IIEF, International Index of Erectile Function; QoL-AGHDA, Quality of Life Assessment of Growth Hormone Deficiency in Adults; SF-36, Short-Form 36; STAI, State-Trait Anxiety Inventory; TAS-20, Toronto Alexithymia Score.

^aAvailable in 19 HCs.

^bOnly in women (11 HHD and 11 HCs).

^cOnly in men (9 HHD and 9 HCs).

transport between the hypothalamus and the pineal gland that may promote interaction between both neurohormones (34, 37, 66). Further, MEL receptors are found in the median

eminence and hypothalamic nuclei (67). (ii) The pineal gland has been shown to have effects on the 24-hour rhythmic variations of the neurohypophysial hormones (68, 69). Indeed, pinealectomy and the exposure to constant light (that inhibits MEL) change the daily rhythm of posterior pituitary hormone secretion (65, 70, 71). (iii) MEL could interfere with the metabolism of some intermediate neurotransmitters (serotonin, glutamate, noradrenaline, dopamine, and prostaglandins) in the hypothalamus that are also involved in the release of neurohypophysial hormones (33). (iv) MEL can cross the brain-blood barrier and, due to its lipophilicity, could also act on cells of the neurohypophysial system through a non-receptor-mediated manner independent of the effect on its receptors (60, 72).

In clinical practice, the evaluation of an endocrine deficiency (ie, GH or ACTH) often requires a provocative test since a single hormone measurement may overlap with physiological states and does not always rule out or confirm a hormone deficiency. Currently, despite the increasing evidence of a possible OXT deficient state in patients with HHD, a validated, safe, and feasible provocative test to rule out OXT-D does not exist. Although we observed a significant increase of OXT concentrations of 26% from T0 to T90 in HCs compared to no change (–1%) in HHD, this increment was mild compared to the reported 650% of OXT increase (at 180 minutes) after MDMA administration or 200% (at 45 minutes) after an IIH test (22, 26) or 50% after hypertonic saline (at 150 minutes) (24). In addition, high interindividual variability in OXT response was observed, indicating that MEL at this dose regimen is not a suitable candidate for a provocative test to unmask OXT-D. Therefore, further research is needed to determine whether different doses or time of MEL administration could induce more powerful increments in OXT release. Importantly, the MEL test is inexpensive, easy to administrate, well tolerated, and accepted by patients and with no risks compared to MDMA or IIH.

Worse psychopathology, sexual function, and QoL have been widely observed in patients with HHD in comparison to healthy individuals (18, 19, 26). Growing evidence suggests that an OXT deficient state could be involved in this poorer mental health and QoL in patients with hypopituitarism and AVP-D (16, 17, 26, 73). Although most of the questionnaires' scores observed in this study were not in the clinical range, we detected some significant negative associations between the OXT dynamics (mean percentage change in OXT levels from T0 to T90) and anxiety symptoms in both groups and depressive symptoms and alexithymia only in the HHD group. One explanation for this lack of association between the mean percentage change in OXT levels from T0 to T90 and depressive symptoms and alexithymia in the HC group could be the lower variability in the questionnaires' scores in comparison to the HHD group. We did not detect any statistically significant association between the OXT dynamics and some psychological, sexual function, and QoL tests in both groups.

Our study had some limitations, but also strengths. This is a proof-of-concept study with a strict protocol testing a potential novel and safe stimulus (MEL) to unmask an OXT deficient state in patients with HHD. While the results confirm prior evidence for an OXT deficient state in HHD and suggest that MEL might be a tool to diagnose OXT-D, the overall effects of MEL on OXT secretion with the current protocol are modest. In addition, the lack of a placebo group does not allow us to state with certitude that MEL was the single driver

of OXT increase. It is possible that different MEL doses may exert a stronger effect on OXT secretion. Therefore, additional doses and regimens of MEL administration should be evaluated to test the efficacy of MEL as an OXT stimulus to warrant its applicability in clinical practice. On the other hand, due to the low molecular weight of OXT and low plasma concentrations in nonpregnant humans, measurement and interpretation of OXT levels are still a major challenge and there is no consensus about the most appropriate assay and sample preparation to use (74). For these reasons, OXT samples were collected following a rigorous and identical protocol for each participant and were measured in a single batch to reduce variability in an experienced laboratory. The sample size, although sufficient to detect relevant differences in OXT release after MEL between HHD and HCs, did not allow us to perform subgroup analysis (sex or pituitary diseases). Finally, some human studies but not all have shown low OXT levels in patients with some psychiatric disorders (ie, posttraumatic stress disorder, depression, and autism spectrum disorders) (75-77). Nevertheless, in our study, OXT levels did not differ at any time point among patients with HHD with or without a psychiatric disorder.

In summary, patients with HHD have a diminished response of plasma OXT concentrations after a low dose of sublingual MEL compared to HCs, supporting the existence of an impaired OXT response at least in a subset of patients with HHD. The associations between OXT changes and psychopathology suggests its role in mood and QoL. These findings support further investigation into MEL's role as a diagnostic tool to address OXT-D.

Acknowledgments

We thank the Center for Drug Research staff and the study participants for their assistance in successfully completing the present study.

Funding

This work has been funded by the grant “Beca Junior Fundación SEEN Proyecto de investigación 2021” to A.A. and by the Instituto de Salud Carlos III through project FIS PI20/01165 (cofunded by the European Regional Development Fund/European Social Fund “A way to make Europe”/“Investing in your future”) and JR19/00013 to A.A. Q.A. received a postresidency fellowship by Fundació de Gestió Sanitària de l'Hospital de la Santa Creu i Sant Pau in 2021.

Disclosures

E.A.L. receives grant support and research study drug from Tonix Pharmaceuticals and receives royalties from UpToDate. E.A.L. and/or immediate family member holds stock in Thermo Fisher Scientific, Zoetis, Danaher Corporation, Intuitive Surgical, Merck, and West Pharmaceutical Services. E.A.L. is an inventor on US provisional patent application number 63/467 980 (Oxytocin-based therapeutics to improve cognitive control in individuals with attention deficit hyperactivity disorder). No other conflicts of interest, financial or otherwise, are declared by the authors.

Data Availability

Some or all data sets generated during and/or analyzed during the present study are not publicly available but are available from the corresponding author on reasonable request.

Clinical Trial Information

Clinical trial registration number NCT05319301 (registered March 17, 2022).

References

- Lynch S, Merson S, Beshyah SA, *et al.* Psychiatric morbidity in adults with hypopituitarism. *J R Soc Med.* 1994;87(8):445-447.
- Fleseriu M, Hashim IA, Karavitaki N, *et al.* Hormonal replacement in hypopituitarism in adults: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab.* 2016;101(11):3888-3921.
- Gimpl G. The oxytocin receptor system: structure, function and regulation. *Physiol Rev.* 2001;81(2):629-683.
- Baribeu. Oxytocin and vasopressin: linking pituitary neuropeptides and their receptors to social neurocircuits. *Front Neurosci.* 2015;9:335.
- Althammer F, Grinevich V. Diversity of oxytocin neurons: beyond magno- and parvocellular cell types? *J Neuroendocrinol.* 2017;30:e12549.
- Leng G, Sabatier N. Oxytocin—the sweet hormone? *Trends Endocrinol Metab.* 2017;28(5):365-376.
- Neumann ID, Slattery DA. Oxytocin in general anxiety and social fear: a translational approach. *Biol Psychiatry.* 2016;79(3):213-221.
- Aulinas A, Pulumo RL, Asanza E, *et al.* Endogenous oxytocin levels in relation to food intake, menstrual phase, and age in females. *J Clin Endocrinol Metab.* 2019;104(4):1348-1356.
- McCormack SE, Blevins JE, Lawson EA. Metabolic effects of oxytocin. *Endocr Rev.* 2020;41(2):121-145.
- Breuil V, Trojani MC, Ez-Zoubir A. Oxytocin and bone: review and perspectives. *Int J Mol Sci.* 2021;22(16):8551.
- Striepen N, Kendrick KM, Maier W, Hurlmann R. Prosocial effects of oxytocin and clinical evidence for its therapeutic potential. *Front Neuroendocrinol.* 2011;32(4):426-450.
- Tamma R, Colaianne G, Zhu LL, *et al.* Oxytocin is an anabolic bone hormone. *Proc Natl Acad Sci U S A.* 2009;106(17):7149-7154.
- Wang P, Wang SC, Yang H, *et al.* Therapeutic potential of oxytocin in atherosclerotic cardiovascular disease: mechanisms and signaling pathways. *Front Neurosci.* 2019;13:454.
- Wang P, Yang HP, Tian S, *et al.* Oxytocin-secreting system: a major part of the neuroendocrine center regulating immunologic activity. *J Neuroimmunol.* 2015;289:152-161.
- Daubenbuchel AM, Hoffmann A, Eveslage M, *et al.* Oxytocin in survivors of childhood-onset craniopharyngioma. *Endocrine.* 2016;54(2):524-531.
- Daughters K, Manstead ASR, Rees DA. Hypopituitarism is associated with lower oxytocin concentrations and reduced empathic ability. *Endocrine.* 2017;57(1):166-174.
- Gebert D, Auer MK, Stieg MR, *et al.* De-masking oxytocin-deficiency in craniopharyngioma and assessing its link with affective function. *Psychoneuroendocrinology.* 2018;88:61-69.
- Aulinas A, Guarda FJ, Yu EW, *et al.* Lower oxytocin levels are associated with lower bone mineral density and less favorable hip geometry in hypopituitary men. *Neuroendocrinology.* 2021;111(1-2):87-98.
- Aulinas A, Plessow F, Asanza E, *et al.* Low plasma oxytocin levels and increased psychopathology in hypopituitary men with diabetes insipidus. *J Clin Endocrinol Metab.* 2019;104(8):3181-3191.
- Nussey SS, Ang VT, Finer N, Jenkins JS. Responses of neurohypophysial peptides to hypertonic saline and insulin-induced hypoglycaemia in man. *Clin Endocrinol (Oxf).* 1986;24(1):97-105.

21. Chiodera P, Volpi R, Capretti L, *et al.* Hypoglycemia-induced arginine vasopressin and oxytocin release is mediated by glucoreceptors located inside the blood-brain barrier. *Neuroendocrinology*. 1992; 55(6):655-659.
22. Chiodera P, Volpi R, Capretti L, *et al.* Oxytocin response to challenging stimuli in elderly men. *Regul Pept*. 1994;51(2):169-176.
23. Volpi R, Chiodera P, Capretti L, *et al.* Effect of residual endogenous insulin secretion on the abnormal oxytocin response to hypoglycaemia in insulin-dependent diabetics. *J Intern Med*. 1998;244(1): 43-48.
24. Sailer CO, Winzeler B, Urwyler SA, *et al.* Oxytocin levels in response to pituitary provocation tests in healthy volunteers. *Eur J Endocrinol*. 2021;185:355-364.
25. Srinivasa S, Aulinas A, O'Malley T, *et al.* Oxytocin response to controlled dietary sodium and angiotensin II among healthy individuals. *Am J Physiol Endocrinol Metab*. 2018;315(4):E671-E675.
26. Atila C, Holze F, Murugesu R, *et al.* Oxytocin in response to MDMA provocation test in patients with arginine vasopressin deficiency (central diabetes insipidus): a single-centre, case-control study with nested, randomised, double-blind, placebo-controlled crossover trial. *Lancet Diabetes Endocrinol*. 2023;11(7):454-464.
27. Atila C, Mekhattu S, Murugesu R, *et al.* Plasma oxytocin levels in response to glucagon in patients with arginine vasopressin deficiency (central diabetes insipidus) and healthy controls. *Endocrine*. 2024;86:774-781.
28. Asla Q, Garrido M, Urgell E, *et al.* Oxytocin levels in response to CRH administration in hypopituitarism and hypothalamic damage: a randomized, crossover, placebo-controlled trial. *Sci Rep*. 2025;15(1):2360.
29. Galbiati F, Plessow F, Plummer L, *et al.* Sex-dependent increases in oxytocin levels in response to intravenous kisspeptin in humans. *Eur J Endocrinol*. 2025;192(3):K24-K29.
30. Arendt J, Aulinas A. Physiology of the pineal gland and melatonin. In: Feingold KR, Anawalt B, Blackman MR, *et al.*, eds. *Endotext [Internet]*. MDText.com, Inc.; 2000:1-25. [Updated October 30, 2022].
31. Wu YH, Zhou JN, Balesar R, *et al.* Distribution of MT1 melatonin receptor immunoreactivity in the human hypothalamus and pituitary gland: colocalization of MT1 with vasopressin, oxytocin, and corticotropin-releasing hormone. *J Comp Neurol*. 2006;499(6): 897-910.
32. Forsling ML. Diurnal rhythms in neurohypophysial function. *Exp Physiol*. 2000;85:179S-186S.
33. Juszczak M, Guzek JW. Hypothalamic and neurohypophysial vasopressin and oxytocin in melatonin-treated pinealectomized male rats. *J Pineal Res*. 1988;5(6):545-552.
34. Juszczak M, Stempniak B, Guzek JW. Melatonin, pinealectomy, and release of neurohypophysial hormones: in vitro studies. *J Pineal Res*. 1992;12(1):1-6.
35. Windle RJ, Luckman SM, Stoughton RP, Forsling ML. The effect of pinealectomy on osmotically stimulated vasopressin and oxytocin release and Fos protein production within the hypothalamus of the rat. *J Neuroendocrinol*. 1996;8(10):747-753.
36. Juszczak M, Guzek JW. The content of vasopressin and oxytocin in the hypothalamus and neurohypophysis of pinealectomized male rats. *Acta Physiol Pol*. 1983;34(1):41-46.
37. Ross MG, Leake RD, Stegner H, Ervin G, Fisher DA. Oxytocin release induced by melatonin in the ewe. *Dev Pharmacol Ther*. 1985;8(4):254-259.
38. Roszczyk M, Juszczak M. Forskolin-stimulated vasopressin and oxytocin release from the rat hypothalamo-neurohypophysial system in vitro is inhibited by melatonin. *Endokrynol Pol*. 2014;65(2):125-131.
39. Forsling ML, Wheeler MJ, Williams AJ. The effect of melatonin administration on pituitary hormone secretion in man. *Clin Endocrinol (Oxf)*. 1999;51(5):637-642.
40. Olcese J, Beesley S. Clinical significance of melatonin receptors in the human myometrium. *Fertil Steril*. 2014;102(2):329-335.
41. Medical A W. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;310(20):2191-2194.
42. Dixon JR, Jr. The international conference on harmonization good clinical practice guideline. *Qual Assur*. 1998;6(2):65-74.
43. Asla Q, Garrido M, Urgell E, *et al.* Data from: Supplementary Material of the "Melatonin as a possible stimulus to unmask an oxytocin deficient state in hypopituitarism and hypothalamic damage" study (NCT05319301). CORA Repositori de Dades de Recerca. <https://doi.org/10.34810/data2092>. Deposited 28 February 2025.
44. Beck AT, Steer RA, Brown GK. *Manual for the Beck Depression Inventory-II*. Psychological Corporation; 1996.
45. Beck AT, Steer RA, Brown GK, Sanz J, Vázquez C. *BDI-II: Inventario de Depresión de Beck-II: Manual*. Pearson Educación; 2011.
46. Vautier S. A longitudinal SEM approach to STAI data: two comprehensive multitrait-multistate models. *J Pers Assess*. 2004;83(2): 167-179.
47. Spielberger CD, Gorsuch R, Lushene RE. *Manual for the State-Trait Anxiety Inventory*. 7th ed. TEA Ediciones; 2008.
48. Spielberger CD, Gorsuch R, Lushene RE, Vagg PR, Jacobs GA. *Manual for the State-Trait Anxiety Inventory*. Consulting Psychologists Press; 1983.
49. Bagby RM, Taylor GJ, Parker JD. The Twenty-item Toronto Alexithymia Scale--II. Convergent, discriminant, and concurrent validity. *J Psychosom Res*. 1994;38(1):33-40.
50. Carver CS WT. Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: the BIS/BAS scales. *J Pers Soc Psychol*. 1994;67(2):319-333.
51. Rosen R, Brown C, Heiman J, *et al.* The Female Sexual Function Index (FSFI): a multidimensional self-report instrument for the assessment of female sexual function. *J Sex Marital Ther*. 2000; 26(2):191-208.
52. Rosen RC, Riley A, Wagner G, Osterloh IH, Kirkpatrick J, Mishra A. The international index of erectile function (IIEF): a multidimensional scale for assessment of erectile dysfunction. *Urology*. 1997; 49(6):822-830.
53. Hays RD, Sherbourne CD, Mazel RM. The RAND 36-item health survey 1.0. *Health Econ*. 1993;2(3):217-227.
54. McKenna SP, Doward LC, Alonso J, *et al.* The QoL-AGHDA: an instrument for the assessment of quality of life in adults with growth hormone deficiency. *Qual Life Res*. 1999;8(4):373-383.
55. Holt-Lunstad J, Birmingham WA, Light KC. Influence of a "warm touch" support enhancement intervention among married couples on ambulatory blood pressure, oxytocin, alpha amylase, and cortisol. *Psychosom Med*. 2008;70(9):976-985.
56. Pruessner JC, Kirschbaum C, Meinlschmid G, Hellhammer DH. Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocrinology*. 2003;28(7): 916-931.
57. Bojanowska E, Forsling ML. The effects of melatonin on vasopressin secretion in vivo: interactions with acetylcholine and prosta-glandins. *Brain Res Bull*. 1997;42(6):457-461.
58. Juszczak M, Guzek JW, Lewy A. The influence of melatonin on the content of vasopressin and oxytocin in the hypothalamus and neurohypophysis in euhydrated and dehydrated male rats. *J Pineal Res*. 1986;3(3):199-211.
59. Yasin SA, Costa A, Besser GM, Hucks D, Grossman A, Forsling ML. Melatonin and its analogs inhibit the basal and stimulated release of hypothalamic vasopressin and oxytocin in vitro. *Endocrinology*. 1993;132(3):1329-1336.
60. Juszczak M, Debeljuk L, Bartke A, Stempniak B. Melatonin inhibits oxytocin and vasopressin release from the neurointermediate lobe of the hamster pituitary. *Neuroreport*. 1995;6(18):2453-2456.
61. Evans JJ, Janmohamed S, Forsling ML. Gonadotrophin-releasing hormone and oxytocin secretion from the hypothalamus in vitro

- during pro-oestrus: the effects of time of day and melatonin. *Brain Res Bull.* 1999;48(1):93-97.
62. Yasin SA, Grossman A, Forsling ML. Diurnal variation in the effect of melatonin on neurohypophysial hormone release from the rat hypothalamus. *Brain Res Bull.* 1996;39(1):1-5.
 63. Kostoglou-Athanassiou I, Treacher DF, Wheeler MJ, Forsling ML. Melatonin administration and pituitary hormone secretion. *Clin Endocrinol (Oxf).* 1998;48(1):31-37.
 64. Juszczak M, Wolak M, Bojanowska E, Piera L, Roszczyk M. The role of melatonin membrane receptors in melatonin-dependent oxytocin secretion from the rat hypothalamo-neurohypophysial system—an in vitro and in vivo approach. *Endokrynol Pol.* 2016; 67(5):507-514.
 65. Chiodera P, Volpi R, Capretti L, Giuliani N, Caffarri G, Coiro V. Melatonin inhibits oxytocin response to insulin-induced hypoglycemia, but not to angiotensin II in normal men. *J Neural Transm (Vienna).* 1998;105(2-3):173-180.
 66. Buijs RM, Pevet P. Vasopressin- and oxytocin-containing fibres in the pineal gland and subcommissural organ of the rat. *Cell Tissue Res.* 1980;205(1):11-17.
 67. Masson-Pevet M, George D, Kalsbeek A, Saboureau M, Lakhdar-Ghazal N, Pevet P. An attempt to correlate brain areas containing melatonin-binding sites with rhythmic functions: a study in five hibernator species. *Cell Tissue Res.* 1994;278(1): 97-106.
 68. Mens WB, Andringa-Bakker EA, Van Wimersma Greidanus TB. Changes in cerebrospinal fluid levels of vasopressin and oxytocin of the rat during various light-dark regimes. *Neurosci Lett.* 1982; 34(1):51-56.
 69. Demaine C, Forsling ML, Kelestimur H, Stoughton RP. Effects of pinealectomy on daily rhythms of neurohypophysial hormone release in the rat. *J Physiol.* 1990;423:P12.
 70. Windle RJ, Forsling ML, Guzek JW. Daily rhythms in the hormone content of the neurohypophysial system and release of oxytocin and vasopressin in the male rat: effect of constant light. *J Endocrinol.* 1992;133(2):283-290.
 71. Forsling ML, Stoughton RP, Zhou Y, Kelestimur H, Demaine C. The role of the pineal in the control of the daily patterns of neurohypophysial hormone secretion. *J Pineal Res.* 1993;14(1):45-51.
 72. Kostoglou-Athanassiou I, Forsling ML. Effect of 5-hydroxytryptamine and pineal metabolites on the secretion of neurohypophysial hormones. *Brain Res Bull.* 1998;46(5):417-422.
 73. Brandi ML, Gebert D, Kopczak A, Auer MK, Schilbach L. Oxytocin release deficit and social cognition in craniopharyngioma patients. *J Neuroendocrinol.* 2020;32(5):e12842.
 74. Tabak BA, Leng G, Szeto A, *et al.* Advances in human oxytocin measurement: challenges and proposed solutions. *Mol Psychiatry.* 2023;28(1):127-140.
 75. Scantamburlo G, Hansenne M, Fuchs S, *et al.* Plasma oxytocin levels and anxiety in patients with major depression. *Psychoneuroendocrinology.* 2007;32(4):407-410.
 76. Frijling JL, van Zuiden M, Nawijn L, *et al.* Salivary oxytocin and vasopressin levels in police officers with and without post-traumatic stress disorder. *J Neuroendocrinol.* 2015;27(10):743-751.
 77. John S, Jaeggi AV. Oxytocin levels tend to be lower in autistic children: a meta-analysis of 31 studies. *Autism.* 2021;25(8):2152-2161.