

Emergent Carotid Artery Stenting in Patients with Acute Ischemic Stroke with Tandem Lesions: 1 year follow-up results from the SECURIS Study

Authors' names, academic degrees and affiliations

Garbiñe Ezcurra-Díaz^{1,2}; Pere Cardona³; Noelia Rodriguez-Villatoro⁴; Antonio Doncel-Moriano Cubero^{5,6}; Belen Flores-Pina⁷; Georgina Laia Figueras-Aguirre⁸; Isabel Fernández-Pérez⁹; Tomas Xuclà-Ferraron¹⁰; Francisco Purroy¹¹; Alan Flores¹²; Marina Guasch-Jiménez¹; Álvaro Lambea-Gil¹; Luis Prats-Sanchez¹; Anna Ramos-Pachón¹; Alejandro Martinez-Domeño¹; Joan Marti-Fabregas¹; Joan Miquel Fernández-Vidal¹; Sònia Abilleira¹³; Mercè Salvat-Plana¹⁴; Ana Núñez-Guillén³; Blanca Lara Rodríguez³; David Rodriguez-Luna¹⁵; David Hernandez¹⁵; Alejandro Rodríguez Vázquez^{5,6}; Andrea Cabero-Arnold^{5,6}; Alex Menéndez Albarracín⁷; David Canovas⁸; Pol Camps-Renom¹

1. Institut de Recerca Sant Pau (IR SANT PAU), Stroke Unit, Department of Neurology, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain

2. Department of Medicine, Universitat Autònoma de Barcelona, Barcelona, Spain

3. Stroke Unit, Department of Neurology, Hospital Universitari de Bellvitge, L'Hospitalet de Llobregat, Spain

4. Stroke Unit Department of Neurology, Hospital Universitari Vall d'Hebron, Barcelona, Spain

5. Stroke Unit, Department of Neurology, Hospital Clínic i Provincial, Barcelona, Spain

6. Fundació de Recerca Clínic Barcelona - Institut d'Investigacions Biomèdiques August Pi i Sunyer

7. Stroke Unit, Department of Neurology, Hospital Universitari Germans Trias i Pujol, Badalona, Spain

8. Stroke Unit Department of Neurology, Hospital Universitari Parc Taulí, Sabadell, Spain

9. Stroke Unit, Department of Neurology, Hospital del Mar, Barcelona, Spain

10. Stroke Unit, Department of Neurology, Hospital Universitari Josep Trueta, Girona, Spain

11. Stroke Unit, Department of Neurology, Hospital Universitari Arnau de Vilanova, Lleida, Spain

12. Stroke Unit, Department of Neurology, Hospital Universitari Joan XXIII, Tarragona, Spain

13. Stroke Programme, Health Ministry of Catalonia, Fundació TIC Salut Social, Barcelona, Spain

14. Stroke Programme, Health Ministry of Catalonia. Healthcare Quality and Evaluation Agency of Catalonia (AQuAS). CIBER Epidemiology and Public Health. Barcelona, Spain

Corresponding Author: Pol Camps-Renom

Abstract

Background and objectives: Management of carotid lesions during endovascular therapy (EVT) in patients with acute ischemic stroke (AIS) due to tandem lesions remains controversial, and long-term prognosis data are scarce. We evaluated 90-days and 1-year follow-up outcomes of emergent carotid artery stenting (eCAS) versus non-eCAS in these patients.

Methods: Data from a prospective mandatory population-based registry of Code Stroke in Catalonia (Spain) between 2017 and 2023 were analyzed. Patients with large vessel occlusion and ipsilateral $\geq 50\%$ atheromatous cervical internal carotid artery (ICA) stenosis undergoing EVT were identified. eCAS and non-eCAS groups were compared. The primary outcome was the shift in the modified Rankin Scale (mRS) score at 90-days and 1-year of follow-up, adjusted using Inverse Probability of Treatment Weighting. Stratified analyses in predefined key subgroups according to sex, age, degree of ICA stenosis, ASPECTS, baseline NIHSS and previous intravenous thrombolysis were performed. Secondary outcomes included successful recanalization (modified Thrombolysis In Cerebral Infarction [mTICI] $\geq 2b$), 90-day favorable functional outcome (mRS 0-2), hemorrhagic transformation (HT) and 90-day mortality.

Results: Among 578 patients included (mean age 70.2 years-old, 140 [24.2%] women), 344 patients (59.5%) underwent eCAS. eCAS was associated with a better functional outcome at 90 days (cOR for a 1-point less on the mRS score 1.47, 95% CI 1.20-1.82, $p < 0.001$) and 1 year of follow up (cOR of 1.47, 95% CI 1.16-1.85), $p = 0.001$), and with higher successful recanalization rates (OR 4.28, 95% CI 2.61-7.0, $p < 0.001$), without increasing HT risk or mortality. The benefit of eCAS was consistent in both moderate (50-69%) and severe ($\geq 70\%$) ICA stenosis subgroups. Among eCAS group, patients under dual antiplatelet therapy at 24 hours showed higher rates of favorable functional outcome (54.6% vs. 22.9%, $p < 0.001$). Potential treatment effect differences were observed for women and older patients.

Discussion: In this population-based study of patients with AIS due to tandem lesions, eCAS was associated with a better mid and long-term functional outcome without safety concerns, even in moderate ICA stenosis. Limitations include the observational design and possible unmeasured

84 confounders. Further studies are needed to explore the mechanisms underlying the differential
85 effects of eCAS in women and older patients.

86 ***Classification of Evidence:*** This study provides Class II evidence that in patients with acute
87 ischemic stroke and tandem lesions undergoing endovascular therapy, emergent carotid artery
88 stenting is associated with favorable functional outcome at 90 days compared to endovascular
89 therapy alone.

90 **Introduction**

91 Acute ischemic stroke (AIS) due to tandem lesions (TL) involving the extracranial Internal
92 Carotid Artery (ICA) represent up to 20% of all AIS undergoing endovascular therapy (EVT)¹.
93 TL are associated with lower recanalization rates, higher mortality and poorer functional
94 prognosis². Although EVT benefits are well established for the intracranial occlusion in these
95 patients¹, there is no consensus among stroke neurologists and neurointerventionalists regarding
96 the best treatment of the ICA lesion^{3,4}, with two mutually exclusive approaches: revascularization
97 via emergent carotid artery stenting (eCAS) during EVT, or conservative management and
98 deferred decision-making regarding carotid revascularization. While eCAS during EVT enables
99 immediate carotid revascularization and stabilization of the ruptured carotid plaque, it may
100 increase the procedure duration⁵ and may be technically difficult in some patients. eCAS also
101 requires early antithrombotic therapy to prevent acute stent occlusion⁶, potentially increasing the
102 risk of hemorrhagic transformation (HT)^{7,8}. Conversely, deferring revascularization might
103 increase the risk of early stroke recurrence and may negatively affect hemodynamics in the
104 affected hemisphere.

105 Some observational studies⁹⁻¹¹ and a recent meta-analysis¹² suggest that, compared to a
106 conservative strategy, eCAS may be associated with higher recanalization rates and a better
107 functional prognosis at 90 days of follow-up with no significantly increased risk of HT. eCAS
108 seems to be safe even when intravenous thrombolysis (IVT) has been priorly administered¹²⁻¹⁴.
109 However, data on long-term benefit of eCAS are scarce and there is heterogeneity among studies
110 in the definition of TL (thresholds for ICA stenosis may vary from 50%, 70%, 90% or complete

occlusion), in the outcome measurement, stroke etiologies included, and methods used to control for confounders.

In addition, most studies were conducted in highly specialized EVT-capable centers, which may reduce their external validity. Therefore, pending the publication of ongoing randomized trials^{15–19}, there is a need to confirm the potential benefit of eCAS in large population-based studies, with standardized follow-up and broad definition of TL. Moreover, information on treatment effect differences in key subgroups is scarce.

This study addresses the following research questions: 1) Does eCAS improve functional outcomes and remain safe in patients with AIS due to atherosclerotic TL? 2) Is the benefit of eCAS sustained at 1-year follow-up? 3) Is eCAS effective and safe across clinically relevant patient subgroups?

Methods

Study design and population

The Stenting Extracranial Carotid artery stenosis dURING endovascular treatment in patients with acute Ischemic Stroke (SECURIS) study was an observational study based on the Code Stroke Registry of Catalonia (CICAT), a region of Spain with a population near 8 million inhabitants. Study design, data collection, and reporting adhered to the STROBE statement guidelines²⁰. CICAT is a prospective, population-based, mandatory registry including data from all consecutive Code Stroke patients in Catalonia. For this study, patients who underwent EVT for AIS in any of the ten comprehensive stroke centers (CSC) that perform EVT in Catalonia between January 2017 and March 2023 were identified. Inclusion criteria were as follows: 1) age ≥ 18 years old; 2) previous modified Rankin Scale (mRS) score 0-3; 3) time from symptom onset or last time seen well (LTSW) ≤ 24 hours; 4) AIS with large vessel occlusion (LVO) of anterior circulation including terminal ICA (TICA), segments M1/M2 of the middle cerebral artery (MCA) or segment A1 of the anterior cerebral artery (ACA); 5) concomitant atheromatous stenosis of cervical ICA $\geq 50\%$; and 6) treatment with any EVT modality, with or without prior IVT administration.

Patients with lack of critical follow-up data (mRS score at 90 days and follow-up computed tomography (CT) within the first 36 hours after treatment) were excluded. Patients with non-atheromatous etiology of ICA stenosis (e. g., dissection) were also excluded, except for those with an indeterminate etiology due to the coexistence of atrial fibrillation and atheromatous ICA stenosis.

Standard Protocol Approvals, Registrations, and Patient Consents

The study protocol was approved by the Ethics Committee of Hospital de la Santa Creu i Sant Pau (IIBSP-SEC-2023-136) and the CICAT scientific committee. The need for written informed consent was waived due to the retrospective nature of the study.

Study variables

Collected directly from the CICAT registry were data as follows: demographic data (age, sex); relevant previous medical history (prior mRS score, hypertension, dyslipidemia, diabetes, ischemic heart disease, antithrombotic treatment, peripheral arteriopathy, previous stroke, smoking); index event characteristics and metrics (baseline National Institute of Health Stroke Scale (NIHSS) score, Alberta Stroke Program Early CT Score (ASPECTS), occlusion site, vital signs, time from onset to hospital arrival, door-to-needle and door-to-groin puncture times); treatment data (IVT, number of passes, EVT-associated complications (arterial perforation, artery dissection or distal embolism), general anesthesia use, final modified Thrombolysis In Cerebral Infarction [mTICI]²¹ score) and presence of HT in the first follow-up CT at 24-36 hours.

Additional data not included in the registry but collected from electronic health records were furnished anonymized by the staff of the 10 CSC: stroke etiology according to TOAST criteria²², degree of ICA stenosis (according to North American Symptomatic Carotid Endarterectomy Trial (NASCET) criteria²³ in the last angiographical series previous to stent decision), HT classification on the follow-up CT if not provided in the CICAT registry and antithrombotic regimen at 24 hours after EVT. The degree of ICA stenosis was determined by investigators of each center based on angiography reports or, when not specified, through a review of source imaging. Investigators were all stroke neurologists not blinded to the eCAS status. Decisions regarding eCAS, prior IVT, antithrombotic regimen, and initiation time were at the discretion of each practitioner. Since

during the study period tenecteplase was not an approved indication for AIS according to the Spanish Agency for Medicines and Medical Devices (AEMPS), it was not used in routine clinical practice. Accordingly, we assume that most of the patients received alteplase.

Study outcomes

The primary outcome was the shift on the mRS score at two time points: 90 days and 1 year of follow-up. At 90-days of follow-up, mRS score was assessed centrally through a structured telephone interview conducted by certified central evaluators who were blinded to the clinical information. The 1-year mRS score was assessed by the treating neurologist, not blinded to eCAS status, during routine follow-up, either in person or by telephone.

Pre-planned stratified analysis in key subgroups for the primary outcome included: sex, degree of ICA stenosis prior to stenting decision (50-69% and $\geq 70\%$), stroke severity according to baseline NIHSS score (mild: ≤ 10 points, moderate-severe: 11-20 points, very severe: >20 points), ASPECTS (0-5 points, 6-8 points, 9-10 points), age (<80 years-old and ≥ 80 years-old) and prior IVT.

Secondary efficacy outcomes were favorable functional outcome at 90 days and 1 year of follow-up defined by a mRS score 0-2 or 3 if the previous mRS was 3; and successful recanalization after EVT defined by final mTICI ≥ 2 b. Secondary safety outcomes were mortality at 90 days of follow-up and the presence of relevant hemorrhagic transformation (HT), defined as the presence of either symptomatic intracerebral hemorrhage (SICH) or any parenchymal hemorrhage (PH) defined according to the European Cooperative Acute Stroke Study-II (ECASS-II) criteria²⁴ in the 24-36 hours follow-up CT scan^{25,26}.

Statistical analysis

Categorical variables were reported using counts (n) and percentages (%), and quantitative variables were reported as means and standard deviations (SD) or medians and interquartile range (IQR) if non-normally distributed according to the Shapiro-Wilk test.

Univariable analyses using the χ^2 test for categorical variables and the Student-t test for continuous variables (or Wilcoxon Rank Sum test when a non-parametric test was required) were performed to compare differences between the eCAS and non-eCAS groups.

To explore the relationship between eCAS and the primary outcome, ordinal logistic regression was used to perform a shift analysis. mRS score was assessed using the full-scale range (from 0 to 6). An Inverse Probability of Treatment Weighting (IPTW) approach using a propensity score for adjusting for possible confounders influencing the decision regarding stent placement was applied. The propensity score was computed using a multivariable logistic regression model, accounting for the following pre-specified variables, selected for their potential association with both the individual decision to perform eCAS and the primary outcome: previous mRS score, age, atrial fibrillation, baseline NIHSS, ASPECTS, time from LTSW to groin puncture, previous IVT and the number of passes. This analysis was repeated in the two time points.

For the graphical depiction of treatment effect in key subgroups as previously described, a forest plot was constructed and a p for interaction was calculated.

Regarding secondary outcomes, favorable functional outcome at 90 days was assessed by dichotomizing the mRS score (as previously described), performing a multivariable logistic regression analysis also adjusted for the IPTW. Number needed to treat (NNT) for achieving a favorable functional outcome was calculated. Successful recanalization, mortality at 90 days of follow-up and relevant HT rates were compared between the eCAS and non-eCAS groups. A multivariable logistic regression analysis adjusted for the pre-specified variables (number of passes, previous IVT, time from LTSW to groin puncture, age, ASPECTS and baseline NIHSS score) was performed.

Additional exploratory analyses were performed in the eCAS group to assess the relationship between antithrombotic regimen at 24 hours from EVT and favorable functional outcome. Predictors of relevant HT in the eCAS group were also assessed.

Analyses were conducted using available data only; no imputation of missing data was performed.

A two-tailed p-value <0.05 was considered significant. All statistical analyses were performed using Stata version 17.0 (StataCorp, College Station, TX).

Data availability

Anonymized data not published within this article will be made available by request from any qualified investigator.

Results

Study population

Of the 6096 EVT performed in Catalonia during the study period, 5656 of them involved anterior circulation, and 937 were initially identified as TL on the first angiographical series. After excluding non-atherothrombotic etiologies and those patients who did not meet inclusion criteria, a total of 578 patients were finally included (Figure 1). Differences between patients included in the analysis and those excluded due to missing follow-up data are detailed in eTable1.

Patient demographic, clinical and radiological characteristics are summarized in Table 1. Mean age was 70.2 years-old (SD=10.8), and 140 patients (24.2%) were women. eCAS was performed in 344 patients (59.5%). Women were less represented in the eCAS group compared to the non-eCAS group (21.2% vs. 28.6%, $p=0.041$). eCAS and non-eCAS groups were similar in key prognostic variables such as age, previous mRS, systolic blood pressure (SBP), glycemia, baseline NIHSS score, ASPECTS, and LTSW-to-groin puncture time. Compared with the non-eCAS group, patients in the eCAS group had higher rates of atrial fibrillation (9.9% vs. 4.7%, $p=0.022$), received less frequently IVT (36.1% vs. 47.9%, $p=0.005$), underwent a smaller number of passes (1 vs. 2, $p<0.001$) and had higher rates of general anesthesia during the procedure (25.6% vs. 14.5%; $p<0.001$). Procedure duration and rates of arterial perforation, dissection and distal embolism during EVT were similar between both groups.

Primary outcome

The median mRS score at 90 days for all of the patients was 3 (IQR 2-5), being significantly lower in the eCAS group compared with the non-eCAS group (3 [1-5] vs. 3 [2-5]; $p = 0.016$). When applying IPTW, the distribution of propensity scores was well balanced between treatment groups, as evidenced by substantial overlap in the density plots (eFigure 1). After adjusting for the IPTW, eCAS was independently associated with better functional outcomes (cOR for a 1-point less in the mRS of 1.47, 95% CI 1.20 -1.82; $p<0.001$). Information of mRS score at 1 year of follow-up was available for 501 patients. The benefit of eCAS persisted at 1 year of follow-up (cOR of 1.47, 95% CI 1.16 – 1.85, $p = 0.001$) (Figure 2).

A graphical depiction of treatment effect in key subgroups is shown in Figure 3. No significant heterogeneity in the treatment effect of eCAS was observed, as assessed in the interaction analysis. However, potential differences on treatment effect were observed for women, older patients and lower ASPECTS. Since the degree of ICA stenosis and sex were not initially included in the IPTW (unlike the other variables used to define the strata), further stratified analyses were performed in these subgroups after adjusting for the IPTW. After adjustment, eCAS was associated also with better functional prognosis in moderate (50-69%) ICA stenosis at 90 days (cOR 3.70, 95% CI 1.49-9.09, $p = 0.04$) and at 1 year of follow-up (cOR 4.17, 95% CI 1.41-12.5, $p=0.01$). Contrariwise, after adjustment, a lack of benefit of eCAS persisted in women (eFigure 2).

Secondary outcomes

At 90 days of follow up, a favorable functional outcome was observed in 244 patients (42.2%): 45.9% in the eCAS group vs. 36.8% in the non-eCAS group, representing a risk ratio for a better functional outcome of 1.25, 95% CI 1.02-1.53; $p=0.028$. In the adjusted multivariable logistic regression analysis, eCAS was independently associated with a higher probability of a 90-day favorable functional outcome (OR 1.43, 95% CI 1.12-1.82; $p=0.004$). The NNT to achieve a favorable outcome was 11. Likewise, at 1 year of follow-up eCAS persisted independently associated with a higher probability of a favorable functional outcome (OR 1.35, 95% CI 1.04-1.76; $p=0.023$).

Final mTICI was available for 575 patients of the 578 included patients. Successful recanalization was observed in 483 patients (84%), 315 (91.7%) in the eCAS group vs. 170 (73.0%) in the non-eCAS group ($p<0.001$). This association remained significant after adjusting for the pre-specified variables (OR 4.28, 95% CI 2.61-7.01; $p<0.001$). Furthermore, this association remained significant when the analysis was repeated only in patients who had received ≥ 1 passes, in order to rule out unsuccessful treatments due to lack of access to the intracranial occlusion.

Relevant HT was observed in 87 patients (15.1%), of whom 51 cases (8.9%) met the criteria for sICH. According to the ECASS-II classification, HT subtypes were distributed as follows: 36 patients had PH1, 36 had PH2, 5 had HI1, 7 had HI2, and 3 had remote PH. A slight but non-

significant trend towards a greater risk of relevant HT was observed in the eCAS group compared to the non-eCAS group (58 (16.9%) vs 29 (12.5%), $p=0.142$).

At 90 days of follow-up, 121 patients had died (20.9%), 20.6% in the eCAS group and 21.4% in the non-eCAS group ($p=0.833$). Mortality rates at 1 year of follow-up were 24% in the eCAS group and 29.9% in the non-eCAS group ($p=0.200$).

Specific eCAS group analyses

The follow-up CT showed relevant HT in 58 (16.9%) of the 344 patients who underwent eCAS. In a multivariable logistic regression analysis, only higher baseline NIHSS scores, greater number of passes and arterial perforation during EVT independently predicted a relevant HT after eCAS (Table 2).

In the eCAS group ($n=344$), specific antithrombotic regimen at 24 hours post-EVT was available for 339 patients. Among them, 227 (67.0%) received dual antiplatelet therapy (DAPT), 64 (18.9%) single antiplatelet therapy (SAPT), 2 (0.6%) anticoagulation therapy only, 5 (1.5%) both anticoagulation and antiplatelet therapy, and 41 (12.1%) any antithrombotic agent. Lower ASPECTS ($p<0.001$) and higher NIHSS scores ($p=0.012$) were the only variables independently associated with DAPT delivery at 24 hours post-EVT. Neither atrial fibrillation nor prior IVT were associated with DAPT delivery at 24 hours post-EVT. Patients receiving anticoagulation therapy at 24 hours post-EVT were excluded from further analyses given their small representation within the study cohort ($n=7$). Of the remaining patients, those under DAPT at 24 hours post-EVT showed a higher proportion of favorable functional outcome at 90 days of follow-up (54.6% vs. 22.9%, $p<0.001$), a lower rate of relevant HT (5.1% vs. 13.8% $p=0.045$) and lower mortality (12.3% Vs 37.1% $p<0.001$) compared to patients receiving other regimens. In adjusted multivariable regression analyses, these associations remained statistically significant.

Overall, this study provides Class II evidence that in patients with AIS and TL undergoing EVT, eCAS is associated with favorable functional outcome at 90 days compared to EVT alone.

Discussion

In this population-based study, eCAS during EVT was associated with a better functional outcome in patients with AIS secondary to TL, with no increased risk of relevant HT. This benefit remained significant at 1 year of follow-up. Our findings support previously reported data on the potential benefit of eCAS in these patients⁹⁻¹² and add valuable information on the safety and effectiveness at long-term follow-up and in key subgroups.

Notably, after adjustment, the cOR of a favorable shift in the mRS score at 90 days in our study was 1.47, closely aligned with the cOR of 1.46 reported in 2023 by Farooqui M. et al⁹. This observation might be explained by the fact that both studies had similar inclusion/exclusion criteria, as well as similar adjustment, and increases the evidence of the potential size effect of eCAS in these patients. However, our study goes further in providing additional data, as it was conducted in a real-life scenario over an extended inclusion period and included all AIS admitted to the ten CSC that currently perform EVT in Catalonia, thereby reducing the risk of hospital selection bias and enhancing the external validity of our results.

Long-term follow-up data of patients with TL treated with eCAS are scarce. One recent observational study found eCAS as an independent predictor of favorable functional outcome at 1 year of follow-up, despite its small sample size and unicenter design²⁷. Our study reinforces this finding and provides more robust evidence supporting the potential long-term benefit of eCAS. Moreover, we observed that the magnitude of this benefit, the cOR, remained consistent at both 90 days and 1 year of follow-up, further strengthening its clinical relevance.

We observed some differences in baseline characteristics between patients who underwent eCAS and those who did not. First, a significantly higher rate of atrial fibrillation was observed in the eCAS group. This may be partly related to a higher prevalence of ischemic heart disease and hypertension, both established risks factors for atrial fibrillation²⁸. However, given the limited number of cases, a chance finding cannot be excluded. Still, as atrial fibrillation was included in the IPTW, we are confident potential confounding was controlled. Second, women were underrepresented in the eCAS group. This disparity may be partly attributed to the perception among treating physicians of a higher procedural risk in women, influenced by their historical underrepresentation in carotid revascularization trials and the higher rates of periprocedural

complications reported in this population²⁹. Regarding the potential benefit of eCAS in key subgroups, no significant modification of the treatment effect was observed across subgroups, although some findings warrant further consideration.

First, a potential benefit was observed not only in patients with severe ($\geq 70\%$) ICA stenosis, but also in patients with moderate (50-69%) ICA stenosis, although they represented only 15.9% of the study population. This finding is important because the definition of TL is highly heterogeneous between studies. It ranges from the strict 90% threshold used in the TITAN registry¹¹ to the ambiguous “extracranial internal carotid occlusive or stenotic lesion” definition used in the STRATIS registry¹⁰, suggesting an underrepresentation of patients with moderate stenosis in these registries. Our results not only reinforce the idea that eCAS should not be dismissed outright in patients with moderate (50%-69%) ICA stenosis, but also highlights the need for a more standardized definition of TL.

After adjustment, we observed a lack of benefit of eCAS in women with TL, although this observation might be unpowered due to the relatively small sample size of women included. Data on sex-specific influences during the acute phase of TL are scarce. Prior trials suggested a reduced benefit of elective carotid endarterectomy in women³⁰, as well as higher rates of restenosis or occlusion following elective CAS³¹. Several factors may contribute to these findings, including older age and greater comorbidity at stroke onset, smaller arterial diameters, and more stable plaque composition, which may reduce the potential benefit of eCAS³². Since these studies were not conducted in the setting of AIS, these observations can only be inferred to our population cautiously. Nevertheless, these sex-based differences highlight the need for tailored therapeutic approaches and further research to understand underlying sex-specific factors influencing treatment outcomes.

As previously reported^{12-14,33}, our results showed that IVT did not impact on the potential benefit of eCAS, and should not be avoided in patients who may be potential candidates to eCAS if there are not contraindications.

In patients with lower ASPECTS, the benefit of eCAS did not reach statistical significance. Recent randomized trials³⁴⁻³⁶ demonstrated the efficacy and safety of EVT in patients with large

ischemic core despite poorer functional outcomes than non-large core patients. Although patients with TL are underrepresented or even excluded from these trials, observational data suggest that patients with TL and a large core may benefit from EVT similarly to non-TL patients³⁷. Moreover, eCAS might provide a slight additional benefit compared to non-eCAS³⁸. However, these findings should be interpreted with caution due to the high risk of bias and the small number of eCAS patients included. Our study was probably underpowered to demonstrate a significant benefit of eCAS in patients with ASPECTS 6-8, and those with ASPECTS 0-5 were highly underrepresented (n=20), making it difficult to draw meaningful conclusions.

Elderly patients did not benefit significantly from eCAS, consistent with previous reports³⁹. Notably, in the context of elective carotid revascularization outside the acute phase, elderly patients who underwent CAS experienced poorer outcomes than those who underwent carotid endarterectomy, largely attributed to an increased risk of periprocedural stroke⁴⁰. The underlying mechanisms of this finding may involve not only their higher comorbidity burden but also anatomical factors (increased vascular tortuosity and calcifications) and a distinct plaque composition that increases their vulnerability to rupture during stenting⁴¹.

In addition to better functional outcome, patients who underwent eCAS showed a 4-time fold increased probability of successful recanalization after EVT, which is also a well-established predictor of favorable functional outcome in patients with TL⁴². However, this association may partly reflect a reverse causality: clinicians may have been more inclined to perform eCAS in cases where successful recanalization had already been achieved. Inversely, poorer recanalization rates in the non-eCAS group may be explained by failed attempts of intracranial recanalization or the impossibility of crossing the extracranial stenosis to perform intracranial thrombectomy.

Information on the EVT approach strategy for the eCAS group (anterograde vs. retrograde) was not available, so this limitation cannot be totally addressed. To partially address potential bias due to the latter scenario, we repeated the analysis in those patients who underwent at least one pass (thus the intracranial occlusion was reached), and the association remained consistent. Nevertheless, the lack of registration of failed intracranial thrombectomies, stenting attempts and EVT approach could have introduced bias into the results.

Several reasons may explain the better outcomes observed in the eCAS group, including improved cerebral hemodynamics of the affected hemisphere, which may lead to smaller infarctions or a reduced risk of early stroke recurrences, among others. Previous randomized trials⁴³⁻⁴⁵ have demonstrated the efficacy of non-emergent symptomatic carotid revascularization in reducing ipsilateral ischemic recurrences, although specific data on recurrence risk following eCAS is lacking.

Regarding antithrombotic therapy following eCAS, we observed better effectiveness and safety outcomes in patients receiving DAPT at 24 hours from EVT. This observation may partly be explained by the fact that these patients had more favorable baseline characteristics, which likely influenced physicians' decisions based on an assumed better prognosis or a decreased perceived risk of bleeding. Furthermore, lower rates of HT in early follow-up CT may have also influenced DAPT initiation based on current guidelines, which recommend DAPT as the optimal approach for elective CAS⁴⁶. While reverse causality cannot be excluded, previous studies have also identified early DAPT initiation as an independent predictor of good outcome in patients with AIS due to TL treated with eCAS⁴⁷. Thus, despite potential confounding, our findings suggest that DAPT seems to be safe, even when prior IVT has been administered.

Finally, we aimed to assess independent predictors of HT in patients receiving eCAS. The only variables that remained associated with relevant HT after the multivariable regression analysis were higher NIHSS score, higher number of passes and arterial rupture during EVT. Interestingly, neither previous IVT nor prior anticoagulation therapy were among them, supporting previous evidence that suggests that IVT should not be withheld in AIS patients with TL who are likely to require eCAS^{13,14}.

The main strength of our study is that it is based on a mandatory, population-based registry that consecutively includes all patients with AIS undergoing EVT in a region of 8 million inhabitants, which not only allowed the inclusion of a large number of patients, but also enhances the external validity of the results and its generalization to a real-life population. Furthermore, the primary outcome (mRS at 90 days of follow-up) was assessed blindly and centrally, thereby limiting observer bias. In addition, using the IPTW as the statistical adjustment method to control for

418 confounders significantly reduced the risk of selection bias in a procedure (eCAS) that is
419 potentially highly influenced by practitioner experience and even CSC internal protocols.

420 We acknowledge some limitations of this study. First, we cannot rule out selection bias due to the
421 exclusion of patients with missing follow-up data. As shown in eTable 1, while baseline
422 characteristics were generally similar, excluded patients underwent a higher number of passes and
423 had lower rates of successful recanalization, suggesting more complications during EVT that may
424 have influenced outcomes and the eCAS decision. Second, although it is based on a prospective
425 registry, some data were collected retrospectively by the participating centers, which may have
426 led to inherent bias. Specially, despite its close concordance with 90-day mRS score, 1-year mRS
427 score was not centrally assessed. Also, radiological image analyses of angiographies and follow-
428 up CT was not centralized nor blinded to eCAS status, potentially introducing inter-observer
429 variability and interpretive bias. Information on the follow-up image modality used (CT or
430 magnetic resonance imaging [MRI]) was not available, which might have influenced the
431 classification of HT. Nevertheless, MRI is exceptionally uncommon as follow-up imaging in our
432 clinical setting due to its higher cost and limited availability. CT is the standard practice, therefore,
433 the risk of bias introduced by this limitation is considered minimal. Third, although the IPTW was
434 employed to mitigate selection bias, decision-making as to stent placement and antithrombotic
435 regimen were ultimately at the discretion of the treating practitioner at each center, leading to a
436 wide array of treatment combinations that are challenging to adjust for accurately. For instance,
437 unmeasured factors, including vascular anatomy, collateral status, or other clinical considerations,
438 may have influenced the decision to perform eCAS. Finally, the absence of data on stroke
439 recurrence during follow-up results in a significant limitation. This limits our ability to assess the
440 potential safety concern of an increased rate of peri-procedural strokes in the eCAS group, as well
441 as the possible efficacy benefit of a reduced incidence of recurrent strokes during follow-up in
442 this group.

443 **Conclusions**

444 In patients with AIS due to TL, eCAS performed during EVT was found to be safe and associated
445 with a better functional outcome, even in a long-term follow-up. DAPT may be safe and

associated with a better functional outcome than other antithrombotic regimens. The benefit of eCAS seems consistent not only in patients with severe ($\geq 70\%$) ICA stenosis, but also in moderate (50-69%) stenosis, and further research is needed to assess potential sex-related or age-related differences on the benefit of emergent carotid revascularization.

References

1. Goyal M, Menon BK, Van Zwam WH, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: A meta-analysis of individual patient data from five randomised trials. *The Lancet*. 2016;387(10029):1723-1731. doi:10.1016/S0140-6736(16)00163-X
2. Rubiera M, Ribo M, Delgado-Mederos R, et al. Tandem internal carotid artery/middle cerebral artery occlusion: An independent predictor of poor outcome after systemic thrombolysis. *Stroke*. 2006;37(9):2301-2305. doi:10.1161/01.STR.0000237070.80133.1d
3. Jacquin G, Poppe AY, Labrie M, et al. Lack of Consensus Among Stroke Experts on the Optimal Management of Patients With Acute Tandem Occlusion. *Stroke*. 2019;50(5):1254-1256. doi:10.1161/STROKEAHA.118.023758
4. Zevallos CB, Farooqui M, Quispe-Orozco D, et al. Proximal Internal Carotid artery Acute Stroke Secondary to tandem Occlusions (PICASSO) international survey. *J Neurointerv Surg*. 2021;13(12):1106-1110. doi:10.1136/NEURINTSURG-2020-017025
5. Stampfl S, Ringleb PA, Möhlenbruch M, et al. Emergency Cervical Internal Carotid Artery Stenting in Combination with Intracranial Thrombectomy in Acute Stroke. *AJNR Am J Neuroradiol*. 2014;35(4):741. doi:10.3174/AJNR.A3763
6. Allard J, Delvoye F, Pop R, et al. 24-Hour Carotid Stent Patency and Outcomes After Endovascular Therapy: A Multicenter Study. *Stroke*. 2023;54(1):124-131. doi: 10.1161/STROKEAHA.122.039797
7. Heck D V., Brown MD. Carotid stenting and intracranial thrombectomy for treatment of acute stroke due to tandem occlusions with aggressive antiplatelet therapy may be associated with a high incidence of intracranial hemorrhage. *J Neurointerv Surg*. 2015;7(3):170-175. doi:10.1136/NEURINTSURG-2014-011224

8. Strbian D, Sairanen T, Meretoja A, et al. Patient outcomes from symptomatic intracerebral hemorrhage after stroke thrombolysis. *Neurology*. 2011;77(4):341-348. doi: 10.1212/WNL.0b013e3182267b8c
9. Farooqui M, Zaidat OO, Hassan AE, et al. Functional and Safety Outcomes of Carotid Artery Stenting and Mechanical Thrombectomy for Large Vessel Occlusion Ischemic Stroke With Tandem Lesions. *JAMA Netw Open*. 2023;6(3):E230736. doi:10.1001/JAMANETWORKOPEN.2023.0736
10. Jadhav AP, Zaidat OO, Liebeskind DS, et al. Emergent Management of Tandem Lesions in Acute Ischemic Stroke: Analysis of the STRATIS Registry. *Stroke*. 2019;50(2):428-433. doi:10.1161/STROKEAHA.118.021893
11. Papanagiotou P, Haussen DC, Turjman F, et al. Carotid Stenting With Antithrombotic Agents and Intracranial Thrombectomy Leads to the Highest Recanalization Rate in Patients With Acute Stroke With Tandem Lesions. *JACC Cardiovasc Interv*. 2018;11(13):1290-1299. doi:10.1016/j.jcin.2018.05.036
12. Zevallos CB, Farooqui M, Quispe-Orozco D, et al. Acute Carotid Artery Stenting Versus Balloon Angioplasty for Tandem Occlusions: A Systematic Review and Meta-Analysis. *J Am Heart Assoc*. 2022;11(2). doi:10.1161/JAHA.121.022335
13. Mujanovic A, Dobrocky T, Pfeilschifter W, et al. Value of intravenous alteplase before thrombectomy among patients with tandem lesions and emergent carotid artery stenting: A subgroup analysis of the SWIFT DIRECT trial. *Eur J Neurol*. 2024;31(6). doi:10.1111/ENE.16256
14. Anadani M, Spiotta AM, Alawieh A, et al. Emergent Carotid Stenting Plus Thrombectomy After Thrombolysis in Tandem Strokes: Analysis of the TITAN Registry. *Stroke*. 2019;50(8):2250-2252. doi:10.1161/STROKEAHA.118.024733
15. Zhu F, Hossu G, Soudant M, et al. Effect of emergent carotid stenting during endovascular therapy for acute anterior circulation stroke patients with tandem occlusion: A multicenter, randomized, clinical trial (TITAN) protocol. *International Journal of Stroke*. 2021;16(3):342-348. doi:10.1177/1747493020929948
16. Endovascular Acute Stroke Intervention - Tandem OCclusion Trial | ClinicalTrials.gov identifier: NCT04261478. Updated November 18, 2023. Accessed May 10, 2025. <https://clinicaltrials.gov/study/NCT04261478>

17. Maes L, Van Elk T, van der Meij A, et al. Carotid Artery Stenting during Endovascular treatment of acute ischemic Stroke (CASES) study protocol for a multicenter randomized clinical trial. *Eur Stroke J*. Published online February 16, 2025. doi:10.1177/23969873251319941
18. Proximal Internal Carotid Artery Acute Stroke Secondary to Tandem or Local Occlusion Thrombectomy Trial (PICASSO). ClinicalTrials.gov. identifier: NCT05611242. Updated December 5, 2024. Accessed May 10, 2025. <https://www.clinicaltrials.gov/study/NCT05611242>
19. Stent Implantation Versus Balloon Dilation for Acute Anterior Circulation Tandem Occlusion (START). ClinicalTrials.gov identifier: NCT05902000. Updated June 28, 2023. Accessed May 10, 2025. <https://www.clinicaltrials.gov/study/NCT05902000>
20. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. 2007;370(9596):1453-1457. doi:10.1016/S0140-6736(07)61602-X
21. Zaidat OO, Yoo AJ, Khatri P, et al. Recommendations on angiographic revascularization grading standards for acute ischemic stroke: A consensus statement. *Stroke*. 2013;44(9):2650-2663. doi: 10.1161/STROKEAHA.113.001972
22. Adams HP, Bendixen BH, Kappelle L Jaap, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. *Stroke*. 1993;24(1):35-41. doi:10.1161/01.STR.24.1.35
23. Barnett HJM. North American Symptomatic Carotid Endarterectomy Trial. Methods, patient characteristics, and progress. *Stroke*. 1991;22(6):711-720. doi:10.1161/01.STR.22.6.711
24. Hacke W, Kaste M, Fieschi C, et al. Randomised double-blind placebo-controlled trial of thrombolytic therapy with intravenous alteplase in acute ischaemic stroke (ECASS II). *Lancet*. 1998;352(9136):1245-1251. doi:10.1016/S0140-6736(98)08020-9
25. Toscano-Prat C, Martínez-González JP, Guasch-Jiménez M, et al. Asymptomatic parenchymal haemorrhage following endovascular treatment: Impact on functional

543 outcome in patients with acute ischaemic stroke. *Eur J Neurol*. 2024;31(2):e16112.
544 doi:10.1111/ENE.16112

545 26. Wang Y, Maeda T, You S, et al. Patterns and Clinical Implications of Hemorrhagic
546 Transformation After Thrombolysis in Acute Ischemic Stroke. *Neurology*.
547 2024;103(11). doi:10.1212/WNL.0000000000210020

548 27. Ferraù L, Giammello F, Tessitore A, et al. Predictors of Good Functional Outcome
549 in Patients with Tandem Occlusion After Revascularization Treatment: Single-
550 Center Experience with 12-Month Follow-Up. *World Neurosurg*. 2024;188:e382-
551 e389. doi:10.1016/J.WNEU.2024.05.120

552 28. Michaud GF, Stevenson WG. Atrial Fibrillation. *New England Journal of*
553 *Medicine*. 2021;384(4). doi:10.1056/NEJMCP2023658

554 29. Howard VJ, Lutsep HL, Mackey A, et al. Influence of sex on outcomes of stenting
555 versus endarterectomy: a subgroup analysis of the Carotid Revascularization
556 Endarterectomy versus Stenting Trial (CREST). *Lancet Neurol*. 2011;10(6):530-
557 537. doi:10.1016/S1474-4422(11)70080-1

558 30. Warlow C, Farrell B, Fraser A, Sandercock P, Slattery J. Randomised trial of
559 endarterectomy for recently symptomatic carotid stenosis: Final results of the MRC
560 European Carotid Surgery Trial (ECST). *Lancet*. 1998;351(9113):1379-1387.
561 doi:10.1016/S0140-6736(97)09292-1

562 31. Lal BK, Beach KW, Roubin GS, et al. Restenosis after carotid artery stenting and
563 endarterectomy: a secondary analysis of CREST, a randomised controlled trial.
564 *Lancet Neurol*. 2012;11(9):755-763. doi:10.1016/S1474-4422(12)70159-X

565 32. Gasbarrino K, Iorio D Di, Daskalopoulou SS. Importance of sex and gender in
566 ischaemic stroke and carotid atherosclerotic disease. *Eur Heart J*. 2022 Feb
567 10;43(6):460-473. doi:10.1093/eurheartj/ehab756

568 33. Cavalcante F, Treurniet K, Kaesmacher J, et al. Intravenous thrombolysis before
569 endovascular treatment versus endovascular treatment alone for patients with large
570 vessel occlusion and carotid tandem lesions: individual participant data meta-
571 analysis of six randomised trials. *Lancet Neurol*. 2025;24(4):305-315.
572 doi:10.1016/S1474-4422(25)00045-6

573 34. Costalat V, Jovin TG, Albucher JF, et al. Trial of Thrombectomy for Stroke with a
574 Large Infarct of Unrestricted Size. *New England Journal of Medicine*.
575 2024;390(18):1677-1689. doi:10.1056/NEJMOA2314063

35. Sarraj A, Hassan AE, Abraham MG, et al. Trial of Endovascular Thrombectomy for Large Ischemic Strokes. *New England Journal of Medicine*. 2023;388(14):1259-1271. doi:10.1056/NEJMOA2214403
36. Bendszus M, Bonekamp S, Herweh C, et al. Endovascular thrombectomy for acute ischaemic stroke with established large infarct: multicentre, open-label, randomised trial. *Lancet*. 2023;402(10414):1753-1763. doi:10.1016/S0140-6736(23)02032-9
37. Galecio-Castillo M, Farooqui M, Guerrero WR, et al. Endovascular Treatment of Patients With Acute Ischemic Stroke With Tandem Lesions Presenting With Low Alberta Stroke Program Early Computed Tomography Score. *J Am Heart Assoc*. 2024;13(22). doi:10.1161/JAHA.124.035977
38. Deliktas Y, Derraz I, Finitis S, et al. Clinical and safety outcomes of acute stenting plus thrombectomy for carotid tandem lesions with large ischemic core. *J Neurointerv Surg*. Published online January 25, 2025. doi:10.1136/JNIS-2024-021517
39. Huber C, Berberat J, Sassenburg R, et al. Age and Stroke Severity Matter Most for Clinical Outcome in Acute Arteriosclerotic Tandem Lesions. *Journal of Clinical Medicine* 2024, Vol 13, Page 2315. 2024;13(8):2315. doi:10.3390/JCM13082315
40. Voeks JH, Howard G, Roubin GS, et al. Age and outcomes after carotid stenting and endarterectomy: The Carotid Revascularization Endarterectomy Versus Stenting Trial. *Stroke*. 2011;42(12):3484-3490. doi:10.1161/STROKEAHA.111.624155
41. Van Lammeren GW, Reichmann BL, Moll FL, et al. Atherosclerotic plaque vulnerability as an explanation for the increased risk of stroke in elderly undergoing carotid artery stenting. *Stroke*. 2011;42(9):2550-2555. doi:10.1161/STROKEAHA.110.607382
42. Rodriguez-Calienes A, Galecio-Castillo M, Hassan A, et al. Association of Acute Carotid Stenting with Functional Outcome Mediated by Successful Reperfusion in Tandem Lesions: Mediation Analysis of a Multicenter Registry (P2-5.006). *Neurology*. 2024;102(17_supplement_1). doi:10.1212/WNL.0000000000206545
43. Collaborators* NASCET. Beneficial Effect of Carotid Endarterectomy in Symptomatic Patients with High-Grade Carotid Stenosis. *New England Journal of Medicine*. 1991;325(7):445-453. doi:10.1056/NEJM199108153250701

44. Warlow C, Farrell B, Fraser A, Sandercock P, Slattery J. Randomised trial of endarterectomy for recently symptomatic carotid stenosis: final results of the MRC European Carotid Surgery Trial (ECST). *The Lancet*. 1998;351(9113):1379-1387. doi:10.1016/S0140-6736(97)09292-1
45. Barnett HJ, Taylor DW, Eliasziw M, et al. Benefit of carotid endarterectomy in patients with symptomatic moderate or severe stenosis. North American Symptomatic Carotid Endarterectomy Trial Collaborators. *N Engl J Med*. 1998;339(20):274-275. doi:10.1056/NEJM199811123392002
46. Schirmer CM, Bulsara KR, Al-Mufti F, Haranhalli N, Thibault L, Hetts SW. Antiplatelets and antithrombotics in neurointerventional procedures: Guideline update. *J Neurointerv Surg*. 2023;15(11):1155-1162. doi:10.1136/JNIS-2022-019844
47. Havlíček R, Šaňák D, Černík D, et al. Predictors of Good Clinical Outcome After Endovascular Treatment for Acute Ischemic Stroke due to Tandem Lesion in Anterior Circulation: Results from the ASCENT Study. *Cardiovasc Intervent Radiol*. 2024;47(2):218-224. doi:10.1007/S00270-023-03649-X

Figure legends

Figure 1. Participant flowchart. Abbreviations: EVT, endovascular therapy; ICA, internal carotid artery.

Figure 2. Modified Rankin Scale score distributions at 90 days of follow-up (A) and 1 year of follow-up (B) for eCAS and non-eCAS groups. Abbreviations: eCAS, emergent carotid artery stenting; mRS, modified Rankin scale.

Figure 3. Stratified analyses of treatment effect of eCAS in key subgroups at 90 days of follow-up (A) and 1 year of follow-up (B). Abbreviations: ASPECTS, Alberta Stroke Program Early Tomography Score; CI, confidence interval, eCAS, emergent carotid artery stenting; ICA, internal carotid artery; mRS, modified Rankin scale; NIHSS, National Institute of Health Stroke Scale; OR, odds ratio.

639 Supplemental Figure 1. Density plot comparing the distribution of eCAS/non eCAS before and
640 after IPTW, demonstrating improved covariate balance.

641 Supplemental Figure 2. Adjusted ordinal regression analysis stratified by sex and stenosis degree
642 prior to stenting decision. Abbreviations: aOR, adjusted Odds Ratio; CI, confidence interval;
643 eCAS, emergent carotid artery stenting; ICA, internal carotid artery; mRS, modified Rankin scale.
644

Tables

Table 1. Patients clinical and radiological characteristics

	Total (n=578)	eCAS (n=344)	Non-eCAS (n=234)	p
Demographics				
Age, mean (SD)	70.2 (10.8)	70.5 (10.2)	69.8 (11.7)	0.438
Sex (female), n (%)	140 (24.2)	73 (21.2)	67 (28.6)	0.041
Previous medical history				
Previous mRS, mdn (IQR)	0 (0-1)	0 (0-1)	0 (0-1)	0.117
Current smoking, n (%)	206 (35.5)	131 (38.2)	74 (31.6)	0.106
Hypertension, n (%)	367 (63.6)	228 (66.5)	139 (59.4)	0.083
Diabetes, n (%)	143 (24.8)	92 (26.8)	51 (21.8)	0.170
Dyslipidemia	283 (49.1)	181 (52.8)	102 (43.6)	0.030
Ischemic heart disease, n (%)	81 (14.0)	59 (17.2)	22 (9.4)	0.008
Atrial fibrillation, n (%)	45 (7.8)	34 (9.9)	11 (4.7)	0.022
Previous ischemic stroke, n (%)	63 (10.9)	40 (11.7)	23 (9.8)	0.488
Previous antiplatelet therapy, n (%)	173 (29.9)	116 (33.7)	57 (24.4)	0.016
Previous anticoagulant therapy, n (%)	46 (8.0)	30 (8.7)	16 (6.8)	0.412
Previous statin therapy, n (%)	222 (38.4)	145 (42.2)	77 (32.9)	0.025
Stroke severity and acute phase variables				
Baseline NIHSS score, mdn (IQR)	16 (11-20)	16 (11-20)	17 (11-20)	0.945
SBP (mmHg), mdn (IQR)	150 (135-169)	153 (135-170)	150 (134-167)	0.156
Glycemia (mg/dL), mdn (IQR)	125 (107-155)	125 (106-155)	124 (108-155)	0.682
ASPECTS, mdn (IQR)	9 (8-10)	9 (8-10)	9 (8-10)	0.434
LTSW-groin puncture (hours), mdn (IQR)	4.3 (2.7-9.3)	4.5 (2.7-9.3)	4.1 (2.6-9.3)	0.705
IVT, n (%)	236 (40.8)	124 (36.1)	112 (47.9)	0.005
Endovascular treatment				
Number of passes, mdn (IQR)	1 (1-3)	1 (1-2)	2 (1-3)	<0.001
mTICI $\geq$ 2b, n (%)	483 (84.0)	313 (91.5)	170 (73.0)	<0.001
General anesthesia, n (%)	122 (21.1)	88 (25.6)	34 (14.5)	0.001
Arterial perforation*, n (%)	13 (2.7)	7 (2.4)	6 (3.1)	0.646
Artery dissection*, n (%)	28 (5.8)	20 (6.9)	8 (4.2)	0.207
Distal embolism*, n (%)	29 (6.0)	20 (6.9)	9 (4.7)	0.307
Groin puncture-recanalization (minutes), mdn (IQR)	62 (40-95)	65 (43-95)	59 (38-95)	0.123
TOAST classification				
Atherothrombotic, n (%)	525 (90.8)	317 (92.2)	208 (88.9)	0.182
Undetermined (two/more causes), n (%)	53 (9.2)	27 (7.9)	26 (11.1)	0.182

Legend: Abbreviations: ASPECTS, Alberta Stroke Program Early CT Score; CI, confidence interval; eCAS, emergent carotid artery stenting; IQR, interquartile range; IVT, intravenous thrombolysis; LTSW, last time seen well; mdn, median; mRS, modified Rankin Scale; mTICI: modified Thrombolysis In Cerebral Infarction; NIHSS, National Institute of Health Stroke Scale; OR, odds ratio; QRS, interquartile range; SBP, systolic blood pressure; SD, standard deviation.

*Information available for 480 patients

Table 2. Analysis of predictors of relevant hemorrhagic transformation in the eCAS group

Bivariable analysis	HT (n=58)	No HT (n=286)	p value
Age, mean (SD)	69.8 (9.7)	70.6 (10.3)	0.570
Sex (female), n (%)	12 (20.7%)	61 (21.3%)	0.914
Previous anticoagulation, n (%) ^a	5 (8.8%)	19 (6.8%)	0.590
Previous atrial fibrillation, n (%)	9 (15.5%)	25 (8.7%)	0.115
Glycemia, mdn (IQR) ^b	137 (117-178)	121 (105-151)	0.005
ASPECTS, mdn (IQR)	8 (7-10)	9 (8-10)	0.001
Baseline NIHSS, mdn (IQR)	19 (15-22)	16 (11-20)	<0.001
Prior IVT, n (%)	22 (37.9%)	102 (35.7%)	0.743
LTSW-groin puncture (hours), mdn (IQR)	4.8 (2.8-9.4)	4.4 (2.6-9.3)	0.391
Number of passes, mdn (IQR)	2 (1-3)	1 (1-2)	0.052
Arterial perforation, n (%) ^c	4 (7.8%)	3 (1.3%)	0.020
Successful recanalization, n (%) ^d	51 (87.9%)	362 (92.3%)	0.282
Multivariable logistic regression analysis	OR	95% CI	p value
Baseline NIHSS	1.08	1.02-1.15	0.009
Number of passes	1.35	1.09-1.68	0.007
Arterial perforation	9.18	1.76-47.93	0.009

Legend: Abbreviations: ASPECTS, Alberta Stroke Program Early Computed Tomography Score; CI, confidence interval; HT, hemorrhagic transformation; IVT, intravenous thrombolysis; LTSW, last time seen well; NIHSS, National Institute of Health Stroke Scale; OR, odds ratio; QRS, interquartile range; SD, standard deviation.

^aAvailable for 338 patients.

^bAvailable for 331 patients.

^cAvailable for 288 patients.

^dAvailable for 342 patients.