
















Country-specific key lifestyle factors and health outcomes for resource allocation in the general population: a network analysis across 29 countries

Jiaying Li^{1,2} , Daniel Yee Tak Fong¹ , Kris Yuet Wan Lok¹ , Janet Yuen Ha Wong³ , Mandy Man Ho¹ , Edmond Pui Hang Choi¹ , Vinciya Pandian² , Patricia M Davidson⁴ , Wenjie Duan⁵ , Marie Tarrant⁶ , Jung Jae Lee¹ , Chia-Chin Lin¹ , Oluwadamilare Akingbade^{7,8} , Khalid M Alabdulwahhab⁹ , Mohammad Shakil Ahmad¹⁰ , Mohamed Alborale¹¹ , Meshari A Alzahrani¹² , Anil S Bilimale¹³ , Sawitree Boonpatcharanon¹⁴ , Samuel Byiringiro³ , Muhammad Kamil Che Hasan¹⁵ , Luisa Clausi Schettini¹⁶, Walter Corzo¹⁷, Josephine M. De Leon¹⁸ , Anjanette S. De Leon¹⁸, Hiba Deek¹⁹ , Fabio Efficace²⁰ , Mayssah A El Nayal²¹, Fathiya El-Raey²² , Eduardo Enseldo-Carrasco²³ , Pilar Escotorin²⁴, Oluwadamilola Agnes Fadodun²⁵, Israel Opeyemi Fawole²⁶ , Yong-Shian Shawn Goh²⁷ , Devi Irawan²⁸, Naimah Ebrahim Khan²⁹ , Binu Koirala² , Ashish Krishna³⁰, Cannas Kwok³¹, Tung Thanh Le³², Daniela Giambruno Leal³³ , Miguel Ángel Lezana-Fernández³⁴ , Emery Manirambona³⁵ , Leandro Cruz Mantoani³⁶ , Fernando Meneses-González³⁴ , Iman Elmahdi Mohamed³⁷ , Madeleine Mukeshimana³⁸, Chinh Thi Minh Nguyen³², Huong Thi Thanh Nguyen³², Khanh Thi Nguyen³², Son Truong Nguyen³², Mohd Said Nurumal¹⁵ , Aimable Nzabonimana³⁹, Nagla Abdelrahim Mohamed Ahmed Omer⁴⁰ , Oluwabunmi Ogungbe² , Angela Chiu Yin Poon⁴¹ , Areli Reséndiz-Rodriguez⁴² , Busayasachee Puang-Ngern¹⁴ , Ceryl G Sagun¹⁸, Riyaz Ahmed Shaik¹⁰, Nikhil Gauri Shankar⁴³, Kathrin Sommer²⁰, Edgardo Toro³³, Hanh Thi Hong Tran³², Elvira L Urgel¹⁸, Emmanuel Uwiringiyimana³⁵ , Tita Vanichbuncha¹⁴, Naglaa Youssef⁴⁴ 

Affiliations are at the end of the article

Correspondence to:

Daniel Yee Tak Fong
School of Nursing, Li Ka Shing Faculty of Medicine, University of Hong Kong
5/F, Academic Building 3 Sassoon Road, Pokfulam, Hong Kong
China
dytfong@hku.hk

Background We aimed to identify the central lifestyle, the most impactful among lifestyle factor clusters; the central health outcome, the most impactful among health outcome clusters; and the bridge lifestyle, the most strongly connected to health outcome clusters, across 29 countries to optimise resource allocation for local holistic health improvements.

Methods From July 2020 to August 2021, we surveyed 16 461 adults across 29 countries who self-reported changes in 18 lifestyle factors and 13 health outcomes due to the pandemic. Three networks were generated by network analysis for each country: lifestyle, health outcome, and bridge networks. We identified the variables with the highest bridge expected influence as central or bridge variables. Network validation included nonparametric and case-dropping subset bootstrapping, and centrality difference tests confirmed that the central or bridge variables had significantly higher expected influence than other variables within the same network.

Results Among 87 networks, 75 were validated with correlation-stability coefficients above 0.25. Nine central lifestyle types were identified in 28 countries: cooking at home (in 11 countries), food types in daily meals (in one country), less smoking tobacco (in two countries), less alcohol consumption (in two countries), less duration of sitting (in three countries), less consumption of snacks (in five countries), less sugary drinks (in five countries), having a meal at home (in two countries), taking alternative medicine or natural health products (in one country). Six central health outcomes were noted among 28 countries: social support received (in three countries), physical health (in one country), sleep quality (in four countries), quality of life (in seven countries), less mental burden (in three countries), less emotional distress (in 13 countries). Three bridge lifestyles were identified in 19 countries: food types in daily meals (in one country), cooking at home (in one country), overall amount of exercise (in 17 countries). The centrality difference test showed the central and bridge variables had significantly higher centrality indices than others in their networks ($P < 0.05$).

Conclusions In 29 countries, cooking at home, less emotional distress, and overall amount of exercise emerged as common central lifestyle, health outcome, and bridge lifestyle factors, respectively. However, notable regional variations necessitate tailored interventions and resource allocations to effectively address unique local key variables and promote holistic health in each locale. The study's cross-sectional design and self-reported data may limit generalisability, emphasising the need for cautious interpretation and further longitudinal research.

Keywords global; across-country comparisons; lifestyle; health outcomes; network analysis

Lifestyle factors – including dietary risks, alcohol and tobacco consumption, sedentary behaviours, and physical inactivity – significantly influence health outcomes both immediately and over the long term. Short-term impacts manifest in changes to physical and psychological quality of life [1], sleep [2], and social well-being [3]. Conversely, persistent unhealthy lifestyle behaviours substantially elevate the risk for non-communicable diseases such as cardiovascular disease [4], diabetes, cancer, and chronic respiratory diseases [5], ultimately affecting life expectancy and mortality [6]. The global prevalence of non-communicable diseases underscores the critical importance of lifestyle in health promotion. However, implementing comprehensive lifestyle interventions that cover all lifestyle aspects at the population level is complex and resource-intensive, necessitating the identification of the most impactful lifestyle for targeted interventions and effective resource allocation for effective holistic health promotion, especially for those low- and middle-income countries with heavier burden and less resources.

Network analysis is a technique that examines the effects of one variable within its network cluster or its connections with another network cluster while considering all other within-network variables as confounders [7]. In network analysis theory, the central lifestyle and health outcome variables are the most impactful in their respective networks, offering the highest predictive value and potential for substantial improvements when modified. The bridge lifestyle, which is the lifestyle factor most strongly connected to the health outcome network, serves a similar function but impacts the entire health network [7]. The presence of most impactful lifestyle is supported by the interconnected relationships identified by previous studies within lifestyles, health outcomes, and between them. For instance, physical inactivity is associated with other lifestyle behaviours, such as smoking, alcohol consumption and an unhealthy diet [8,9] Additionally, physical inactivity affects various interim health outcomes, including mental health and sleep quality [10]. Moreover,

behaviours like smoking, drinking, and unhealthy eating also adversely impact other health outcomes, such as mental health and sleep quality [11]. Building on these intertwined relationships, a study utilised network analysis to identify central lifestyle factors, health outcomes, and bridge lifestyles using a sample drawn from various countries [12]. However, global approaches may overlook crucial cross-country variations in lifestyle patterns and cultural norms, limiting the effectiveness of interventions in specific regions. Therefore, identifying central lifestyle factors at the country level is essential to tailor interventions to local contexts.

The assumption that central lifestyle factors vary across countries is supported by multiple drivers, including socioeconomic and cultural influences, which affect the coexistence and strength of associations among lifestyle behaviours. For instance, limited access to healthy foods is more prevalent in low-income countries [13], which may result in insufficient food availability and reduced variety of food options. In contrast, high-income countries often experience higher rates of sedentary behaviours, such as prolonged sitting and increased screen time [14], due to factors like widespread use of digital technology, the prevalence of desk-based and office jobs, and greater reliance on automobiles for transportation. Additionally, cultural factors, such as the American preference for sweet flavours [15,16] and alcohol prohibition in some areas of Saudi Arabia due to religious and legal restrictions [15,16], can increase or decrease frequency of certain lifestyles. Hong Kong's high-density urban living may limit physical activity and increase stress levels [17]. In contrast, Rwanda's predominantly rural environment promotes higher levels of physical activity [18]. The connection between these differences and country-specific key variables can be explained by the pronounced presence of specific lifestyle factors and health outcomes. These factors may strengthen their associations with other coexisting lifestyles or health outcomes, thereby increasing their centrality within a network and making each country's key variables distinct. Identifying these diverse key lifestyles is crucial for developing effective, tailored local health interventions, yet remains poorly understood.

Therefore, this study aims to assess the interplay between lifestyle factors and health outcomes across 29 countries, and to identify the key variables (central lifestyle, central health outcome, and bridge lifestyle). We hypothesise that each country has distinct central lifestyle factors, health outcomes, and bridge lifestyles. The findings are expected to provide country-specific guidance for local resource allocation for effective holistic health improvements.

METHODS

Study design

This was an international cross-sectional study.

Study settings

This study spanned six regions defined by the World Health Organization, covering 29 countries after excluding Spain due to a limited sample size of 51. Recruitment was conducted primarily through online platforms to optimise reach and facilitate voluntary participation in preferred languages. The published protocol provides detailed information on the study design [19].

Participants and sample size

We recruited participants aged 18 and older using convenience and snowball sampling techniques. While these methods enabled efficient participant recruitment across diverse geographical locations, they also introduce potential selection biases, particularly affecting the representativeness of the socioeconomic and rural/urban diversity within the sample populations. Sample sizes for each network – lifestyle (18 nodes), health outcomes (13 nodes), and bridge (31 nodes) – were determined based on the maximum number of edges: 153, 78, and 465, respectively, adhering to the guideline of at least three participants per parameter [20]. Consequently, the required sam-

ple sizes for each network were 459, 234, and 1134 participants, respectively. Sample sizes across countries ranged 150–2238, with most meeting these criteria. Notably, centrality measures in network analysis can be considered reliable even if the sample size falls short of initial estimates, provided they pass the stability test via case-dropping subset bootstrap [20]. To ensure reliability, stability tests were conducted for all 87 networks, with only those networks that passed these tests being reported. We recognise the limitations of these sampling methods and the sample size in some countries, and their potential impact on the generalisability of our findings. To mitigate these effects, stability tests via case-dropping subset bootstrap were conducted for all 87 networks, and only those networks that passed these tests were reported.

Measures

Sociodemographics

The sociodemographic variables included in this study were gender, age, country of residence, marital status, highest educational level attained, and employment status.

Measurement of lifestyle factors and health outcomes

Item generation for the study questionnaire was developed for assessing a thorough literature review and collaborative discussions with public health professionals, nurses, and nutritionists. The chosen variables for lifestyle and health outcomes reflect dimensions commonly acknowledged in public health research, focusing on key lifestyle dimensions, including diet, physical activity, and substance use, as well as health outcomes across physical, psychological, social, and financial well-being. Specifically, 18 lifestyle factors include:

- dietary (types of food in daily meals, consumption of fruits and vegetables, frozen food/food products, snacks, soft drinks, juices, or other sugary drinks, meals at home, cooking at home, takeout food, traditional Chinese medicine or natural health products, and oral supplements/vitamins)
- exercise (frequency, duration, type, and overall amount)
- sedentary behaviours (sitting and screen time duration)
- addiction behaviours (smoking tobacco and alcohol consumption).

The 13 health-related outcomes include:

- physical well-being (weight, appetite, sleep quality, perceived physical health, and quality of life)
- psychological well-being (mental burden, emotional distress)
- social well-being (family disputes, social support provided, social support received, and social activities)
- financial well-being (income and economic burden).

The initial English version was created and face validity ensured through rigorous reviews that refined the questionnaire by enhancing clarity, eliminating redundancies, and structuring the flow of questions to facilitate ease of completion. To ensure cultural relevance, consultations with domain experts from various countries were conducted, followed by translations into multiple languages using a standard forward-backwards method. Lifestyle factors and health outcomes were assessed using single items, making formal psychometric evaluation inapplicable [19].

Participants rated the impact of COVID19 on 18 lifestyle factors and 13 general health outcomes using a 5-point Likert scale, with 1 indicating ‘substantially reduced’, 3 ‘no change’, and 5 ‘substantially increased’. To ensure consistent item alignment reflecting healthier lifestyles and improved health outcomes, certain items were reverse coded by appending a ‘less’ prefix.

Data collection

Data were collected via online survey platforms and supplemented with offline electronic PDF forms in areas with limited internet access. Both methods were standardised in content and structure to ensure comparability and maintain data quality across different collection environments. Rigorous pre-testing confirmed the equivalence of the two formats, allowing for consistent data collection across all regions. To encourage participation, one Hong Kong dollar was donated to the Red Cross for each completed questionnaire. The response rate was recorded at 75.2%.

Validation and rigor

To enhance internal validity, validation questions were incorporated into the questionnaire. Participants were asked ‘Where does the sun rise every day?’ Additionally, to ensure cultural relevance in Nigeria, the question ‘Where is your STATE capital?’ was used. Additionally, each country’s questionnaire was pilot-tested with a minimum of 10 participants prior to administration. The results from these pilot tests indicated that participant responses were within expected parameters and no confusion regarding the questions was reported; therefore, no modifications to the questionnaire were deemed necessary. Moreover, to mitigate social desirability bias, the survey was conducted anonymously. Furthermore, we performed data quality checks before analysis to remove any responses indicating misreporting or misunderstanding. These measures collectively aim to reduce potential biases and enhance the accuracy and reliability of our data set.

Statistical analysis

Data collected for our study were transferred to a Microsoft Excel database to facilitate stringent quality control. This included removing incomplete or duplicate responses to ensure data integrity. We also addressed data inconsistencies by discarding responses that did not align with validation questions, such as incorrect details regarding sunset times or capital cities, thereby enhancing the accuracy of our data set and the validity of our analysis. Subsequent analyses were performed using R Statistical Software (v4.1.1; R Core Team, Vienna, Austria). The network analyses encompassed four domains: topological overlap assessment, network estimation, stability testing, and the computation of centrality and bridge centrality indices.

Checking topological overlap

We used the ‘networktools’ R package’s ‘goldbricker’ function to identify unique variables to prevent artificial relationships among similar variables in a network. A significance threshold of 0.25 for inclusion and a significance level of 0.01 were applied [21].

Network estimation

For each country, three networks were analysed: one comprising all remaining lifestyle factors, another encompassing all remaining health outcomes, and a bridge network connecting the two. Pairwise associations were estimated using partial correlation analysis to control for confounding effects from other nodes. The least absolute shrinkage and selection operator was applied to minimise edges and nullify small correlations, while the extended Bayesian information criterion helped select the tuning parameter, resulting in a more interpretable and sparse network [22]. Network estimation and visualisation were performed using the ‘bootnet’ and ‘qgraph’ R packages, respectively [22]. In visual representations, nodes symbolised network items and edges their relationships, with edge thickness indicating the strength of association – blue for positive and red for negative associations.

Network stability

Network stability was evaluated using the ‘bootnet’ package [22]. Edge weight stability was assessed with 95% confidence intervals (CIs) derived from nonparametric bootstrapping; nar-

lower CIs indicated greater network credibility [22]. Centrality stability was measured by the correlation stability coefficient (CS-C), calculated via case-dropping subset bootstrap. A CS-C value above 0.25 was considered acceptable, with values above 0.5 being preferable [22].

Central node, centrality, bridge node, and bridge centrality

A central node significantly influences a network through its connections with other nodes, while bridge nodes connect closely to different clusters within another network [23,24]. In network analysis, betweenness centrality measures how frequently a node occurs on the shortest paths between other pairs, underscoring its intermediary role. Closeness centrality calculates a node's average distance to all others, indicating network accessibility. Strength centrality totals the edge weights connected to a node, showing its connectivity. Differently, expected influence centrality sums edge weights without absolute conversions, capturing dominant influences in networks with both positive and negative edges. Given our objective to identify nodes that exert significant cumulative effects, particularly in networks with negative interactions, we focus on expected influence centrality to pinpoint central nodes and to identify bridge nodes effectively [25]. The 'qgraph' package in R calculated the expected influence index, which accounts for both positive and negative edges to identify central nodes with the highest values. Bridge nodes were assessed using the 'networktools' package in R, which utilises the bridge expected influence (one-step) index that aggregates a node's edges to nodes in other networks [23]. Centrality differences between two nodes were evaluated using Wilcoxon tests with 1000 bootstrapped indices, generated through the 'bootnet' package in R. Multiple comparisons were adjusted for using Holm-Bonferroni corrections.

RESULTS

Demographics and item description

After screening 19 094 responses, 16 461 were eligible for analysis. Exclusions included blank or incomplete responses (1940), duplicates (116), inconsistent entries (450), responses from non-participating countries (126), and records missing age or gender information (1). **Table 1** outlines the sociodemographic details, means, standard deviations, and abbreviations of lifestyle factors and health outcomes, while Table S1 in the **Online Supplementary Document** displays the breakdown by country.

Non-redundant items remained for network analysis

Redundancy analysis via the Goldbricker method excluded L15 (**Table 1**), L16, and L17 for overlapping with L18, representing exercise habits more accurately. The final network comprised 15 lifestyle factors and 13 health outcomes with no additional redundancies.

Lifestyle networks across each country

Network stability

The bootstrapped 95% CI analysis confirmed the accuracy of edge weights in the lifestyle network, with narrow intervals suggesting reliable estimates (Figure S1 in the **Online Supplementary Document**). Besides, CS-C values for expected influence, derived from a case-dropping subset bootstrap, ranged 0.21–0.75 (**Table 2**, Figure S2 in the **Online Supplementary Document**). Excluding the United States, these values surpassed the 0.25 threshold, affirming the interpretability of 28 country-specific lifestyle networks.

Network structure and central lifestyle

Using Mainland China as an example, **Figure 1**, Panel A shows the lifestyle network with 49.5% of its 150 edges being nonzero, indicating strong connectivity. The strongest connections were

Table 1. Sociodemographic of respondents from 29 countries (n=16 461)

Variables	N/mean	%/SD	Variables	N/mean	%/SD
Age group			Macau	250	1.5
18–24	4845	29.4	Mainland China	667	4.1
25–29	2343	14.2	Malaysia	535	3.3
30–34	1930	11.7	Mexico	1016	6.2
35–39	1850	11.2	Nigeria	590	3.6
40–44	1417	8.6	Philippines	457	2.8
45–49	1147	7.0	Republic Of Sudan	538	3.3
50–54	971	5.9	Rwanda	150	0.9
55–59	663	4.0	Saudi Arabia	631	3.8
60–64	699	4.2	Singapore	237	1.4
≥65	596	3.6	South Africa	198	1.2
Gender			South Korea	2238	13.6
Female	10 317	62.7	Thailand	723	4.4
Male	6045	36.7	UK	212	1.3
Non-binary	99	0.6	USA	213	1.3
Marital status			Vietnam	419	2.5
Married/cohabitation/common-law	7244	44.0	Lifestyle factors (abbreviations)		
Separated/divorced/widowed	730	4.4	Food types in daily meals (L1)	3.01	0.87
Single	8486	51.6	Consumption of fruits and vegetables (L2)	3.15	0.90
Education			Less consumption of frozen food/food products (L3)	3.01	0.97
Primary or below	405	2.5	Less consumption of snacks (L4)	2.96	1.00
Secondary	2618	15.9	Less soft drinks/juices/other sugary drinks (L5)	2.80	1.03
Associate degree	1573	9.6	Having a meal at home (L6)	3.86	0.99
College	2246	13.6	Cooking at home (L7)	3.80	0.98
Bachelor	6495	39.5	Less eating takeout food (L8)	2.95	1.19
Graduate	2952	17.9	Taking alternative medicine or natural health products (L9)	2.89	0.91
Missing value	172	1.0	Taking oral supplements/vitamins (L10)	3.06	0.93
Employment			Less smoking tobacco (L11)	2.63	0.94
Job seeking	883	5.4	Less alcohol consumption (L12)	2.62	0.96
Laid off	170	1.0	Less duration of sitting (L13)	3.65	0.98
Not in workforce	990	6.0	Less duration of screen time (L14)	3.74	0.98
Retired	612	3.7	Frequency of exercise (L15)	2.81	1.08
Self-employed	1300	7.9	Duration of exercise (L16)	2.78	1.07
Student	4577	27.8	Type of exercise (L17)	2.77	1.02
Working (≥40 h/week)	5183	31.5	Overall amount of exercise (L18)	2.77	1.07
Working (1–39 h/week)	2746	16.7	Health outcomes (abbreviations)		
Country stay			Lose weight (H1)	3.21	0.90
Australia	639	3.9	Appetite (H2)	3.13	0.83
Brazil	553	3.4	Physical health (H3)	2.91	0.80
Burundi	369	2.2	Sleep quality (H4)	2.86	0.96
Canada	368	2.2	Quality of life (H5)	2.71	0.98
Chile	342	2.1	Less mental burden (H6)	3.40	1.07
Egypt	461	2.8	Less emotional distress (H7)	3.36	1.04
Guatemala	229	1.4	Family disputes (H8)	3.10	0.87
Hong Kong	2127	12.9	Social support provided (H9)	3.09	0.86
India	529	3.2	Social support received (H10)	2.97	0.84
Indonesia	482	2.9	Social activities (H11)	2.36	1.06
Italy	203	1.2	Income (H12)	2.65	0.93
Lebanon	440	2.7	Less economic burden (H13)	3.24	1.01
Libya	645	3.9			

H – health outcome, L – lifestyle, SD – standard deviation

between (the edge connected Lifestyle 7 and Lifestyle 6 (L7–L6), partial correlation coefficient=0.85), L11–L12 (partial correlation coefficient=0.75), and L13–L14 (partial correlation coefficient=0.66) (Table 2). The central node, cooking at home (L7), was confirmed by centrality index plots and bootstrapped difference tests ($P<0.05$) (Figure 1, Panel B). For the other 27 countries,

Table 2. Summary of network measures across 29 countries (n=16461)

Country	Lifestyle network						Health outcome network						Bridge network			
	Nonzero/ total edges = 105 (%)	Largest edge (PC-C)	Second largest edge (PC-C)	Third largest edge (PC-C)	CS-C value	Central lifestyle	Nonzero/ total edges = 78 (%)	Largest edge (PC-C)	Second largest edge (PC-C)	Third largest edge (PC-C)	CS-C value	Central health outcome	Nonzero/ total edges = 378 (%)	CS-C value	Bridge lifestyle	Bridge edge (PC-C)
Australia	68 (64.8)	L9-L2 (0.37)	L9-L1 (0.24)	L10-L6 (0.24)	0.59	L9	52 (66.7)	H13-H12 (0.64)	H5-H4 (0.34)	H6-H3 (0.26)	0.21	NA	124 (32.8)	0.13	NA	NA
Brazil	50 (47.6)	L7-L6 (0.63)	L14-L13 (0.61)	L5-L4 (0.40)	0.67	L4	21 (26.9)	H10-H9 (0.52)	H7-H6 (0.51)	H2-H1 (0.38)	0.75	H5	60 (15.9)	0.75	L18	L18-H3 (0.34)
Burundi	46 (43.8)	L2-L1 (0.43)	L7-L6 (0.26)	L4-L3 (0.24)	0.28	L1	46 (59.0)	H8-H7 (0.34)	H7-H6 (0.28)	H4-H3 (0.26)	0.52	H3	83 (22.0)	0.28	L18	L18-H4 (0.12)
Canada	35 (33.3)	L14-L13 (0.68)	L7-L6 (0.64)	L2-L1 (0.45)	0.67	L13	40 (51.3)	H7-H6 (0.75)	H10-H9 (0.58)	H5-H4 (0.31)	0.67	H7	64 (16.9)	0.52	L18	L18-H3 (0.25)
Chile	42 (40.0)	L14-L13 (0.67)	L7-L6 (0.48)	L10-L9 (0.36)	0.28	L4	25 (32.1)	H7-H6 (0.70)	H2-H1 (0.47)	H5-H4 (0.40)	0.75	H5	32 (8.5)	0.59	L18	L18-H3 (0.25)
Egypt	64 (61.0)	L7-L6 (0.68)	L12-L11 (0.66)	L14-L13 (0.49)	0.67	L6=L7	34 (43.6)	H2-H1 (0.40)	H7-H6 (0.39)	H8-H7 (0.37)	0.67	H4	125 (33.1)	0.00	NA	NA
Guatemala	16 (15.2)	L7-L6 (0.58)	L12-L11 (0.47)	L14-L13 (0.42)	0.59	L7	22 (28.2)	H7-H6 (0.47)	H2-H1 (0.38)	H5-H4 (0.30)	0.52	H4	23 (6.1)	0.44	L18	L18-H3 (0.24)
Hong Kong	72 (68.6)	L7-L6 (0.60)	L5-L4 (0.57)	L14-L13 (0.57)	0.75	L5	46 (59.0)	H7-H6 (0.51)	H10-H9 (0.37)	H5-H4 (0.29)	0.75	H7	169 (44.7)	0.67	L18	L18-H11 (0.14)
India	59 (56.2)	L12-L11 (0.76)	L7-L6 (0.60)	L14-L13 (0.58)	0.75	L5	26 (33.3)	H7-H6 (0.63)	H10-H9 (0.44)	H2-H1 (0.40)	0.67	H5=H7	85 (22.5)	0.59	L18	L18-H3 (0.23)
Indonesia	68 (64.8)	L12-L11 (0.71)	L7-L6 (0.54)	L14-L13 (0.47)	0.67	L11	53 (68.0)	H7-H6 (0.63)	H10-H9 (0.55)	H13-H12 (0.49)	0.75	H10	176 (46.6)	0.44	L18	L18-H5 (0.08)
Italy	25 (23.8)	L14-L13 (0.64)	L7-L6 (0.49)	L5-L4 (0.32)	0.21	NA	17 (21.8)	H5-H4 (0.41)	H7-H6 (0.40)	H2-H1 (0.36)	0.52	H5	37 (9.8)	0.13	NA	NA
Lebanon	59 (56.2)	L14-L13 (0.68)	L7-L6 (0.60)	L10-L9 (0.29)	0.52	L7	42 (53.9)	H7-H6 (0.54)	H2-H1 (0.53)	H8-H7 (0.37)	0.67	H7	127 (33.6)	0.20	NA	NA
Libya	64 (61.0)	L7-L6 (0.71)	L12-L11 (0.59)	L14-L13 (0.53)	0.67	L5=L6	37 (47.4)	H2-H1 (0.55)	H7-H6 (0.42)	H8-H7 (0.36)	0.75	H7	86 (22.8)	0.36	L1	L1-H2 (0.11)
Macau	43 (41.0)	L7-L6 (0.72)	L12-L11 (0.64)	L14-L13 (0.57)	0.52	L7	14 (18.0)	H7-H6 (0.60)	H2-H1 (0.35)	H5-H4 (0.35)	0.67	H7	23 (6.1)	0.00	NA	NA

Table 2. continued

Country	Lifestyle network						Health outcome network						Bridge network			
	Nonzero/ total edges = 105 (%)	Largest edge (PC-C)	Second largest edge (PC-C)	Third largest edge (PC-C)	CS-C value	Central lifestyle	Nonzero/ total edges = 78 (%)	Largest edge (PC-C)	Second largest edge (PC-C)	Third largest edge (PC-C)	CS-C value	Central health outcome	Nonzero/ total edges = 378 (%)	CS-C value	Bridge lifestyle	Bridge edge (PC-C)
Mainland China	52 (49.5)	L7-L6 (0.85)	L12-L11 (0.75)	L14-L13 (0.66)	0.75	L7	40 (51.3)	H7-H6 (0.67)	H5-H4 (0.47)	H2-H1 (0.42)	0.75	H5	144 (38.1)	0.44	L18	L18-H1 (0.15)
Malaysia	59 (56.2)	L12-L11 (0.78)	L7-L6 (0.62)	L14-L13 (0.61)	0.67	L11	46 (59.0)	H7-H6 (0.66)	H10-H9 (0.60)	H2-H1 (0.47)	0.75	H10	117 (31.0)	0.44	L18	L18-H3 (0.16)
Mexico	53 (50.5)	L14-L13 (0.68)	L7-L6 (0.65)	L12-L11 (0.54)	0.75	L4=L5	44 (56.4)	H7-H6 (0.59)	H2-H1 (0.46)	H5-H4 (0.38)	0.75	H5	107 (28.3)	0.75	L18	L18-H3 (0.34)
Nigeria	50 (47.6)	L12-L11 (0.66)	L7-L6 (0.60)	L14-L13 (0.45)	0.75	L7	39 (50.0)	H7-H6 (0.51)	H10-H9 (0.43)	H5-H4 (0.26)	0.75	H7	140 (37.0)	0.36	L7	L7-H5 (0.07)
Philippines	53 (50.5)	L7-L6 (0.74)	L12-L11 (0.66)	L14-L13 (0.60)	0.75	L7	44 (56.4)	H7-H6 (0.74)	H10-H9 (0.57)	H2-H1 (0.43)	0.75	H10	139 (36.8)	0.44	L18	L18-H3 (0.20)
Republic of Sudan	42 (40.0)	L12-L11 (0.62)	L7-L6 (0.61)	L14-L13 (0.32)	0.67	L7	34 (43.6)	H7-H6 (0.41)	H2-H1 (0.38)	H4-H3 (0.24)	0.67	H7	101 (26.7)	0.21	NA	NA
Rwanda	34 (32.4)	L14-L13 (0.55)	L2-L1 (0.53)	L7-L6 (0.50)	0.36	L12= L13=L7	7 (9.0)	H7-H6 (0.38)	H2-H1 (0.18)	H12-H11 (0.15)	0.44	H7=H6	22 (5.8)	0.00	NA	NA
Saudi Arabia	67 (63.8)	L7-L6 (0.67)	L14-L13 (0.58)	L10-L9 (0.34)	0.59	L7	43 (55.1)	H7-H6 (0.51)	H2-H1 (0.50)	H5-H4 (0.40)	0.75	H4	125 (33.1)	0.21	NA	NA
Singapore	36 (34.3)	L14-L13 (0.61)	L7-L6 (0.54)	L12-L11 (0.46)	0.28	L13	24 (30.8)	H7-H6 (0.70)	H10-H9 (0.45)	H5-H4 (0.33)	0.44	H7=H6	83 (22.0)	0.13	NA	NA
South Africa	19 (18.1)	L7-L6 (0.53)	L12-L11 (0.39)	L14-L13 (0.39)	0.44	L7	19 (24.4)	H7-H6 (0.58)	H10-H9 (0.43)	H2-H1 (0.41)	0.60	H6	54 (14.3)	0.00	NA	NA
South Korea	71 (67.6)	L7-L6 (0.71)	L5-L4 (0.52)	L14-L13 (0.52)	0.75	L4	39 (50.0)	H7-H6 (0.78)	H10-H9 (0.59)	H4-H3 (0.37)	0.75	H7	133 (35.2)	0.75	L18	L18-H3 (0.28)
Thailand	58 (55.2)	L12-L11 (0.74)	L14-L13 (0.52)	L7-L6 (0.43)	0.67	L5	44 (56.4)	H7-H6 (0.53)	H13-H12 (0.42)	H2-H1 (0.41)	0.75	H4	135 (35.7)	0.28	L18	L18-H11 (0.10)
U K	24 (22.9)	L7-L6 (0.58)	L14-L13 (0.49)	L5-L4 (0.24)	0.36	L4	12 (15.4)	H7-H6 (0.60)	H10-H9 (0.32)	H4-H3 (0.27)	0.44	H7	36 (9.5)	0.28	L18	L18-H3 (0.16)
USA	29 (27.6)	L7-L6 (0.63)	L14-L13 (0.57)	L10-L9 (0.42)	0.21	NA	26 (33.3)	H7-H6 (0.72)	H2-H1 (0.43)	H10-H9 (0.42)	0.60	H5	37 (9.8)	0.44	L18	L18-H3 (0.22)
Vietnam	49 (46.7)	L12-L11 (0.76)	L7-L6 (0.68)	L14-L13 (0.53)	0.75	L12	39 (50.0)	H10-H9 (0.73)	H7-H6 (0.56)	H8-H7 (0.38)	0.75	H7	145 (38.4)	0.52	L18	L18-H11 (0.12)

CS-C – correlation stability coefficient, H – health outcome, L – lifestyle, NA – not applicable, PC-C – partial correlation coefficient




Figure 1. Network structure and centrality difference test of lifestyles (Panels A–B), health outcomes (Panels C–D), and combined (Panels E–F) in mainland China. The abbreviations of nodes in Panels A, C, and E can be found in Table 1. In Panels B, D, and F, a grey cell indicates that there is no significant difference between the corresponding two variables. A dark cell indicates that there is a significant difference between the corresponding two variables at 5% level of significance. A white cell displays the value of the expected influence or bridge expected influence.

Figure S3 in the [Online Supplementary Document](#) presents the network structures, with non-zero edge percentages ranging from 15.2 in Guatemala to 68.6% in Hong Kong (Table 2). Figure S4 in the [Online Supplementary Document](#) contains centrality index plots and results of centrality bootstrapped difference tests. Table S2 in the [Online Supplementary Document](#) contains the partial correlation coefficients for all edges across all countries.

Health outcome networks across countries

Network stability

Narrow CIs across countries confirmed the precision of estimated edge weights (Figure S5 in the [Online Supplementary Document](#)). Besides, CS-C values ([Table 2](#); Figure S6 in the [Online Supplementary Document](#)), varied 0.21–0.75. Australia was the only country below the 0.25 threshold, resulting in 28 interpretable health outcome networks.

Network structure and central health outcome

For Mainland China ([Figure 1](#), Panel C) displays a network structure with 51.3% (40 out of 78) of edges being nonzero. The strongest connections were between H7–H6 ([Table 1](#)) and H5–H4 (both 0.67), and H2–H1 (0.42) ([Table 2](#)). Quality of life (H5) was identified as the most influential health outcome, confirmed by centrality index and bootstrapped difference tests ($P < 0.05$) ([Figure 1](#), Panel D). For the other 27 countries, Figure S3 in the [Online Supplementary Document](#) illustrates the network structures, with nonzero edge percentages ranging from 15.4 in the UK to 68.0% in Indonesia ([Table 2](#)). Figure S4 in the [Online Supplementary Document](#) contains centrality index plots and results of centrality bootstrapped difference tests. Table S3 in the [Online Supplementary Document](#) provides the partial correlation matrix for all edges across all networks.

Bridge networks across countries

Network stability

Figure S7 in the [Online Supplementary Document](#) illustrates the precision of estimated edge weights for each country with narrow CIs, indicating high accuracy. Besides, CS-C values for bridge networks varied between 0.00–0.75 ([Table 2](#); Figure S8 in the [Online Supplementary Document](#)). Ten countries, including Australia, Egypt, Italy, Lebanon, Macau, Republic of Sudan, Rwanda, Saudi Arabia, Singapore, and South Africa, fell below the 0.25 threshold, yielding 19 interpretable bridge networks.

Network structure and bridge lifestyle

[Figure 1](#), Panel E shows Mainland China's network with 144 of 378 edges being nonzero. The overall amount of exercise (L18) emerged as the most influential bridge lifestyle, significantly outperforming other nodes as shown by centrality bootstrapped difference tests ($P < 0.05$) ([Figure 1](#), Panel F). The corresponding bridge edge is L18–H1 (lose weight). For the other 18 countries, Figure S3 in the [Online Supplementary Document](#) displays network structures, with nonzero edge percentages ranging from 6.1 in Guatemala to 46.6% in Indonesia ([Table 2](#)). Figure S4 in the [Online Supplementary Document](#) contains centrality index plots and results of centrality bootstrapped difference tests. Table S4 in the [Online Supplementary Document](#) contains the partial correlation matrix for all networks.

Summary of the central and bridge variables in networks across 29 countries

Nine types of central lifestyle were identified among 28 countries:

- cooking at home (L7) in Guatemala, Lebanon, Macau, Mainland China, Nigeria, Philippines, Republic of Sudan, Rwanda, Saudi Arabia, South Africa, and Egypt
- food types in daily meals (L1) in Burundi
- less smoking tobacco (L11) in Indonesia and Malaysia
- less alcohol consumption (L12) in Rwanda and Vietnam
- less duration of sitting (L13) in Canada, Rwanda, and Singapore

- less consumption of snacks (L4) in Brazil, Chile, Mexico, South Korea, and UK
- less sugary drinks (L5) in Hong Kong, India, Libya, Mexico, and Thailand
- having a meal at home (L6) in Egypt and Libya
- taking alternative medicine or natural health products (L9) in Australia.

Six types of central health outcome were identified among 28 countries:

- social support received (H9) in Indonesia, Malaysia, and Philippines
- physical health (H3) in Burundi
- sleep quality (H4) in Egypt, Guatemala, Saudi Arabia, and Thailand
- quality of life (H5) in Brazil, Chile, India, Italy, Mainland China, Mexico, and USA
- less mental burden (H6) in Rwanda, Singapore, and South Africa
- less emotional distress (H7) in Canada, Hong Kong, India, Lebanon, Libya, Macau, Nigeria, Republic of Sudan, Rwanda, Singapore, South Korea, UK, and Vietnam.

Three types of bridge lifestyle were identified among 19 countries:

- food types in daily meals (L1) in Libya
- cooking at home (L7) in Nigeria
- overall amount of exercise (L18) in Brazil, Burundi, Canada, Chile, Guatemala, Hong Kong, India, Indonesia, Mainland China, Malaysia, Mexico, Philippines, South Korea, Thailand, United Kingdom, United States, and Vietnam.

Figure 2 visually summarises the distribution of central lifestyles, health outcomes, and bridge lifestyles across all participating countries.

DISCUSSION

This study employed network analysis to examine complex relationships between lifestyle factors and health outcomes across 29 countries, pinpointing key variables for use as potential predictive markers or intervention targets. We identified four central lifestyle categories: diet-related behaviours, natural health product usage, substance use, and sedentary habits; and four key health outcomes: physical health, emotional distress, social support, and quality of life. Additionally, three bridge lifestyle factors – food types in daily meals, home cooking, and overall exercise amount – were recognised as focal points for targeted resource allocation to enhance network-wide health outcomes, given the established causal relationship between lifestyle and health. Modifying central lifestyle factors can enhance all within-network lifestyles and improve long-term health outcomes. Central health outcomes highlight opportunities for beyond-lifestyle interventions, such as better health care service utilisation. Bridge lifestyles offer specific pathways for optimising overall health through targeted adjustments. Shared central or bridge variables across regions highlight the potential for collaborative strategies that leverage common strengths and address shared challenges.

In 28 lifestyle networks, four central lifestyle categories emerged: diet-related behaviours, natural health product usage, substance use, and sedentary habits, reflecting complex cultural, economic, and public health interactions. Specifically, Australia's focus on natural health products as a central lifestyle, likely driven by strong cultural acceptance and a preference for alternative health practices [26], enhances preventive lifestyles such as diet and exercise. In contrast, central alcohol consumption in Indonesia and Malaysia is influenced by significant binge drinking in Malaysia and rising harm from illicit home-produced alcohol in Indonesia [27,28]. Similarly, tobacco consumption was prominent in Rwanda and Vietnam, likely due to cultural norms and high smoking prevalence [29,30]. Reduced sitting time, significant in Rwanda and Singapore, corresponds with high urbanisation and pandemic-related increases in sedentary behaviour [31,32].




Figure 2. Central lifestyles, central health outcomes, and bridge lifestyles by country.

Moreover, diet-related behaviours such as specific food choices, reduced consumption of snacks and sugary drinks, home cooking, and having meals at home are pivotal in Rwanda and 23 other countries. Diet's centrality within the lifestyle network may be explained by its integral role in daily life and its deep connections to cultural and economic conditions [33,34], along with its coexistence with other lifestyle factors through behavioural and psychological mechanisms, such as self-regulation and discipline. These central behaviours not only predict overall lifestyle patterns but also serve as catalysts for broader modifications, suggesting that interventions targeting these areas could significantly influence entire lifestyle networks. Globally, effective strategies to address these dietary challenges could include enhancing nutritional education and awareness, improving food accessibility and affordability, and subsidising healthy foods or taxing unhealthy ones to leverage diet's centrality for holistic improvements in overall healthy lifestyles.

In 28 health outcome networks, we identified four primary categories of central health outcomes: physical health (including sleep quality), emotional distress, social support, and overall quality of life. Social support received was central in Indonesia, Malaysia, and the Philippines, reflecting their collective cultural values [35,36]. Physical health was prominent in Burundi, highlighting direct concerns with bodily well-being, while sleep quality was central in Egypt, Guatemala, Saudi Arabia, and Thailand, possibly due to hot climates affecting sleep [37]. Emotional distress or mental burden emerged as central health concerns in countries including Canada, Hong Kong,

India, Lebanon, Libya, Macau, Nigeria, Republic of Sudan, Rwanda, Singapore, South Africa, South Korea, UK, and Vietnam, driven by diverse factors. Specifically, political turmoil and economic instability significantly impacted Lebanon, Libya, Sudan, and Nigeria [38], while rapid urbanisation and economic stress affected Singapore, Hong Kong, South Korea, Macau, and others [39]. Quality of life, a multidimensional outcome encompassing physical health, mental health, and social relationships [40], was central in Brazil, Chile, India, Italy, Mainland China, Mexico, and the USA. This highlights the varied needs for health outcome interventions in each country-targeting physical, psychological, and social factors beyond lifestyle changes-to effectively improve holistic health.

In 19 countries with an interpretable bridge network, our study identified three key bridge lifestyles significantly affecting overall health outcomes: daily food types in Libya, home cooking in Nigeria, and exercise in other nations. In the context of network analysis, bridge lifestyle variables serve as critical connectors between the lifestyle network and the health outcomes network. These variables exhibit strong influence within the lifestyle network while simultaneously linking strongly to health outcome networks. Targeting these bridges in interventions can produce widespread, synergistic benefits by improving both the variables themselves and associated lifestyle behaviours, leading to enhanced overall health outcomes. Libya's reliance on specific meal types, influenced by its desert climate, heavy reliance on food imports, and political instability, highlights the critical role of dietary choices in health outcomes [41,42]. In Nigeria, home cooking may influence health through nutritional control over diets and economic relief via cost-effectiveness [43,44]. All remaining 17 countries or regions had exercise as their bridge lifestyle. The well-documented benefits of exercise on mental and physical health may explain this consistency across various nations [45], suggesting that strategic enhancements in exercise accessibility could lead to substantial health improvements. Globally, effective strategies to improve exercise accessibility include enhancing public infrastructure, implementing workplace wellness programmes, promoting community initiatives, integrating physical activity into urban planning, conducting education campaigns, ensuring inclusivity, partnering with health care providers, leveraging digital platforms, and supporting research and monitoring. Given the well-established causal relationship between lifestyle factors and health outcomes, targeted resource allocation and intensified interventions on bridge lifestyles are crucial for optimising these modifications across diverse settings.

This study identifies target points requiring increased resources and targeted interventions but does not specify which kinds of strategies are most effective. It assumes countries will adopt context-proven strategies. While comprehensive interventions for all lifestyle factors are ideal, they may only be feasible in resource-rich settings. Network theory suggests that prioritising central lifestyles, key health outcomes, and bridging lifestyles achieves greater health improvements than addressing other factors alone, which is particularly useful for resource-limited settings. For example, high-income Macau, middle-income mainland China, and low-income Sudan all identify home cooking as a central lifestyle factor. In resource-rich Macau, allocating substantial resources to home cooking can enhance other lifestyles, while additional resources can directly target other factors. In low-resource Sudan, dedicating a higher proportion of resources to home cooking can maximise impact. Similarly, targeting central health outcomes and bridging lifestyles can holistically improve health outcomes. Policymakers should design cost-effective programmes focused on these key variables and adopt context-specific interventions.

Limitations

Several limitations were noted. First, our online recruitment might cause selection bias, under-representing individuals with low socio-economic status and limited digital literacy, possibly skewing the relationships between lifestyle and health outcomes. Future studies should use stratified sampling and offline data collection to improve representativeness and validate findings. Second, our reliance on self-reported data subject to recall bias, social desirability bias, and inaccurate self-assessment, potentially distorting the observed relationships between variables in the

network. Although mitigated by validation questions and strict data quality controls, these limitations necessitate caution when interpreting the network structures and their implications. Future research should incorporate more accurate assessments, such as wearable technology or 24-hour recall, to enhance validity. Third, methodological limitations inherent in network analysis, such as assumptions of sparsity and potential overfitting, may affect the stability and generalisability of the identified network structures. The network estimation techniques rely on statistical models that may not fully capture the complex interplay between variables, and small changes in data can lead to different network configurations. Therefore, our network analysis results should be interpreted with consideration of these methodological constraints. Fourth, partial correlations within networks were based on individual perceptions of lifestyle and health outcome changes driven by the external factor of the COVID-19 pandemic. Although we anticipate these changes will revert in equal magnitude but opposite direction, preserving the relationships between variables post-pandemic, the findings require future validation during non-pandemic periods. Fifth, the cross-sectional design of our study during the COVID-19 pandemic limits our findings to associative insights and precludes the analysis of temporal network dynamics. By measuring outcomes relative to pre-pandemic conditions rather than their current status, the study's applicability to the post-pandemic period may be constrained. However, if we posit a rebound effect of equal magnitude and opposite direction, the findings could retain their relevance in a post-pandemic context. To substantiate and expand upon these insights, we recommend conducting longitudinal studies and randomised controlled trials post-pandemic. Lastly, small sample sizes in several countries limited network interpretability and robust comparisons, especially concerning bridge lifestyles. Increasing sample sizes in future research could address this issue.

CONCLUSIONS

This study delineated central lifestyle, health outcome, and bridge lifestyle variables across 29 countries. Modifications in these identified variables hold the potential to significantly improve holistic lifestyle and holistic health outcomes, as well as holistic health outcomes via lifestyle, respectively. Their substantial impact supports the need for targeted interventions and increased resource allocation to achieve effective, comprehensive health improvements. The variation in these variables across different countries-shaped by diverse cultural, socio-demographic, political, and economic factors-emphasises the importance of customised strategies that exploit the unique effects of central or bridge lifestyles. Additionally, the similarities observed among some countries suggest opportunities for collaborative initiatives to promote global health.

¹School of Nursing, Li Ka Shing Faculty of Medicine, University of Hong Kong, Hong Kong SAR, China

²School of Nursing, Johns Hopkins University, Baltimore, Maryland, USA

³School of Nursing and Health Studies, Hong Kong Metropolitan University, Hong Kong SAR, China

⁴Vice-Chancellor and Principal, University of Wollongong, Wollongong, Australia

⁵East China University of Science and Technology, Department of Social Work, Shanghai, China

⁶School of Nursing, The University of British Columbia, Kelowna, British Columbia, Canada

⁷The Nethersole School of Nursing, The Chinese University of Hong Kong, Hong Kong, China

⁸Institute of Nursing Research, Osogbo, Osun State, Nigeria

⁹College of Medicine, Majmaah University, Al Majmaah, Saudi Arabia

¹⁰Department of Family & Community Medicine, College of Medicine, Majmaah University, Majmaah, Saudi Arabia

¹¹Department of Internal Medicine, Al-Azhar University, Cairo, Egypt

¹²Department of Urology, College of Medicine, Majmaah University, Al Majmaah, Saudi Arabia

¹³School of Public Health, JSS Medical College, JSS AHER, Mysuru, India

¹⁴Department of Statistics, Chulalongkorn Business School, Bangkok, Thailand

¹⁵Kulliyyah of Nursing, International Islamic University, Kuantan, Malaysia

¹⁶Italian Association against Leukaemia, Lymphoma and Myeloma, Rome, Italy

- ¹⁷Diálogos Guatemala, Guatemala, Guatemala
- ¹⁸School of Nursing, Centro Escolar University, Manila, Philippines
- ¹⁹Nursing Department, Faculty of Health Science, Beirut Arab University, Lebanon
- ²⁰Data Centre and Health Outcomes Research Unit, Italian Group for Adult Hematologic Disease, Rome, Italy
- ²¹Department of Psychology, Beirut Arab University, Beirut, Lebanon
- ²²Department of hepatogastroenterology and infectious diseases, Damietta faculty of medicine, Al-Azher University, Kairo, Egypt
- ²³Ergonomics Research Centre, University of Guadalajara, Jalisco, Mexico
- ²⁴Laboratory of Applied Prosocial Research, Department of Basic, Developmental and Educational Psychology, Autonomous University of Barcelona, Barcelona, Spain
- ²⁵Faculty of Health Sciences, University of Lethbridge, Lethbridge, Alberta, Canada
- ²⁶Faculty of Nursing, Ladoke Akintola University of Technology, Ogbomosho, Nigeria
- ²⁷Alice Lee Centre for Nursing Studies, National University of Singapore, Singapore, Singapore
- ²⁸School of Nursing, Wijaya Husada Health Institute, Bogor, Indonesia
- ²⁹Department of Optometry, University of Kwazulu-Natal, Durban, South Africa
- ³⁰Ecove, Ghaziabad, India
- ³¹School of Nursing, Paramedicine and Health Care Science, Charles Sturt University, New South Wales, Australia
- ³²Nam Dinh University of Nursing, Nam Dinh, Vietnam
- ³³Pontificia Universidad Católica de Valparaíso, School of Social Work, Valparaíso, Chile
- ³⁴Research Department, National Commission for Medical Arbitration, Mexico, Mexico
- ³⁵College of Medicine and Health Sciences, University of Rwanda, Kigali, Rwanda
- ³⁶Department of Physiotherapy, Presidente Prudente, Faculty of Science and Technology, São Paulo State University (UNESP), São Paulo, Brazil
- ³⁷Pharmacology and Toxicology Department, Faculty of Pharmacy, Benghazi University, Benghazi, Libya
- ³⁸School of Nursing and Midwifery, College of Medicine and Health Sciences, University of Rwanda, Kigali, Rwanda
- ³⁹Centre for Language Enhancement, College of Arts and Social Sciences, University of Rwanda, Huye, Rwanda
- ⁴⁰Faculty of Medicine, Alzaïem Alazhari University, Khartoum North, Sudan
- ⁴¹Faculty of Health Sciences and Sports, Macao Polytechnic University, Macao, China
- ⁴²National Autonomous University of Mexico, Mexico City, Mexico
- ⁴³Mental Health and Learning division, Wrexham Maelor Hospital, Wrexham, UKyy
- ⁴⁴Medical-surgical Nursing Department, Faculty of Nursing, Cairo University, Cairo, Egypt

Acknowledgements: We would like to thank Miss Bobo Chan for her tremendous administrative support of the project.

Ethics statement: This study was approved by the Institutional Review Board of the University of Hong Kong-the Hospital Authority Hong Kong West Cluster (reference no: UW 20-272). Informed consent was obtained from all participants, ensuring they were aware of the study's purpose, data collection methods, and their rights. To protect data privacy and handle sensitive health information confidentially, all participants completed the survey anonymously, and data storage and analysis were conducted in strict compliance with the Declaration of Helsinki, upholding the highest standards of data security and confidentiality.

Data availability: Data are available upon reasonable request from the corresponding author.

Funding: This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Authorship contributions: JL analysed the data and drafted the manuscript. DYTF is the principal investigator of the study and is responsible for conducting the study overall. DYTF, JYHW, MH, KYWL and EPHC conceived the study. DYTF, MH, KYWL, JYHW, EPHC, WD, JJJ and CCL contributed to the design of the study. All authors contributed to data acquisition, critically appraised and approved the manuscript, and assume responsibility for the contents of the manuscript.

Disclosure of interest: The authors completed the ICMJE Disclosure of Interest Form (available upon request from the corresponding author) and disclose no relevant interests.

Additional material

Online Supplementary Document

- 1 Pisinger C, Ladelund S, Glümer C, Toft U, Aadahl M, Jørgensen T. Five years of lifestyle intervention improved self-reported mental and physical health in a general population: the Inter99 study. *Prev Med.* 2009;49:424–8. **Medline:19664653 doi:10.1016/j.ypmed.2009.07.020**
- 2 Wong VW-H, Ho FY-Y, Wong YS-H, Chung K-F, Yeung W-F, Ng CH, et al. Efficacy of lifestyle medicine on sleep quality: A meta-analysis of randomized controlled trials. *J Affect Disord.* 2023;330:125–38. **Medline:36863476 doi:10.1016/j.jad.2023.02.111**
- 3 Parsamehr M, Rasoulinezhad SP. The study of the relationship between lifestyle and social health among people of Talesh city. *Quarterly Journal of Social Development.* 2015;10:35–66.
- 4 Neupane D, McLachlan CS, Mishra SR, Olsen MH, Perry HB, Karki A, et al. Effectiveness of a lifestyle intervention led by female community health volunteers versus usual care in blood pressure reduction (COBIN): an open-label, cluster-randomised trial. *Lancet Glob Health.* 2018;6:e66–73. **Medline:29241617 doi:10.1016/S2214-109X(17)30411-4**
- 5 Nyberg ST, Singh-Manoux A, Pentti J, Madsen IEH, Sabia S, Alfredsson L, et al. Association of Healthy Lifestyle With Years Lived Without Major Chronic Diseases. *JAMA Intern Med.* 2020;180:760–8. **Medline:32250383 doi:10.1001/jamainternmed.2020.0618**
- 6 Sun Q, Yu D, Fan J, Yu C, Guo Y, Pei P, et al. Healthy lifestyle and life expectancy at age 30 years in the Chinese population: an observational study. *Lancet Public Health.* 2022;7:e994–1004. **Medline:35926549 doi:10.1016/S2468-2667(22)00110-4**
- 7 Hevey D. Network analysis: a brief overview and tutorial. *Health Psychol Behav Med.* 2018;6:301–28. **Medline:34040834 doi:10.1080/21642850.2018.1521283**
- 8 Noble N, Paul C, Turon H, Oldmeadow C. Which modifiable health risk behaviours are related? A systematic review of the clustering of Smoking, Nutrition, Alcohol and Physical activity (‘SNAP’) health risk factors. *Prev Med.* 2015;81:16–41. **Medline:26190368 doi:10.1016/j.ypmed.2015.07.003**
- 9 Corrales-Gutierrez I, Baena-Antequera F, Gomez-Baya D, Leon-Larios F, Mendoza R. Relationship between eating habits, physical activity and tobacco and alcohol use in pregnant women: Sociodemographic inequalities. *Nutrients.* 2022;14:557. **Medline:35276912 doi:10.3390/nu14030557**
- 10 Kramer A. An overview of the beneficial effects of exercise on health and performance. *Adv Exp Med Biol.* 2020;1228:3–22. **Medline:32342447 doi:10.1007/978-981-15-1792-1_1**
- 11 Vajdi M, Farhangi MA. A systematic review of the association between dietary patterns and health-related quality of life. *Health Qual Life Outcomes.* 2020;18:337. **Medline:33046091 doi:10.1186/s12955-020-01581-z**
- 12 Li J, Fong DYT, Lok KYW, Wong JYH, Ho MM, Choi EPH, et al. Key lifestyles and interim health outcomes for effective interventions in general populations: A network analysis of a large international observational study. *J Glob Health.* 2023;13:04125. **Medline:37861130 doi:10.7189/jogh.13.04125**
- 13 Qiao J, Lin X, Wu Y, Huang X, Pan X, Xu J, et al. Global burden of non-communicable diseases attributable to dietary risks in 1990–2019. *J Hum Nutr Diet.* 2022;35:202–13. **Medline:33834556 doi:10.1111/jhn.12904**
- 14 Katzmarzyk PT, Friedenreich C, Shiroma EJ, Lee I-M. Physical inactivity and non-communicable disease burden in low-income, middle-income and high-income countries. *Br J Sports Med.* 2022;56:101–6. **Medline:33782046 doi:10.1136/bjsports-2020-103640**
- 15 Alexander Bentley R, Ruck DJ, Fouts HN. US obesity as delayed effect of excess sugar. *Econ Hum Biol.* 2020;36:100818. **Medline:31540873 doi:10.1016/j.ehb.2019.100818**
- 16 Al-Ansari B, Thow AM, Day CA, Conigrave KM. Extent of alcohol prohibition in civil policy in Muslim majority countries: the impact of globalization. *Addiction.* 2016;111:1703–13. **Medline:26508526 doi:10.1111/add.13159**
- 17 Li J, Fong DYT, Ho MM, Choi EPH, Lok KYW, Lee JJ, et al. Prevalence and Factors Associated With Willingness to Sustain Pandemic-Induced Digital Work in the General Population and Moderating Effects of Screen Hours: Cross-Sectional Study. *J Med Internet Res.* 2024;26:e53321. **Medline:38805704 doi:10.2196/53321**
- 18 Li J, Fong DYT, Lok KYW, Wong JYH, Man Ho M, Choi EPH, et al. Global impacts of COVID-19 on lifestyles and health and preparation preferences: An international survey of 30 countries. *J Glob Health.* 2023;13:06031. **Medline:37565394 doi:10.7189/jogh.13.06031**
- 19 Lok KY-W, Ko RWT, Fan HSL, Wong JY, Choi EP, Shek NWM, et al. International survey on fear and childbirth experience in pregnancy and the postpartum period during the COVID-19 pandemic: study protocol. *BMJ Open.* 2021;11:e050132. **Medline:34344684 doi:10.1136/bmjopen-2021-050132**
- 20 Li J, Fong DYT, Lok KYW, Wong JYH, Ho MM, Choi EPH, et al. Key lifestyles and health outcomes across 16 prevalent chronic diseases: A network analysis of an international observational study. *J Glob Health.* 2024;14:04068. **Medline:38606605 doi:10.7189/jogh.14-04068**
- 21 Li J, Fong DYT, Lok KYW, Wong JYH, Ho MM, Choi EPH, et al. Key lifestyles and interim health outcomes for effective interventions in general populations: A network analysis of a large international observational study. *J Glob Health.* 2023;13:04125. **Medline:37861130 doi:10.7189/jogh.13.04125**

- 22 Epskamp S, Borsboom D, Fried EI. Estimating psychological networks and their accuracy: A tutorial paper. *Behav Res Methods*. 2018;50:195–212. **Medline:28342071** doi:10.3758/s13428-017-0862-1
- 23 Fonseca-Pedrero E. [Network analysis in psychology]. *Pap Psicol*. 2018;39:1–12. Spanish.
- 24 Jones PJ, Ma R, McNally RJ. Bridge centrality: A network approach to understanding comorbidity. *Multivariate Behav Res*. 2021;56:353–67. **Medline:31179765** doi:10.1080/00273171.2019.1614898
- 25 Hevey D. Network analysis: a brief overview and tutorial. *Health Psychol Behav Med*. 2018;6:301–28. **Medline:34040834** doi:10.1080/21642850.2018.1521283
- 26 Harnett JE, McIntyre E, Steel A, Foley H, Sibbritt D, Adams J. Use of complementary medicine products: a nationally representative cross-sectional survey of 2019 Australian adults. *BMJ Open*. 2019;9:e024198. **Medline:31315853** doi:10.1136/bmjopen-2018-024198
- 27 Hatta M, Balkish M, Rozanim K, Hamizatul A, Norsiah A. How Severe is Binge Drinking in Malaysia and Who are at Risk. *J Alcohol Drug Depend*. 2013;1:2.
- 28 Nurprasetia AN. Alcohol policies, consumption and government revenue: Indonesia. *World Customs J*. 2022;16:15–34. doi:10.55596/001c.116319
- 29 Nguyen TNP, Hunsberger M, Löve J, Duong TA, Phan TH, Luong NK, et al. Patterns and determinants of tobacco purchase behaviors among male cigarette smokers in Vietnam: A latent class analysis. *Tob Induc Dis*. 2024;22:1–10. **Medline:38835515** doi:10.18332/tid/187869
- 30 Brathwaite R, Addo J, Smeeth L, Lock K. A systematic review of tobacco smoking prevalence and description of tobacco control strategies in sub-Saharan African countries; 2007 to 2014. *PLoS One*. 2015;10:e0132401. **Medline:26162085** doi:10.1371/journal.pone.0132401
- 31 Lau JH, Nair A, Abdin E, Kumarasan R, Wang P, Devi F, et al. Prevalence and patterns of physical activity, sedentary behaviour, and their association with health-related quality of life within a multi-ethnic Asian population. *BMC Public Health*. 2021;21:1939. **Medline:34696751** doi:10.1186/s12889-021-11902-6
- 32 Nganabashaka JP, Niyibizi JB, Umwali G, Rulisa SM, Bavuma C, Byiringiro JC, et al. The effects of COVID-19 mitigation measures on physical activity (PA) participation among adults in Rwanda: An online cross-sectional survey. *PLoS One*. 2023;18:e0293231. **Medline:37943889** doi:10.1371/journal.pone.0293231
- 33 Desbouys L, Méjean C, De Henauw S, Castetbon K. Socio-economic and cultural disparities in diet among adolescents and young adults: a systematic review. *Public Health Nutr*. 2020;23:843–60. **Medline:31466544** doi:10.1017/S1368980019002362
- 34 Lobcowska K, Banik A, Forberger S, Kaczmarek K, Kubiak T, Neumann-Podczaska A, et al. Social, economic, political, and geographical context that counts: meta-review of implementation determinants for policies promoting healthy diet and physical activity. *BMC Public Health*. 2022;22:1055. **Medline:35619065** doi:10.1186/s12889-022-13340-4
- 35 Quah S. *Families in Asia: Home and kin*. Milton Park, Abingdon, Oxon: Routledge; 2008.
- 36 HOFSTEDE G. The 6-D model of national culture 2024. Available from: <https://geerthofstede.com/culture-geert-hofstede-gert-jan-hofstede/6d-model-of-national-culture/>. Accessed: 2 January 2025.
- 37 Tawatsupa B, Yiengprugsawan V, Kjellstrom T, Seubsman S-a, Sleigh A, Team TCS. Heat stress, health and well-being: findings from a large national cohort of Thai adults. *BMJ Open*. 2012;2:e001396. **Medline:23135538** doi:10.1136/bmjopen-2012-001396
- 38 Mohamed MS, Panda DS, Fadul FA, Saadeldin AA, Idriss MT, Khan YH, et al. Assessment of the impact of COVID-19 and political instability on mental health of university students in Sudan. *East Mediterr Health J*. 2024;30:272–82. **Medline:38808403** doi:10.26719/2024.30.4.272
- 39 Ventriglio A, Torales J, Castaldelli-Maia JM, De Berardis D, Bhugra D. Urbanization and emerging mental health issues. *CNS Spectr*. 2021;26:43–50. **Medline:32248860** doi:10.1017/S1092852920001236
- 40 Schalock RL. The concept of quality of life: what we know and do not know. *J Intellect Disabil Res*. 2004;48:203–16. **Medline:15025663** doi:10.1111/j.1365-2788.2003.00558.x
- 41 Nouh F. Prevalence of food insecurity in eastern part of Libya: a study of associated factors. *Scholars Acad J Biosci*. 2021;8:192-8. doi:10.36347/sajb.2021.v09i08.002
- 42 Hameed M, Moradkhani H, Ahmadalipour A, Moftakhari H, Abbaszadeh P, Alipour A. A review of the 21st century challenges in the food-energy-water security in the Middle East. *Water*. 2019;11:682. doi:10.3390/w11040682
- 43 Petrikova I, Bhattacharjee R, Fraser PD. The 'Nigerian Diet' and its evolution: review of the existing literature and household survey data. *Foods*. 2023;12:443. **Medline:36765972** doi:10.3390/foods12030443
- 44 Mbah P, Olabisi B. Promoting household food preparation as a strategy to combating nutritional problems in Nigeria: Key to sustainable development. *Education and Science Journal of Policy Review and Curriculum Development*. 2015;5:82–92.
- 45 Kelly RS, Kelly MP, Kelly P. Metabolomics, physical activity, exercise and health: A review of the current evidence. *Biochim Biophys Acta Mol Basis Dis*. 2020;1866:165936. **Medline:32827647** doi:10.1016/j.bbadis.2020.165936