

Available online at [ScienceDirect](https://www.sciencedirect.com)

Resuscitation

journal homepage: www.elsevier.com/locate/resuscitation

Original paper

Defibrillation in patients with accidental hypothermia and core temperatures $\leq 30^{\circ}\text{C}$ – A retrospective observational study



Evelien Cools^a, Robert Blasco^b, Peter Paal^c, Delphine S. Courvoisier^d, Ken Zafren^e, Hermann Brugger^f, Simon Rauch^{f,g}, Beat Walpoth^{h,*}

Abstract

Background: Defibrillation often fails in patients in hypothermic cardiac arrest with shockable rhythms. In patients with core temperatures (T_c) $\leq 30^{\circ}\text{C}$ the success rate, optimal number, timing of defibrillations, and factors including rewarming associated with successful defibrillation are not well known.

Aims: To determine the success rate of defibrillation in patients in hypothermic cardiac arrest with shockable rhythms. We studied T_c at defibrillation attempt, number of shocks, and factors facilitating defibrillation.

Methods: This was a retrospective observational study from the International Hypothermia Registry of patients with hypothermic cardiac arrest ($T_c \leq 30^{\circ}\text{C}$) and shockable rhythms undergoing defibrillation attempts before and during rewarming with or without extracorporeal life support. We performed multivariate binomial logistic regression to analyse the probability of defibrillation success according to the defibrillation temperature and ECLS rewarming.

Results: Thirty-seven patients with cardiac arrest, shockable rhythm, and $T_c \leq 30^{\circ}\text{C}$ were included. Defibrillation was attempted in 20 patients before rewarming and in 17 during rewarming. The overall success rate was 22/37 (59%). Both the success rate of defibrillation (100% [17/17] vs. 25% [5/20]; $P < 0.001$) and T_c at the time of the defibrillation attempt (29.0°C [28.0–30.0°C] vs 25.8°C [24.0–26.2°C]; $P < 0.001$) were higher if defibrillation was performed during rewarming. No defibrillation attempt was successful at $T_c < 24.8^{\circ}\text{C}$.

Conclusions: Defibrillation attempts can be successful before rewarming at $T_c \leq 30^{\circ}\text{C}$. The success rate is higher during rewarming and at a higher T_c . T_c is a strong and independent predictor of defibrillation success. Defibrillation at $T_c < 25^{\circ}\text{C}$ is unlikely. These findings require confirmation in larger studies.

Introduction

Accidental hypothermia (core temperature (T_c) $< 35^{\circ}\text{C}$) is a life-threatening condition that may occur in urban and nonurban environments.^{1,2} Hypothermia can provoke arrhythmias and cardiac arrest (CA).^{3–6} A hypothermic heart with a $T_c < 30^{\circ}\text{C}$ may be resistant to defibrillation. At $T_c < 30^{\circ}\text{C}$, ventricular fibrillation (VF) often recurs

despite initial successful defibrillation.^{7,8} Repeated defibrillations can cause myocardial injury.⁹ If VF persists after three shocks, the 2021 and 2025 guidelines of the European Resuscitation Council (ERC) recommend delaying further defibrillation attempts until T_c is $> 30^{\circ}\text{C}$.^{10,11} In contrast to the ERC guidelines, the 2025 Special Circumstances guidelines of the American Heart Association (AHA) state that it may be reasonable to perform one defibrillation attempts according to the standard algorithm and to withhold administration of

* Corresponding author.

E-mail addresses: Evelien.Cools@hopitalrivierachablais.ch (E. Cools), roblasc.rb@gmail.com (R. Blasco), Peter.Paal@icloud.com (P. Paal), Delphine.Courvoisier@hug.ch (D.S. Courvoisier), kenzafren@gmail.com (K. Zafren), hermann.brugger@eurac.edu (H. Brugger), simon.rauch@eurac.edu (S. Rauch), Beat.Walpoth@unige.ch (B. Walpoth).

<https://doi.org/10.1016/j.resuscitation.2026.110971>

Received 28 October 2025; Received in Revised form 22 December 2025; Accepted 31 December 2025

epinephrine until the core Tc is $>30^{\circ}\text{C}$.¹² Similarly to the AHA 2025 special circumstances guidelines, the 2019 guidelines of the Wilderness Medical Society (WMS) recommend that a single shock at a maximum power be given for patients with a temperature $<30^{\circ}\text{C}$.¹³ This discrepancy in guidelines can be explained by the different interpretation of poor-quality data from animal studies, case reports and case series in humans.^{7,14–16}

We aimed to determine the incidence of successful defibrillation in patients in cardiac arrest with shockable rhythms with Tc $\leq 30^{\circ}\text{C}$. We evaluated Tc at the time of successful defibrillation, number of shocks necessary for successful defibrillation, and factors associated with successful defibrillation. We hypothesised that rewarming patients in hypothermic cardiac arrest on Extracorporeal Life Support (ECLS) would be associated with higher defibrillation success, because of improved myocardial oxygen support, making the heart more responsive to defibrillation.

Methods

This study was approved by the ethical committee of Geneva, Switzerland (2023-01087). We extracted the data on September 1, 2023. The study was registered in clinicaltrials.gov (number NCT06131892). This research adhered to the STROBE guidelines.¹⁷

We performed a retrospective observational study using data from the International Hypothermia Registry (IHR). The IHR collects data of adults and children with accidental hypothermia (Tc $<35^{\circ}\text{C}$).¹⁸ The registry was created in 2008 (protocol No. 08-040R, Geneva, 23 September 2008). On August 31, 2023, 28 centres in 10 countries (Austria, Denmark, Germany, Italy, Japan, Poland, Spain, Switzerland, the UK, and the USA) were participating in the registry. We included all patients entered into the registry by August 31, 2023, with attempted defibrillation at Tc $\leq 30^{\circ}\text{C}$. We studied two groups: those with attempted defibrillation before rewarming and those with attempted defibrillation during rewarming. Rewarming with extracorporeal life support (ECLS) refers to rewarming using cardiopul-

monary bypass (CPB) or extracorporeal membrane oxygenation (ECMO) (venoarterial or venovenous). Non-ECLS rewarming included all other invasive and noninvasive rewarming techniques. The non-ECLS rewarming techniques that can be selected in the International Hypothermia Registry are haemodialysis, continuous venovenous haemofiltration, intravenous catheter rewarming, continuous arteriovenous rewarming, thoracic lavage (thoracotomy), thoracic lavage (thoracoscopy), mediastinal lavage sternotomy, peritoneal lavage, peritoneal dialysis, bladder lavage, gastric lavage, warm IV fluids, and other.

We extracted the following data: age, sex, country, cause of hypothermia (mountain accidents, including avalanche, outdoor exposure, and crevasse accidents; water-related accidents (drowning and immersion); and rural or urban accidents (outdoor exposure in patients experiencing homelessness and other vulnerable populations), presence of asphyxia (submersion in water or avalanche with burial of the head under the snow), lowest measured Tc, first rhythm (prehospital or inhospital), Tc at the time of successful defibrillation or the time of the last shock if attempted defibrillation was unsuccessful, initial cardiac rhythm, and whether CA was witnessed. We extracted the type of defibrillation (i.e. electrical or spontaneous) and recurrence of VF after successful defibrillation.

The primary outcome was the incidence of successful defibrillation with persistent return of spontaneous circulation (ROSC) in patients in hypothermic cardiac arrest with shockable rhythms and Tc $\leq 30^{\circ}\text{C}$.¹⁹ We defined persistent ROSC as sustained spontaneous circulation for ≥ 30 s. We defined successful defibrillation as either spontaneous or electrically successful defibrillation with persistent ROSC. Secondary outcomes were the influence of Tc on defibrillation success, the number of shocks necessary to obtain successful defibrillation, the influence of ECLS rewarming, the use of epinephrine and other factors associated with defibrillation success. Although the registered trial protocol included recurrent ventricular fibrillation and cardiac dysfunction as secondary outcomes, these measures were not consistently collected throughout the study.




Fig. 1 – Flowchart of the study population.

Table 1 – Baseline characteristics of patients stratified by timing of defibrillation.

Variable	Overall	Defibrillation before rewarming	Defibrillation during rewarming	p-value
Number of patients	N = 37	N = 20	N = 17	
Patient characteristics				
Sex male (n, %)	25 (68%)	14 (70%)	11 (65%)	0.99
Age (mean, $\pm$ SD)	43 $\pm$ 18	42 $\pm$ 20	44 $\pm$ 17	0.78
Cardiac arrest characteristics				
Shockable rhythm prehospitally (n, %)	15 (58%)	10 (67%) Missing = 5	5 (45%) Missing = 6	0.43
Lowest prehospital temp. (median, IQR)	24.6°C (22.3–26)	25.8°C (24–26.4)	22.5°C (20.7–24.7)	0.006
Core temp. at first defibrillation (median, IQR)	26.8°C (25–29)	25.8°C (24–26.2)	29°C (28–30)	<0.001
Accident site (n, %)				0.009
– Mountain	13 (35%)	5 (25%)	8 (47%)	
– Water	11 (30%)	10 (50%)	1 (6%)	
– Rural/Urban	13 (35%)	5 (25%)	8 (47%)	
Rescue collapse (n, %)	11 (48%)	6 (43%), Missing = 6	5 (56%), Missing = 8	0.68
Asphyxia (n, %)	15 (43%)	7 (39%), Missing = 2	8 (47%)	0.73
Witnessed CA (n, %)	19 (66%)	12 (71%), Missing = 3	7 (58%), Missing = 5	0.69
pH (mean, $\pm$ SD)	6.9 $\pm$ 0.2	6.9 $\pm$ 0.2, Missing = 8	6.9 $\pm$ 0.2, Missing = 3	0.47
Lactate (mean, $\pm$ SD)	14.7 $\pm$ 7.8	17.4 $\pm$ 9, Missing = 9	12.7 $\pm$ 6.5, Missing = 2	0.13
Potassium (median, IQR)	3.9 (3.1–4.57)	3.7 (2.80–6.29), Missing = 6	4 (3.32–4.43), Missing = 1	0.98
Treatment provided				
Epinephrine (n, %)	22 (59%)	16 (80%)	6 (35%)	0.008
ECLS rewarming (n, %)	25 (73.5%)	9 (53%), Missing = 3	16 (94%)	0.017
CPR duration (median, IQR)	119 (75–159)	122 (64–164), Missing = 8	119 (76–157), Missing = 3	0.73
Number of shocks given (median, IQR)	2 (1–3.5)	3 (1.25–4.75)	2 (1–2)	0.15
Energy defibrillation (mean, SD)	255 (93)	230 (82)	274 (99)	0.27
Low flow time (median, IQR)	94 (66–145)	82 (39–122), Missing = 4	140 (75–155), Missing = 4	0.07

(continued on next page)

Table 1 (continued)

Variable	Overall	Defibrillation before rewarming	Defibrillation during rewarming	p-value
Outcomes				
Successful defibrillation (n, %)	22 (59%)	5 (25%)	17 (100%)	<0.001
HOPE score (mean, SD)	55 (27)	41 (26), Missing = 10	67 (23), Missing = 6	0.026
CPC ICU discharge (n, %)		Missing = 9	Missing = 1	0.09
– CPC 1	11 (41%)	2 (18%)	9 (56%)	
– CPC 2	4 (15%)	3 (27%)	1 (6%)	
– CPC 3	1 (4%)	0	1 (6%)	
– CPC 4	0	0	0	
– CPC 5	11 (41%)	6 (54%)	5 (31%)	

CA denotes cardiac arrest, CPC cerebral performance category, ECLS extracorporeal life support, ICU intensive care unit, IQR interquartile range, SD standard deviation.

Statistical analysis

We report general characteristics of the study population using descriptive statistics. We expressed normally distributed data as mean $\pm$ standard deviation (SD) and non-normally distributed data as median $\pm$ interquartile range (IQR). We assessed normal distribution using the Shapiro-Wilk test and Q-Q plots. We tested homogeneity of variances using Levene's test. We compared defibrillation characteristics of patients (defibrillation during or before rewarming) and those with or without successful defibrillation using Fisher's exact test for categorical characteristics and Student's *t*-test or Welch's *t*-test for continuous variables if normality was satisfied, or the Mann-Whitney *U* test for non-normal distributions. We set the level of significance at <0.05 . Analyses were performed as complete-case analyses (Supplementary Table S1 for missing data by variable); defibrillation success was available for all patients, while variables with missing values (lactate, potassium, CPC) were handled using complete-case approach in respective analyses. We assessed univariate associations with defibrillation success using binomial logistic regression. We performed multivariate binomial logistic regression to analyse the probability of defibrillation success according to the defibrillation Tc and ECLS rewarming. Because of quasi-complete separation in the ECLS rewarming variable and the sample size, we complemented the analysis with Firth's penalised logistic regression to obtain more stable estimates. We used a bootstrap with 5000 replicates to assess the robustness and stability of the penalised regression results. To assess temporal confounding over the 40-year study period, we stratified analyses by pre-2010 vs post-2010, comparing ECLS utilisation, success rates, and temperatures across periods. We included period as a binary variable in our multivariable model using Firth's penalised regression to control for temporal changes in ECLS availability.

Statistical analysis was performed using R version 4.4.3 (2025-02-28).

Results

Of the 37 patients, 20 had attempted defibrillation before rewarming and 17 during rewarming (Fig. 1). Twenty-six patients were excluded: 9 patients because the temperature at defibrillation was $>30^{\circ}\text{C}$, 11 patients because temperature at defibrillation was not known. Six patients had defibrillation attempts both before and during rewarming (Supplementary Table S2 compares the main characteristics of included and excluded patients).

Patients came from Switzerland,¹⁵ Poland,⁹ France,⁴ Austria,⁵ Italy,² Spain,¹ and the United Kingdom.¹ The cases occurred between August 5, 1982 and December 6, 2022.

Patients were mostly male (25/37, 68%), with a mean age of 43 ± 18 years (Table 1). Accidents were more likely to be water-related in patients with defibrillation before rewarming (10/20 (50%) versus 1/17 (6%)). The initial prehospital median core temperature was lower in patients who underwent defibrillation during rewarming compared to those who were defibrillated before rewarming (22.5°C [$20.7\text{--}24.7^{\circ}$] vs. 25.8°C [$24.0\text{--}26.4^{\circ}$], $P = 0.006$). Patients with successful defibrillation before rewarming required ECLS to be rewarmed less often than patients who were not defibrillated in the prehospital phase (53% vs 94%, $p = 0.017$) (Table 1).

Primary outcome

The overall success rate of electrical and spontaneous defibrillation was 22 of 37 (59%) (Table 1). Three of the 22 patients had spontaneous defibrillation and ROSC before rewarming. Defibrillation was more successful when done during, rather than before, rewarming (17/17 (100%) vs 5/20 (25%), $p < 0.001$) (Table 1).

Secondary outcomes

The median temperature at the time of the first defibrillation attempt, successful or unsuccessful, was higher in the group with defibrillation during rewarming than in the group with defibrillation before rewarm-

Table 2 – Differences between patients with successful or not successful defibrillation.

Variable	Overall	Successful defibrillation	Not successful defibrillation	p-value
Number of patients	N = 37	N = 22 (59.5%)	N = 15 (40.5%)	
Patient characteristics				
Age (mean, $\pm$ SD)	43 $\pm$ 18.4	45 $\pm$ 19.7	39 $\pm$ 16.1	0.29
Sex male (n, %)	25 (68%)	14 (64%)	11 (73%)	0.72
Cardiac arrest characteristics				
Shockable rhythm prehospitally (n, %)	15 (58%)	8 (53%) Missing = 7	7 (64%) Missing = 4	0.7
Prehospital lowest temperature (median, IQR)	24.6°C (22.3–26)	23.6°C (21.5–25.5)	25°C (23.6–26)	0.25
Core temperature at first defibrillation (median, IQR)	26.8°C (25–29)	28°C (26.8–29.9)	25°C (23.6–26)	<0.001
Accident site (n, %)				0.59
– Mountain	13 (35%)	8 (36%)	5 (33%)	
– Water	11 (30%)	5 (23%)	6 (40%)	
– Rural/Urban	13 (35%)	9 (41%)	4 (27%)	
Rescue collapse (n, %)	11 (48%)	7 (54%), Missing = 9	4 (40%), Missing = 5	0.68
Asphyxia (n, %)	15 (43%)	10 (45%)	5 (38%), Missing = 2	0.73
Witnessed CA (n, %)	19 (66%) Missing = 8	12 (71%) Missing = 5	7 (58%) Missing = 3	0.69
pH (mean, $\pm$ SD)	6.9 $\pm$ 0.2	6.9 $\pm$ 0.18, Missing = 5	6.87 $\pm$ 0.21, Missing = 6	0.48
Lactate (mean, $\pm$ SD)	14.7 $\pm$ 7.8	12.2 $\pm$ 5.9, Missing = 3	21.3 $\pm$ 8.8, Missing = 8	0.006
Potassium (median, IQR)	3.9 (3.1–4.57)	3.75 (3–4.4), Missing = 2	5 (3.15–7.6), Missing = 5	0.08
Treatment provided				
Epinephrine (n, %)	22 (59%)	10 (45%)	12 (80%)	0.047
CPR duration (median, IQR)	119 min (75–159)	102 min (69–155) Missing = 5	153 min (94–175) Missing = 6	0.17
Low flow time (median, IQR)	94 min (66–145)	98.5 min (67–153) Missing = 4	94 min (67–125) Missing = 4	0.73
Number of shocks given (median, IQR)	2 (1–3.5)	2 (1–3) Missing = 5	3 (1.25–5.75) Missing = 1	0.16
Energy defibrillation (mean, $\pm$ SD)	255 $\pm$ 93.1	276 $\pm$ 96 Missing = 8	222 $\pm$ 82.7 Missing = 6	0.18
ECLS rewarming (n, %)	25 (73.5%)	18 (85%) Missing = 1	7 (53%), Missing = 2	0.05

(continued on next page)

Table 2 (continued)

Variable	Overall	Successful defibrillation	Not successful defibrillation	p-value
– Before rewarming		2 (50%)	7 (54%)	
– During rewarming		16 (94%)	0	
Outcomes				
HOPE score (mean, $\pm$ SD)	55.0 $\pm$ 27.3 Missing = 16	65.3 $\pm$ 24.6 Missing = 8	34.6 $\pm$ 21.1 Missing = 8	0.011
CPC at ICU discharge (n, %)		Missing = 3	Missing = 7	0.17
– CPC 1	11 (41%)	10 (53%)	1 (12%)	
– CPC 2	4 (15%)	2 (10%)	2 (25%)	
– CPC 3	1 (4%)	1 (5.3%)	0	
– CPC 4	0	0	0	
– CPC 5	11 (41%)	6 (31.6%)	5 (62.5%)	

CA denotes cardiac arrest, CPR cardiopulmonary resuscitation, ECLS extracorporeal life support, IQR interquartile range, SD standard deviation. HOPE: Hypothermia Outcome Prediction after Extracorporeal life support.

ing (29.0°C (28.0–30°C) vs 25.8°C (24.0–26.2°C), $P < 0.001$). No defibrillation with $T_c < 24.8^\circ\text{C}$ was successful. A higher median core temperature was associated with higher defibrillation success (28°C (26.8–29.9) vs 25°C (23.6–26); $P < 0.001$) (Table 2). Each additional 1°C in core temperature at the time of shock was associated with greater odds of successful defibrillation (OR 2.68; $P = 0.003$) (Table 3). The T_c at defibrillation was positively associated with defibrillation success, with adjusted odds ratios increasing from 2.68 (95% CI: 1.39–5.14, $P = 0.003$) in univariate analysis to 4.03 (95% CI: 1.3–12.3, $P = 0.015$) in multivariate analysis, adjusting for ECLS rewarming (Table 4, Fig. 2). At the median cohort temperature of 26.8°C, the difference in the probability of success between patients managed with or without extracorporeal life support (ECLS) was 31% (78% vs. 47%) (Fig. 2). The discriminative ability of this model was high (AUC 0.94). The application of Firth's penalised regression produced more stable estimates than the standard model, with defibrillation temperature remaining a stable predictor of success (OR 2.83; CI 95% 1.54–9.56, $P < 0.001$). Analyses using bootstrapping confirmed the robustness of the temperature effect.

The number of shocks was reported in 31 patients (Fig. 3, Supplementary Fig. 2). Seventeen of the 31 patients (55%) were successfully defibrillated. Of the 17 patients successfully defibrillated, 5 (29%) were successfully defibrillated with 1 shock, 7 (41%) with 2 shocks, 3 (1%) with 3 shocks and 2 (12%) with 4 to 5 shocks (Fig. 3, Supplementary Fig. 2).

Patients who received rewarming with ECLS had greater than 5 times higher odds of successful defibrillation than those who did not receive ECLS rewarming (OR 5.4; $P = 0.05$) (Table 3). The absolute number of ECLS-treated patients increased from pre-2010 ($n = 3$) to post-2010 ($n = 22$), reflecting likely changes in ECLS availability. The proportion of patients receiving ECLS was similar between periods (75% pre-2010 vs 73.3% post-2010), suggesting consistent selection criteria. Pre-2010 patients had higher median core temperatures at defibrillation (27.5°C vs 26.9°C) yet lower suc-

cess rates (50% vs 63.3%), supporting the hypothesis that improved outcomes were driven by temperature effects rather than increased ECLS availability. In the multivariable model adjusted for period (pre-2010 vs post-2010), temperature retained strong significance (OR 2.45, 95% CI 1.47–7.82, $p < 0.001$), while the period variable was not significant (OR 0.15, 95% CI 0–12.39, $p = 0.53$), indicating that the association between temperature and success is consistent across eras and confirming the absence of temporal confounding (Supplementary Fig. S1).

More epinephrine was used in the patients defibrillated before than during rewarming (16/20 (80%) versus 6/17 (35%), $P = 0.008$) (Table 1). Epinephrine administration was associated with a lower success rate (10/22 (45%) vs 12/15 (80%); $P = 0.05$).

Higher mean lactate levels were associated with a lower probability of success (21.4 $\pm$ 8.9 mmol/L vs 12.2 $\pm$ 5.9 mmol/L) vs. $p = 0.006$). Patients with witnessed cardiac arrest had a trend to higher chance of a successful defibrillation that was not statistically significant (OR 1.7, 95%CI 0.5–8.1, $p = 0.5$). Each 1 mEq/L of higher potassium was associated with a 41% decrease in the odds of successful defibrillation (OR 0.58; $P = 0.046$) (Table 3).

Discussion

We assessed the incidence of successful defibrillation in 37 patients in hypothermic cardiac arrest with shockable rhythms. In 25%, defibrillation was successful before rewarming compared to 100% during rewarming. Three of 20 patients (15%) experienced spontaneous defibrillation with ROSC at core temperatures below 30°C prior to rewarming.

In patients in hypothermic cardiac arrest, the likelihood of successful defibrillation is strongly dependent on T_c . Our findings suggest that achieving a T_c of $\geq 25^\circ\text{C}$ prior to a shock is key to successful defibrillation. This observation should be treated as

Table 3 – Univariate binomial logistic regression for predictors of success in defibrillation.

	Unit of change	OR	IC	p-value	AUC
Age	per 1 year	1.02	0.98–1.06	0.29	0.61
Witnessed CA	Binary	1.7	0.36–8.1	0.50	0.56
Low flow time	per 1 min	1	0.99–1.01	0.78	0.54
Lactate	per 1 mmol/L	0.83	0.71–0.98	0.02	0.81
pH	per 0.1 units	1.18	0.75–1.86	0.47	0.59
K	per 1 mmol/L	0.58	0.34–0.99	0.05	0.63
Epinephrine	Binary	0.21	0.05–0.95	0.04	0.67
CPR duration	per 1 min	0.99	0.98–1	0.31	0.67
Prehospital temperature	per 1°C	0.91	0.72–1.15	0.44	0.61
ECLS	Binary	5.14	1.00–26.5	0.05	0.66
Shocks given	per 1 shock	0.68	0.45–1.03	0.07	0.68
Defibrillation energy (per shock)	per 1 Joule	1.007	0.99–1.02	0.18	0.65
Temperature at defibrillation	per 1°C	2.68	1.39–5.14	0.003	0.89
HOPE score	per 1%	1.06	1.00–1.14	0.04	0.88

AUC denotes area under the curve, CA cardiac arrest, CPR cardiopulmonary resuscitation, ECLS extracorporeal life support, IQR interquartile range, IR information coefficient, OR odds ratio. HOPE: Hypothermia Outcome Prediction after Extracorporeal life support.

Table 4 – Multivariate logistic regression for the probability of defibrillation success according to the defibrillation temperature during defibrillation and the ECLS rewarming and a comparative for outcome predictors using different models.

Model	Variable	OR	CI 95%	p-value
Multivariable classic logistic regression	Defibrillation temperature (°C)	4.03	1.32–12.3	0.015
	ECLS rewarming (Yes/No)	5.42	0.51–57	0.15
Firth's penalised logistic regression	Defibrillation temperature (°C)	2.83	1.54–9.56	<0.001
	ECLS rewarming (Yes/No)	3.85	0.53–37.4	0.18
Bootstrap after firth	Defibrillation temperature (°C)	3.74	1.98–45	–
	ECLS rewarming (Yes/No)	5.07	0.56–109	–

Classic model: AUC 0.94, Sensibility 0.905, Specificity 0.846. Cox & Snell R square 0.518, Nagelkerke R square 0.704). AUC denotes area under the curve, CA cardiac arrest, CI confidence interval, ECLS extracorporeal life support, OR odds ratio.

hypothesis-generating and should be corroborated by further studies providing additional data. Successful defibrillation is possible in patients with $T_c \leq 30^\circ\text{C}$ ^{14,15,20,21} but it is unlikely at a $T_c < 25^\circ\text{C}$ despite a few reported cases with successful defibrillation with lower T_c .^{14,16}

Our data suggest that most patients can be successfully defibrillated at T_c under 30°C during ECLS rewarming (Fig. 2). This finding has significant implications for clinical practise. Based on our data, defibrillation during ECLS rewarming should be attempted with a $T_c > 25^\circ\text{C}$ and should not be withheld until T_c reaches 30°C . Other




Fig. 2 – Probability of successful defibrillation among patients rewarmed or not using ECLS and the temperature at the first defibrillation.

ECLS = extracorporeal life support.

factors for successful defibrillation in hypothermic cardiac arrest are immediate high-quality cardiopulmonary resuscitation (CPR), and a higher core temperature. These results reinforce the importance of achieving a T_c of $\geq 25^{\circ}\text{C}$ before attempting defibrillation in hypothermic cardiac arrest patients. Although achieving a temperature of at least 25°C may improve the chances of success, attempts at defibrillation should not be deterred in patients with lower temperatures. These findings inform expectations of defibrillation success and could help to refine resuscitation protocols. Based on the accident sites, all the patients analysed had access to an ECLS centre in less than 6 h.

It is not clear whether the higher likelihood of successful defibrillation during ECLS rewarming can be attributed solely to the higher T_c observed during ECLS rewarming or whether improved myocar-

dial perfusion, oxygenation and metabolic state also play a role.²² Our data support continuing CPR even if ECLS is not available, e.g. in case of bad weather, long distances, low-resource settings, because there is a 25% chance of successful defibrillation and ROSC using standard CPR and non-ECLS rewarming with a T_c $>25^{\circ}\text{C}$. When transfer to an ECLS centre is not possible, active rewarming should be started as early as possible. With a T_c $<25^{\circ}\text{C}$ cerebral metabolism is reduced by $>70\%$, allowing CPR to be intermittent during a difficult evacuation,²³ but with a T_c between 28 and 30°C , even short interruptions of CPR are associated with a high risk of hypoxic cerebral injury. Above a T_c of 28, successful defibrillation may be life-saving.²³ In the defibrillation before rewarming group, 8 patients were first transferred to a non-ECLS hospital. Only one patient wasn't transferred to an ECLS centre. Eight of the patients



Fig. 3 – Number of shocks versus core temperature at defibrillation.

of the ECLS group were first admitted in a non-ECLS hospital, before arriving in an ECLS centre.

Our data support the ERC 2021 and 2025 guidelines to deliver more than one shock to patients in cardiac arrest with $T_c \leq 30^\circ\text{C}$ and shockable rhythms.^{10,11} Using the WMS 2019 and the AHA 2025 guidelines that recommend only one defibrillation attempt, would have reduced the chances of achieving a successful defibrillation in 67% of patients in the present study.^{12,13} No patient in our study had a successful defibrillation after more than 5 shocks. All patients requiring 6 or more shocks ($n = 4$) failed to achieve defibrillation success despite continued resuscitation efforts.

Higher lactate and potassium levels were associated with a lower probability of successful defibrillation, suggesting that poor tissue perfusion and a worse metabolic state reduce the efficacy of defibrillation. These associations likely reflect the severity and duration of cardiac arrest rather than being causal. Epinephrine administration was also associated with a lower success rate but, we were unable to determine if this was a causal effect. It is still unclear whether epinephrine improves outcome in hypothermic cardiac arrest. Because of potential detrimental effects of epinephrine and the risk of accumulation in hypothermic cardiac arrest^{24–26} the 2021 European Resuscitation guidelines did not recommend its use¹⁰ and the 2025 European Resuscitation guidelines¹¹ recommend only 1 mg during hypothermic cardiac arrest. The WMS 2019 and the AHA 2025 do not recommend any epinephrine use at all.^{12,13} Some experimental studies have suggested that the rate of successful defibrillation is higher after administration of epinephrine but, so far, no clinical studies have confirmed this.^{24, 25} The discrepancy in the AHA, ERC and WMS guidelines may be explained with a different interpretation of data and their clinical applicability and efficiency. Certainly, the differences in these guidelines are decreasing over time.

The HOPE score was lower in the unsuccessful defibrillation group ($35 \pm 21\%$) vs $65 \pm 25\%$) but was far above the 10% cut-off below which patients should not be rewarmed with ECLS. The pre-hospital HOPE score should be interpreted as being associated with the hospital-based HOPE score, rather than as a predictor of pre-hospital outcomes.

Limitations

Of 251 patients in the IHR, few patients ($n = 37$) were included in this study (15% of inclusion rate). The geographic representation is biased, which limits generalisability and information about severity of trauma was not reported. The largest human series on defibrillation in hypothermic cardiac arrest includes four patients.¹⁶ The data were collected retrospectively with a significant rate of missing data (17%). A selection bias may have led to preferential inclusion of patients with better outcomes, such as higher rates of defibrillation success or survival. Differences in baseline characteristics between patients defibrillated before and during ECLS rewarming may have confounded the defibrillation success variable. We do not have access to injury severity scores (ISS) or other detailed injury characteristics that would enable us to adjust for injury severity as a confounding factor. Furthermore, we lack documented criteria for ECLS selection decisions made by clinical teams during the study period. We are not aware of any study, which has included a similarly large number of patients to assess this study question.

Conclusion

Defibrillation attempts can be successful before rewarming at $T_c \leq 30^\circ\text{C}$. The success rate is higher during rewarming and at a higher T_c . T_c is a strong and independent predictor of defibrillation success. Defibrillation at $T_c < 25^\circ\text{C}$ is unlikely. These findings require confirmation in larger studies.

CRedit authorship contribution statement

Evelien Cools: Visualization, Resources, Project administration, Methodology, Investigation, Formal analysis, Data curation, Conceptualization. **Robert Blasco:** Writing – review & editing, Writing – original draft, Visualization, Validation, Software, Methodology, Investigation, Formal analysis, Data curation. **Peter Paal:** Writing – review & editing, Writing – original draft, Visualization, Validation, Supervision, Methodology, Investigation, Conceptualization. **Delphine S. Courvoisier:** Software, Methodology, Formal analysis, Data curation. **Ken Zafren:** Writing – review & editing, Validation. **Hermann Brugger:** Writing – review & editing, Validation, Methodology. **Simon Rauch:** Writing – review & editing. **Beat Walpoth:** Writing – review & editing, Writing – original draft, Validation, Supervision, Resources, Conceptualization.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of competing interest

All authors declare to have no conflicts of interest.

Acknowledgements

We thank the centres participating in the IHR and the rescuers for collecting the prehospital data.

The centres participating in the International Hypothermia Registry are Medical University of Innsbruck, Innsbruck Austria, Krankenhaus Barmherzige Brüder, Salzburg Austria, Rigshospitalet, København, Denmark, Immenstadt Hospital, Immenstadt im Allgäu, Germany, Krankenhaus Bruneck, Brunico, Italy, Institute of Mountain Emergency Medicine, Eurac Research, Bozen/Bolzano, Italy, Ospedale di Bolzano, Bozen/Bolzano, Italy, Krankenhaus Brixen, Brixen/Bressanone, Italy, Krankenhaus F. Tappeiner Meran, Meran, Italy, Krankenhaus Innichen, Innichen, Italy, Ospedale Santa Chiara, Santa Chiara, Italy, ASST Bergamo Est, Seriate, Italy, Krankenhaus Schlanders, Silandro, Italy, Krankenhaus Sterzing, Vipiteno, Italy, Hokkaido Ohno Memorial Hospital, Sapporo, Hokkaido, Japan, Medical University of Silesia, Katowice, Poland, Val d'Hebron University Hospital, Barcelona, Spain, Hospital de Cerdanya, Girona, Spain, Universitätsspital Basel, Basel, Switzerland, Hôpitaux Universitaires Genève, Geneva, Switzerland, Kantonspital Graubünden, Chur,

Switzerland, Luzerner Kantonspital, Luzerne, Switzerland, Ospedale Regionale di Lugano, Lugano, Switzerland, University Hospital Zürich, Zürich, Switzerland, Patterdale Mountain Rescue Team, Patterdale, UK, University of Portsmouth, Portsmouth, UK

Appendix A. Supplementary material

Supplementary material to this article can be found online at <https://doi.org/10.1016/j.resuscitation.2026.110971>.

Author details

^aDepartment of Anaesthesiology, Hôpital Riviera-Chablais, Route du Vieux-Sequoia 20, 1847 Rennaz, Air Zermatt, Spissstrasse 107, 3920 Zermatt, Switzerland ^bFaculty of Medicine. Autonomous University of Barcelona. Department of Anaesthesiology, Vall d'Hebron University Hospital, Barcelona, Spain ^cDepartment of Anaesthesiology and Intensive Care Medicine, St. John of God Hospital, Paracelsus Medical University, Salzburg, Austria ^dUniversity Hospital Geneva, Rue Gabrielle-Perret-Gentil 4, 1205 Genève, Switzerland ^eDepartment of Emergency Medicine, Stanford University Medical Center, Palo Alto, CA, USA ^fInstitute of Mountain Emergency Medicine, Eurac Research, 39100 Bolzano, Italy ^gDepartment of Anaesthesiology and Intensive Care Medicine, Hospital of Merano 39012 Merano, Italy ^hEmeritus, Dept. of Cardiovascular Surgery, University Hospital & University of Geneva, Switzerland

REFERENCES

- Walpoth BH, Walpoth-Aslan BN, Mattle HP, Radanov BP, Schroth G, Schaeffler L, et al. Outcome of survivors of accidental deep hypothermia and circulatory arrest treated with extracorporeal blood warming. *N Engl J Med* 1997;337(21):1500–5.
- Caroselli C, Gabrieli A, Pisani A, Bruno G. Hypothermia: an underestimated risk. *Intern Emerg Med* 2009;4(3):227–30.
- Paal P, Gordon L, Strapazzon G, Brodmann Maeder M, Putzer G, Walpoth B, et al. Accidental hypothermia-an update : the content of this review is endorsed by the International Commission for Mountain Emergency Medicine (ICAR MEDCOM). *Scand J Trauma Resusc Emerg Med* 2016;24(1):111.
- Turk EE. Hypothermia. *Forensic Sci Med Pathol* 2010;6(2):106–15.
- Iyer A, Rajkumar V, Sadasivan D, Bruce J, Gilfillan I. No one is dead until warm and dead. *J Thorac Cardiovasc Surg* 2007;134(4):1042–3.
- Walpoth BH, Maeder MB, Courvoisier DS, Meyer M, Cools E, Darocha T, et al. Hypothermic cardiac arrest – retrospective cohort study from the international hypothermia registry. *Resuscitation* 2021;167:58–65.
- Kuhnke M, Albrecht R, Schefold JC, Paal P. Successful resuscitation from prolonged hypothermic cardiac arrest without extracorporeal life support: a case report. *J Med Case Rep* 2019;13(1):354.
- Schwarz B, Mair P, Raedler C, Deckert D, Wenzel V, Lindner KH. Vasopressin improves survival in a pig model of hypothermic cardiopulmonary resuscitation. *Crit Care Med* 2002;30(6):1311–4.
- Gazmuri RJ, Deshmukh S, Shah PR. Myocardial effects of repeated electrical defibrillations in the isolated fibrillating rat heart. *Crit Care Med* 2000;28(8):2690–6.
- Lott C, Truhlar A, Alfonso A, Barelli A, Gonzalez-Salvado V, Hinkelbein J, et al. European Resuscitation Council guidelines 2021: cardiac arrest in special circumstances. *Resuscitation* 2021;161:152–219.
- Lott C, Karageorgos V, Abelairas-Gomez C, Alfonso A, Bierens J, Cantellow S, et al. European Resuscitation Council guidelines 2025 special circumstances in resuscitation. *Resuscitation* 2025;215(Suppl 1):110753.
- Cao D, Arens AM, Chow SL, Easter SR, Hoffman RS, Lagina 3rd AT, et al. Part 10: adult and pediatric special circumstances of resuscitation: 2025 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2025;152(16_suppl_2):S578–672.
- Dow J, Giesbrecht GG, Danzl DF, Brugger H, Sagalyn EB, Walpoth B, et al. Wilderness Medical Society Clinical Practice guidelines for the out-of-hospital evaluation and treatment of accidental hypothermia: 2019 update. *Wilderness Environ Med* 2019;30(4S):S47–69.
- Kosinski S, Drzewiecka A, Pasquier M, Golba KS, Podsiadlo P, Drwila R, et al. Successful defibrillation at a core temperature of 18.2 °C. *Wilderness Environ Med* 2020;31(2):230–4.
- Thomas R, Cahill CJ. Successful defibrillation in profound hypothermia (core body temperature 25.6 °C). *Resuscitation* 2000;47(3):317–20.
- Mair P, Gasteiger L, Mair B, Stroehle M, Walpoth B. Successful defibrillation of four hypothermic patients with witnessed cardiac arrest. *High Alt Med Biol* 2019;20(1):71–7.
- von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007;370(9596):1453–7.
- Anonymous. International Hypothermia Registry; 2025. Available from: <https://hypothermia-registry.org/>.
- Cummins RO, Chamberlain DA, Abramson NS, Allen M, Baskett PJ, Becker L, et al. Recommended guidelines for uniform reporting of data from out-of-hospital cardiac arrest: the Utstein style. A statement for health professionals from a task force of the American Heart Association, the European Resuscitation Council, the Heart and Stroke Foundation of Canada, and the Australian Resuscitation Council. *Circulation* 1991;84(2):960–75.
- Marimuthu V, Balaji GA, Chandrasekar N, Mathai K. Successful defibrillation/resuscitation in accidental profound hypothermia. *Indian J Anaesth* 2013;57(2):213–4.
- Clift J, Munro-Davies L. Best evidence topic report. Is defibrillation effective in accidental severe hypothermia in adults?. *Emerg Med J* 2007;24(1):50–1.
- Podsiadlo P, Darocha T, Kosinski S, Salapa K, Zietkiewicz M, Sanak T, et al. Severe hypothermia management in mountain rescue: a survey study. *High Alt Med Biol* 2017;18(4):411–6.
- Gordon L, Paal P, Ellerton JA, Brugger H, Peek GJ, Zafren K. Delayed and intermittent CPR for severe accidental hypothermia. *Resuscitation* 2015;90:46–9.
- Kornberger E, Lindner KH, Mayr VD, Schwarz B, Rackwitz KS, Wenzel V, et al. Effects of epinephrine in a pig model of hypothermic cardiac arrest and closed-chest cardiopulmonary resuscitation combined with active rewarming. *Resuscitation* 2001;50(3):301–8.
- Kondratiev TV, Myhre ES, Simonsen O, Nymark TB, Tveita T. Cardiovascular effects of epinephrine during rewarming from hypothermia in an intact animal model. *J Appl Physiol* 1985;100(2):457–64.
- Dietrichs ES, Schanche T, Kondratiev T, Gaustad SE, Sager G, Tveita T. Negative inotropic effects of epinephrine in the presence of increased beta-adrenoceptor sensitivity during hypothermia in a rat model. *Cryobiology* 2015;70(1):9–16.