

Linking high-frequency lacustrine sequences to orbitally-induced cyclicity (Lower Cretaceous, Iberian Basin)

Natalia Illueca Carlos L. Liesa Ana R. Soria

Dpto. de Ciencias de la Tierra, Grupo GEOTransfer - Instituto Universitario de Investigación en Ciencias Ambientales de Aragón (IUCA), Universidad de Zaragoza

Pedro Cerbuna 12, 50009 Zaragoza, Spain

Illueca E-mail: nillueca@unizar.es, Illueca ORCID: 0009-0008-9396-5915

Liesa E-mail: carluis@unizar.es, Liesa ORCID: 0000-0002-9130-117X

Soria E-mail: anasoria@unizar.es, Soria ORCID: 0000-0003-2963-8422

ABSTRACT

The lacustrine El Castellar Formation in the Castillo de Aliaga section (Early Cretaceous Galve sub-basin, eastern Spain) features two carbonate successions of marl and limestone, separated by a mudstone and marl interval. Sequence analysis revealed small-scale (44 elementary sequences), medium-scale (ten complete and two incomplete parasequences), and large-scale (three sets of parasequences, one incomplete), high-frequency lacustrine sequences, with an average thickness of 2.9, 12.4 and 54m, respectively. These sequences start with a sudden facies change (deepening) and exhibit a shallowing-upward trend with features of subaerial exposure (bioturbation, oxidation, or brecciation) at the top, indicating phases of climate-modulated lake expansion and retraction. The temporal framework of the lacustrine sequences is further characterized by the correlation of these sequences with sedimentary-cycle periodicities of 3.3, 13.2 and 57.3m, attributed, respectively, to the long precession cycle (22.4kyr) and the short (95kyr) and long (405kyr) eccentricity cycles of the Earth's orbit. The hierarchical stacking of sequences aligns with orbitally driven cyclicity, with thickness variations interpreted as tectonic subsidence effects (accommodation) resulting from normal fault slip in a rift system. The three sets of parasequences (SPS-1 to SPS-3) align with stages of lake system evolution. SPS-1 represents deposition in a low-energy shallow carbonate lake. SPS-2 indicates a significant lake expansion and deepening, linked with clastic input and a mixed lake system, and correlates with a major increase in accommodation, over ~200kyr, involving fault-induced local tilting. SPS-3 represents deposition in a high-energy carbonate lake. The parasequences identified show variations in cyclic thickness tied to a >700kyr tectonic cycle. The elementary sequences, mainly corresponding to marl-limestone bundles, exhibit thickness changes probably due to shorter-term tectonic pulses. Accelerated tectonic activity resulted in increased accommodation in shorter (40-50kyr) periods, followed by longer (>100kyr) periods of progressive deceleration.

KEYWORDS | Lacustrine facies. Sequence stratigraphy. Climatic forcing. Milankovitch cycles. Tectonic forcing.

INTRODUCTION

Lacustrine systems are depositional settings that are quite sensitive to environmental changes, forced primarily by variations in climate but also in tectonics (e.g. Argylian and Forman, 2003; Armenteros and Huerta, 2006; Gierlowski Kordesch *et al.*, 1991; Jin *et al.*, 2020; Meléndez *et al.*, 2009; Moreno *et al.*, 2012; Sáez *et al.*, 2009; Scheidt *et al.*, 2020; Wei *et al.*, 2018). These factors mainly affect the level and depth of the lake's water, the terrestrial input, the productivity of carbonates, and the flow of groundwater, thus modulating the environmental evolution of the lake and consequently the sedimentary pattern recorded in the lake basin (Carroll and Bohacs, 1999; de Wet *et al.*, 2015; Gómez-Fernández and Meléndez, 1994; Kovács *et al.*, 2021; Ming *et al.*, 2020; Rits *et al.*, 2017; Woolway *et al.*, 2020). Nevertheless, disentangling the signal and response of each of these factors, or their interaction and relative contribution, in relation to the environmental changes recorded in the sedimentation, is not an easy task.

In this regard, cyclostratigraphy is a helpful tool. By conducting spectral analysis of various paleoclimatic proxies in the lake record (e.g. lithology, carbonate content, magnetic susceptibility, the gamma-ray signal, color parameters of the sediments, and distality-ranked lacustrine facies), orbitally induced climate cycles (Milankovitch cycles; Milankovitch, 1941) related to changes in the Earth's insolation can be detected. Interpreting the recorded cyclicity can thus provide valuable insights into past climate variations and trends (e.g. Aziz *et al.*, 2000; Ezquerro *et al.*, 2022a; Juhász *et al.*, 1997; Hoorn *et al.*, 2022; Lepre and Quinn, 2022; Schnyder *et al.*, 2009; Valero *et al.*, 2014; Zhong *et al.*, 2022).

The astronomical forcing of lake sedimentary records has been assessed in both recent (Anselmetti *et al.*, 2006; Camuera *et al.*, 2018; Coianiz *et al.*, 2019; Guitér *et al.*, 2003; Kuzucuoğlu *et al.*, 1997; Tian *et al.*, 2017; Wei *et al.*, 2020; Zhong *et al.*, 2015) and ancient lakes (Gomes *et al.*, 2020; Huang *et al.*, 2020; Ielpi, 2012; Li *et al.*, 2018; Tang *et al.*, 2023; Wei *et al.*, 2023; Xu *et al.*, 2023). Climatic cyclicity is often well recorded even in lakes with active tectonics capable of modifying the accommodation and, therefore, the rate of sediment accumulation (Azenoud *et al.*, 2023; Deino *et al.*, 2006; Luzón *et al.*, 2002; Morrill *et al.*, 2001; Soria *et al.*, 2012; Steenbrink *et al.*, 2006; Walzer and Hendel, 2023; Wang *et al.*, 2012). This is common, for example, in the Cretaceous lacustrine synrift basins of Iberia (Angulo and Muñoz, 2013; Casas *et al.*, 2024; Liesa *et al.*, 2023; Muñoz *et al.*, 2020; Navarrete, 2015; Soria *et al.*, 2012).

Sequence stratigraphy, originally developed to interpret changes in relative sea level in marine successions (Boulila

et al., 2011; Catuneanu, 2022; Emery and Meyers, 2009; Posamentier and Vail, 1988), has also been applied with good results to the study of lake successions. Lake sequences are commonly associated with variations in lake levels driven by climatic and environmental changes, providing evidence of external forcing (Azenoud *et al.*, 2023; Changsong *et al.*, 2001; Keighley *et al.*, 2003; Nutz *et al.*, 2017; Oviatt *et al.*, 1994; Wei *et al.*, 2017; Zavala *et al.*, 2022).

In this study, we present a sequence stratigraphy analysis of the lacustrine synrift record of the Early Cretaceous El Castellar Formation in the Galve sub-basin (western Maestrazgo Basin, eastern Spain). By characterizing lacustrine facies and facies associations, we not only interpret the sedimentary environment and the overall evolutionary trend of the lake, but also analyze high-frequency lacustrine sequences at different scales, along with the behavior and patterns of these sequences throughout the lacustrine succession. These results are then integrated with the orbitally forced climatic cycles (long and short eccentricity, and long precession) recently described by Illueca *et al.* (2023) for the studied deposits. This allows us to link the high-frequency lacustrine sequences of different orders to climate changes induced by variations in the Earth's orbit (Milankovitch cycles), despite the significant influence of tectonics expected in this synrift unit. Variations in the thickness of the lacustrine sequences and cycles are interpreted as tectonically forced changes in accommodation and are used to analyse the sequence stacking and describe patterns of tectonic activity recorded in the lacustrine succession.

GEOLOGICAL SETTING

The Cretaceous Galve sub-basin is located in the eastern Iberian Chain (eastern Spain) (Fig. 1A). This intraplate chain was formed in the context of the Cenozoic collision between the Eurasian and African plates, resulting from the tectonic inversion of a Mesozoic rifting area known as the Iberian Basin (Capote *et al.*, 2002; Liesa *et al.*, 2018, 2023). The Mesozoic rifting occurred in several stages, driven primarily by the opening and spreading of the Tethys Sea and the central-north Atlantic (Capote *et al.*, 2002; Salas and Casas, 1993; Salas *et al.*, 2001). During the Late Jurassic–Early Cretaceous rifting stage, the Iberian Basin became compartmentalized, leading to the development of smaller basins and sub-basins (Martín-Chivelet *et al.*, 2019). The Galve sub-basin was one of these sub-basins that developed at the western margin of the Maestrazgo Basin (Fig. 1A–B). The sedimentary record of the Maestrazgo Basin and its sub-basins comprises two synrift sequences separated by a regional unconformity (Liesa *et al.*, 2019): synrift




FIGURE 1. A) Geological map of the Aliaga region in western Maestrazgo, showing the location (star) of the Castillo de Aliaga section of the El Castellar Formation (modified from Meléndez et al., 2009). The inset displays the location of the studied section within the Iberian Chain, in eastern Iberian Peninsula (after Liesa et al., 2018). B) Sedimentary succession of the Lower Cretaceous El Castellar Formation in the Galve sub-basin, along with the evolutionary stages of the unit (after Meléndez et al., 2009). Section 1 corresponds to the Castillo de Aliaga section studied here. C) Cartography on aerial imagery displaying the limestone packages (bed lines shown as dashed white lines) of the El Castellar Formation and the normal faults (solid white lines) that displace them. The location of the Castillo de Aliaga section logged in this study is shown with solid yellow lines, with numbers indicating the metres in the stratigraphic series. The Middle Surface (MS; dashed pink line) is used to differentiate the evolutionary stages of this formation. Also depicted are its stratigraphic contact with the overlying Camarillas Formation (dashed yellow line) and the boundaries of the parasequences sets (SPS-1 to SPS-3) distinguished in this study.

1 sequence 1 (mid Tithonian to Berriasian) and synrift
2 sequence 2 (Valanginian to early Albian).

3
4 The El Castellar Formation (Salas, 1987), dating
5 from the late Hauterivian to early Barremian, represents
6 the first unit of synrift sequence 2 in the Galve sub-basin
7 (Aurell *et al.*, 2016; Liesa *et al.*, 2019, 2023). This unit
8 is mainly lacustrine with some alluvial deposits and it
9 lies unconformably over synrift sequence 1 and locally on
10 earlier Jurassic rocks (Fig. 1B) (García-Penas and Aurell,
11 2017; Liesa *et al.*, 2004, 2006; Meléndez *et al.*, 2009).
12 Its upper boundary is marked by the input of coarse- to
13 medium-grained white sandstones from the fluvial and
14 estuarine Barremian Camarillas Formation (Navarrete,
15 2015; Soria, 1997; Soria *et al.*, 2023). The thickness of
16 the El Castellar Formation varies significantly (Fig. 1B)
17 due to contemporaneous normal faults trending NNW-SSE
18 and ENE-WSW, which controlled subsidence at different
19 scales, influencing both sedimentation patterns and
20 paleogeography (Liesa *et al.*, 2006, 2019, 2023; Meléndez
21 *et al.*, 2009). This unit consists of a lower alluvial section
22 and an upper lacustrine carbonate section (Fig. 1B), with
23 facies associations belonging to alluvial and palustrine
24 plains, palustrine fringes, a low-energy shallow lake, a
25 mixed siliciclastic-carbonate lake, and a high-energy
26 lake (Aurell *et al.*, 2016; Liesa *et al.*, 2006; Meléndez *et*
27 *al.*, 2009; Soria, 1997; Soria *et al.*, 2001). In addition to
28 the typical biota of Cretaceous lakes (e.g. charophytes,
29 ostracods, and bivalves), this unit also contains dinosaur
30 remains and footprints (e.g. Castanera *et al.*, 2022; Gasca
31 *et al.*, 2009; Ruiz Omeñaca, 2006).

32
33 The sedimentary evolution of the El Castellar
34 Formation in the Galve sub-basin has been related to the
35 interaction and linkage of normal faults that controlled
36 both the basin structure and the subsidence pattern
37 (Liesa *et al.*, 2006). The alluvial and lacustrine sections
38 correspond to different phases of the rifting process:
39 the alluvial deposits represent the initial rifting stage,
40 whereas the lacustrine deposits mark the onset of the
41 rifting climax (Liesa *et al.*, 2006; Meléndez *et al.*, 2009;
42 Soria *et al.*, 2001). During the initial stage, the basin
43 exhibited a highly heterogeneous distribution of alluvial
44 and palustrine facies, driven by the activity of individual
45 listric faults (Liesa *et al.*, 2006; Soria *et al.*, 2001). In
46 contrast, the subsequent stage is characterized by facies
47 homogenization, marked by the development of a large
48 carbonate lake that covered the entire Galve sub-basin
49 (Meléndez *et al.*, 2009). The transition between these
50 stages is notable for the presence of diagenetic gypsum
51 crystals, which are associated with the fault linkage
52 process. This process likely facilitated the circulation
53 and ascent of highly mineralized water from the Triassic
54 Keuper facies, which contains gypsum (Liesa *et al.*, 2006;
55 Meléndez *et al.*, 2009) (Fig. 1B).

1 Recently, Illueca *et al.* (2023) conducted a
2 cyclostratigraphic analysis of lithological and magnetic
3 susceptibility data, identifying three main periodicities
4 (~57.3, ~13.2 and ~3.3m). The authors attributed these
5 periodicities respectively to the long eccentricity (405kyr),
6 short eccentricity (95kyr) and long precession (22.4kyr)
7 cycles of the Earth's orbit, suggesting that these orbitally
8 induced cycles correspond to periodic climatic and
9 environmental variations that influenced sedimentation
10 patterns. Despite the area's highly active extensional
11 context, these variations led to the formation of sedimentary
12 sequences that are both recurrent and recognizable.

13
14 Regarding the paleoclimatic context, the Cretaceous has
15 traditionally been considered an archetypical supergreenhouse
16 period in the Earth's history (e.g. Barron and Washington,
17 1984; Moriya, 2011; O'Connor *et al.*, 2019; Wang *et al.*,
18 2014). The Maestrazgo Basin (eastern Iberian Plate) has been
19 located in the subtropical zone (latitude: 26-35°North) during
20 the Barremian (e.g. Blakey, 2008; Dercourt *et al.*, 1986;
21 Scotese, 2016; Stampfli and Borel, 2002). Deposition has thus
22 traditionally been assumed to occur under subtropical climatic
23 conditions (Buscalioni and Fregenal-Martínez, 2010; Föhlmi,
24 2012; Haywood *et al.*, 2004; Martín-Chivelet *et al.*, 2019;
25 Ziegler *et al.*, 1987) that fall within the temperate, humid
26 subtropical (Ca) climatic zone in the Hauterivian-Barremian
27 paleo-Köppen climate map (Burgener *et al.*, 2023); that is, the
28 climate was fairly warm, humid and seasonal and conducive
29 to significant paleoweathering (Yuste *et al.*, 2015). However,
30 our growing understanding of the global paleoclimate and
31 sea-level changes (e.g. Sames *et al.*, 2016; Hasegawa *et al.*,
32 2022; Rodríguez-López *et al.*, 2022; Wang *et al.*, 2023, and
33 references therein) has revealed more complex and variable
34 paleoclimatic conditions during the Cretaceous, with an
35 episodically active Early Cretaceous cryosphere, both at
36 Palearctic latitudes (Vickers *et al.*, 2019) and paleoplateau
37 altitudes (Rodríguez-López *et al.*, 2022), and with more
38 changing paleoclimates, also in Iberia (e.g. Hay and Floegel,
39 2012; Laita *et al.*, 2020, 2021). The recent discovery of
40 Hauterivian glacial deposits in the Cameros Basin (northern
41 Iberia) (Rodríguez-López *et al.*, 2024), together with recent
42 paleogeographic reconstructions (e.g. King *et al.*, 2021;
43 Rosembaum *et al.*, 2002; Tugend *et al.*, 2015; Vissers and
44 Meijers, 2012) and the high estimated inclinations (55-60°) of
45 the Valanginian-Barremian magnetic vector in the Lusitanian
46 Basin (Moreau *et al.*, 1997) and the Pyrenean Organyà Basin
47 (Gong *et al.*, 2008), point to a latitude of up to 42-50°N for
48 Iberia during the Early Cretaceous, which lies within the
49 temperate climatic zone.

METHODS

50
51 The carbonate interval of the El Castellar Formation
52 (129m) was logged bed by bed at a 1:100 scale in the

1 Castillo de Aliaga section (40°40'37"N and 0°42'26"W).
2 This logging sought to characterize facies and facies trends
3 and to identify high-frequency sequences (Fig. 1B). Given
4 the abundant metric- to hectometric-scale faults with
5 metric to decametric offsets, the outcrop was studied to
6 analyze fault displacements. A detailed aerial orthoimage
7 (1:5000) was also used for this purpose (Fig. 1C). As a
8 result, the section was logged with lateral displacement
9 along the outcrop to ensure a continuous record and
10 prevent repetition. The field study involved describing and
11 documenting the lithology, sedimentary facies, sedimentary
12 structures, and fossil content using standard field tools.
13 Each meter of the section was marked in the outcrop (using
14 markers or paint) to facilitate subsequent observations.
15 During the logging process, 146 samples were collected,
16 comprising 98 rock samples and 48 sediment samples,
17 with at least one sample taken per meter of the section to
18 aid in the comprehensive description of the sedimentary
19 facies and fossil identification. From the collected samples,
20 98 polished sections and 64 selected thin sections (30 μ m
21 thick) of rocks were produced and studied using a binocular
22 loupe and a polarizing petrographic microscope with
23 camera (model OPO 185, KERN Optics). High-resolution
24 images were also captured with an Epson V100 photo
25 scanner (model J231B) and a Plustek OpticFilm scanner
26 (model 8300iSE). Sediment samples were levigated using
27 a sieve shaker machine with 1000, 500 and 100 μ m mesh
28 openings, and the residue was examined with a binocular
29 loupe.

30 The concept of high-frequency sequences of different
31 scales is used here from a descriptive perspective, implying
32 a hierarchical organization where smaller sequences are
33 nested within larger ones (e.g. Mitchum and Van Wagoner,
34 1991). The identification of these sequences, including
35 elementary sequences, parasequences, and parasequence
36 sets, relies on recognizing discontinuity surfaces that
37 indicate breaks in sedimentation and/or vertical facies trends
38 within sequences (e.g. Bosence *et al.*, 2009; Sevillano *et al.*,
39 2020). The lacustrine sequences at different scales mainly
40 correspond to shallowing-upward successions, with features
41 of exposure (oxidation, brecciation and root bioturbation)
42 toward the top of the sequences. The time frame for each
43 sequence type is based on the cyclostratigraphy from Illueca
44 *et al.* (2023), who undertook a spectral analysis of the time
45 series (depth domain) of the magnetic susceptibility and
46 lithological data (data every 0.5m) of the stratigraphic
47 section studied here. Smoothed curves for significant
48 periodicities (95kyr and 22.4kyr cycles) were obtained by
49 applying a Gaussian filter using PAST software (Hammer
50 *et al.*, 2001). The analysis of variations in accommodation
51 over time, following Schwarzacher (2005), involved
52 constructing a graph of accumulated thickness deviations
53 for each stratigraphic sequence or cycle in relation to their
54 respective average thicknesses.

RESULTS

The Castillo de Aliaga section

1 The strata of the El Castellar Formation in the Castillo
2 de Aliaga section exhibit NW-SE to E-W strikes and high
3 dips toward the NE and N. Numerous faults, ranging from
4 meter- to hectometer-scale, cut through strata, resulting
5 in thickness variations in discrete beds of a clearly
6 synsedimentary character (Fig. 1C). The logged section
7 includes three different lithological groups (stratigraphic
8 sections I, II, and III in Fig. 2). The lowest section (I) is
9 27m thick and mainly consists of limestone with a few
10 marl beds. The middle section (II), which is 42m thick, is
11 dominated by variegated marls and clays with a much lower
12 proportion of limestone. Finally, the upper section (III) is
13 58m thick and mainly consists of a succession of marl,
14 marlstone and limestone.

18 The El Castellar Formation is made up of decimeter-
19 to meter-thick tabular strata of marl, marlstone, marly
20 limestone, and limestone, varying in predominance
21 throughout the unit (Fig. 2). Claystone is present to a
22 lesser extent. Occasionally, more detrital beds, such as
23 silts, fine sands, or conglomerates, and mixed siliciclastic-
24 carbonate lithologies (silty/sandy marls and limestones)
25 are intercalated. The limestones and marlstones
26 frequently display oxidation (Fe oxides), bioturbation
27 and brecciation, especially near the top of the beds. The
28 limestones are mainly mudstones and wackestones, with
29 some packstones present. Sedimentary structures such
30 as parallel and cross-lamination, cross-bedding, and
31 symmetric or asymmetric ripples are common in the
32 upper part of the section. The marls are mostly green and
33 gray, and are usually massive and sometimes laminated.
34 The clays are also massive and exhibit a range of colors,
35 including red, violet, ocher and gray. The fossil content is
36 abundant, with a predominance of charophytes (oogonia
37 and stalks), along with bivalve fragments, ostracods, and
38 scarce gastropods and fish teeth.

Facies associations

39 Nine lacustrine facies associations are distinguished
40 (Fig. 3): Low-Energy Shallow Lake (LESL)-1 and LESL-
41 2; Mixed Siliciclastic-Carbonate Lake (MSCL)-1, MSCL-
42 2, and MSCL-3 and High-Energy Lake (HEL)-1, HEL-2,
43 HEL-3 and HEL-4.

44 The LESL facies associations are made up of limestone
45 sequences occasionally found overlying marls and clastic
46 sequences. The limestone evolves from micritic mudstone
47 or wackestone to bioclastic packstone, typically ending
48 with features of subaerial exposure such as brecciation,
49 oxidation, and/or bioturbation.




FIGURE 2. The El Castellar Formation in the Castillo de Aliaga section. Lithologic and textural log including the sedimentary structures, fossil content, subaerial exposure features, and diagenetic structures, as well as the three different stratigraphic sections (I, II and III) identified in the studied section.




FIGURE 3. Facies associations of the lacustrine El Castellar Formation in the Castillo de Aliaga section. A) Distribution of facies associations in the stratigraphic log. B) Ideal stratigraphical sections of the facies associations recognized in the lake system, from proximal to distal areas, evolving over time from a Low-Energy Shallow Lake (LESL), through a Mixed Siliciclastic-Carbonate Lake (MSCL), to a High-Energy Lake (HEL).

LESL-1 is made up of facies sequences from 2 to 4m thick and mainly comprises limestones (Fig. 3B). When present, the marl is gray to green and forms massive, dm-scale tabular beds containing abundant ostracods and charophytes. Occasionally, a lower siliciclastic term occurs, featuring a fining-upward sequence that includes a cm-thick conglomerate with lenticular geometry and homometric, sub-rounded calcareous pebbles, a dm-thick fine-grained sandstone, and a gray tabular silt with parallel lamination. The upper limestones consist of dm- to m-thick tabular beds of mudstone-wackestone textures with charophytes, ostracods, and bivalve fragments (Fig. 4A), transitioning to wackestone to wackestone-packstone textures and occasionally packstones with charophytes (Fig. 4B-C), ostracods, bivalve fragments, and fewer gastropods. The preservation of the fossils is generally good, although some fossils may be fragmented. The tops of these limestone sequences frequently show evidence of subaerial exposure, including oxidation (Fe oxides), but also root bioturbation or brecciation (Fig. 4D). This facies succession mainly occurs in the lower section (I) and occasionally in the middle section (II) of the stratigraphic profile and is vertically related to the LESL2 facies association (Fig. 3A).

LESL-2 consists of 2 to 3m-thick successions of limestones that often overlie dm- to m-scale tabular beds of massive gray to green marls containing abundant ostracods and charophytes. The limestones are dark gray, occur in dm-thick tabular layers, and range from mudstone to wackestone or wackestone-packstone, containing charophytes, bivalve fragments, ostracods, and gastropods (Fig. 3B). The preservation of fossils, especially of charophytes, can be exceptional (Fig. 4E). At the top of the sequences, these limestones occasionally show features of subaerial exposure such as brecciation and oxidation surfaces (Fig. 4F), but without root bioturbation. This facies succession is particularly prevalent in the lower section (I) and occasionally in the middle section (II) of the stratigraphic profile and is related to the LESL-1 and MSCL-3 facies associations (Fig. 3A).

The MSCL facies associations are characterized by mixed siliciclastic and carbonate facies. There is a gradual increase in carbonate content from clays or silts at the base to marls, marlstones, and occasionally limestones at the top, with possible intermediate stages of sandy or silty marls (Fig. 5A). Generally, the marls, marlstones, and limestones also contain a high proportion of terrigenous material.

MSCL-1 forms 3-4m-thick successions that consist of up to 1.5m of massive silts, sandy or silty marls (both with scarce fossil content), and mudstone-wackestone to wackestone-packstone limestones arranged in dm-thick tabular beds, with bivalve fragments, charophytes and ostracods, and locally oncoids (Figs. 3B; 5A, B).

The limestone also exhibits sedimentary features such as brecciation, bioturbation, and oxidation at the top of the beds (Fig. 5D-E). This facies association is identified in the middle part (II) of the section and is vertically related to the MSCL-2 and MSCL-3 facies associations.

MSCL-2 forms 3-4m-thick successions that begin with massive clays, which are red and versicolored on the surface and greenish to greenish-gray in fresh section. These clays often contain displaceable, isolated crystals or aggregates of gypsum, ranging from millimeters to centimeters in size, which are transparent or white (red on the surface). The sequence then transitions to sandy marls, clayey-marls, and finally, marlstones (Fig. 5C) or to charophyte wackestone-packstone limestones, with less oxidation and bioturbation at the top of the beds. This facies association is identified in the middle part (II) of the section and is vertically related to the LESL-2, MSCL-1, and MSCL-3 facies associations.

MSCL-3 consists of m-thick sequences of marl, marlstone, and limestone, with marl being the main lithology (Fig. 2C). The marl and marlstone are green and gray, rich in charophytes and less abundant in ostracods, and are present as massive and sometimes laminated tabular bodies on a meter to decimeter scale. The limestone is gray and exhibits mudstone and wackestone textures with charophytes and ostracods, and is arranged in dm-thick tabular beds. Evidence of subaerial exposure is rare, with only occasional oxidation. This facies association is identified in the middle part (II) of the section and is vertically related to the LESL-1, LESL-2, and especially to the MSCL-1 and MSCL-2 facies associations.

The HEL facies associations consist of a succession of marls, marlstones, and slabby limestones, frequently exhibiting cross-bedding, internal lamination, and ripples (Figs. 3B; 6). Charophytes are the predominant fossils, accompanied by ostracods and bivalves. In general, the fossils are more fragmented and less well preserved than in the LESL facies associations, with wackestone and packstone textures being common (Fig. 6B-C). Features of subaerial exposure are also common at the top of the limestone beds.

HEL-1 consists of 1.5-2.5m-thick successions of green marls, sandy marlstones, and decimeter-thick beds of mudstone to wackestone with charophytes and ostracods (Fig. 3B). Centimeter- to decimeter-thick layers of charophyte mudstone-wackestone are intercalated within the marl. Early diagenetic processes are particularly intense, evidenced by strong mottling due to oxidation, brecciation in the upper parts of the sequences, and especially root bioturbation, which is abundant across all the limestone facies, with roots penetrating deep (up to 20-30cm) into the marls (Fig. 6D). This facies association is located in the




FIGURE 4. Low-Energy Shallow Lake (LESL) facies association. A) Field view of the charophyte mudstone-wackestone of LESL-1 facies association (elementary sequence ES3). B) Thin section (parallel nicols) of mudstone-wackestone facies (LESL-1). C) Thin section (parallel nicols) of packstone facies (uppermost of LESL-1). D) Field view of limestones at the top of the LESL-1 facies association showing root bioturbation, oxidation and brecciation. E) Thin section (parallel nicols) of wackestone facies of LESL-2, with a good preservation of charophyte stems and oogonia (upper left corner). F) Polished section of a wackestone-packstone limestone from an upper level of LESL2, with oxidation and brecciation.

lower part of stratigraphic section III, where it is vertically related to HEL-3 and HEL-4 (Fig. 3A).

HEL-2 typically consists of a succession up to 3.5m thick of gray marls, marlstones, and limestones (Fig. 3B). The marls, up to dm-thick, are massive, contain charophytes and ostracods, and occasionally exhibit nodulization at the top. The marlstones, also up to dm-thick, are laminated, contain charophytes, ostracods, and bivalve fragments,

and occasionally display nodulization as well. Some beds have molds of entire large bivalves (articulated and disarticulated). Locally, there are cm- to dm-thick tabular layers of highly oxidized calcareous sandstones with bivalve molds. The limestones of HEL-2 are micritic, ranging from mudstone to mudstone-wackestone, with charophytes and ostracods, and abundant molds of entire bivalves (Fig. 6E), along with occasional fish teeth. These limestone beds often have oxidized (Fe oxides) tops. HEL-2 is situated




FIGURE 5. Mixed siliciclastic-carbonate lake (MSCL) facies associations. A) Field view of elementary sequences and parasequences at the boundary between stratigraphic sections I and II. Yellow dashed line shows the boundary between parasequence sets SPS-1 and SPS-2 (PS-2–PS-3 boundary). Note the local tilting and erosion of underlaying PS-2 parasequence. B) Thin section (parallel nicols) of an oncolytic limestone (MSCL-1). C) Thin section (parallel nicols) of marlstone (upper term of MSCL-2). D) Field view of the stratigraphic surface of the boundary between the PS-2 and PS-3 parasequences, also the boundary between SPS-1 and SPS-2 (see location in Fig. 5A), showing the significant subaerial exposure, with oxidation, brecciation, and bioturbation. E) Detail of brecciated and oxidized facies shown in D.

in the upper half of stratigraphic section III, where it is vertically related to HEL-3 (Fig. 3A).

HEL-3 forms sequences up to 3.5m thick, consisting of a succession of gray to green marls, marlstones, marly limestones, and bioclastic limestones (Fig. 3B). The marls, arranged in dm-thick tabular beds, are massive or laminated and contain abundant charophytes and

ostracods. Locally, they intercalate with cm-thick beds with wackestone-packstone textures containing charophytes and ostracods. The marlstones and marly limestones contain charophytes, ostracods, and bivalve fragments, and are arranged in dm-thick beds, generally laminated and rarely massive. They are sometimes oxidized and may present ripples, with intercalated cm-thick layers of bioclastic packstone. The limestones, which are




FIGURE 6. High-Energy Lake (HEL) facies associations. A) Field view of elementary sequences and parasequences of the base of stratigraphic section III and the boundary between SPS-2 and SPS-3 (dashed yellow line). B) and C) Thin sections (parallel nicols) show charophyte packstone facies from the HEL facies association, with occasional ostracods. D) Field view of palustrine HEL-1 facies, characterized by highly oxidized limestones with vertical root bioturbation extending into the marl. E) Field view of the limestones term of HEL-2 facies association, containing abundant entire bivalve moulds. F) and G) Field views of high-energy littoral bar facies from the HEL-3 facies association, exhibiting lamination, ripples, and cross-bedding.

occasionally silty, are mainly bioclastic wackestones and packstones, with abundant charophytes, ostracods, bivalve fragments, and a lower content of gastropods (Fig. 6B-C). These limestones are frequently laminated or slabby and are arranged in decimeter- to meter- thick tabular bundles, exhibiting internal cross-lamination and current or oscillation ripples (Fig. 6F). Toward the top, they often show signs of oxidation and bioturbation, mainly vertical root traces, and occasionally horizontal bioturbation, with less frequent brecciation. The sequence ends with dm- to m-thick tabular beds of bioclastic limestones, ranging from wackestone to packstone with charophytes, ostracods, and bivalves, which may intercalate with cm-thick marl beds. The facies are intensely laminated and slabby, with dm-scale cross-stratification and abundant ripples (Fig. 6G). The limestones also show frequent oxidation, bioturbation, or brecciation on their upper surfaces. Facies association HEL-3 is located in the lower half of stratigraphic section III and is vertically related to HEL-2, as well as to HEL-1 and HEL-4 (Fig. 3A).

HEL-4 consists of a thick lower unit (3 to 5m thick) of green massive marls with charophytes, and a thin bed of limestone (wackestone) with charophytes, ostracods, and bivalve fragments, showing some oxidation at the top (Fig. 3B). Facies association HEL-4 is located at the base of stratigraphic section III and is vertically related to MSCL-2, HEL-1 and HEL-3 (Fig. 3A).

Interpretation of the facies associations and sedimentary systems

The nine lacustrine facies associations that are distinguished are interpreted as representing deposition in three different lake systems that evolved over time (Fig. 3): a LESL, MSCL and HEL.

Low-Energy Shallow Lake (LESL)

The facies characteristics and vertical relationships of the LESL facies associations indicate that they represent low-energy, shallowing lacustrine sequences interrupted by occasional, more pronounced deepening events (Fig. 3B). These sequences were the result of the progressive infilling of a very shallow, hard water lake where charophytes were the main carbonate source, under conditions similar to those described by Murphy and Wilkinson (1980). LESL-1 and LESL-2 correspond to progressively more distal carbonate facies (Fig. 3B). In this context, the marl lithology represents sediment settling in the more distal parts of the lake system, whereas the micritic carbonate facies (mudstone and wackestone) suggest deposition under low-energy conditions in the proximal littoral or sublittoral areas. The charophyte and ostracod packstone facies, containing fragments of bivalves and gastropods,

are indicative of deposition in littoral zones during more energetic episodes. These episodes, likely associated with storms or floods, would have caused the fragmentation and accumulation of bioclasts and introduced sediments from the distal alluvial-fluvial system. This process contributed to the formation of cm-thick conglomerates, fine-grained sandstones, and laminated silts in the littoral lacustrine areas.

The LESL facies associations correspond to sequences of infilling in the low-gradient margins of a shallow, poorly developed lake (Platt and Wright, 1991). The predominance of mudstone/wackestone, the good preservation of fossils, and the absence of wave or current structures suggest that these were isolated hard water lakes with low-energy margins. In such environments, the thickest accumulation of carbonate occurs in littoral areas due to the high productivity of benthic organisms (Platt and Wright, 1991; Alonso Zarza *et al.*, 2006; Meléndez *et al.*, 2009). These lakes are highly sensitive to minor fluctuations in lake level, which can lead to subaerial exposure of more proximal deposits and the development of palustrine-type facies, such as oxidized, brecciated, or bioturbated limestones (Alonso Zarza and Wright, 2010). However, the extent of subaerial exposure is relatively limited (see the exposure index given by Platt and Wright, 1992), indicating a stable system with minimal sedimentary interruptions or erosive episodes. Consequently, features such as paleosols, microkarsts or calcrites are absent (Alonso Zarza and Wright, 2010; Luzón *et al.*, 2002).

Mixed Siliciclastic-Carbonate Lake (MSCL)

The three distinct facies associations that are identified represent deposition in different settings within a mixed siliciclastic-carbonate lacustrine system (Fig. 3B): proximal (MSCL-1), intermediate (MSCL-2), and distal (MSCL-3). This lacustrine system mainly occurs in the middle part (section II) of the stratigraphic profile and the distinct facies associations exhibit a vertical relationship with one another (Fig. 3A).

The MSCL facies associations are interpreted as shallowing-upward lacustrine sequences, with an increasing carbonate content in a lake environment characterized by higher siliciclastic input and/or lower carbonate production. During periods of high sediment supply to the lake, terrigenous facies (silts and sandy marls) were mainly deposited in the proximal areas of the lacustrine system (MSCL-1), whereas clays settled by decantation in the intermediate areas (MSCL-2). In the central parts of the lake, further from the terrigenous sources, marl deposition dominated (MSCL-3). As the sediment supply diminished, carbonate production within the lake resumed, leading to the deposition of marls, followed by limestones, in both the

marginal and central lake areas. Accordingly, this sequence reflects a progressive decrease in terrigenous input to the lake. The proportions of limestone relative to marl, along with the indicators of subaerial exposure (such as oxidation and bioturbation) in the limestones at the top of the lacustrine sequences, are greater in the proximal areas (MSCL-1) than in the intermediate (MSCL-2) and distal areas (MSCL-3) of the lake. The gypsum crystals found in the massive greenish-gray clays of MSCL-2, as suggested by Liesa *et al.* (2006), are interpreted as late diagenetic in origin, having formed in sediments saturated with highly mineralized water.

High-Energy Lake (HEL)

The four identified HEL facies associations characterize a high-energy lake system (Fig. 3B), which occurs predominantly in the upper part (section III) of the stratigraphic profile (Figs. 3A; 6A).

The HEL facies associations represent deposition in different subenvironments within this lake system (Fig. 3B), ranging from the palustrine fringe (HEL-1), through the high-energy littoral bar (HEL-3), to the sublittoral environment (HEL-4). HEL-2 reflects sedimentation in restricted areas protected by the littoral bar. HEL-1 is interpreted as deposition in a low-gradient palustrine belt characterized by vegetated marshes or ponds, where marls and limestones formed and where the vegetation bioturbated and mottled the sediments. The sandy facies are attributed to detrital inputs from alluvial flows entering the palustrine belt. This facies association constitutes a marginal upward-shallowing sequence, in which the palustrine area gradually expands until the sediments are fully exposed, leading to the development of paleosols during prolonged periods of vegetation colonization. HEL-2 is interpreted as having formed under low-energy conditions in restricted and protected areas where mud facies were deposited, giving rise to significant accumulations and preservation of bivalves, along with some siliciclastic inputs from nearby areas. HEL-3 corresponds to the progradational sequence of a lacustrine littoral bar that developed in high-energy settings. Massive or laminated marls and thin layers of tabular (sometimes laminated) bioclastic limestone represent bottomset deposits, whereas dm-scale cross-bedded bioclastic limestones correspond to the foreset migration of bioclastic bars with superimposed ripples, which developed in high-energy lake environments. Finally, HEL-4 represents the most distal facies associated with this high-energy system, characterized by thick deposits of marls that reflect the settling of suspended sediments in more central lacustrine areas. The few limestones at the top of the sequences, showing local subaerial exposure, represent sedimentation during periods of low lake levels,

indicating a shallowing upward sequence very similar to those in the low-energy systems (*i.e.* MSCL-3).

The HEL facies associations are interpreted as representing lacustrine sedimentation in a shallow, extensive, high-energy, low-gradient (ramp-type) lake (Platt and Wright, 1991). Large-scale bars and ripples would have developed in littoral zones under high-energy conditions characterized by currents and wave action, as described in other lake systems from the Lower Cretaceous of the Iberian Basin (Fregenal-Martínez and Meléndez, 2000; Meléndez *et al.*, 2000). Fluctuations in lake level would have controlled the deposition and stacking of different sequences, such as the superimposition of palustrine environments over lacustrine ones, or the sudden deepening of the system. Therefore, this HEL system, although more developed and extensive than the systems described above, would have still been a shallow lake sensitive to these oscillations in water level, often undergoing subaerial exposure and root bioturbation during periods of low lake level.

High-frequency sequences

The analysis of bounding surfaces and facies trends makes it possible to recognize hierarchically stacked stratigraphic sequences at three distinct scales within the lacustrine evolution of the El Castellar Formation (Fig. 7A–B): the Elementary Sequence (ES), the parasequence (PS), and the Set of Parasequences (SPS).

Elementary Sequences (ES) are high-frequency, small-scale sequences with an average thickness of ~2.91m (ranging from 1.25 to 5.25m; mode: 2.75-3.24m) (Fig. 8A). A total of 44 elementary sequences (ES-1 to ES-44) are identified (Fig. 7B). The ESs are bounded by distinct surfaces, typically characterized by a sudden deepening of the lacustrine facies that overlie the shallower facies of the preceding elementary sequence. These previous facies often exhibit features of subaerial exposure, such as brecciation, oxidation, and/or root bioturbation. Differentiating elementary sequences in distal facies composed of marls and clays is challenging, so some sequence boundaries are defined tentatively (ES-16 to ES-17). ESs mainly correspond to the facies associations described in the previous section, representing sequences of shallowing lacustrine settings.

Parasequences (PS) are medium-scale, high-frequency sequences with an average thickness of ~12.4m (ranging from 9.75 to 17.5m) (Fig. 8B). A total of ten complete parasequences (PS-1 to PS-10) and two incomplete ones (PS-0 and PS-11) are recognized (Fig. 7B). The complete parasequences typically consist of four elementary sequences (PS-1, PS-2, PS-3, PS-5, PS-8, PS-9, PS-10, and probably PS-4) or occasionally five (PS-6 and PS-7), with




FIGURE 7. Linking high-frequency sequences of different orders with orbitally induced cyclicity in the lacustrine El Castellar Formation. A) Stratigraphic record. B) Facies associations and high-frequency sequences of small (elementary sequences), medium (parasequences) and large scale (sets of parasequences) recognized and their vertical facies trend (sequences numbered in correlative order). C) Raw and smoothed curves from cyclostratigraphic time series data, including Magnetic Susceptibility (MS) and lithology, as well as the 95kyr cycles identified in the stratigraphic section (after Illueca et al., 2023). Smoothing was carried out using a 25 data (12.5m) or 6 data (3m) Gaussian filter for the 95kyr and 22.4kyr filter, respectively. Note the inverse correlation between the two curves and the eleven maxima in the 95kyr MS filter (orange bands), which Illueca et al. (2023) used to identify ten complete and two incomplete cycles throughout the profile.

an average of 4.2 ESs per PS. Their thickness generally increases from PS-2 to PS-6 and then progressively decreases. The parasequence boundaries are marked by clear stratification surfaces defined by sharp lithological changes, corresponding to deeper facies overlying the shallower facies of the previous parasequence, often exhibiting signs of subaerial exposure (such as brecciation, oxidation, and/or root bioturbation). Most parasequences show a vertical progradational evolution, characterized by a shallowing-upward trend (PS-3 to PS-7 and PS-10). However, some show a significant aggradational component with a relatively progressive deepening at the base of the parasequence (PS-1, PS-2, PS-8, and PS-9) (Fig. 7B). The shallowing-upward trend is represented by the succession of distal LESL-2 and proximal LESL-1 facies (PS-1 and PS-2) in the LESL (stratigraphic section I); by the succession, whether complete or incomplete, of distal carbonate (MSCL-3), distal mixed (MSCL-2), and proximal mixed (MSCL-1) facies (PS-4 and PS-5), and locally of LESL-2 and LESL-1 (PS-3), in the MSCL (section II); and by the succession of sublittoral (HEL-4), littoral bar (HEL-3) and palustrine (HEL-1) facies (PS-6 and PS-7) or the succession of littoral bar (HEL-3) and restricted area (HEL-2) facies (PS-8 to PS-10) in the HEL (section III).

Sets of Parasequences (SPS) are large-scale, high-frequency sequences each about 54m in thickness. Three SPSs are identified (SPS-1 to SPS-3), with SPS-1 being incomplete (Fig. 7B). These sets correspond to three distinct lithological sections bounded by key flooding surfaces. SPS-1, which is 27m thick and incomplete, includes two parasequences (PS-1 and PS-2) and is made up of limestone from the proximal areas of a low-energy lake system. SPS-2 marks the beginning of significant clastic supply and the development of a low-energy mixed siliciclastic-carbonate lake system. SPS-2 is 54m thick, comprises four stacked parasequences (PS-3 to PS-6), and is essentially composed of mixed siliciclastic-carbonate lake shales and marls. SPS-3 is 48m thick, includes four parasequences (PS-7 to PS-10) and the elementary sequence ES-44, and is mainly composed of limestone sedimented in a lacustrine littoral bar and proximal areas of a high-energy lacustrine system. SPS-3 marks the onset of sublittoral facies over the palustrine environments of SPS-2 in the high-energy lake system. Its upper limit is defined by a sharp change of facies from the El Castellar Formation to the fluvial deposits of the Camarillas Formation. The sedimentary evolution is aggradational for SPS-1 and shallowing for the two other sets, reflecting an evolution consistent with the individual parasequences within each set. It is worth noting that even though SPS-1, SPS-2, and SPS-3 correspond approximately to lithological groups I, II, and III (Fig. 7A-B), there are discrepancies. For example, SPS-2 extends up to meter 81 instead of 68.5 (the top of II) and includes parasequence PS-6. The upper limit of SPS-3

is based primarily on a significant flooding event, marked by the transition from sublittoral to palustrine facies.

DISCUSSION

Linking high-frequency lacustrine sequences to orbitally-forced cyclicity

The high-frequency sequences within the lacustrine El Castellar Formation mainly constitute shallowing-upward sequences, stacked hierarchically at different scales (elementary sequences, parasequences, and sets of parasequences). These stratigraphic units are characterized by clear bounding surfaces that indicate abrupt changes in sedimentation and facies trends. The boundaries between different sequence orders mainly correspond to lacustrine flooding surfaces, which often overlie shallower facies exhibiting signs of subaerial exposure such as brecciation, oxidation surfaces, and root bioturbation. The development of the sequences and their facies trend is closely linked to changes in lake levels driven by climatic variations that controlled the supply of water and clastics to the hard water lake, where bioinduced carbonate production occurred.

Furthermore, in our case study, these stratigraphic sequences of varying orders can be correlated with different orbital cycles. The average thickness of elementary sequences (~2.9m; range 1.25 to 5.25m), parasequences (~12.4m; range 9.75 to 17.5m), and sets of parasequences (~54m) in the lacustrine deposits (Fig. 8) corresponds closely to the main periodicities of ~3.3, ~13.2 and ~57.3m, respectively. These periodicities were inferred from the spectral analysis of the time series of the magnetic susceptibility and lithology of this same section of the El Castellar Formation (Illueca *et al.*, 2023) (Fig. 7C). Accordingly, these high-frequency lacustrine sequences can be linked to this cyclicity, which was driven by orbital parameters (Illueca *et al.*, 2023). The elementary sequences are thus associated with the cyclical climate control of sedimentation induced by the long precession cycle (22.4kyr), which affects the orientation of the Earth's rotation axis. Similarly, parasequences and sets of parasequences are sedimentary cycles forced respectively by the short (95kyr) and long (405kyr) cycles of the Earth's orbit around the Sun (Fig. 7B-C).

The link between high-frequency lacustrine sequences and orbitally forced cyclicity not only provides a temporal framework for the former but also offers an additional line of evidence for correlating sedimentary cycles and orbital cycles. This is achieved by examining the number of stacked sequences relative to the corresponding time frame. For instance, each parasequence (95kyr) comprises four elementary sequences (22.4kyr) in eight cases

and five in two cases, resulting in an average ratio of 4.2 elementary sequences per parasequence (Figs. 7; 8). This ratio aligns the time span of these medium- and small-scale high-frequency sequences with the expected relationship between the orbital cycles (4.24 precession cycles per short eccentricity cycle). Similarly, the sets of parasequences (405kyr) stack four parasequences (95kyr) in SPS-2, or four parasequences plus one elementary sequence in SPS-3, which also aligns with the ratio of 4.26 between the eccentricity orbital cycles.

An additional constraint can be established by comparing the lacustrine parasequences with the 95kyr cycles defined from the smoothed data curves (25 data points, 12.5m intervals) of the magnetic susceptibility and lithology data series (Fig. 7C). Illueca *et al.* (2023) used the maxima in magnetic susceptibility (and minima in lithology) in these smoothed curves as references to distinguish short eccentricity cycles (orange bars). They identified ten complete 95kyr cycles with an average thickness of 12.3m, which is nearly identical to the average thickness of the parasequences (12.4m) (Fig. 8A-C). The correlation between the parasequences and the cycles established from the smoothed curves is strong both in terms of the number of cycles and the distribution of their thicknesses (Fig. 7B-C). For instance, the thickest parasequence, PS-6, corresponds to the thickest cycle, cycle-6, whereas the thinner PS-9 and PS-10 align with the thinner cycle-9 and cycle-10, respectively.

However, the boundaries of the parasequences appear slightly out of phase with the magnetic susceptibility maxima used to define the 95kyr cycles (Illueca *et al.*, 2023), with only three instances where they practically coincide (PS-1, PS-2, and PS-10). Generally, the boundaries of the parasequences coincide with or are close to the inflection points of the curves. This is consistent with observations in the study of sequences and cyclic behavior, such as sequence stratigraphy (Christie-Blick, 1991; Posamentier and Vail, 1988; Wright and Marriott, 1993). In dynamic systems such as sedimentary environments, the inflection point marks the change in behavior, defining a sequence boundary (Simmons, 2016). In sequence stratigraphy, for example, bounding unconformities are related to the inflection points of eustatic falls (Posamentier and Vail, 1988), or boundaries of seismic sequences are located near the inflection points of eustatic sea-level fluctuations, where the rates of fall or rise are maximized (Jervey, 1988). The small observed mismatches may result from variations in the thickness of sedimentary sequences and from the analytical procedure itself, which may introduce minor errors during curve smoothing using a Gaussian filter.

Finally, the small-scale (elementary sequence) and medium-scale (parasequence) high-frequency lacustrine

sequences, like the recorded long precession (22.4kyr) and short eccentricity (95kyr) cycles, exhibit variations in thickness over time within the stratigraphic series (Figs. 7; 8). These variations in thickness are comparable, which also highlights the link between high-frequency lacustrine sequences and climatically driven, orbitally induced cyclicity (Fig. 8).

Consequently, the high-frequency shallowing-upward sequences represent phases of lacustrine expansion and retraction caused by cyclic variations in orbital parameters, specifically the Earth's orbital eccentricity and precession cycles. These sequences may therefore be associated with climatic factors that correspond to changes in lake levels related to water supply, aridity, or variations in the water table (Bohacs *et al.*, 2000; Gierlowski-Kordesch *et al.*, 1991; Gómez-Fernández and Meléndez, 1994; Soria *et al.*, 2008, 2012). According to Morrill *et al.* (2001), orbital forcing can significantly impact the water balance of lake systems, especially in the absence of large salinity changes, as observed in our section. This impact is primarily driven by changes in lake evaporation and runoff coefficients, both of which are influenced by climatic factors such as short-wave radiation, temperature, and precipitation.

It is not clear how the effects of these orbital variations were transferred through the climate system and into the lacustrine sedimentary record (e.g. Morrill *et al.*, 2001; Ruddiman, 2006). Periods of tilt and precession force changes in insolation and greenhouse gases responsible for variations in climate and in the evolution of sedimentary systems. Despite negligible insolation forcing at the period of eccentricity, it has been suggested that the 100kyr periods are the result of internal resonance and feedback between the tilt and precession periods (Ruddiman, 2006). In our opinion, latitudinal variations in the climatic zones induced by orbital forcing cannot be ruled out, which may be responsible for cyclic sedimentary and climatic changes over long periods of time. This is particularly relevant for the study area, situated in a mid-latitude zone that may fall within the influence of either temperate or subtropical climatic belts. The paleogeographic changes in land and sea masses caused by the spreading of the southern North Atlantic during the Early Cretaceous (King *et al.*, 2021; Tugend *et al.*, 2015) could have also influenced regional climatic changes. Whatever the case, a detailed study of paleoclimatic proxies (e.g. conventional or clumped $d^{18}\text{O}$ and $d^{13}\text{C}$ carbonate isotopes) is necessary to characterize the relevance of these paleoclimatic changes, *i.e.* to resolve the question of whether the orbital cycles resulted in changes to the point of completely resetting the dominant climate, or whether some variables changed but the same climate persisted.




FIGURE 8. Analysis of thickness variation in high-frequency sequences, 95kyr cycles, and tectonic forcing on stacked sequences. A) Scatter plots and histograms illustrating the thicknesses of elementary sequences and parasequences, with the average thickness for each sequence type also indicated. B) Cumulative thickness deviation diagram (based on Schwarzacher, 2005; see methods) for elementary sequences and parasequences, depicting variations in the creation (mainly tectonic) of accommodation. In this graph, a drop in the curve indicates that the sequence/cycle is thinner than the average, while a rise indicates a thickness greater than the average. Thus, a positive slope indicates an increase in accommodation (accelerated tectonic subsidence), whereas a negative slope indicates a decrease (decelerated tectonic subsidence). C) Plots showing thickness and cumulative thickness deviation for short eccentricity cycles (95kyr) identified from the smoothed magnetic susceptibility curve (see Fig. 7C). Note the similarity among the different curves in relation to changes in thickness.

1 Tectonic control on sequence stacking

2 Changes in the thickness of sedimentary sequences
 3 and cyclicity of different temporal orders indicate changes
 4 in the rate of sediment accumulation over time (Fig. 8A–
 5 C). These variations in thickness within a synrift unit
 6 (Liesa *et al.*, 2004, 2006; Salas, 1987; Soria, 1997)
 7 are interpreted as having been strongly controlled by the
 8 generation of accommodation associated with tectonic
 9 processes, particularly tectonic subsidence. Variations in
 10 fault displacement likely caused changes in subsidence
 11 rates, thereby affecting the thickness of the recorded cycles,
 12 as also observed in other lacustrine systems (e.g. Casas *et*
 13 *al.*, 2024; Muñoz *et al.*, 2020; Soria *et al.*, 2012).

16 The creation of accommodation driven by tectonics is
 17 consistent with the significant faulting associated with the
 18 deposition of the El Castellar Formation in the Galve sub-
 19 basin (Aurell *et al.*, 2016; Liesa *et al.*, 2004, 2006, 2019;
 20 Meléndez *et al.*, 2009; Soria, 1997). Numerous normal faults
 21 in the study area displaced and altered the thickness of the
 22 sedimentary series (Fig. 1C). For example, the stratigraphic
 23 surface between SPS-1 and SPS-2 (the PS-2–PS-3 boundary)
 24 is marked by a major episode of subaerial exposure resulting
 25 from tilting and local erosion (Fig. 5A). Here, bed tilting
 26 was caused by the sliding of normal faults on a metric to
 27 decametric scale, which also led to the thickening of the
 28 clayey elementary sequences ES-10 and ES11 (Fig. 1C).

30 In this context, variations in the thickness of orbital
 31 cycles recorded in stratigraphic sequences can serve as
 32 a tool for analyzing tectonic activity associated with
 33 normal fault displacement. Such analysis relies on two
 34 key assumptions: first, that sediments fill the depositional
 35 space generated by fault displacement; and second, that
 36 most shallowing-upward lacustrine sequences culminate
 37 in facies indicating subaerial exposure, meaning the
 38 accommodation available was filled. Accordingly, the
 39 observed increase in parasequence thickness (and 95kyr
 40 cycles) from PS-2 to PS-6 (cycle 2 to cycle 6), followed by
 41 the progressive decrease until PS-10 (Fig. 8A–C), outlines
 42 a cycle of rising and then declining tectonic activity. This
 43 tectonic cycle spans approximately 30 elementary sequences
 44 (from ES-15 to ES-44), covering around 700kyr, which is
 45 comparable to certain tectonic cycles documented in the
 46 Teruel rift basin (Ezquerro, 2017; Ezquerro *et al.*, 2022b).
 47 Given the average thickness (without decompression) of
 48 parasequences (12.4m) and elementary sequences (2.9m),
 49 along with their respective time spans (95 and 22.4kyr), the
 50 average accommodation rate in the lake system was 130m/
 51 Myr. However, the accommodation rate varied between 102
 52 and 184m/Myr for different parasequences, and between 67
 53 and 245m/Myr for elementary sequences. These values are
 54 typical of rift systems (Casas *et al.*, 2024; Ezquerro *et al.*,
 55 2020; Friedmann and Burbank, 1995; Leeder, 1991).

1 The deviations in the accumulated thickness observed
 2 in each elementary sequence, parasequence (Fig. 8B), and
 3 95kyr cycle (Fig. 8C), relative to their respective average
 4 thicknesses, reveal that the increase in accommodation
 5 mainly occurred during the PS-5 to PS-7 parasequences
 6 (95kyr cycles 5 to 7). This period of enhanced subsidence
 7 was both preceded and followed by periods characterized
 8 by reduced tectonic subsidence. These periods roughly
 9 correlate with, and could be responsible for, the three sets
 10 of parasequences (SPS-1, SPS-2 and SPS-3) and the three
 11 stages distinguished in the evolution of the lake system:
 12 LESL, MSCL and HEL.

13 Thus, LESL facies associations are characteristic of
 14 overfilled lake basins where the sedimentation rate exceeds
 15 the available accommodation (Alonso Zarza *et al.*, 2006;
 16 Carroll and Bohacs, 1999; Meléndez *et al.*, 2009). This
 17 scenario is consistent with the low subsidence expected
 18 during the initial rifting phase (Liesa *et al.*, 2006; Meléndez
 19 *et al.*, 2009). Subsequently, the lake system evolved into a
 20 mixed lake (MSCL) due to an increase in clastic supply
 21 and/or a decrease in carbonate production, associated with
 22 an increase in tectonic subsidence that marks the transition
 23 to the rifting climax (Liesa *et al.*, 2006). This phase was
 24 responsible for the abrupt deepening associated with SPS-
 25 2. The increased clastic supply could have been driven by
 26 a rejuvenation of relief caused by tectonic uplift during the
 27 rifting period. However, the influence of climatic factors
 28 cannot be ruled out, as increased humidity and runoff
 29 could also have led to an increase in surface water reaching
 30 the lake, thereby increasing clastic supply and causing
 31 further lake deepening. Furthermore, the supply of sulfate
 32 from groundwater into the lake might have altered the
 33 environmental conditions, disrupting carbonate production.
 34 Finally, as tectonic subsidence slowed, the clastic input to
 35 the basin decreased, and carbonate production resumed in
 36 a larger, more energetic carbonate lake. During this stage,
 37 typical lacustrine littoral facies characteristic of a balanced-
 38 fill lake basin developed, as suggested by the more complete
 39 lacustrine sequences identified in this phase (Meléndez *et*
 40 *al.*, 2009).

41 The results derived from the analysis of elementary
 42 sequence thicknesses are consistent with those observed
 43 in the parasequences and 95kyr cycles (Fig. 8B–C), further
 44 supporting the idea that tectonic forcing influenced
 45 subsidence changes across different time scales. However,
 46 the elementary sequences also exhibit a more pulsating
 47 variation, with thicker sequences (3.5–4.0m; e.g. ES-17,
 48 ES-19, ES-33, ES-35 or ES-36) alternating with thinner
 49 ones (1.5–2.0m; e.g. ES-15, ES-18, ES-34 or ES-37). This
 50 probably indicates that, over shorter time periods, tectonic
 51 subsidence occurred in a more discontinuous manner
 52 due to the pulsating activity of local normal faults. This
 53 could explain the periodicity of ~1.5m also identified by

1 Illueca *et al.* (2023). It is plausible that this periodicity
 2 corresponds to the precession cycle during periods of lower
 3 sedimentation rates, rather than a half-precession cycle
 4 (~10kyr; Berger and Loutre, 1997; Boulila *et al.*, 2010; Park
 5 *et al.*, 1993) as proposed by Illueca *et al.* (2023). Similar
 6 instances of multiple periodicities linked to a single orbital
 7 cycle, associated with changes in fault activity, have been
 8 documented in other lacustrine synrift deposits (Casas *et*
 9 *al.*, 2024; Muñoz *et al.*, 2020; Soria *et al.*, 2012).

11 Examined in detail, the cumulative thickness deviation
 12 curve of elementary sequences reveals two periods of
 13 enhanced tectonic subsidence: a main period from ES-
 14 15 to ES-23 (PS-4 to PS-6) and an initial, shorter phase
 15 of accommodation creation during ES-4 and ES-5 (PS-
 16 1). Both phases were followed by periods of reduced
 17 accommodation creation (Fig. 8B). These findings suggest
 18 that the cycles of increasing and decreasing tectonics are
 19 asymmetric, with shorter durations for accelerating tectonic
 20 subsidence compared to longer periods of deceleration.
 21 Accelerated tectonic periods typically span two consecutive
 22 elementary sequences (40-50kyr), whereas decelerating
 23 tectonic periods are longer, usually exceeding 100kyr.
 24 Exceptions occur when several accelerating tectonic periods
 25 cluster to form a higher-order period, such as the one
 26 identified between ES-15 and ES-23. In such cases, brief
 27 periods of quiescence separate two periods of increasing
 28 tectonic activity. This pattern mirrors the slip history of
 29 faults observed in recent rift systems (e.g. Ezquerro *et al.*,
 30 2015, 2016; Simón *et al.*, 2016).

32 Despite the significant influence of tectonics as the
 33 main factor generating accommodation and driving
 34 changes in the thicknesses of the sedimentary sequences
 35 and cycles across various time scales, the sedimentary and
 36 climatic cyclicity was not entirely obscured. This study has
 37 not accounted for other potential factors that might disrupt
 38 cyclicity, such as diagenetic processes (*i.e.* cementation,
 39 dissolution), sediment compaction, or variations in
 40 carbonate production and clastic supply during basin filling.
 41 Even though these processes can distort and complicate
 42 an assessment of astronomical influence (Meyers, 2019),
 43 they are interpreted as having less impact on the thickness
 44 and stacking patterns of lacustrine sequences compared to
 45 extensional tectonics. It is likely that their effects were more
 46 pronounced in smaller-scale sequences.

47 CONCLUSIONS

48 The nine lacustrine facies associations distinguished
 49 in the synrift El Castellar Formation mainly correspond to
 50 shallowing-upward sequences that developed in different
 51 subenvironments within a shallow lacustrine system. This
 52 system evolved through three stages over time: a low-

53 energy shallow lake, a mixed siliciclastic-carbonate lake,
 54 and a high-energy lake. Three hierarchically stacked high-
 55 frequency lacustrine sequences are distinguished, resulting
 56 in the identification of 44 elementary sequences, ten
 57 complete parasequences (and two incomplete ones), and
 58 three sets of parasequences (one of them incomplete). The
 59 average thicknesses of these stratigraphic sequences are
 60 2.9m (ranging from 1.25 to 5.25m), 12.4m (9.75 to 17.5m),
 61 and approximately 54m, respectively. The stratigraphic
 62 sequences are bounded by clear stratification surfaces that
 63 reflect flooding events following episodes of subaerial
 64 exposure. The elementary sequences, parasequences, and
 65 sets of parasequences align with the main periodicities of
 66 3.3, 13.2 and 57m, corresponding respectively to the long
 67 precession cycle (22.4kyr) and to the short (95kyr) and
 68 long (405kyr) eccentricity cycles of the Earth's orbit. These
 69 correlations provide a robust linkage between the high-
 70 frequency lacustrine sequences and climate forcing driven
 71 by the Earth's orbital variations.

72 Variations in the thicknesses of elementary sequences
 73 and parasequences, as well as 95kyr cycles, are attributed
 74 to changes in accommodation related to tectonic activity
 75 associated with normal faults. An asymmetric cycle of
 76 >700kyr, characterized by alternating phases of increasing
 77 and decreasing tectonics, marks the transition from rift
 78 initiation to rifting climax stages, with an accelerated
 79 tectonic phase lasting around 200kyr. The changes in
 80 accommodation exhibit brief periods of increased tectonic
 81 subsidence lasting about 40-50kyr followed by longer
 82 (>100kyr) periods of decreased subsidence. The major
 83 phase of accelerated tectonics contributed to the deepening
 84 of the lake, an increased clastic supply, and the development
 85 of a mixed siliciclastic-carbonate lake system.

86 DECLARATION OF COMPETING INTEREST

87 The authors declare that they have no known competing
 88 financial interests or personal relationships that could have
 89 appeared to influence the work reported in this paper.

90 ACKNOWLEDGMENTS

91 The authors thank Marian Fregenal-Martínez, an anonymous
 92 reviewer, and editor Telm Bover-Arnal for their valuable
 93 comments and suggestions, and Rupert Glasgow for improving
 94 the English text. We would like to acknowledge the Servicio
 95 General de Apoyo a la Investigación-SAI from the Universidad
 96 de Zaragoza. This research has been financed by the Agencia
 97 Estatal de Investigación (MCIU/AEI/10.13039/501100011033)
 98 of the Spanish Government (grants PID2019-108705-GB-I00 and
 99 PID2023-148949NB-I00) and the Gobierno de Aragón (grant
 100 E32_23R of the GEOTransfer Research Group). N.I. benefits

1 from an INVESTIGO Program contract within the Recovery,
 2 Transformation and Resilience, financed by the European Union-
 3 NextGenerationEU.

6 REFERENCES |

7 Alonso-Zarza, A.M., Wright, V.P., 2010. Palustrine carbonates.
 8 In: Alonso-Zarza, A.M., Tanner, L.H. (eds.). *Carbonates in
 9 Continental Settings: Geochemistry, Diagenesis and
 10 Applications. Developments in Sedimentology*, 61, 103-132.
 11 DOI: [https://doi.org/10.1016/S0070-4571\(09\)06102-0](https://doi.org/10.1016/S0070-4571(09)06102-0)

12 Alonso-Zarza, A.M., Dorado-Valiño, M., Valdeolmillos-
 13 Rodríguez, A., Ruiz-Zapata, B., 2006. A recent analogue for
 14 palustrine carbonate environments: The Quaternary deposits
 15 of Las Tablas de Daimiel wetlands, Ciudad Real, Spain.
 16 Geological Society of America, 416 (Special Paper), 153-168.
 17 DOI: [https://doi.org/10.1130/2006.2416\(10\)](https://doi.org/10.1130/2006.2416(10))

18 Angulo, A., Muñoz, A., 2013. Análisis de la periodicidad climática
 19 de baja frecuencia registrada en los sedimentos lacustres del
 20 Grupo Enciso (Cretácico inferior de la Cuenca de Cameros,
 21 La Rioja). Aplicaciones a la correlación y datación de la serie.
 22 Boletín Geológico y Minero de España, 124(2), 203-219.

23 Anselmetti, F.S., Ariztegui, D., Hodell, D.A., Hillesheim, M.B.,
 24 Brenner, M., Gilli, A., McKenzie, J.A., Mueller, A.D., 2006.
 25 Late Quaternary climate-induced lake level variations in Lake
 26 Petén Itzá, Guatemala, inferred from seismic stratigraphic
 27 analysis. *Palaeogeography, Palaeoclimatology, Palaeoecology*,
 28 230(1-2), 52-69. DOI: <https://doi.org/10.1016/j.palaeo.2005.06.037>

29 Argylian, E.P., Forman, S.T., 2003. Lake level response to seasonal
 30 climatic variability in the Lake Michigan-Huron System from
 31 1920 to 1995. *Journal of Great Lakes Research*, 29(3), 488-
 32 500. DOI: [https://doi.org/10.1016/S0380-1330\(03\)70453-5](https://doi.org/10.1016/S0380-1330(03)70453-5)

33 Armenteros, I., Huerta, P., 2006. The role of clastic sediment influx
 34 in the formation of calcrete and palustrine facies: a response
 35 to paleographic and climatic conditions in the southeastern
 36 Tertiary Duero basin (northern Spain). In: Alonso-Zarza,
 37 A.M., Tanner, L.H. (eds.). *Paleoenvironmental Record*
 38 and Applications of Calcretes and Palustrine Carbonates.
 39 Geological Society of America, 416 (Special Paper), 119-132.
 40 DOI: [https://doi.org/10.1130/2006.2416\(08\)](https://doi.org/10.1130/2006.2416(08))

41 Aurell, M., Bádenas, B., Gasca, J.M., Canudo, J.I., Liesa, C., Soria,
 42 A.R., Moreno-Azanza, M., Najes, L., 2016. Stratigraphy
 43 and evolution of the Galve sub-basin (Spain) in the middle
 44 Tithonian-early Barremian: implications for the setting and
 45 age of some dinosaur fossil sites. *Cretaceous Research*, 65,
 46 138-162. DOI: <https://doi.org/10.1016/j.cretres.2016.04.020>

47 Azzennoud, K., Baali, A., Mesquita-Joanes, F., El Asmi, H., Ait-
 48 Brahim, Y., 2023. Disentangling orbital, sub-orbital, and
 49 tectonic signatures from lacustrine sediments developed upon
 50 a half-graben (Lake Ifrah Basin, Northwest Africa): Insights
 51 into lowest-rank T-R sequences in low accommodation
 52 basins. *Sedimentary Geology*, 449, 106376. DOI: <https://doi.org/10.1016/j.sedgeo.2023.106376>

53 Aziz, H.A., Hilgen, F., Krijgsman, W., Sanz, E., Calvo, J.P., 2000.
 54 Astronomical forcing of sedimentary cycles in the middle to
 55 late Miocene continental Calatayud Basin (NE Spain). *Earth
 1 and Planetary Science Letters*, 177(1-2), 9-22. DOI: [https://doi.org/10.1016/S0012-821X\(00\)00035-2](https://doi.org/10.1016/S0012-821X(00)00035-2)

2 Barron, E.J., Washington, W.M., 1984. The role of geographic
 3 variables in explaining paleoclimates: Results from
 4 Cretaceous climate model sensitivity studies. *Journal of
 5 Geophysical Research: Atmospheres*, 89(D1), 1267-1279.
 6 DOI: <https://doi.org/10.1029/JD089iD01p01267>

7 Berger, A., Loutre, M.F., McIntyre, A., 1997. Intertropical latitudes
 8 and precessional and half-precessional cycles/Response.
 9 *Science*, 278(5342), 1476-1478. Last accessed: 12 September
 10 2024. Website: <https://www.jstor.org/stable/2894768>

11 Blakey, R., 2008. Gondwana palaeogeography from assembly to
 12 breakup – a 500m.y. odyssey. In: Fielding, C.R., Frank, T.D.,
 13 Isbell, J.L. (eds.). *Resolving the Late Paleozoic Ice Age in
 14 Time and Space*. Geological Society of America, 441 (Special
 15 Paper), 1-28. DOI: [https://doi.org/10.1130/2008.2441\(01\)](https://doi.org/10.1130/2008.2441(01))

16 Bohacs, K.M., Carroll, A.R., Neal, J.E., Mankiewicz, P.J., 2000.
 17 Lake-basin type, source potential, and hydrocarbon character:
 18 an integrated sequence-stratigraphic geochemical framework.
 19 In: Gierlowski-Kordesch, E.H., Kelts, K.R. (eds.). *Lake Basins
 20 through Space and Time*. American Association of Petroleum
 21 Geologists Studies in Geology, 46, 3-33. DOI: <https://doi.org/10.1306/St46706C1>

22 Bosence, D.W.J., Procter, E., Aurell, M., Bel Kahla, A., Boudagher-
 23 Fadel, M., Casaglia, F., Cirilli, S., Mehdié, M., Nieto, L., Rey,
 24 J., Scherreiks, R., Soussi, M., Waltham, D., 2009. A tectonic
 25 signal in high-frequency, peritidal carbonate cycles? A
 26 regional analysis of Liassic platforms from western Tethys.
 27 *Journal of Sedimentary Research*, 79, 389-415. DOI: <https://doi.org/10.2110/jsr.2009.038>

28 Boulila, S., Galbrun, B., Hinnov, L.A., Collin, P.Y., Ogg, J.G.,
 29 Fortwengler, D., Marchand, D., 2010. Milankovitch and sub-
 30 Milankovitch forcing of the Oxfordian (Late Jurassic) Terres
 31 Noires Formation (SE France) and global implications. *Basin
 32 Research*, 22, 717-732. DOI: [https://doi.org/10.1111/j.1365-2117.2009.00429.x](https://doi.org/10.1111/j.1365-

 33 2117.2009.00429.x)

34 Boulila, S., Galbrun, B., Miller, K.G., Pekar, S.F., Browning, J.V.,
 35 Laskar, J., Wright, J.D., 2011. On the origin of Cenozoic and
 36 Mesozoic “third-order” eustatic sequences. *Earth-Science
 37 Reviews*, 109, 94-112. DOI: [https://doi.org/10.1016/j.earscirev.2011.09.003](https://doi.org/10.1016/j.

 38 earscirev.2011.09.003)

39 Burgener, L., Hyland, E., Reich, B.J., Scotes, C., 2023. Cretaceous
 40 climates: Mapping palaeo-Köppen climatic zones using a
 41 Bayesian statistical analysis of lithologic, paleontologic, and
 42 geochemical proxies. *Palaeogeography, Palaeoclimatology,
 43 Palaeoecology*, 613, 11373. DOI: [https://doi.org/10.1016/j.palaeo.2022.111373](https://doi.org/10.1016/j.

 44 palaeo.2022.111373)

45 Buscalioni, A.D., Fregenal Martínez, M.A., 2010. A holistic
 46 approach to the palaeoecology of Las Hoyas Konservat-
 47 Lagerstätte (La Huérguina Formation, Lower Cretaceous,
 48 Iberian Ranges, Spain). *Journal of Iberian Geology*, 36(2),
 49 297-326. DOI: [10.5209/rev_JIGE.2010.v36.n2.13](https://doi.org/10.5209/rev_JIGE.2010.v36.n2.13)

50

1 Camuera, J., Jiménez-Moreno, G., Ramos-Román, M.J., García-
 2 Alix, A., Toney, J.L., Anderson, R.S., Jiménez-Espejo, F.,
 3 Kaufman, D., Bright, J., Webster, C., Yanes, Y., Carrión, J.S.,
 4 Ohkouchi, N., Suga, H., Yamame, M., Yokoyama, Y., Martínez-
 5 Ruiz, E., 2018. Orbital-scale environmental and climatic
 6 changes recorded in a new \square 200,000-year-long multiproxy
 7 sedimentary record from Padul, southern Iberian Peninsula.
 8 *Quaternary Science Reviews*, 198, 91-114. DOI: <https://doi.org/10.1016/j.quascirev.2018.08.014>

9 Capote, R., Muñoz, J.A., Simón, J.L., Liesa, C.L., Arlegui, L.E.,
 10 2002. Alpine Tectonics I: The Alpine System North of the
 11 Betic Cordillera. In: Gibbons, W., Moreno, T. (eds.). *The*
 12 *Geology of Spain*. London, The Geological Society, 367-400.

13 Carroll, A.R., Bohacs, K.M., 1999. Stratigraphic classification
 14 of ancient lakes: Balancing tectonic and climatic controls.
 15 *Geology*, 27(2), 99-102. DOI: [https://doi.org/10.1130/0091-7613\(1999\)027<0099:SCOALB>2.3.CO;2](https://doi.org/10.1130/0091-7613(1999)027<0099:SCOALB>2.3.CO;2)

16 Casas, A.M., Muñoz, A., Tella, A., Liesa, C.L., 2024. Magnetic
 17 surveying as a proxy for defining cyclicity in thick sedimentary
 18 fillings: Application to the Cretaceous Cameros Basin (N
 19 Spain). *Cretaceous Research*, 154, 105736. DOI: <https://doi.org/10.1016/j.cretres.2023.105736>

20 Castanera, D., Bádenas, B., Aurell, M., Canudo, J.I., Gasca, J.M.,
 21 2022. New ornithopod tracks from the Lower Cretaceous
 22 El Castellar Formation (Spain): Implications for track
 23 preservation and evolution of ornithopod footprints.
 24 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 591,
 25 110866. DOI: <https://doi.org/10.1016/j.palaeo.2022.110866>

26 Catuneanu, O., 2022. *Principles of sequence stratigraphy*. Elsevier
 27 Science, 2nd Edition, 486pp. DOI: <https://doi.org/10.1016/C2009-0-19363-5>

28 Changsong, L., Eriksson, K., Sitian, L., Yongxian, W., Jianye,
 29 R., Yanmei, Z., 2001. Sequence architecture, depositional
 30 systems, and controls on development of lacustrine basin
 31 fills in part of the Erlian Basin, northeast China. *American
 32 Association of Petroleum Geologists Bulletin*, 85, 2017-
 33 2043. DOI: <https://doi.org/10.1306/8626D0DB-173B-11D7-8645000102C1865D>

34 Christie-Blick, N., 1991. Onlap, offlap, and the origin of
 35 unconformity-bounded depositional sequences. *Marine
 36 Geology*, 97, 35-56. DOI: [https://doi.org/10.1016/0025-3227\(91\)90018-Y](https://doi.org/10.1016/0025-3227(91)90018-Y)

37 Coianiz, L., Bialik, O.M., Ben-Avraham, Z., Lazar, M., 2019. Late
 38 Quaternary lacustrine deposits of the Dead Sea basin: high
 39 resolution sequence stratigraphy from downhole logging data.
 40 *Quaternary Science Reviews*, 210, 175-189. DOI: <https://doi.org/10.1016/j.quascirev.2019.03.009>

41 de Wet, C.B., Godfrey, L., de Wet, A.P., 2015. Sedimentology and
 42 stable isotopes from a lacustrine-to-palustrine limestone
 43 deposited in an arid setting, climatic and tectonic factors:
 44 Miocene–Pliocene Opache Formation, Atacama Desert, Chile.
 45 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 426, 46-
 46 67. DOI: <https://doi.org/10.1016/j.palaeo.2015.02.039>

47 Deino, A.L., Kingston, J.D., Glen, J.M., Edgar, R.K., Hill, A.,
 48 2006. Precessional forcing of lacustrine sedimentation in
 49 the late Cenozoic Chemeron Basin, Central Kenya Rift, and
 50 calibration of the Gauss/Matuyama boundary. *Earth and
 51 Planetary Science Letters*, 247(1-2), 41-60. DOI: <https://doi.org/10.1016/j.epsl.2006.04.009>

52 Dercourt, J., Zonenshain, L.P., Ricou, L.E., Kazmin, V.G., Le
 53 Pichon, X., Knipper, A.L., Grandjacquet, C., Sbortshikov,
 54 I.M., Geyssant, J., Leprevier, C., Pechersky, D.H., Boulin,
 55 J., Sibuet, J.C., Savostin, L.A., Sorokhtin, O., Westphal, M.,
 56 Bazhenov, M.L., Lauer, J.P., Biju-Duval, B., 1986. Geological
 57 evolution of the Tethys belt from the Atlantic to the Pamir
 58 since the Lias. *Tectonophysics*, 123, 241-315. DOI: [https://doi.org/10.1016/0040-1951\(86\)90199-X](https://doi.org/10.1016/0040-1951(86)90199-X)

59 Emery, D., Myers, K. (eds.), 2009. *Sequence stratigraphy*. New
 60 Jersey, John Wiley & Sons, 309pp.

61 Ezquerro, L., 2017. El sector norte de la cuenca neógena de Teruel:
 62 tectónica, clima y sedimentación. PhD Thesis. Zaragoza,
 63 Universidad de Zaragoza, 494pp. Last accessed: 21 October
 64 2024. Website: <http://zaguan.unizar.es/record/77098#>

65 Ezquerro, L., Moretti, M., Liesa, C.L., Luzón, A., Simón, J.L.,
 66 2015. Seismites from a well core of palustrine deposits
 67 as a tool for reconstructing the palaeoseismic history of
 68 a fault. *Tectonophysics*, 655, 191-205. DOI: <http://dx.doi.org/10.1016/j.tecto.2015.05.025>

69 Ezquerro, L., Moretti, M., Liesa, C.L., Luzón, A., Pueyo, E.L.,
 70 Simón, J.L., 2016. Controls on space–time distribution of soft-
 71 sediment deformation structures: Applying palaeomagnetic
 72 dating to approach the apparent recurrence period of
 73 paleoseisms at the Conud Fault (eastern Spain). *Sedimentary
 74 Geology*, 344, 91-111. DOI: <http://dx.doi.org/10.1016/j.sedgeo.2016.06.007>

75 Ezquerro, L., Simón, J.L., Luzón, A., Liesa, C.L., 2020. Segmentation
 76 and increasing activity in the Neogene-
 77 Quaternary Teruel Basin rift (Spain) revealed by
 78 morphotectonic approach. *Journal of Structural Geology*, 135,
 79 104043. DOI: <https://doi.org/10.1016/j.jsg.2020.104043>

80 Ezquerro, L., Muñoz, A., Liesa, C.L., Simón, J.L., Luzón, A.,
 81 2022a. Late Neogene to early Quaternary climate evolution
 82 in southwestern Europe from a continental perspective.
 83 *Global and Planetary Change*, 211, 103788. DOI: <https://doi.org/10.1016/j.gloplacha.2022.103788>

84 Ezquerro, L., Luzón, A., Simón, J.L., Liesa, C.L., 2022b. A review of the European
 85 Neogene Mammal zones from integration of litho-, bio-
 86 and magnetostratigraphy in the Teruel Basin. *Earth-Science
 87 Reviews*, 234, 104223. DOI: <https://doi.org/10.1016/j.earscirev.2022.104223>

88 Földmi, K.B., 2012. Early Cretaceous life, climate and anoxia.
 89 *Cretaceous Research*, 35, 230-257. DOI: <https://doi.org/10.1016/j.cretres.2011.12.005>

90 Fregenal-Martínez, M.A., Meléndez, N., 2000. The lacustrine
 91 fossiliferous deposits of the Las Hoyas Subbasin (Lower
 92 Cretaceous, Serranía de Cuenca, Iberian Ranges, Spain). In:
 93 Gierlowski-Kordesch, E.H., Kelts, K.R. (eds.). *Lake Basins
 94 through Space and Time*. American Association of Petroleum
 95 Geologists Studies in Geology, 46, 303-313. DOI: <https://doi.org/10.1306/St46706C25>

1 Friedmann, S.J., Burbank, D.W., 1995. Rift Basins and
2 supradetachment basins: intracontinental extensional end-
3 members. *Basin Research*, 7(2), 109-127. DOI: <https://doi.org/10.1111/j.1365-2117.1995.tb00099.x>

4 García-Penas, A., Aurell, M., 2017. Tectono-sedimentary evolution
5 around the Jurassic-Cretaceous transition in Galve (Aguilar
6 del Alfambra Formation, Teruel, Iberian Chain). *Revista de la
7 Sociedad Geológica de España*, 30(2), 79-90.

8 Gasca, J.M., Moreno-Azanza, M., Canudo, J.I., 2009. The dinosaur
9 assemblage of the El Castellar Formation (upper Hauterivian-
10 lowemost Barremian, Teruel, Spain). In: Godefroit, P.,
11 Lambert, O. (eds.). *Tribute to Charles Darwin to Bernissart*
12 *Iguanodonts: New perspectives on Vertebrate Evolution and*
13 *Early Cretaceous Ecosystems*. Brussels (Belgium),
14 Programme, Abstracts and Field Trips Guidebook, 43pp.

15 Gierlowski-Kordesch, E.H., Gómez-Fernández, J.C., Meléndez,
16 N., 1991. Carbonate and coal deposition in an alluvial-
17 lacustrine setting: Lower Cretaceous (Weald) in the Iberian
18 Range (east-central Spain). In: Anadón, P., Cabrera, Ll.,
19 Kelts, K.R. (eds.). *Lacustrine Facies Analysis*. International
20 Association of Sedimentologists, 13 (Special Publication),
21 109-125. DOI: <https://doi.org/10.1002/9781444303919.ch6>

22 Gomes, J.P.B., Bunevich, R.B., Tonietto, S.N., Alves, D.B., Santos,
23 J.F., Whitaker, F.F., 2020. Climatic signals in lacustrine deposits
24 of the Upper Yacoraite Formation, Western Argentina: evidence
25 from clay minerals, analcime, dolomite and fibrous calcite. *Sedimentology*, 67(5), 2282-2309. DOI: <https://doi.org/10.1111/sed.12700>

26 Gómez-Fernández, J.C., Meléndez, N., 1994. Climatic control on
27 Lower Cretaceous sedimentation in a playa-lake system of a
28 tectonically active basin, Huérteles alloformation, Cameros
29 Basin, (north-central Spain). *Journal of Paleolimnology*, 11,
30 91-107. DOI: <https://doi.org/10.1007/BF00683272>

31 Gong, Z., Langereis, C.G., Mullender, T.A.T., 2008. The rotation of
32 Iberia during the Aptian and the opening of the Bay of Biscay. *Earth and Planetary Science Letters*, 273, 80-93. DOI: <https://doi.org/10.1016/j.epsl.2008.06.016>

33 Guiter, F., Andrieu-Ponel, V., de Beaulieu, J.L., Cheddadi, R.,
34 Calvez, M., Ponel, P., Reille, M., Keller, T., Goeury, C., 2003. The last climatic cycles in Western Europe: a comparison
35 between long continuous lacustrine sequences from France and other terrestrial records. *Quaternary International*, 111(1),
36 59-74. DOI: [https://doi.org/10.1016/S1040-6182\(03\)00015-6](https://doi.org/10.1016/S1040-6182(03)00015-6)

37 Hammer, Ø., Harper, D.A.T., Ryan, P.D., 2001. PAST: Paleontological statistics software package for education and data analysis. *Paleontologia Electronica*, 4(1), 1-9. Last accessed: 21 October 2024. Website: <http://palaeo-electronica.org>

38 Hasegawa, H., Katsuta, N., Muraki, Y., Heimhofer, U., Ichinnorov,
39 N., Asahi, H., Ando, H., Yamamoto, K., Murayama, M., Ohta, T.,
40 Yamamoto, M., Ikeda, M., Ishikawa, K., Kuma, R., Hasegawa,
41 T., Hasebe, N., Nishimoto, S., Yamaguchi, K., Abe, F., Tada, R.,
42 Nakagawa, T., 2022. Decadal-centennial-scale solar-linked
43 climate variations and millennial-scale internal oscillations
44 during the Early Cretaceous. *Scientific Reports*, 12, 21894.

45 DOI: <https://doi.org/10.1038/s41598-022-25815-w>

46 Hay, W.W., Floegel, S., 2012. New thoughts about the Cretaceous
47 climate and oceans. *Earth-Science Reviews*, 115(4), 262-272.
48 DOI: <https://doi.org/10.1016/j.earscirev.2012.09.008>

49 Haywood, A.M., Valdes, P.J., Markwick, P.J., 2004. Cretaceous
50 (Wealden) climates: a modelling perspective. *Cretaceous Research*, 25(3), 303-311. DOI: <https://doi.org/10.1016/j.cretres.2004.01.005>

51 Hoorn, C., Kukla, T., Bogotá-Ángel, G., van Soelen, E., González-
52 Arango, C., Wesselingh, F.P., Vonhof, H., Val, P., Morcote-
53 Rios, G., Roddaz, M., Luiz Dantas, E., Ventura Santos, R.,
54 Sinninghe Damsté, J.S., Kim, J.-H., Morley, R.J., 2022. Cyclic
55 sediment deposition by orbital forcing in the Miocene wetland
of western Amazonia? New insights from a multidisciplinary
approach. *Global and Planetary Change*, 210, 103717. DOI:
<https://doi.org/10.1016/j.gloplacha.2021.103717>

56 Huang, C., Ogg, J.G., Kemp, D.B., 2020. Cyclostratigraphy and
57 astrochronology: Case studies from China. *Palaeogeography,
58 Palaeoclimatology, Palaeoecology*, 560, 110017. DOI: <https://doi.org/10.1016/j.palaeo.2020.110017>

59 Ielpi, A., 2012. Orbitally-driven climate forcing in late Pliocene
60 lacustrine siderite-rich clastic rhythms (Upper Valdarno
61 Basin, Northern Apennines, Italy). *Palaeogeography,
62 Palaeoclimatology, Palaeoecology*, 331, 119-135. DOI: <https://doi.org/10.1016/j.palaeo.2012.03.004>

63 Illueca, N., Liesa, C.L., Soria, A.R., 2023. Ciclicidad climática en
64 sedimentos lacustres de la Formación El Castellar (Cretácico
65 inferior, Cordillera Ibérica). *Geogaceta*, 74, 3-6. DOI: <https://doi.org/10.55407/geogaceta98398>

66 Jersey, M.T., 1988. Quantitative geological modelling of
67 siliciclastic rock sequences and their seismic expression. In:
68 Wilgus, C.K., Hastings, B.S., Kendall, C.G.St.C., Posamentier,
69 H.W., Ross, C.A., Van Wagoner, J.C. (eds.). *Sea Level Changes – An Integrated Approach*. Society of Economic Paleontologists
70 and Mineralogists (SEPM), 42 (Special Publication), 47-69.
71 DOI: <https://doi.org/10.2110/pec.88.01.0047>

72 Jin, Z., Yu, J., Zhang, F., Qiang, X., 2020. Glacial-interglacial
73 variation in catchment weathering and erosion paces the
74 Indian summer monsoon during the Pleistocene. *Quaternary
75 Science Reviews*, 248, 106619. DOI: <https://doi.org/10.1016/j.quascirev.2020.106619>

76 Juhász, E., Kovács, L.Ó., Müller, P., Tóth-Makk, A., Phillips, L.,
77 Lantos, M., 1997. Climatically driven sedimentary cycles
78 in the Late Miocene sediments of the Pannonian Basin,
79 Hungary. *Tectonophysics*, 282(1-4), 257-276. DOI: [https://doi.org/10.1016/S0040-1951\(97\)00222-9](https://doi.org/10.1016/S0040-1951(97)00222-9)

80 Keighley, D., Flint, S., Howell, J., Moscariello, A., 2003. Sequence
81 stratigraphy in lacustrine basins: A model for part of the
82 Green River Formation (Eocene), Southwest Uinta Basin,
83 Utah, U.S.A. *Journal of Sedimentary Research*, 73(6), 987-
84 1006. DOI: <https://doi.org/10.1306/050103730987>

85 King, M.T., Welford, J.K., Cadenas, P., Tugend, J., 2021. Investigating
86 the plate kinematics of the Bay of Biscay using deformable
87 plate tectonic models. *Tectonics*, 40, e2020TC006467. DOI:
<https://doi.org/10.1029/2020TC006467>

1 Kovács, Á., Balázs, A., Špelić, M., Sztanó, O., 2021. Forced or
 2 normal regression signals in a lacustrine basin? Insights from
 3 3D stratigraphic forward modeling in the SW Pannonian
 4 Basin. *Global and Planetary Change*, 196, 103376. DOI:
 5 <https://doi.org/10.1016/j.gloplacha.2020.103376>

6 Kuzucuoğlu, C., Karabiyikoğlu, M., Fontugne, M., Pastre,
 7 J.F., Ercan, T., 1997. Environmental changes in Holocene
 8 lacustrine sequences from Karapınar in the Konya plain
 9 (Turkey). In: Dalfes, H.N., Kukla, G., Weiss, H. (eds.). *Third*
 10 *Millennium BC Climate Change and Old World Collapse*.
 11 Berlin, Heidelberg, Springer, NATO ASI Series, 49, 451-463.
 12 DOI: https://doi.org/10.1007/978-3-642-60616-8_18

13 Laita, E., Bauluz, B., Aurell, M., Bádenas, B., Canudo, J.I., Yuste,
 14 A., 2020. A change from warm/humid to cold/dry climate
 15 conditions recorded in lower Barremian clay-dominated
 16 continental successions from the SE Iberian Chain (NE
 17 Spain). *Sedimentary Geology*, 403, 105673. DOI: <https://doi.org/10.1016/j.sedgeo.2020.105673>

18 Laita, E., Bauluz, B., Aurell, M., Bádenas, B., Yuste, A., 2022.
 19 Weathering events recorded in uppermost Hauterivian–lower
 20 Barremian clay-dominated continental successions from the
 21 NW Iberian Range: climatic vs. tectonic controls. *Journal of*
 22 *Iberian Geology*, 48, 45-63. DOI: <https://doi.org/10.1007/s41513-021-00181-0>

23 Leeder, M.R., 1991. Denudation, vertical crustal movements and
 24 sedimentary basin fill. *Geologische Rundschau*, 80, 441-458.
 25 DOI: 10.1007/bf01829376

26 Lepre, C.J., Quinn, R.L., 2022. Aridification and orbital forcing
 27 of eastern African climate during the Plio-Pleistocene.
 28 *Global and Planetary Change*, 208, 103684. DOI: <https://doi.org/10.1016/j.gloplacha.2021.103684>

29 Li, M., Kump, L.R., Hinnov, L.A., Mann, M.E., 2018. Tracking
 30 variable sedimentation rates and astronomical forcing in
 31 Phanerozoic paleoclimate proxy series with evolutionary
 32 correlation coefficients and hypothesis testing. *Earth and*
 33 *Planetary Science Letters*, 501, 165-179. DOI: <https://doi.org/10.1016/j.epsl.2018.08.041>

34 Liesa, C.L., Casas, A.M., Soria, A.R., Simón, J.L., Meléndez, A.,
 35 2004. Estructura extensional cretácea e inversión terciaria
 36 en la región Aliaga-Montalbán. In: Colombo, F., Liesa, C.L.,
 37 Meléndez, G., Pocoví, A., Sancho C., Soria, A.R. (eds.). *Itinerarios*
 38 *Geológicos por Aragón*. Zaragoza, Sociedad
 39 Geológica de España, 151-180.

40 Liesa, C.L., Soria, A.R., Meléndez, N., Meléndez, A., 2006.
 41 Extensional fault control on the sedimentation patterns in a
 42 continental rift basin: El Castellar Formation, Galve subbasin,
 43 Spain. *Journal of the Geological Society*, 163(3), 487-498.
 44 DOI: <https://doi.org/10.1144/0016-764904-169>

45 Liesa, C.L., Casas, A.M., Simón, J.L., 2018. La tectónica de
 46 inversión en una región intraplaca: la Cordillera Ibérica.
 47 *Revista de la Sociedad Geológica de España*, 31(2), 23-50.

48 Liesa, C.L., Soria, A.R., Casas, A., Aurell, M., Meléndez, N.,
 49 Bádenas, B., Fregenal-Martínez, M., Navarrete, R., Peropadre,
 50 C., Rodríguez-López, J.P., 2019. The South-Iberian, Central
 51 Iberian and Maestrazgo Basins. In: Oliveira, J.T., Quesada,
 52 C. (eds.). *The Geology of Iberia: a Geodynamic Approach*,
 53 Vol. 3 (The Alpine Cycle), (Chapter 5) (Late Jurassic-Early
 54 Cretaceous rifting). Springer Nature, *Regional Geology*
 55 *Reviews*, 214-228. DOI: https://doi.org/10.1007/978-3-030-11295-0_5

56 Liesa, C.L., Casas, A.M., Aurell, M., Simón, J.L., Soria, A.R.,
 57 2023. Salt tectonics vs. inversion tectonics: The anticlines
 58 of the western Maestrazgo revisited (eastern Iberian Chain,
 59 Spain). *Basin Research*, 35, 295-335. DOI: <https://doi.org/10.1111/bre.12713>

60 Luzón, A., González, A., Muñoz, A., Sánchez-Valverde, B.,
 61 2002. Upper Oligocene-Lower Miocene shallowing-
 62 upward lacustrine sequences controlled by periodic and
 63 non-periodic processes (Ebro Basin, northeastern Spain).
 64 *Journal of Paleolimnology*, 28, 441-456. DOI: <https://doi.org/10.1023/A:1021675227754>

65 Martín-Chivelet, J., López-Gómez, J., Aguado, R., Arias, C.,
 66 Arribas, J., Arribas, M.E., Aurell, M., Bádenas, B., Benito,
 67 M.I., Bover-Arnal, T., Casas-Sainz, A., Castro, J.M., Coruña,
 68 F., de Gea, G.A., Fornós, J.J., Fregenal-Martínez, M., García-
 69 Senz, J., Garofano, D., Gelabert, B., Giménez, J., González-
 70 Acebrón, J., Guimerà, J., Liesa, C.L., Mas, R., Meléndez, N.,
 71 Molina, J.M., Muñoz, J.A., Navarrete, R., Nebot, M., Nieto,
 72 L.M., Omodeo-Salé, S., Pedrera, A., Peropadre, C., Quijada,
 73 I.E., Quijano, M.L., Reolid, M., Robador, A., Rodríguez-
 74 López, J.P., Rodríguez-Perea, A., Rosales, I., Ruiz-Ortiz, P.A.,
 75 Sàbat, F. Salas, R., Soria, A.R., Suárez-González, P., Vilas,
 76 L., 2019. The Late Jurassic-Early Cretaceous Rifting. In:
 77 Quesada, C., Oliveira, J.T. (eds.). *The Geology of Iberia: A*
 78 *Geodynamic Approach, The Alpine Cycle*. Springer Nature,
 79 *Regional Geology Reviews*, vol. 5, 169-249. DOI: https://doi.org/10.1007/978-3-030-11295-0_5

80 Meléndez, M.N., Liesa, C.L., Soria, A.R., Meléndez, A.,
 81 2009. Lacustrine system evolution during early rifting:
 82 El Castellar Formation (Galve sub-basin, Central Iberian
 83 Chain). *Sedimentary Geology*, 222, 64-77. DOI: <https://doi.org/10.1016/j.sedgeo.2009.05.019>

84 Meléndez, M.N., Soria, A.R., Meléndez, A., Aurell, M., Liesa,
 85 C.L., 2000. Early Cretaceous lacustrine systems of the
 86 Aguilón Subbasin (Central Iberian range, NE Spain): tectonic
 87 evolution. In: Gierlowski-Kordesch, E.H., Kelts, K.R. (eds.).
 88 *Lake Basins through Space and Time*. American Association
 89 of Petroleum Geologists Studies in Geology, 46, 285-294.
 90 DOI: <https://doi.org/10.1306/St46706C23>

91 Meyers, S.R., 2019. Cyclostratigraphy and the problem of
 92 astrochronologic testing. *Earth-Science Reviews*, 190, 190-
 93 223. DOI: <https://doi.org/10.1016/j.earscirev.2018.11.015>

94 Milankovitch, M.M., 1941. Canon of insolation and the iceage
 95 problem. *Königlich Serbische Akademie*, 484pp.

96 Ming, G., Zhou, W., Wang, H., Cheng, P., Shu, P., Xian, F., Fu, Y.,
 97 2020. Moisture variations in Lacustrine-eolian sequence
 98 from the Hunshandake sandy land associated with the East
 99 Asian Summer Monsoon changes since the late Pleistocene.
 100 *Quaternary Science Reviews*, 233, 106210. DOI: <https://doi.org/10.1016/j.quascirev.2020.106210>

1 Mitchum, R.M., Van Wagoner, J.C., 1991. High-frequency
2 sequences and their stacking patterns: sequence-stratigraphic
3 evidence of high-frequency eustatic cycles. *Sedimentary
4 Geology*, 70, 131-160. DOI: [https://doi.org/10.1016/0037-0738\(91\)90139-5](https://doi.org/10.1016/0037-0738(91)90139-5)

5 Moreau, M.G., Berthou, J.Y., Malod, J.A., 1997. New paleomagnetic
6 Mesozoic data from the Algarve (Portugal): fast rotation of
7 Iberia between the Hauterivian and the Aptian. *Earth and
8 Planetary Science Letters*, 146, 689-701. DOI: [https://doi.org/10.1016/S0012-821X\(96\)00239-7](https://doi.org/10.1016/S0012-821X(96)00239-7)

9 Moreno, A., González-Sampériz, P., Morellón, M., Valero-Garcés,
10 B.L., Fletcher, W.J., 2012. Northern Iberian abrupt climate
11 change dynamics during the last glacial cycle: a view from
12 lacustrine sediments. *Quaternary Science Reviews*, 36, 139-
13 153. DOI: <https://doi.org/10.1016/j.quascirev.2010.06.031>

14 Moriya, K., 2011. Development of the Cretaceous greenhouse
15 climate and the oceanic thermal structure. *Paleontological
16 Research*, 15(2), 77-88. DOI: <https://doi.org/10.2517/1342-8144-15.2.077>

17 Morrill, C., Small, E.E., Sloan, L.C., 2001. Modeling orbital
18 forcing of lake level change: Lake Gosiute (Eocene), North
19 America. *Global and Planetary Change*, 29, 57-76. DOI:
20 [https://doi.org/10.1016/S0921-8181\(00\)00084-9](https://doi.org/10.1016/S0921-8181(00)00084-9)

21 Muñoz, A., Angulo, A., Liesa, C.L., Luzón, M.A., Mayayo, M.J.,
22 Pérez, A., Soria, A.R., Val, V., Yuste, A., 2020. Climatic
23 periodicities and astrochronological dating of the Enciso
24 Group in the eastern Cameros Basin (N of Spain). *Boletín
25 Geológico y Minero*, 131(2), 243-268. DOI: <https://doi.org/10.21701/bolgeomin.131.2.003>

26 Murphy, D.H., Wilkinson, B.H., 1980. Carbonate deposition
27 and facies distribution in a central Michigan marl
28 lake. *Sedimentology*, 27, 123-135. DOI: <https://doi.org/10.1111/j.1365-3091.1980.tb01164.x>

29 Navarrete, R., 2015. Controles alocíclicos de la sedimentación
30 barremiense en la Subcuenca de Galve (Fm. Camarillas,
31 margen occidental de la Cuenca del Maestrazgo). PhD Thesis.
32 Zaragoza, Universidad de Zaragoza, 448pp.

33 Nutz, A., Schuster, M., Boës, X., Rubino, J.L., 2017. Orbitally-
34 driven evolution of Lake Turkana (Turkana Depression, Kenya,
35 EARS) between 1.95 and 1.72 Ma: A sequence stratigraphy
36 perspective. *Journal of African Earth Sciences*, 125, 230-243.
37 DOI: <https://doi.org/10.1016/j.jafrearsci.2016.10.016>

38 O'Connor, L., Robinson, S.A., Naafs, B.D.A., Jenkyns,
39 H.C., Henson, S., Clarke, M., Pancost, R.D., 2019. Late
40 Cretaceous temperature evolution of the southern high
41 latitudes: A TEX86 perspective. *Paleoceanography* and
42 *Paleoclimatology*, 34, 436-454. DOI: <https://doi.org/10.1029/2018PA003546>

43 Oviatt, Ch., McCoy, W.D., Nash, W.P., 1994. Sequence stratigraphy
44 of lacustrine deposits: A Quaternary example from the
45 Bonneville basin, Utah. *Geological Society of America
46 Bulletin*, 106(1), 133-144. DOI: [https://doi.org/10.1130/0016-7606\(1994\)106<0133:SSOLDA>2.3.CO;2](https://doi.org/10.1130/0016-7606(1994)106<0133:SSOLDA>2.3.CO;2)

47 Park, J., d'Hont, S.L., King, J.W., Gibson, C., 1993. Late
48 Cretaceous precessional cycles in double time: A warm-Earth
49 Milankovitch response. *Science*, 261(5127), 1431-1434. DOI:
50 <https://doi.org/10.1126/science.261.5127.1431>

51 Platt, N.H., Wright, V.P., 1991. Lacustrine carbonates: facies
52 models, facies distribution and hydrocarbon aspects. In:
53 Anadón, P., Cabrera, L., Kelts, K. (eds.). *Lacustrine Facies
54 Analysis*. Special Publication International Association of
55 Sedimentologists, 13, 57-74.

56 Platt, N.H., Wright, V.P., 1992. Palustrine carbonates and the
57 Florida Everglades: towards and exposure index for the fresh-
58 water environment? *Journal Sedimentary Petrology*, 62, 1058-
59 1071. DOI: <https://doi.org/10.1306/D4267A4B-2B26-11D7-8648000102C1865D>

60 Posamentier, H.W., Vail, P.R., 1988. Eustatic controls on clastic
61 deposition II: Sequence and systems tract models. In: Wilgus,
62 C.K., Hastings, B.S., Posamentier, H., Van Wagoner, J., Ross,
63 C.A., Kendall, C.G.St.C. (eds.). *Sea Level Changes: An
64 Integrated Approach*. Society of Economic Paleontologists
65 and Mineralogists (SEPM), 42 (Special Publication), 125-
66 154. DOI: <https://doi.org/10.2110/pec.88.01.0125>

67 Rits, D.S., van Balen, R.T., Prins, M.A., Zheng, H., 2017. Evolution
68 of the alluvial fans of the Luo River in the Weihe Basin,
69 central China, controlled by faulting and climate change-A
70 reevaluation of the paleogeographical setting of Dali Man site.
71 *Quaternary Science Reviews*, 166, 339-351. DOI: <https://doi.org/10.1016/j.quascirev.2017.01.013>

72 Rodríguez-López, J.P., Wu, C., Vishnivetskaya, T.A., Murton,
73 J.B., Tang, W., Ma, C., 2022. Permafrost in the Cretaceous
74 supergreenhouse. *Nature Communications*, 13, 7946. DOI:
75 <https://doi.org/10.1038/s41467-022-35676-6>

76 Rodríguez-López, J.P., Liesa, C.L., Luzón, A., Muñoz, A., Mayayo,
77 M.J., Murton, J.B., Soria, A.R., 2024. Ice rafted dropstones at
78 midlatitudes in the Cretaceous of continental Iberia. *Geology*,
79 52, 33-38. DOI: <https://doi.org/10.1130/G51725.1>

80 Rosenbaum, G., Lister, G.S., Duboz, C., 2002. Relative motions
81 of Africa, Iberia and Europe during Alpine orogeny.
82 *Tectonophysics*, 359, 117-129. DOI: [https://doi.org/10.1016/S0040-1951\(02\)00442-0](https://doi.org/10.1016/S0040-1951(02)00442-0)

83 Ruddiman, W.F., 2006. Orbital changes and climate. *Quaternary
84 Science Reviews*, 25, 3092-3112. DOI: <https://doi.org/10.1016/j.quascirev.2006.09.001>

85 Ruiz Omeñaca, J.I., 2006. Restos directos de dinosaurios
86 (Saurischia, Ornithischia) en el Barremiense (Cretácico
87 Inferior) de la Cordillera Ibérica en Aragón (Teruel, España).
88 PhD Thesis. Zaragoza, Universidad de Zaragoza, 432pp.

89 Sáez, A., Valero-Garcés, B.L., Giralt, S., Moreno, A., Bao,
90 R., Pueyo, J.I., Hernández, A., Casas, D., 2009. Glacial to
91 Holocene climate changes in the SE Pacific. The Raraku
92 lake sedimentary record (Easter Island, 27 S). *Quaternary
93 Science Reviews*, 28(25-26), 2743-2759. DOI: <https://doi.org/10.1016/j.quascirev.2009.06.018>

94 Salas, R., 1987. El Malm i el Cretaci inferior entre el Massís de
95 Garraf i la Serra d'Espadà: anàlisi de conca. PhD Thesis.
96 Barcelona, Universitat de Barcelona, 345pp.

97 Salas, R., Casas, A., 1993. Mesozoic extensional tectonics,
98 stratigraphy and crustal evolution during the Alpine cycle of

1 the eastern Iberian basin. *Tectonophysics*, 228, 33-55. DOI:
2 [https://doi.org/10.1016/0040-1951\(93\)90213-4](https://doi.org/10.1016/0040-1951(93)90213-4)

3 Salas, R., Guimerà, J., Mas, R., Martín-Closas, C., Meléndez, A.,
4 Alonso, A., 2001. Evolution of the Mesozoic central Iberian
5 Rift System and its Cainozoic inversion (Iberian Chain). In: Ziegler, P.A., Cavazza, W., Robertson, A.H.F., Crasquin-
6 Soleau, S. (eds.). *Peri-Tethyan Rift/Wrench Basins and*
7 *Passive Margins. Peri-Tethys Memoir 6: Peri-Tethyan Rift/*
8 *Wrench Basins and Passive Margins*. Paris, Muséum national
9 d'Histoire naturelle, 186, 145-185.

10 Sames, B., Wagreich, M., Wendler, J.E., Haq, B.U., Conrad, C.P.,
11 Melinte-Dobrinescu, M.C., Hu, X., Wendler, I., Wolfgring, E.,
12 Yilmaz, I.Ö., Zorina, S.O., 2016. Review: Short-term seal-level
13 changes in a greenhouse world – A review from the Cretaceous.
14 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 441, 393-
15 411. DOI: <https://doi.org/10.1016/j.palaeo.2015.10.045>

16 Scheidt, S., Hambach, U., Hao, Q., Rolf, C., Wennrich, V., 2020.
17 Environmental signals of Pliocene-Pleistocene climatic
18 changes in Central Europe: Insights from the mineral magnetic
19 record of the Heidelberg Basin sedimentary infill (Germany).
20 *Global and Planetary Change*, 187, 103112. DOI: <https://doi.org/10.1016/j.gloplacha.2020.103112>

21 Schnyder, J., Dejax, J., Keppens, E., Tu, T.T.N., Spagna, P., Boulila,
22 S., Galbrun, B., Riboulleau, A., Tshibangu, J.P., Yans, J.,
23 2009. An Early Cretaceous lacustrine record: organic
24 matter and organic carbon isotopes at Bernissart (Mons
25 Basin, Belgium). *Palaeogeography, Palaeoclimatology,*
26 *Palaeoecology*, 281(1-2), 79-91. DOI: <https://doi.org/10.1016/j.palaeo.2009.07.014>

27 Schwarzbacher, W., 2005. The stratification and cyclicity of the
28 Dachstein Limestone in Lofer, Leogang and Steinernes
29 Meer (Northern Calcareous Alps, Austria). *Sedimentary*
30 *Geology*, 181, 93-106. DOI: <https://doi.org/10.1016/j.sedgeo.2005.07.001>

31 Scotese, C.R., 2016. Tutorial: PALEOMAP Paleoatlas for GPlates
32 and the PaleoData Plotter Program. Tutor, PALEOMAP
33 Project. Last accessed: 12 September 2024. Website: <http://www.earthbyte.org/paleomap-paleoatlas-for-gplates/>

34 Sevillano, A., Bádenas, B., Rosales, I., Barnolas, A., López-García,
35 J.M., 2020. Orbital cycles, differential subsidence and internal
36 factors controlling the high-frequency sequence architecture
37 in a Sinemurian shallow carbonate platform (Mallorca Island,
38 Spain). *Sedimentary Geology*, 407, 105729. DOI: <https://doi.org/10.1016/j.sedgeo.2020.105729>.

39 Simmons, G.F., 2016. *Differential Equations with Applications*
40 and *Historical Notes*. Boca Raton, CRC Press, 740pp.

41 Simón, J.L., Arlegui1, L.E., Ezquerro, L., Lafuente, P., Liesa, C.L.,
42 Luzón, A., 2016. Enhanced palaeoseismic succession at the
43 Concord Fault (Iberian Chain, Spain): new insights for seismic
44 hazard assessment. *Natural Hazards*, 80, 1967-1993. DOI:
45 <https://doi.org/10.1007/s11069-015-2054-6>

46 Soria, A.R., 1997. Estudio estratigráfico-sedimentológico y
47 tectónico del Cretácico inferior en la parte occidental de la
48 cuenca del Maestrazgo, subcuenca de Las Parras y Galve. La
49 sedimentación en las cuencas marginales del surco ibérico
50 durante el Cretácico Inferior y su control estructural. PhD
51 Thesis. Zaragoza, Universidad de Zaragoza, 363pp.

52 Soria, A.R., Liesa, C.L., Meléndez, A., Meléndez, M.N., 2001.
53 Sedimentación sintectónica de la Fm. El Castellar (Cretácico
54 Inferior) en la Subcuenca de Galve (Cuenca Ibérica).
55 *Geotemas*, 3, 257-260.

56 Soria, A.R., Muñoz, A., Liesa, C.L., Meléndez, A., Meléndez,
57 M.N., Soto, R., 2008. Ciclicidad climática en una unidad
58 lacustre cretácica: la Fm. Villanueva de Huerva en la
59 Subcuenca de Aguilón (Cordillera Ibérica). *Geotemas*, 10,
60 1487-1490.

61 Soria, A.R., Muñoz, A., Liesa, C.L., Luzón, A., Meléndez, A.,
62 Meléndez, M.N., 2012. Climate-driven cyclicity in an Early
63 Cretaceous synrift lacustrine series (Aguilón subbasin,
64 NE Spain). *Terra Nova*, 24, 407-416. DOI: <https://doi.org/10.1111/j.1365-3121.2012.01080.x>

65 Soria, A.R., Liesa, C.L., Navarrete, R., Rodríguez-López, J.P., 2023.
66 Sedimentology and stratigraphic architecture of Barremian
67 synrift barrier island-estuarine depositional systems from
68 blended field and drone-derived data. *Sedimentology*, 70,
69 1812-1855. DOI: <https://doi.org/10.1111/sed.13097>.

70 Stampfli, G.M., Borel, G.D., 2002. A plate tectonic model for
71 the Paleozoic and Mesozoic constrained by dynamic plate
72 boundaries and restored synthetic oceanic isochrons. *Earth*
73 and *Planetary Science Letters*, 196, 17-33. DOI: [https://doi.org/10.1016/S0012-821X\(01\)00588-X](https://doi.org/10.1016/S0012-821X(01)00588-X)

74 Steenbrink, J., Hilgen, F.J., Krijgsman, W., Wijbrans, J.R.,
75 Meulenkamp, J.E., 2006. Late Miocene to Early Pliocene
76 depositional history of the intramontane Florina-Ptolemais-
77 Servia Basin, NW Greece: Interplay between orbital
78 forcing and tectonics. *Palaeogeography, Palaeoclimatology,*
79 *Palaeoecology*, 238(1-4), 151-178. DOI: <https://doi.org/10.1016/j.palaeo.2006.03.023>

80 Tang, Y., He, W., Wang, R., Ren, H., Jin, Z., Yang, Z., Zhang, Y.,
81 2023. Cyclostratigraphy of Lower Permian alkaline lacustrine
82 deposits in the Mahu Sag, Junggar basin and its stratigraphic
83 implication. *Frontiers in Earth Science*, 11, 1232418. DOI:
84 <https://doi.org/10.3389/feart.2023.1232418>

85 Tian, F., Wang, Y., Liu, J., Tang, W., Jiang, N., 2017. Late Holocene
86 climate change inferred from a lacustrine sedimentary
87 sequence in southern Inner Mongolia, China. *Quaternary*
88 *International*, 452, 22-32. DOI: <https://doi.org/10.1016/j.quaint.2017.01.029>

89 Tugend, J., Manatschal, G., Kusznir, N.J., 2015. Spatial and temporal
90 evolution of hyperextended rift systems: Implication for the
91 nature, kinematics, and timing of the Iberian-European plate
92 boundary. *Geology*, 43, 15-18. DOI: <https://doi.org/10.1130/G36072.1>

93 Valero, L., Garcés, M., Cabrera, L., Costa, E., Sáez, A., 2014. 20
94 Myr of eccentricity paced lacustrine cycles in the Cenozoic
95 Ebro Basin. *Earth and Planetary Science Letters*, 408, 183-
96 193. DOI: <https://doi.org/10.1016/j.epsl.2014.10.007>

97 Vickers, M.L., Price, G.D., Jerrett, R.M., Sutton, M., Watkinson,
98 M.P., FitzPatrick, M., 2019. The duration and magnitude of
99 Cretaceous cold events: Evidence from the northern high

1 latitudes. *Geological Society of America Bulletin*, 131, 1979-
2 1994. DOI: <https://doi.org/10.1130/B35074.1>

3 Vissers, R.L.M., Meijer, P.Th., 2012. Mesozoic rotation of Iberia:
4 Subduction in the Pyrenees? *Earth-Science Reviews*, 110, 93-
5 110. DOI: <https://doi.org/10.1016/j.earscirev.2011.11.001>

6 Walzer, U., Hendel, R., 2023. Natural climate change and
7 glaciations. *Earth-Science Reviews*, 241, 104435. DOI:
8 <https://doi.org/10.1016/j.earscirev.2023.104435>

9 Wang, Y., Deng, T., Flynn, L., Wang, X., Yin, A., Xu, Y., Parker,
10 W., Lochner, E., Zhang, Ch., Biasatti, D., 2012. Late Neogene
11 environmental changes in the central Himalaya related
12 to tectonic uplift and orbital forcing. *Journal of Asian
13 Earth Sciences*, 44, 62-76. DOI: <https://doi.org/10.1016/j.jseas.2011.05.020>

14 Wang, Y., Huang, C., Sun, B., Quan, C., Wu, J., Lin, Z., 2014.
15 Paleo-CO₂ variation trends and the Cretaceous greenhouse
16 climate. *Earth-Science Reviews*, 129, 136-147. DOI: <https://doi.org/10.1016/j.earscirev.2013.11.001>

17 Wang, T., He, S., Zhang, Q., Ding, L., Farnsworth, A., Cai, F., Wang,
18 C., Xie, J., Li, G., Sheng, J., Yue, Y., 2023. Ice Sheet Expansion
19 in the Cretaceous Greenhouse World. *Fundamental Research*,
20 1-9. DOI: <https://doi.org/10.1016/j.fmre.2023.05.005>

21 Wei, W., Lu, Y., Xing, E., Liu, Z., Pan, L., Algeo, T.J., 2017.
22 Sedimentary facies associations and sequence stratigraphy
23 of source and reservoir rocks of the lacustrine Eocene
24 Niubao Formation (Lunpola Basin, central Tibet). *Marine
25 and Petroleum Geology*, 86, 1273-1290. DOI: <https://doi.org/10.1016/j.marpetgeo.2017.07.032>

26 Wei, Z., Zhong, W., Shang, S., Ye, S., Tang, X., Xiaowen, T., Xue, J.,
27 Jibin, X., Jun, O., Smol, J.P., 2018. Lacustrine mineral magnetic
28 record of postglacial environmental changes from Dahu
29 Swamp, southern China. *Global and Planetary Change*, 170,
30 62-75. DOI: <https://doi.org/10.1016/j.gloplacha.2018.08.010>

31 Wei, X., Yan, D., Luo, P., Jiang, P., Wang, H., Zhou, J., Cong, F.,
32 Yang, X., Niu, X., Li, T., Liu, L., Liu, E., 2020. Astronomically
33 forced climate cooling across the Eocene-Oligocene transition
34 in the Pearl river Mouth basin, northern South China Sea.
35 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 558,
36 109945. DOI: <https://doi.org/10.1016/j.palaeo.2020.109945>

37 Wei, R., Zhang, R., Li, M., Wang, X., Jin, Z., 2023. Obliquity
38 forcing of lake-level changes and organic carbon burial
39 during the Late Paleozoic Ice Age. *Global and Planetary
40 Change*, 223, 104092. DOI: <https://doi.org/10.1016/j.gloplacha.2023.104092>.

41 Woolway, R.I., Kraemer, B.M., Lenters, J.D., Merchant, C.J.,
42 O'Reilly, C.M., Sharma, S., 2020. Global lake responses to
43 climate change. *Nature Reviews Earth & Environment*, 1,
44 388-403. DOI: <https://doi.org/10.1038/s43017-020-0067-5>

45 Wright, V.P., Marriott, S.B., 1993. The sequence stratigraphy of
46 fluvial depositional systems: the role of floodplain sediment
47 storage. *Sedimentary Geology*, 86, 203-210. DOI: [https://doi.org/10.1016/0037-0738\(93\)90022-W](https://doi.org/10.1016/0037-0738(93)90022-W)

48 Xu, T.-Y., Peng, J., Yu, L.-D., Han, H.-D., Yang, Y.-M., Zeng, Y.,
49 Wang, Y.B., 2023. The control of astronomical cycles on
50 lacustrine fine-grained event Sedimentation—A case study of
51 the Chunshang sub-member of the upper Es4 in the Dongying
52 Sag. *Petroleum Science*, 20(3), 1395-1410. DOI: <https://doi.org/10.1016/j.petsci.2022.11.023>

53 Yuste, A., Bauluz, B., Mayayo, M.J., 2015. Genesis and mineral
54 transformations in Lower Cretaceous karst bauxites (NE
55 Spain): climatic influence and superimposed processes.
56 *Geological Journal*, 50(6), 839-857. DOI: <https://doi.org/10.1002/gj.2604>

57 Zavala, C., Liu, H., Li, X., Arcuri, M., Di Meglio, M., Zorzano,
58 A., Otharán, G., Hao, B., Wang, Y., 2022. Lacustrine sequence
59 stratigraphy: New insights from the study of the Yanchang
60 Formation (Middle-Late Triassic), Ordos Basin, China. In:
61 Renchao Yang, A.J., Van Loon, T. (eds.). *The Ordos Basin. Sedimentological Research for Hydrocarbons Exploration*,
62 309-335. DOI: <https://doi.org/10.1016/B978-0-323-85264-7.00012-6>

63 Zhong, W., Cao, J., Xue, J., Ouyang, J., 2015. A 15,400-year record
64 of climate variation from a subalpine lacustrine sedimentary
65 sequence in the western Nanling Mountains in South China.
66 *Quaternary Research*, 84(2), 246-254. DOI: <https://doi.org/10.1016/j.yqres.2015.06.002>

67 Zhong, Q., Zhang, J.G., Tang, D., Jiang, J.H., Shen, J.J., Cai,
68 M.X., Li, P.X., 2022. Research status of lacustrine mudrock
69 deposition constrained from astronomical forcing. *Journal of Palaeogeography*, 11(3), 315-331. DOI: <https://doi.org/10.1016/j.jop.2022.05.003>

70 Ziegler, A.M., Raymond, A.L., Gierlowski, T.C., Horrell, M.A.,
71 Rowley, D.B., Lottes, A.L., 1987. Coal, climate and terrestrial
72 productivity: the present and early Cretaceous compared.
73 London, Geological Society, 32(1, Special Publications), 25-
74 49. DOI: <https://doi.org/10.1144/GSL.SP.1987.032.01.04>

Manuscript received June 2024;
revision accepted August 2024;
published Online November 2024.