

Nota briològica

Two nomenclatural changes in *Entosthodon*

Montserrat Brugués

Universitat Autònoma de Barcelona
Botànica. Facultat de Ciències
08193 Bellaterra (Barcelona). Spain
mbrugues@uab.es

Manuscript received in September 2000

Funaria convexa Spruce and *F. pulchella* Philib. are two species that Crundwell & Nyholm (1974) related to *F. muhlenbergii* Turn. According to Fife (1985, 1996), the genus *Funaria* is restricted to species with capsules inclined, asymmetric and sulcate when dry, double peristome with strongly sigmoid teeth, which are fused at apices, and a compound and revolute annulus; species with inclined and asymmetric but not sulcate capsules when dry, more or less sigmoid teeth not fused, and simple, non-revolute annulus, like *E. muhlenbergii* (Tur.) Fife, belong to *Entosthodon* subg. *Plagiodus*. Consequently, the following new combinations are necessary.

***Entosthodon convexus* (Spruce) Brugués, comb. nov.**

Basionym: *Funaria convexa* Spruce, *Musc. Pyr.* 149, 1847.

***Entosthodon pulchellus* (Philib.) Brugués, comb. nov.**

Basionym: *Funaria pulchella* Philib., *Rev. Bryol.*: 11, 41. 1884.

References

- Crundwell, A.C.; Nyholm, E. 1974. *Funaria muhlenbergii* and related European species. *Lindbergia* 2: 222-229.
- Fife, A.J. 1985. A generic revision of the Funariaceae (Bryophyta: Musci). Part I. The Journal of the Hattori Botanical Laboratory 58: 149.
- 1996. Lectotypification of *Weissia radians* Hedw., a neglected name for a widespread Australian moss species, with notes on the status of *Funaria glabra* Tayl. and the typification of some other Hedwigian names. Journal of Bryology 19: 39-48.