

SEMIGROUP ANALYSIS OF STRUCTURED POPULATIONS WITH DISTRIBUTED STATES-AT-BIRTH ARISING IN MATHEMATICAL AQUACULTURE

JÓZSEF Z. FARKAS, DARREN GREEN, AND PETER HINOW

ABSTRACT. Motivated by the modelling of structured parasite populations in aquaculture we consider a class of physiologically structured population models, where individuals may be recruited into the population at different sizes in general. That is, we consider a size-structured population model with distributed states-at-birth. The mathematical model which describes the evolution of such a population is a first order nonlinear partial integro-differential equation of hyperbolic type. First, we use positive perturbation arguments and utilise results from the spectral theory of semigroups to establish conditions for the existence of a positive equilibrium solution of our model. Then we formulate conditions that guarantee that the linearised system is governed by a positive quasicontraction semigroup on the biologically relevant state space. We also show that the governing linear semigroup is eventually compact, hence growth properties of the semigroup are determined by the spectrum of its generator. In case of a separable fertility function we deduce a characteristic equation and investigate the stability of equilibrium solutions in the general case using positive perturbation arguments.

1. Introduction

In this paper we study the following partial integro-differential equation

$$\begin{aligned} \frac{\partial}{\partial t} p(s, t) + \frac{\partial}{\partial s} (\gamma(s, P(t)) p(s, t)) \\ = -\mu(s, P(t)) p(s, t) + \int_0^m \beta(s, y, P(t)) p(y, t) dy, \end{aligned} \quad (1.1)$$

$$\gamma(0, P(t)) p(0, t) = 0, \quad (1.2)$$

$$p(s, 0) = p_0(s), \quad P(t) = \int_0^m p(s, t) ds. \quad (1.3)$$

Here the function $p = p(s, t)$ denotes the density of individuals of size (or other developmental stage) s at time t , thus, $P(t)$ is the total population at time t .

Date: June 22, 2009.

1991 Mathematics Subject Classification. 92D25, 47D06, 35B35.

Key words and phrases. Mathematical aquaculture; Quasicontraction semigroups, Positivity, Spectral methods; Stability.

$\mu \geq 0$ and $\gamma \geq 0$ denote the mortality and growth rates of individuals, respectively, and both of these vital rates depend on both size s and on the total population size $P(t)$. It is assumed that individuals may have different sizes at birth and therefore $\beta(s, y, \cdot)$ denotes the rate at which individuals of size y “produce” individuals of size s . That is the non-local integral term in (1.1) represents the recruitment of individuals into the population. We make the following general assumptions on the model ingredients

$$\begin{aligned} \mu &\in C^1([0, m] \times [0, \infty)), & \beta &\in C^1([0, m] \times [0, m] \times [0, \infty)) \\ \beta, \mu &\geq 0, & \gamma &\in C^2([0, m] \times [0, \infty)), & \gamma &> 0. \end{aligned} \tag{1.4}$$

Our motivation to investigate model (1.1)-(1.3) is the modelling of structured populations in aquaculture. In particular we are interested in farmed and wild salmonid fish that have particular relevance both industrially and commercially to the UK. These species are subject to parasitism from a number of copepod parasites of the family Caligidae. These organisms are well studied with a large literature: below we draw attention to some of the key review papers. Sea lice cause reduced growth and appetite, wounding, and susceptibility to secondary infections [5], resulting in significant damage to crops and therefore they are economically important. For salmon, louse burden in excess of 0.1 lice per gram of fish can be considered pathogenic [5]. The best studied species is *Lepeophtheirus salmonis*, principally a parasite of salmonids and frequent parasite on British Atlantic salmon (*Salmo salar*) farms [22]. It also infects sea trout (*Salmo trutta*) and rainbow trout (*Oncorhynchus mykiss*). The life history of the parasite is direct, with no requirement for intermediate hosts. It involves a succession of 10 distinct developmental stages, separated by moults, from egg to adult. Initial *naupliar* and *copepodid* stages are free living and planktonic. Following attachment of the infective copepodid to a host, the parasite passes through four *chalimus* stages that are firmly attached to the host, before entering sexually dimorphic *pre-adult* and *adult* stages where the parasite can once again move over the host surface and transfer to new hosts.

The state of the art for population-level modelling of *L. salmonis* is represented by Revie *et al.* [20]. These authors presented a series of delay differential equations to model different life history stages and parameterised the model using data collected around Scottish salmon farms. A similar compartmental model was proposed by Tucker *et al.* [21]. The emphasis of these papers was not however, in analytical study, but on numerical simulation and parameterisation using field [20] and lab [21] data. An earlier model by Heuch & Mo [13] investigated the infectivity, in term of *L. salmonis* egg production, posed by the Norwegian salmon industry, using a simple deterministic model. Other authors have considered the potential for long-distance dispersal of mobile parasite stages through sea currents [18], looking at Loch Shielraig in NW Scotland, a long-term study site for sea lice research.

In this paper we focus on the dynamics of individuals at the chalimus stage. At this stage individuals are not yet sexually differentiated. Individuals entering the chalimus stage from the non-feeding planktonic stage are distributed over different sizes, hence we have the zero influx boundary condition (1.2) and the recruitment term in (1.1). Of course, modelling the whole sea lice population will involve a rather complex model and system (1.1)-(1.3) would just describe the dynamics at the chalimus stage. In that setting the vital rates will be more specific and for example the fertility β would refer to the fertility of gravid females in the population. In fact, our aim here is to present a preliminary step towards the analysis of this rather complex model by giving a mathematical treatment of a quite general scramble competition model with distributed states-at-birth. We use the words scramble competition to describe the scenario when individuals in the population have the same chance when competing for resources such as food (see e.g. [6]). Therefore in our present model all the vital rates, i.e. growth, fertility and mortality depend on the total population size. In other populations, such as a tree population or a cannibalistic population, there is a natural hierarchy among individuals of different sizes, which results in mathematical models incorporating infinite dimensional nonlinearities, see e.g. [10, 11]. We also note that the analysis presented in this paper could be extended to these type of models and also to other models, for example to ones which involve a different type of recruitment term.

In the present work we are particularly interested in the asymptotic behaviour of solutions of model (1.1)-(1.3). Our analysis will be based on linearisation around equilibrium solutions (see e.g. [10, 19]) and we will utilise well-known results from linear operator theory which can be found for example in the excellent monographs [1, 4, 9]. We will also utilise some novel ideas on positive perturbations of linear operators. For basic concepts and results from the theory of structured population dynamics we refer the interested reader to [6, 14, 17, 23].

Traditionally, structured population models have been formulated as partial differential equations for population densities. However, the recent unified approach of Diekmann *et al.*, making use of the rich theory of delay and integral equations, has been resulted in significant results. In fact, the Principle of Linearized Stability has been proven in [7, 8] for a wide class of physiologically structured population models formulated as delay equations (or abstract integral equations). We shall also note that it is not clear yet whether the models formulated in [7, 8] as delay equations are equivalent to those formulated as partial differential equations.

In the remarkable paper [3], Calsina and Saldaña studied well-posedness of a very general size-structured model with distributed states-at-birth. They established global existence and uniqueness of solutions utilising results from the theory of nonlinear evolution equations. Model (1.1)-(1.3) is a special case of the general model treated in [3], however, in [3] qualitative questions were not addressed. In contrast to [3], our paper focuses on the existence and local asymptotic stability

of equilibrium solutions of system (1.1)-(1.3) with particular regards to the effects of the distributed states-at-birth compared to more simple models we addressed previously e.g. in [10]. First, we will establish conditions in theorem 2.6 that guarantee the existence of equilibrium solutions, in general. Then we show in Theorem 3.2 that a positive quasicontraction semigroup describes the evolution of solutions of the system linearized at an equilibrium solution. Then we establish a further regularity property in Theorem 3.6 of the governing linear semigroup that allows in principle to address stability questions of positive equilibrium solutions of (1.1)-(1.3). We use rank one perturbations of the general recruitment term to arrive at stability/instability conditions for equilibria. Finally we briefly discuss some issues regarding positivity of the governing linear semigroup.

2. Existence of equilibrium solutions

Model (1.1)-(1.3) admits the trivial solution. If we look for positive time-independent solutions of (1.1)-(1.3) we arrive at the following integro-differential equation

$$\gamma(s, P_*)p'_*(s) + (\gamma_s(s, P_*) + \mu(s, P_*))p_*(s) = \int_0^m \beta(s, y, P_*)p_*(y) dy \quad (2.1)$$

$$\gamma(0, P_*)p_*(0) = 0, \quad P_* = \int_0^m p_*(s) ds. \quad (2.2)$$

2.1 Separable fertility function

In the special case of

$$\beta(s, y, P) = \beta_1(s, P)\beta_2(y), \quad s, y \in [0, m], \quad P \in (0, \infty), \quad (2.3)$$

equation (2.1) reduces to

$$\gamma(s, P_*)p'_*(s) + (\gamma_s(s, P_*) + \mu(s, P_*))p_*(s) = \beta_1(s, P_*)\bar{P}_*, \quad (2.4)$$

where

$$\bar{P}_* = \int_0^m \beta_2(y)p_*(y) dy.$$

The solution of (2.4) satisfying the initial condition in (2.2) is readily obtained as

$$p_*(s) = \bar{P}_*F(s, P_*) \int_0^s \frac{\beta_1(y, P_*)}{F(y, P_*)\gamma(y, P_*)} dy, \quad (2.5)$$

where

$$F(s, P_*) = \exp \left\{ - \int_0^s \frac{\gamma_s(y, P_*) + \mu(y, P_*)}{\gamma(y, P_*)} dy \right\}.$$

Multiplying equation (2.5) by β_2 and integrating from 0 to m yields the following necessary condition for the existence of a positive equilibrium solution

$$1 = \int_0^m \beta_2(s)F(s, P_*) \int_0^s \frac{\beta_1(y, P_*)}{F(y, P_*)\gamma(y, P_*)} dy ds. \quad (2.6)$$

Therefore we define a net reproduction function R as follows

$$R(P) = \int_0^m \int_0^s \frac{\beta_1(y, P)\beta_2(s)}{\gamma(s, P)} \exp \left\{ - \int_y^s \frac{\mu(z, P)}{\gamma(z, P)} dz \right\} dy ds. \quad (2.7)$$

On the other hand it is straightforward to show that for every positive value P_* for which $R(P_*) = 1$ holds formula (2.5) yields a unique positive stationary solution p_* , where \bar{P}_* may be determined from equation (2.5) as

$$\bar{P}_* = \frac{P_*}{\int_0^m F(s, P_*) \int_0^s \frac{\beta_1(y, P_*)}{F(y, P_*)} dy ds}.$$

Then it is straightforward to establish the following result.

Proposition 2.1. *Assume that the fertility function β satisfies (2.3) and that the following conditions hold true:*

$$\int_0^m \exp \left\{ - \int_0^s \frac{\mu(y, 0)}{\gamma(y, 0)} dy \right\} ds < m - 1, \quad \text{and} \quad \beta(s, y, 0) > \mu(s, 0) \quad (2.8)$$

for $s, y \in [0, m]$, $P \in (0, \infty)$; and

$$\int_0^m \beta_1(s, P) ds \rightarrow 0 \quad \text{as} \quad P \rightarrow \infty, \quad \text{and} \quad 0 < \gamma^* \leq \gamma(s, P), \quad (2.9)$$

for $s \in [0, m]$, $P \in (0, \infty)$. Then model (1.1)-(1.3) admits at least one positive equilibrium solution.

Proof. Condition (2.8) implies

$$\begin{aligned} R(0) &= \int_0^m \exp \left\{ - \int_0^s \frac{\mu(y, 0)}{\gamma(y, 0)} dy \right\} \int_0^s \frac{\beta_2(s)\beta_1(y, 0)}{\gamma(y, 0)} \exp \left\{ \int_0^y \frac{\mu(z, 0)}{\gamma(z, 0)} dz \right\} dy ds \\ &> \int_0^m \exp \left\{ - \int_0^s \frac{\mu(y, 0)}{\gamma(y, 0)} dy \right\} \int_0^s \left(\exp \left\{ \int_0^y \frac{\mu(z, 0)}{\gamma(z, 0)} dz \right\} \right)' dy ds \\ &> 1. \end{aligned} \quad (2.10)$$

On the other hand, condition (2.9) and the growth behaviour of the functions involved in (2.7) imply that

$$\lim_{P \rightarrow +\infty} R(P) = 0,$$

hence the claim holds true on the grounds of the Intermediate Value Theorem. \square

2.2 The general case

For a fixed $P \in (0, \infty)$ we define the operator \mathcal{B}_P by

$$\mathcal{B}_P u = - \frac{\partial}{\partial s} (\gamma(\cdot, P)u) - \mu(\cdot, P)u + \int_0^m \beta(\cdot, y, P)u(y) dy,$$

$$\text{Dom}(\mathcal{B}_P) = \{u \in W^{1,1}(0, m) \mid u(0) = 0\}. \quad (2.11)$$

Our goal is to show, that there exists a P_* such that the operator \mathcal{B}_{P_*} has eigenvalue 0 with a corresponding unique positive eigenvector. To this end, first we establish that \mathcal{B}_P is the generator of a positive semigroup. Then we determine conditions that guarantee that it generates an irreducible semigroup. We also establish that the governing linear semigroup is eventually compact, which implies that the Spectral Mapping Theorem holds true for the semigroup and its generator, and the spectrum of the generator may contain only isolated eigenvalues of finite algebraic multiplicity. It then follows that the spectral bound is a dominant (real) eigenvalue λ_P of geometric multiplicity one with a corresponding positive eigenvector [4, Chapter 9]. Finally we need to establish conditions which imply that there exist a $P^+ \in (0, \infty)$ such that the spectral bound $s(\mathcal{B}_{P^+})$ is negative and therefore the dominant eigenvalue $\lambda_{P^+} = s(\mathcal{B}_{P^+})$ is also negative; and a $P^- \in (0, \infty)$ such that this dominant eigenvalue $\lambda_{P^-} = s(\mathcal{B}_{P^-})$ is positive. Then it follows from standard perturbation results on eigenvalues (see e.g. [15]) that there exists a zero eigenvalue. A similar strategy was employed in [2] to establish existence and uniqueness of an equilibrium solution of a cyclin structured cell population model.

Lemma 2.2. *For every $P \in (0, \infty)$ the semigroup $\mathcal{T}(t)$ generated by the operator \mathcal{B}_P is positive.*

Proof. We rewrite (2.11) as, $\mathcal{B}_P = \mathcal{A}_P + \mathcal{C}_P$, where

$$\begin{aligned} \mathcal{A}_P u &= -\frac{\partial}{\partial s} (\gamma(\cdot, P)u) - \mu(\cdot, P)u \\ \text{Dom}(\mathcal{A}_P) &= \{u \in W^{1,1}(0, m) \mid u(0) = 0\}, \\ \mathcal{C}_P u &= \int_0^m \beta(\cdot, y, P)u(y) dy, \\ \text{Dom}(\mathcal{C}_P) &= L^1(0, m). \end{aligned} \tag{2.12}$$

For $0 \leq f \in L^1(0, m)$ the solution of the resolvent equation

$$(\lambda \mathcal{I} - \mathcal{A}_P)u = f,$$

is

$$u(s) = \int_0^s \exp \left\{ - \int_y^s \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} f(y) dy.$$

This shows that the resolvent operator $\mathcal{R}(\lambda, \mathcal{A}_P)$ is a positive bounded operator, hence \mathcal{A}_P generates a positive semigroup. Since \mathcal{C}_P is a positive and bounded operator, the statement follows. \square

Lemma 2.3. *The linear semigroup $\mathcal{T}(t)$ generated by the operator \mathcal{B}_P is eventually compact.*

Proof. We note that \mathcal{A}_P generates a nilpotent semigroup, while it is easily shown that \mathcal{C}_P is a compact operator if conditions (1.4) hold true. (For more details see also Theorem 3.6.) \square

Lemma 2.4. *Assume that for every $P \in (0, \infty)$ there exists an $\varepsilon_0 > 0$ such that for all $0 < \varepsilon \leq \varepsilon_0$*

$$\int_0^\varepsilon \int_{m-\varepsilon}^m \beta(s, y, P) \, dy \, ds > 0. \quad (2.13)$$

Then the linear semigroup $\mathcal{T}(t)$ generated by the operator \mathcal{B}_P is irreducible.

Proof. We only need to show that under condition (2.13) for every $p_0 \in L^1_+(0, m)$ there exists a t_0 such that

$$\text{supp } \mathcal{T}(t_0)p_0 = [0, m],$$

for all $t \geq t_0$. Since $\gamma > 0$, there exist t_* such that

$$\text{supp } \mathcal{T}(t)p_0 \cap \text{supp } \beta(s, \cdot) \neq \emptyset$$

for every $t_* \leq t$ and every $s \in (0, \varepsilon]$. By assumption (2.13), $\mathcal{T}(t)p_0(s) > 0$ for $t_* \leq t$ and $s \in (0, \varepsilon]$. After this, eventually the support of the solution $\mathcal{T}(t_0)p_0$ will cover the entire size space $[0, m]$. \square

Lemma 2.5. *Assume that there exists a $\beta^-(s, y, P) = \beta_1^-(s, P)\beta_2^-(y)$ and a $P^- \in (0, \infty)$ such that*

$$\beta_1^-(s, P^-)\beta_2^-(y) \leq \beta(s, y, P^-), \quad s, y \in [0, m], \quad (2.14)$$

and

$$\int_0^m \int_0^s \frac{\beta_1^-(y, P^-)\beta_2^-(s)}{\gamma(y, P^-)} \exp \left\{ - \int_y^s \frac{\gamma_s(z, P^-) + \mu(z, P^-)}{\gamma(z, P^-)} \, dz \right\} \, dy \, ds > 1, \quad (2.15)$$

and a $\beta^+(s, y, P) = \beta_1^+(s, P)\beta_2^+(y)$ and a $P^+ \in (0, \infty)$ such that

$$\beta(s, y, P^+) \leq \beta_1^+(s, P^+)\beta_2^+(y), \quad (2.16)$$

and

$$\int_0^m \int_0^s \frac{\beta_1^+(y, P^+)\beta_2^-(s)}{\gamma(y, P^+)} \exp \left\{ - \int_y^s \frac{\gamma_s(z, P^+) + \mu(z, P^+)}{\gamma(z, P^+)} \, dz \right\} \, dy \, ds < 1. \quad (2.17)$$

Then the operator \mathcal{B}_{P^-} has a dominant real eigenvalue $\lambda_{P^-} > 0$ and the operator \mathcal{B}_{P^+} has a dominant real eigenvalue $\lambda_{P^+} < 0$, with corresponding positive eigenvectors.

Proof. First assume, that there exists a $\beta^-(s, y, P) = \beta_1^-(s, P)\beta_2^-(y)$ and a P^- such that condition (2.14) and (2.15) hold true. Let \mathcal{B}_{P^-} denote the operator

that corresponds to the fertility β^- and the constant P^- . The solution of the eigenvalue problem

$$\mathcal{B}_{P^-}^- u = \lambda u, \quad u(0) = 0 \quad (2.18)$$

is

$$u(s) = \int_0^m \beta_2^-(s) u(s) ds \int_0^s \frac{\beta_1^-(y, P^-)}{\gamma(y, P^-)} \exp \left\{ - \int_y^s \frac{\lambda + \gamma_s(z, P^-) + \mu(z, P^-)}{\gamma(z, P^-)} dz \right\} dy. \quad (2.19)$$

We multiply equation (2.19) by β_2^- and integrate from 0 to m to arrive at the characteristic equation

$$1 = \int_0^m \beta_2^-(s) \int_0^s \frac{\beta_1^-(y, P^-)}{\gamma(y, P^-)} \exp \left\{ - \int_y^s \frac{\lambda + \gamma_s(z, P^-) + \mu(z, P^-)}{\gamma(z, P^-)} dz \right\} dy ds. \quad (2.20)$$

It is clear that equation (2.20) admits a unique dominant real solution $\lambda_{P^-}^- > 0$ if condition (2.15) holds true. Since $\mathcal{B}_{P^-}^-$ is a generator of a positive semigroup and $(\mathcal{B}_{P^-} - \mathcal{B}_{P^-}^-)$ is a positive (and bounded) operator by condition (2.14), it follows that \mathcal{B}_{P^-} has a dominant real eigenvalue $\lambda_{P^-} \geq \lambda_{P^-}^- > 0$, see e.g. [9, Corollary VI.1.11].

In a similar way, let us assume that there exists a $\beta^+(s, y, P) = \beta_1^+(s, P)\beta_2^+(y)$ and a P^+ such that condition (2.16) and (2.17) hold true. Let $\mathcal{B}_{P^+}^+$ denote the operator which corresponds to the fertility β^+ and the constant P^+ . The solution of the eigenvalue problem

$$\mathcal{B}_{P^+}^+ u = \lambda u, \quad u(0) = 0 \quad (2.21)$$

is now

$$u(s) = \int_0^m \beta_2^+(s) u(s) ds \int_0^s \frac{\beta_1^+(y, P^+)}{\gamma(y, P^+)} \exp \left\{ - \int_y^s \frac{\lambda + \gamma_s(z, P^+) + \mu(z, P^+)}{\gamma(z, P^+)} dz \right\} dy. \quad (2.22)$$

We multiply equation (2.22) by β_2^+ and integrate from 0 to m to arrive at the characteristic equation

$$1 = \int_0^m \beta_2^+(s) \int_0^s \frac{\beta_1^+(y, P^+)}{\gamma(y, P^+)} \exp \left\{ - \int_y^s \frac{\lambda + \gamma_s(z, P^+) + \mu(z, P^+)}{\gamma(z, P^+)} dz \right\} dy ds. \quad (2.23)$$

It is clear that equation (2.23) admits a unique dominant real solution $\lambda_{P^+}^+ < 0$ if condition (2.17) holds true. Since \mathcal{B}_{P^+} is a generator of a positive semigroup and $(\mathcal{B}_{P^+}^+ - \mathcal{B}_{P^+})$ is a positive operator by condition (2.16), it follows that \mathcal{B}_{P^+} has a dominant real eigenvalue $\lambda_{P^+} \leq \lambda_{P^+}^+ < 0$.

In both cases, the positivity of the corresponding eigenvector follows from the irreducibility of the semigroup $\mathcal{T}(t)$, see [4, Theorem 9.11]. \square

Theorem 2.6. *Assume that conditions (2.13), (2.14)-(2.17) are satisfied. Then system (1.1)-(1.3) admits at least one positive equilibrium solution.*

Proof. Let $P^* > 0$ be such that $s(\mathcal{B}_{P^*}) = 0$. Then, since the spectrum consists only of isolated eigenvalues we have $\lambda_{P^*} = s(\mathcal{B}_{P^*}) = 0$ and there exists a corresponding positive eigenvector p_* . Then $\frac{P^*}{\|p_*\|_1} p_*$ is the desired equilibrium solution with total population size P^* . \square

3. The linearised semigroup and its regularity

First we note that when we say linearised semigroup, we mean the linear semigroup governing the linearised system. However, since it was proved in [3] that model (1.1)-(1.3) is well-posed, there exists a semigroup of nonlinear operators $\{\Sigma(t)\}_{t \geq 0}$ defined via $\Sigma(t)p(s, 0) = p(s, t)$. We also note that it was proven in [8] that if the nonlinearities are smooth enough (namely, the vital rates are differentiable) then this nonlinear semigroup $\Sigma(t)$ is Frechét differentiable and the Frechét derivative around an equilibrium solution p_* defines a semigroup of bounded linear operators. In this section we will establish the existence of this semigroup and at the same time arrive at a condition which guarantees that it is positive, as well.

Given a positive stationary solution p_* of system (1.1)-(1.3), we introduce the perturbation $u = u(s, t)$ of p by making the ansatz $p = u + p_*$. Then we are using Taylor series expansions of the vital rates to arrive at the linearised problem (see e.g. [10])

$$\begin{aligned} u_t(s, t) = & -\gamma(s, P_*) u_s(s, t) - (\gamma_s(s, P_*) + \mu(s, P_*)) u(s, t) \\ & - (\gamma_{sP}(s, P_*) p_*(s) + \mu_P(s, P_*) p_*(s) + \gamma_P(s, P_*) p_*'(s)) U(t) \\ & + \int_0^m u(y, t) \left(\beta(s, y, P_*) + \int_0^m \beta_P(s, z, P_*) p_*(z) dz \right) dy, \end{aligned} \quad (3.1)$$

$$\gamma(0, P_*) u(0, t) = 0 \quad (3.2)$$

where we have set

$$U(t) = \int_0^m u(s, t) ds. \quad (3.3)$$

Eqs. (3.1)–(3.2) are accompanied by the initial condition

$$u(s, 0) = u_0(s). \quad (3.4)$$

Our first objective is to establish conditions which guarantee that the linearised system is governed by a positive semigroup. To this end, we cast the linearised system (3.1)-(3.4) in the form of an abstract Cauchy problem on the state space $\mathcal{X} = L^1(0, m)$ as follows

$$\frac{d}{dt} u = (\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}) u, \quad u(0) = 0, \quad (3.5)$$

where

$$\mathcal{A}u = -\gamma(\cdot, P_*)u_s \quad \text{with domain } \text{Dom}(\mathcal{A}) = \{u \in W^{1,1}(0, m) \mid u(0) = 0\}, \quad (3.6)$$

$$\mathcal{B}u = -(\gamma_s(\cdot, P_*) + \mu(\cdot, P_*))u \quad \text{on } \mathcal{X}, \quad (3.7)$$

$$\begin{aligned} \mathcal{C}u &= -(\gamma_{sP}(\cdot, P_*)p_* + \mu_P(\cdot, P_*)p_* + \gamma_P(\cdot, P_*)p'_*) \int_0^m u(s) ds \\ &= -\rho_*(\cdot) \int_0^m u(s) ds \quad \text{on } \mathcal{X}, \end{aligned} \quad (3.8)$$

$$\mathcal{D}u = \int_0^m u(y) \left(\beta(\cdot, y, P_*) + \int_0^m \beta_P(\cdot, z, P_*)p_*(z) dz \right) dy \quad \text{on } \mathcal{X}, \quad (3.9)$$

where ρ_* is defined via equation (3.8). Our aim is to establish that the linear operator $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}$ is a generator of a quasicontraction semigroup. To this end first we recall (see e.g. [1, 4, 9]) some basic concepts from the theory of linear operators acting on Banach spaces. Let \mathcal{O} be a linear operator defined on the real Banach space \mathcal{Y} with norm $\|\cdot\|$. \mathcal{O} is called dissipative if for every $\lambda > 0$ and $x \in \text{Dom}(\mathcal{O})$,

$$\|(\mathcal{I} - \lambda\mathcal{O})x\| \geq \|x\|.$$

Furthermore, a function $f : \mathcal{Y} \rightarrow \mathbf{R}$ is called sublinear if

$$\begin{aligned} f(x + y) &\leq f(x) + f(y), \quad x, y \in \mathcal{Y} \\ f(\lambda x) &= \lambda f(x), \quad \lambda \geq 0, \quad x \in \mathcal{Y}. \end{aligned}$$

If also $f(x) + f(-x) > 0$ holds true for $x \neq 0$ then f is called a half-norm on \mathcal{Y} . The linear operator \mathcal{O} is called f -dissipative if

$$f(x) \leq f(x - \lambda\mathcal{O}x), \quad \lambda \geq 0, \quad x \in \text{Dom}(\mathcal{O}).$$

An operator \mathcal{O} which is p -dissipative with respect to the half norm

$$p(x) = \|x^+\|,$$

is called dispersive, where $x^+ = x \vee 0$ (and $x^- = (-x)^+$). Finally a C_0 semigroup $\{\mathcal{T}(t)\}_{t \geq 0}$ is called quasicontractive if

$$\|\mathcal{T}(t)\| \leq e^{\omega t}, \quad t \geq 0,$$

for some $\omega \in \mathbf{R}$, and it is called contractive if $\omega \leq 0$. We recall the following characterization theorem from [4].

Theorem 3.1. *Let \mathcal{Y} be a Banach lattice and let $\mathcal{O} : \text{Dom}(\mathcal{O}) \rightarrow \mathcal{Y}$ be a linear operator. Then, the following statements are equivalent.*

- (i) \mathcal{O} is the generator of a positive contraction semigroup.
- (ii) \mathcal{O} is densely defined, $\text{Rg}(\lambda\mathcal{I} - \mathcal{O}) = \mathcal{Y}$ for some $\lambda > 0$, and \mathcal{O} is dispersive.

We also recall that \mathcal{O} is dispersive if for every $x \in \text{Dom}(\mathcal{O})$ there exists $\phi \in \mathcal{Y}^*$ with $0 \leq \phi$, $\|\phi\| \leq 1$ and $(x, \phi) = \|x^+\|$ such that $(\mathcal{O}x, \phi) \leq 0$, where (\cdot, \cdot) is the natural pairing between elements of \mathcal{Y} and its dual \mathcal{Y}^* .

Theorem 3.2. *The operator $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}$ generates a positive strongly continuous (C_0 for short) quasicontraction semigroup $\{\mathcal{T}(t)\}_{t \geq 0}$ of bounded linear operators on \mathcal{X} if the following condition holds true*

$$\rho_*(s) \leq \beta(s, y, P_*) + \int_0^m \beta_P(s, y, P_*) p_*(y) dy, \quad s, y \in [0, m], \quad (3.10)$$

where ρ_* is defined via equation (3.8).

Proof. Our aim is to apply the previous characterization theorem for the perturbed operator $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} - \omega \mathcal{I}$, for some $\omega \in \mathbf{R}$. To this end, for every $u \in \text{Dom}(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} - \omega \mathcal{I})$ we define $\phi_u \in \mathcal{X}^*$ by

$$\phi_u(s) = \frac{u^+(s)}{|u(s)|}, \quad s \in [0, m], \quad u(s) \neq 0, \quad (3.11)$$

if $u(s) = 0$ then let $\phi_u(s) = 0$. Then

$$\|\phi_u\|_\infty \leq 1,$$

and clearly

$$(u, \phi_u) = \int_0^m u(s) \phi_u(s) ds = \|u^+\|_1.$$

Making use of condition (3.10) we obtain the following estimate.

$$\begin{aligned} & ((\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} - \omega \mathcal{I})u, \phi_u) \\ &= - \int_0^m \mathbf{1}_{u^+}(s) (\gamma(s, P_*)u(s))_s ds - \int_0^m \mathbf{1}_{u^+}(s) \mu(s, P_*)u(s) ds - \int_0^m \mathbf{1}_{u^+}(s) \omega u(s) ds \\ & \quad + \int_0^m \mathbf{1}_{u^+}(s) \int_0^m u(y) \left(\beta(s, y, P_*) + \int_0^m \beta_P(s, z, P_*) p_*(z) dz - \rho_*(s) \right) dy ds \\ & \leq - \int_0^m \mathbf{1}_{u^+}(s) (\gamma(s, P_*)u(s))_s ds - \omega \|u^+\|_1 - \inf_{s \in [0, m]} \mu(s, P_*) \|u^+\|_1 \\ & \quad + \|u^+\|_1 \left\| \sup_{y \in [0, m]} \left(\beta(s, y, P_*) + \int_0^m \beta_P(s, z, P_*) p_*(z) dz - \rho_*(s) \right) \right\|_\infty \\ & \leq -\omega \|u^+\|_1 - (\gamma(m, P_*)u(m)) \mathbf{1}_{u^+}(m) \\ & \quad + \|u^+\|_1 \left\| \sup_{y \in [0, m]} \left(\beta(s, y, P_*) + \int_0^m \beta_P(s, z, P_*) p_*(z) dz - \rho_*(s) \right) \right\|_\infty \\ & \leq 0, \end{aligned} \quad (3.12)$$

for some $\omega \in \mathbf{R}$ large enough, hence the operator $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} - \omega\mathcal{I}$ is dispersive. The operator $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} - \omega\mathcal{I}$ is clearly densely defined. We observe that the equation

$$(\lambda I - \mathcal{A})u = h \quad (3.13)$$

for $h \in \mathcal{X}$ and $\lambda > 0$ sufficiently large has a unique solution $u \in \text{Dom}(\mathcal{A})$, given by

$$u(s) = \exp \left\{ - \int_0^s \frac{\lambda}{\gamma(y, P_*)} dy \right\} \int_0^s \exp \left\{ \int_0^y \frac{\lambda}{\gamma(z, P_*)} dz \right\} \frac{h(y)}{\gamma(y, P_*)} dy. \quad (3.14)$$

The fact that $u \in \text{Dom}(\mathcal{A})$ is well defined by (3.14) follows from

$$\begin{aligned} |u'(s)| &\leq \left| \frac{h(s)}{\gamma(s, P_*)} \right| + \frac{\lambda}{\gamma(s, P_*)} \int_0^m \exp \left\{ - \int_y^s \frac{\lambda}{\gamma(z, P_*)} dz \right\} \frac{|h(y)|}{\gamma(y, P_*)} dy \\ &\leq \left| \frac{h(s)}{\gamma(s, P_*)} \right| + M_\lambda, \end{aligned}$$

for λ large enough for some $M_\lambda < \infty$, that is $u \in W^{1,1}(0, m)$. Since $\mathcal{B} + \mathcal{C} + \mathcal{D} - \omega\mathcal{I}$ is bounded, the range condition is satisfied. Theorem 3.1 gives that $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} - \omega\mathcal{I}$ is a generator of a positive contraction semigroup. Since the operator $\omega\mathcal{I}$ is positive (clearly if the dispersivity estimate holds true with an $\omega < 0$ than it holds true with any other $\omega^* > \omega$) a well-known perturbation result (see e.g. [9]) yields that $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}$ is a generator of a positive quasicontraction semigroup \mathcal{T} which obeys

$$\|\mathcal{T}(t)\| \leq e^{\omega t}, \quad t \geq 0. \quad \square$$

Remark 3.3 The proof of Theorem 3.1 shows that if

$$\inf_{s \in [0, m]} \mu(s, P_*) > \left\| \sup_{y \in [0, m]} \left(\beta(s, y, P_*) + \int_0^m \beta_P(s, z, P_*) p_*(z) dz - \rho_*(s) \right) \right\|_\infty$$

holds, then the growth bound ω_0 of the semigroup is negative, hence the semigroup $\{\mathcal{T}(t)\}_{t \geq 0}$ is uniformly exponentially stable (see e.g. [9]), i.e. the equilibrium p_* is locally asymptotically stable.

Remark 3.4 We note that the operator $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}$ is in general a generator of a C_0 quasicontraction (but not positive) semigroup. The proof of this would utilise the Lumer-Phillips Theorem (see e.g. [1, 4, 9]) and goes along similar lines, obtaining a dissipativity estimate in terms of u rather than u^+ , see e.g. [11]. This implies that the linearized problem (3.1)-(3.2) is well-posed.

Remark 3.5 Note that if $\beta = \beta(s, y)$, $\mu = \mu(s)$, $\gamma = \gamma(s)$, i.e. model (1.1)-(1.3) is a linear one, then the biologically relevant conditions $\mu, \beta \geq 0$ and $\gamma > 0$ imply that it is governed by a positive quasicontraction semigroup.

Theorem 3.6. *The semigroup $\{\mathcal{T}(t)\}_{t \geq 0}$ generated by the operator $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}$ is eventually compact.*

Proof. \mathcal{C} is a rank one operator hence it is compact on $\mathcal{X} = L^1(0, m)$. \mathcal{D} is linear and bounded. Hence in view of the Fréchet-Kolmogorov compactness criterion in L^p (see e.g. [24]) we need to show that

$$\lim_{t \rightarrow 0} \int_0^m |\mathcal{D}u(t+s) - \mathcal{D}u(s)| ds = 0, \quad \text{uniformly in } u,$$

for $u \in B$, where B is the unit sphere of $L^1(0, m)$. But this follows from the regularity assumptions we made on β based on the following estimate

$$|\mathcal{D}u(s_1) - \mathcal{D}u(s_2)| \leq \|u\|_1 \left\| \beta(s_1, y, P_*) + \int_0^m \beta_P(s_1, z, P_*) p_*(z) dz - \beta(s_2, y, P_*) + \int_0^m \beta_P(s_2, z, P_*) p_*(z) dz \right\|_\infty.$$

Therefore, it suffices to investigate the operator $\mathcal{A} + \mathcal{B}$. To this end, we note that the abstract differential equation

$$\frac{d}{dt} u = (\mathcal{A} + \mathcal{B}) u \tag{3.15}$$

corresponds to the partial differential equation

$$u_t(s, t) + \gamma(s, P_*) u_s(s, t) + (\gamma_s(s, P_*) + \mu(s, P_*)) u(s, t) = 0, \tag{3.16}$$

subject to the boundary condition (3.2). We solve easily equation (3.16) for example using the method of characteristics. For $t > \Gamma(m)$ we arrive at

$$u(s, t) = u(0, t - \Gamma(s)) \exp \left\{ - \int_0^s \frac{\gamma_s(y, P_*) + \mu(y, P_*)}{\gamma(y, P_*)} dy \right\} = 0, \tag{3.17}$$

where

$$\Gamma(s) = \int_0^s \frac{1}{\gamma(y, P_*)} dy.$$

This means that the semigroup $\{\mathcal{T}(t)\}_{t \geq 0}$ generated by $\mathcal{A} + \mathcal{B}$ is nilpotent. In particular it is compact for $t > \Gamma(m)$ and the claim follows. \square

Remark 3.7 Theorem 3.6 implies that the Spectral Mapping Theorem holds true for the semigroup $\{\mathcal{T}(t)\}_{t \geq 0}$ with generator $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}$ and that the spectrum $\sigma(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D})$ contains only isolated eigenvalues of finite multiplicity (see e.g. [9]).

4. (In) Stability

Our aim in this section is to study the stability of positive equilibrium solutions by studying the point spectrum of the linearised operator $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}$. The main difficulty is, that the eigenvalue equation

$$(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} - \mathcal{I})\lambda = 0,$$

cannot be solved explicitly, since the operator \mathcal{D} has infinite rank, in general. We encountered this problem previously in case of hierarchical size-structured population models, see e.g. [11, 12]. In [11] and [12] we used the dissipativity approach, presented in the previous section, to establish conditions which guarantee that the spectral bound of the linearized semigroup is negative. However, as we can see from remark 3.3 this approach gives a rather restrictive stability condition. Therefore, here we devise a different approach, which uses positive perturbation arguments.

Theorem 4.1. *Assume that there exists an $\varepsilon > 0$ such that*

$$\beta(s, y, P_*) - \rho_*(s) - \varepsilon + \int_0^m \beta_P(s, y, P_*) p_*(y) dy \geq 0, \quad s, y \in [0, m], \quad (4.1)$$

and

$$\varepsilon \int_0^m \int_0^s \exp \left\{ - \int_y^s \frac{\gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} dy ds > 1. \quad (4.2)$$

Then the stationary solution $p_(s)$ of model (1.1)-(1.3) is linearly unstable.*

Proof. Let $\varepsilon > 0$, and define the operator \mathcal{F}_ε on \mathcal{X} as

$$\mathcal{F}_\varepsilon u = \varepsilon \int_0^m u(s) ds = \varepsilon \bar{u}.$$

We first find the solution of the eigenvalue equation

$$(\mathcal{A} + \mathcal{B} + \mathcal{F}_\varepsilon)u = \lambda u$$

as

$$\begin{aligned} u(s) = & \varepsilon \bar{u} \exp \left\{ - \int_0^s \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} \\ & \times \int_0^s \exp \left\{ \int_0^y \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} dy. \end{aligned} \quad (4.3)$$

Next we integrate the solution (4.3) over $[0, m]$ to obtain

$$\begin{aligned} \bar{u} = & \varepsilon \bar{u} \int_0^m \left[\exp \left\{ - \int_0^s \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} \right. \\ & \left. \times \int_0^s \exp \left\{ \int_0^y \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} dy \right] ds. \end{aligned} \quad (4.4)$$

We note that, if $\bar{u} = 0$ then equation (4.3) shows that $u(s) \equiv 0$, hence we have a non-trivial eigenvector if and only if $\bar{u} \neq 0$ and λ satisfies the following characteristic equation

$$\begin{aligned} 1 = K(\lambda) \stackrel{\text{def}}{=} & \varepsilon \int_0^m \left[\exp \left\{ - \int_0^s \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} \right. \\ & \left. \times \int_0^s \exp \left\{ \int_0^y \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} dy \right] ds. \end{aligned} \quad (4.5)$$

It is easily shown that

$$\lim_{\lambda \rightarrow +\infty} K(\lambda) = 0,$$

therefore it follows from condition (4.2), on the grounds of the Intermediate Value Theorem, that equation (4.5) has a positive (real) solution. Hence we have

$$0 < s(\mathcal{A} + \mathcal{B} + \mathcal{F}_\varepsilon),$$

where $s(\mathcal{A} + \mathcal{B} + \mathcal{F}_\varepsilon)$ stands for the spectral bound of the operator $\mathcal{A} + \mathcal{B} + \mathcal{F}_\varepsilon$. Next, for a fixed $0 \leq f \in \mathcal{X}$, we obtain the solution of the resolvent equation

$$(\lambda \mathcal{I} - (\mathcal{A} + \mathcal{B} + \mathcal{F}_\varepsilon)) u = f,$$

as

$$\begin{aligned} u(s) = & \exp \left\{ - \int_0^s \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} \\ & \times \int_0^s \exp \left\{ \int_0^y \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} (\varepsilon \bar{u} + f(y)) dy. \end{aligned} \quad (4.6)$$

We integrate equation (4.6) from 0 to m to obtain

$$\bar{u} = \frac{\int_0^m \exp \left\{ - \int_0^s \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} \int_0^s \exp \left\{ \int_0^y \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} f(y) dy}{1 - \varepsilon \int_0^m \exp \left\{ - \int_0^s \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} \int_0^s \exp \left\{ \int_0^y \frac{\lambda + \gamma_s(\sigma, P_*) + \mu(\sigma, P_*)}{\gamma(\sigma, P_*)} d\sigma \right\} dy} \quad (4.7)$$

It follows from the growth behaviour of the exponential function and from assumptions (1.4), that \bar{u} is well-defined and non-negative for any $0 \leq f \in \mathcal{X}$ and λ large enough. Hence the resolvent operator

$$\mathcal{R}(\lambda, \mathcal{A} + \mathcal{B} + \mathcal{F}_\varepsilon) = (\lambda - (\mathcal{A} + \mathcal{B} + \mathcal{F}_\varepsilon))^{-1}$$

is positive, for λ large enough, which implies that $\mathcal{A} + \mathcal{B} + \mathcal{F}_\varepsilon$ generates a positive semigroup (see e.g. [9]).

Finally, we note that condition (4.1) guarantees that the operator $\mathcal{C} + \mathcal{D} - \mathcal{F}_\varepsilon$ is positive, hence we have for the spectral bound (see e.g. Corollary VI.1.11 in [9])

$$0 < s(\mathcal{A} + \mathcal{B} + \mathcal{F}_\varepsilon) \leq s(\mathcal{A} + \mathcal{B} + \mathcal{F}_\varepsilon + \mathcal{C} + \mathcal{D} - \mathcal{F}_\varepsilon) = s(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}),$$

and the result follows. \square

Next we show that for a separable fertility function we can indeed explicitly characterize the point spectrum of the linearised operator.

Theorem 4.2. *Assume that $\beta(s, y, P) = \beta_1(s, P)\beta_2(y) \stackrel{\text{def}}{=} \tilde{\beta}(s, y, P)$, $s, y \in [0, m]$, $P \in (0, \infty)$. Then for any $\lambda \in \mathbb{C}$, we have $\lambda \in \sigma(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D})$ if and only if λ satisfies the equation*

$$K_{\tilde{\beta}}(\lambda) \stackrel{\text{def}}{=} \det \begin{pmatrix} 1 + a_1(\lambda) & a_2(\lambda) \\ a_3(\lambda) & 1 + a_4(\lambda) \end{pmatrix} = 0, \quad (4.8)$$

where

$$\begin{aligned}
a_1(\lambda) &= - \int_0^m F(\lambda, s, P_*) \int_0^s \frac{g(y)}{F(\lambda, y, P_*)} dy ds, \\
a_2(\lambda) &= - \int_0^m F(\lambda, s, P_*) \int_0^s \frac{\beta_1(y, P_*)}{\gamma(y, P_*)F(\lambda, y, P_*)} dy ds, \\
a_3(\lambda) &= - \int_0^m \beta_2(s)F(\lambda, s, P_*) \int_0^s \frac{g(y)}{F(\lambda, y, P_*)} dy ds, \\
a_4(\lambda) &= - \int_0^m \beta_2(s)F(\lambda, s, P_*) \int_0^s \frac{\beta_1(y, P_*)}{\gamma(y, P_*)F(\lambda, y, P_*)} dy ds, \tag{4.9}
\end{aligned}$$

and

$$\begin{aligned}
g(s) &= \frac{\beta_{1P}(s, P_*) \int_0^m \beta_2(y)p_*(y) dy - \rho_*(s)}{\gamma(s, P_*)}, \quad s \in [0, m], \\
F(\lambda, s, P_*) &= \exp \left\{ - \int_0^s \frac{\lambda + \gamma_s(y, P_*) + \mu(y, P_*)}{\gamma(y, P_*)} dy \right\}, \quad s \in [0, m].
\end{aligned}$$

Proof. To characterize the point spectrum of $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}$ we consider the eigenvalue problem

$$(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} - \lambda \mathcal{I})U = 0, \quad U(0) = 0. \tag{4.10}$$

The solution of (4.10) is found to be

$$U(s) = \bar{U}F(\lambda, s, P_*) \int_0^s \frac{g(y)}{F(\lambda, y, P_*)} dy + \tilde{U}F(\lambda, s, P_*) \int_0^s \frac{\beta_1(y, P_*)}{\gamma(y, P_*)F(\lambda, y, P_*)} dy, \tag{4.11}$$

where

$$\bar{U} = \int_0^m U(s) ds, \quad \tilde{U} = \int_0^m \beta_2(s)U(s) ds.$$

We integrate equation (4.11) from zero to m and multiply equation (4.11) by $\beta_2(s)$ and then integrate from zero to m to obtain

$$\bar{U}(1 + a_1(\lambda)) + \tilde{U}a_2(\lambda) = 0, \tag{4.12}$$

$$\bar{U}a_3(\lambda) + \tilde{U}(1 + a_4(\lambda)) = 0. \tag{4.13}$$

If $\lambda \in \sigma(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D})$ then the eigenvalue equation (4.10) admits a non-trivial solution U hence there exists a non-zero vector (\bar{U}, \tilde{U}) which solves equations (4.12)-(4.13). On the other hand, if (\bar{U}, \tilde{U}) is a non-zero solution of equations (4.12)-(4.13) for some $\lambda \in \mathbb{C}$ then (4.11) yields a non-trivial solution U . This is

because the only scenario for U to vanish would yield

$$\bar{U}F(\lambda, s) \int_0^s \frac{g(y)}{F(\lambda, y)} dy = -\tilde{U}F(\lambda, s) \int_0^s \frac{\beta_1(y, P_*)}{\gamma(y, P_*)F(\lambda, y)} dy, \quad s \in [0, m].$$

This however, together with equations (4.12)-(4.13) would imply $\bar{U} = \tilde{U} = 0$, a contradiction, hence the proof is completed. \square

Theorem 4.3. *Assume that condition (3.10) holds true for some stationary solution p_* . Moreover, assume that there exists a function $\tilde{\beta}(s, y, P) = \beta_1(s, P)\beta_2(y)$ such that $\beta(s, y, P_*) \leq \tilde{\beta}(s, y, P_*)$ for $s, y \in [0, m]$ and the characteristic equation $K_{\tilde{\beta}}(\lambda) = 0$ does not have a solution with non-negative real part. Then the equilibrium solution p_* is linearly asymptotically stable.*

Proof. We need to establish that the spectral bound of the linearised operator $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D}$ is negative. To this end, we rewrite the operator \mathcal{D} as a sum of two operators, namely $\mathcal{D} = \mathcal{G} + \mathcal{H}_\beta$, where

$$\begin{aligned} \mathcal{G}u &= \bar{u} \int_0^m \beta_P(\cdot, z, P_*)p_*(z) dz, \quad \text{on } \mathcal{X}, \\ \mathcal{H}_\beta u &= \int_0^m u(y)\beta(\cdot, y, P_*) dy, \quad \text{on } \mathcal{X}. \end{aligned}$$

Condition (3.10) guarantees that $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{G} + \mathcal{H}_\beta$ is a generator of a positive semigroup, while the eventual compactness of the linearised semigroup assures that the spectrum of $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{G} + \mathcal{H}_{\tilde{\beta}}$ contains only eigenvalues and that the Spectral Mapping Theorem holds true. Since $\mathcal{H}_{\tilde{\beta}} - \mathcal{H}_\beta$ is a positive and bounded operator we have

$$\begin{aligned} s(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{G} + \mathcal{H}_\beta) &\leq s(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{G} + \mathcal{H}_\beta + \mathcal{H}_{\tilde{\beta}} - \mathcal{H}_\beta) \\ &= s(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{G} + \mathcal{H}_{\tilde{\beta}}) < 0, \end{aligned} \quad (4.14)$$

and the proof is completed. \square

Example 4.4 As we can see from equations (4.8)-(4.9) the characteristic function $K_{\tilde{\beta}}(\lambda)$ is rather complicated, in general. Therefore, here we only present a special case when it is straightforward to establish that the point spectrum of the linear operator $\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{G} + \mathcal{H}_{\tilde{\beta}}$ does not contain any element with non-negative real part. In particular, we make the following specific assumption

$$\beta_2(\cdot) \equiv \beta_2.$$

In this case we can cast the characteristic equation (4.8) in the simple form

$$\int_0^m \int_0^s \exp \left\{ - \int_y^s \frac{\lambda + \gamma_s(r, P_*) + \mu(r, P_*)}{\gamma(r, P_*)} dr \right\} \\ \times \left(\frac{g(y)\gamma(y, P_*) + \beta_1(y, P_*)\beta_2}{\gamma(y, P_*)} \right) dy ds = 1. \quad (4.15)$$

We note that, if

$$g(y)\gamma(y, P_*) + \beta_1(y, P_*)\beta_2 \geq 0, \quad y \in [0, m],$$

which is equivalent to the positivity condition (3.10), then equation (4.15) admits a dominant unique (real) solution. On the other hand, it is easily shown that this dominant eigenvalue is negative if

$$\int_0^m \int_0^s \exp \left\{ - \int_y^s \frac{\gamma_s(r, P_*) + \mu(r, P_*)}{\gamma(r, P_*)} dr \right\} \\ \times \left(\frac{g(y)\gamma(y, P_*) + \beta_1(y, P_*)\beta_2}{\gamma(y, P_*)} \right) dy ds < 1. \quad (4.16)$$

It is easy too see, making use of equation (2.7), that (4.16) is satisfied if

$$\int_0^m \frac{1}{\gamma(s, P_*)} \int_0^s \exp \left\{ - \int_y^s \frac{\mu(z, P_*)}{\gamma(z, P_*)} dz \right\} g(y) dy ds < 0,$$

holds true. In this case, we obtain for the growth bound of the semigroup ω_0

$$\omega_0 = s(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{G} + \mathcal{H}_{\tilde{\beta}}) < 0,$$

see e.g. Th.1.15 in Chapter VI of [9], which implies that the equilibrium solution is linearly stable.

5. Concluding remarks

In this paper, using linear semigroup methods, we analysed the asymptotic behaviour of a size-structured scramble competition model. We were motivated by the modelling of structured macro-parasites in aquaculture, where the population dynamics at the first chalimus stage of certain sea lice species is driven by a similar set of model equations. First we studied existence of equilibrium solutions of our model. In the case when the fertility function is separable, we easily established monotonicity conditions on the vital rates which guarantee the existence of a steady state (Proposition 2.1). In the general case we used positive perturbation arguments to establish criteria that guarantee the existence of at least one positive equilibrium solution. Next, we established conditions for the existence of a positive quasicontraction semigroup which governs the linearized problem. Then we established a further regularity property of the governing linear semigroup which in principle allows to study stability of equilibria via the point spectrum of its generator. In the special case of separable fertility function we

explicitly deduced a characteristic function in equation (4.8) whose roots are the eigenvalues of the linearized operator. Then we formulated stability/instability results, where we used once more finite rank lower/upper bound estimates of the very general recruitment term. We note that it would be also straightforward to formulate conditions which guarantee that the governing linear semigroup exhibits asynchronous exponential growth. However, this is not very interesting from the application point of view, since the linearised system is not necessarily a population equation anymore.

It is interesting to note that characterization of positivity using dispersivity resulted in much more relaxed conditions than the ones we obtained in [10] for a more simple size-structured model with 1-state-at-birth by characterizing positivity via the resolvent of the semigroup generator. This although is probably due to the different recruitment terms in the two model equations. Positivity is often crucial for our stability studies, as this was demonstrated in Section 3, indeed, more relaxed positivity conditions result in the much wider applicability (i.e. for a larger set of vital rates) of our analytical stability results.

We also note that in spaces where the positive cone has a non-empty interior there is a much more efficient way of characterizing positivity. Indeed, the positive minimum principle (see e.g. [1]) would potentially yield more relaxed positivity conditions than the ones we obtained in here. The difficulty to characterize positivity lies in the fact that the natural state space L^1 contains in a certain sense too many (non-positive) elements. In fact the interior of the positive cone of L^1 is empty. There is another result to characterize positivity in spaces where the positive cone has empty interior, namely the generalized (or abstract) Kato-inequality (see e.g. [1]). In our setting the abstract Kato-inequality reads

$$S_u (\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D})u \leq (\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D})|u|, \quad (5.1)$$

for $u \in \text{Dom} (\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D})$, where S_u is the signum operator, that is

$$S_u = \frac{u}{|u|}.$$

It is straightforward to show that inequality (5.1) requires

$$\begin{aligned} S_u \int_0^m u(y) \left(\beta(s, y, P_*) + \int_0^m \beta(s, z, P_*) p_*(z) dz - \rho_*(s) \right) dy \\ \leq \int_0^m |u(y)| \left(\beta(s, y, P_*) + \int_0^m \beta(s, z, P_*) p_*(z) dz - \rho_*(s) \right) dy, \quad s \in [0, m], \end{aligned} \quad (5.2)$$

which holds true for every $u \in \text{Dom} (\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D})$ indeed when condition (3.10) is satisfied.

As we have seen previously in Section 3, since the linearised system is not a population model anymore, the governing semigroup is not positive anymore unless some additional condition is satisfied. However, it was proven in [16]

that every quasicontraction semigroup on an L^1 space has a minimal dominating positive semigroup, called the modulus semigroup, which itself is quasicontractive. Hence, in principle, one can prove stability results even in the case of a non-positive governing semigroup, by perturbing the semigroup generator with a positive operator such that the perturbed generator does indeed generate a positive semigroup.

ACKNOWLEDGMENTS

JZF is thankful to the Centre de Recerca Matemàtica and to the Department of Mathematics, Universitat Autònoma de Barcelona for their hospitality while being a participant in the research programme "Mathematical Biology: Modelling and Differential Equations" during 01/2009-06/2009. PH thanks the University of Stirling for its hospitality. We also thank the Edinburgh Mathematical Society for financial support.

REFERENCES

- [1] W. Arendt, A. Grabosch, G. Greiner, U. Groh, H. P. Lotz, U. Moustakas, R. Nagel, F. Neubrander and U. Schlotterbeck, *One-Parameter Semigroups of Positive Operators*, Springer-Verlag, Berlin, (1986).
- [2] R. Borges, À. Calsina and S. Cuadrado, Equilibria of a cyclin structured cell population model, *Discrete Contin. Dyn. Syst., Ser. B* **11** (2009), 613–627.
- [3] À. Calsina and J. Saldaña, Basic theory for a class of models of hierarchically structured population dynamics with distributed states in the recruitment, *Math. Models Methods Appl. Sci.* **16** (2006), 1695–1722.
- [4] Ph. Clément, H. J. A. M. Heijmans, S. Angenent, C. J. van Duijn, and B. de Pagter, *One-Parameter Semigroups*, North-Holland, Amsterdam 1987.
- [5] M. J. Costello, Ecology of sea lice parasitic on farmed and wild fish, *Trends in Parasitol.* **22** (2006), 475–483.
- [6] J. M. Cushing, *An Introduction to Structured Population dynamics*, SIAM, Philadelphia (1998).
- [7] O. Diekmann and M. Gyllenberg, *Abstract delay equations inspired by population dynamics*, in "Functional Analysis and Evolution Equations" (Eds. H. Amann, W. Arendt, M. Hieber, F. Neubrander, S. Nicaise and J. von Below), Birkhäuser, (2007), 187–200.
- [8] O. Diekmann, Ph. Getto and M. Gyllenberg, Stability and bifurcation analysis of Volterra functional equations in the light of suns and stars, *SIAM J. Math. Anal.* **39** (2007), 1023–1069.
- [9] K.-J. Engel and R. Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Springer, New York 2000.
- [10] J. Z. Farkas and T. Hagen, Stability and regularity results for a size-structured population model, *J. Math. Anal. Appl.* **328** (2007), 119–136.
- [11] J. Z. Farkas and T. Hagen, Asymptotic analysis of a size-structured cannibalism model with infinite dimensional environmental feedback, to appear in *Commun. Pure Appl. Anal.*, <http://arxiv.org/abs/0812.1369>
- [12] J. Z. Farkas and T. Hagen, Hierarchical size-structured populations: The linearized semi-group approach, submitted, <http://arxiv.org/abs/0812.1367>

- [13] P. A. Heuch and T. A. Mo, A model of salmon louse production in Norway: effects of increasing salmon production and public management measures, *Dis. Aquat. Org.* **45** (2001), 145–152.
- [14] M. Iannelli, *Mathematical Theory of Age-Structured Population Dynamics*, Giardini Editori, Pisa (1994).
- [15] T. Kato, *Perturbation Theory for Linear Operators*, Springer, New York, 1966.
- [16] Y. Kubokawa, Ergodic theorems for contraction semi-groups, *J. Math. Soc. Japan* **27** (1975), 184–193.
- [17] J. A. J. Metz and O. Diekmann, *The Dynamics of Physiologically Structured Populations*, Springer, Berlin, 1986.
- [18] A. G. Murray and P. A. Gillibrand, Modelling salmon lice dispersal in Loch Torridon, Scotland, *Marine Pollution Bulletin* **53** (2006), 128–135.
- [19] J. Prüß, Stability analysis for equilibria in age-specific population dynamics, *Nonlin. Anal. TMA* **7** (1983), 1291–1313.
- [20] C. W. Revie, C. Robbins, G. Gettinby, L. Kelly and J. W. Treasurer, A mathematical model of the growth of sea lice, *Lepeophtheirus salmonis*, populations on farmed Atlantic salmon, *Salmo salar* L., in Scotland and its use in the assessment of treatment strategies, *J. Fish Dis.* **28** (2005), 603–613.
- [21] C. S. Tucker, R. Norman, A. Shinn, J. Bron, C. Sommerville and R. Wootten, A single cohort time delay model of the life-cycle of the salmon louse *Lepeophtheirus salmonis* on Atlantic salmon *Salmo salar*, *Fish Path.* **37** (2002), 107–118.
- [22] O. Tully and D. T. Nolan, A review of the population biology and host-parasite interactions of the sea louse *Lepeophtheirus salmonis* (Copepoda: Caligidae), *Parasitology* **124** (2002), S165–S182.
- [23] G. F. Webb, *Theory of Nonlinear Age-Dependent Population Dynamics*, Marcel Dekker, New York, (1985).
- [24] K. Yosida, *Functional analysis*, Springer, Berlin, (1995).

JÓZSEF Z. FARKAS

DEPARTMENT OF COMPUTING SCIENCE AND MATHEMATICS
 UNIVERSITY OF STIRLING
 STIRLING, FK9 4LA, UNITED KINGDOM
E-mail address: jzf@maths.stir.ac.uk

DARREN GREEN

INSTITUTE OF AQUACULTURE
 UNIVERSITY OF STIRLING
 STIRLING, FK9 4LA, UNITED KINGDOM
E-mail address: darren.green@stir.ac.uk

PETER HINOW

INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS
 UNIVERSITY OF MINNESOTA
 114 LIND HALL, MINNEAPOLIS
 MN 55455-0134, USA
E-mail address: hinow@ima.umn.edu